

Switzerland's Greenhouse Gas Inventory 1990–2012

National Inventory Report 2014
including reporting elements under the Kyoto Protocol

Submission of 15 April 2014
under the United Nations Framework Convention on Climate
Change and under the Kyoto Protocol



Schweizerische Eidgenossenschaft
Confédération suisse
Confederazione Svizzera
Confederaziun svizra

Swiss Confederation

Federal Office for the Environment FOEN

Publisher

Federal Office for the Environment FOEN, Climate Division, 3003 Bern, Switzerland
www.bafu.admin.ch/climate
www.climatereporting.ch

Inventory Project Leader

Paul Filliger FOEN, Climate Division, Bern

National Inventory Compiler

Anouk-Aimée Bass FOEN, Air Pollution Control and Chemicals Division, Bern

NIR Lead Authors

| | |
|---------------|------------------------------|
| Jürg Heldstab | INFRAS, Zürich |
| Denise Fussen | Ernst Basler+Partner, Zürich |
| Beat Rihm | Meteotest, Bern |

KP-LULUCF

Nele Rogiers FOEN, Forest Division, Bern

QA/QC Officer

Regine Röthlisberger FOEN, Climate Division, Bern

National Registry Administrator

Christine Kieffer FOEN, Climate Division, Bern

Bern, 15 April 2014

Switzerland's Greenhouse Gas Inventory 1990–2012

National Inventory Report 2014

Including reporting elements under the Kyoto Protocol

Submission of 15 April 2014

under the United Nations Framework Convention on Climate
Change and under the Kyoto Protocol

| NIR Chapter / Sector | Responsible Persons |
|---|---|
| Executive Summary | Author: Jürg Heldstab (INFRAS) |
| 1. Introduction | Lead Author: Jürg Heldstab (INFRAS) Authors: Mario Betschart (INFRAS), Martin Herren (INFRAS), Denise Fussen (EBP; Key Categories, uncert. analysis Tier 1), Beat Müller (FOEN; KCA analysis Tier 1/2), Regine Röthlisberger (FOEN; QA/QC), Bettina Schäppi (INFRAS, uncert. Anal. Tier 2) |
| 2. Trends in GHG Emissions and Removals | Lead Author: Jürg Heldstab (INFRAS) Authors: Mario Betschart (INFRAS), Martin Herren (INFRAS) |
| 3. Energy | Lead Authors: Jürg Heldstab (INFRAS), Denise Fussen (EBP) Authors: Anouk-Aimée Bass (FOEN; EMIS data base operation, QA/QC checks), Mario Betschart (INFRAS; Overview, Civil and Military Aviation, Bunker Fuels), Denise Fussen (EBP; Energy Stationary), Martin Herren (INFRAS; Transport), Beat Müller (FOEN; Reference Approach), Benedikt Notter (INFRAS; Transport, Off-road transport) Sector expert for Civil Aviation: Theo Rindlisbacher (FOCA) Technical contributor (Transport): Mario Keller (INFRAS) |
| 4. Industrial Processes | Lead Authors: Jürg Heldstab (INFRAS; F-gases), Denise Fussen (EBP) Authors: Christina Seyler (EBP), Martin Herren (INFRAS; F-gases), Sabine Schenker (FOEN; EMIS data base operation, QA/QC checks) Technical contributor (fluorinated gases): Cornelia Stettler (Carbotech) |
| 5. Solvent and Other Product Use | Lead Author: Denise Fussen (EBP) Author: Christina Seyler (EBP), Sabine Schenker (FOEN; EMIS data base operation, QA/QC checks) |
| 6. Agriculture | Lead Author: Jürg Heldstab (INFRAS) Authors: Benedikt Notter (INFRAS) Sector expert: Daniel Bretscher (Agroscope) |
| 7. LULUCF | Project Leader: Andreas Schellenberger (FOEN) Lead Author: Beat Rihm (Meteotest) Authors: Armin Keller (Agroscope), Jens Leifeld (Agroscope), Reto Meuli (Agroscope), Nele Rogiers (FOEN) Sector experts: Daniel Bretscher (Agroscope), Esther Thürig (WSL), Richard Volz (Meteotest) Technical contributors: Markus Didion (WSL), Christoph Könitzer (Sigmaplan), Lukas Mathys (Sigmaplan), Stefan Müller (Meteotest), Felix Weibel (SFSO) |

| NIR Chapter / Sector | Responsible Persons (continued) |
|-------------------------------------|--|
| 8. Waste | Lead Author: Denise Fussen (EBP) Author: Rainer Kegel (FOEN; EMIS data base operation, QA/QC checks), Maya Wolfensberger (EBP) |
| 9. Other (IPCC sector 7) | Authors: Martin Herren (INFRAS) |
| 10. Recalculations | Authors: Jürg Heldstab (INFRAS), Mario Betschart (INFRAS), Denise Fussen (EBP), Martin Herren (INFRAS), Benedikt Notter (INFRAS), Nele Rogiers, (FOEN), Andreas Schellenberger (FOEN), Christina Seyler (EBP), Maya Wolfensberger (EBP) |
| 11. KP-LULUCF | Author: Nele Rogiers (FOEN) Sector experts: Esther Thürig (WSL), Richard Volz (Meteotest) |
| 12. Accounting of Kyoto units | Authors: Paul Filliger (FOEN), Stefan Meier (FOEN) Sector experts: Yvan Keckeis (FOEN), Matthias Kohler (FOEN) |
| 13. Changes in National System | Author: Regine Röthlisberger (FOEN) |
| 14. Changes in National Registry | Author: Stefan Meier (FOEN) |
| 15. Minimization of adverse effects | Authors: Marianne Abt (seco), Paul Filliger (FOEN) |
| Annex 1 | Author: Denise Fussen (EBP) |
| Annex 2 | Authors: Jürg Heldstab (INFRAS), Mario Betschart (INFRAS) Technical contributor: Anouk-Aimée Bass (FOEN) |
| Annex 3 | Authors: Mario Betschart (INFRAS), Denise Fussen (EBP), Jürg Heldstab (INFRAS), Martin Herren (INFRAS), Stefan Kessler (INFRAS), Benedikt Notter (INFRAS) |
| Annex 4, 5, 6, 7 and 8 | Authors: Denise Fussen (EBP), Jürg Heldstab (INFRAS), Martin Herren (INFRAS), Stefan Reimann (EMPA) |

Table of Contents

| | |
|--|-----|
| Table of Contents | 5 |
| Glossary | 9 |
| Executive Summary | 13 |
| Acknowledgements | 27 |
| PART 1 | 29 |
| 1 Introduction | 29 |
| 1.1 Background Information on Swiss Greenhouse Gas Inventories, Climate Change and Supplementary Information of the Kyoto Protocol (KP) | 29 |
| 1.2 Institutional Arrangements for Inventory Preparation | 31 |
| 1.3 Process for Inventory Preparation | 35 |
| 1.4 Methodologies and Data Sources | 37 |
| 1.5 Description of Key Categories | 39 |
| 1.6 Quality Assurance and Quality Control (QA/QC) | 48 |
| 1.7 Uncertainty Evaluation | 56 |
| 1.8 Completeness Assessment | 69 |
| 2 Trends in Greenhouse Gas Emissions and Removals | 71 |
| 2.1 Aggregated Greenhouse Gas Emissions 2012 (UNFCCC) | 71 |
| 2.2 Emission Trends by Gas | 73 |
| 2.3 Emission Trends by Sources and Sinks | 75 |
| 2.4 Emission Trends for Indirect Greenhouse Gases and SO ₂ | 81 |
| 2.5 Emission Trends (Kyoto Protocol) | 83 |
| 3 Energy | 87 |
| 3.1 Overview | 87 |
| 3.2 Source Category 1A – Fuel Combustion Activities | 91 |
| 3.2.1 Comparison of the Sectoral Approach with the Reference Approach | 91 |
| 3.2.2 International Bunker Fuels | 93 |
| 3.2.3 Feedstocks and Non-Energy Use of Fuels | 96 |
| 3.2.4 CO ₂ Capture from Flue Gases and Subsequent CO ₂ Storage | 97 |
| 3.2.5 Country-specific issues | 98 |
| 3.2.6 Source Category 1A1 - Energy Industries | 116 |
| 3.2.7 Source Category 1A2 - Manufacturing Industries and Construction | 127 |
| 3.2.8 Source Category 1A3 - Transport | 144 |
| 3.2.9 Source Category 1A4 - Other Sectors (Commercial/Institutional, Residential, Agriculture/Forestry/Fisheries) | 164 |
| 3.2.10 Source Category 1A5b - Military | 172 |
| 3.3 Source Category 1B – Fugitive Emissions from Fuels | 175 |

| | | |
|-------|---|-----|
| 3.3.1 | Source Category 1B1 - Solid Fuels | 175 |
| 3.3.2 | Source Category 1B2a - Oil..... | 175 |
| 3.3.3 | Source Category 1B2b - Natural Gas | 177 |
| 3.3.4 | Source Category 1B2c - Venting | 180 |
| 4 | Industrial Processes | 183 |
| 4.1 | Overview | 183 |
| 4.2 | Source Category 2A – Mineral Products..... | 186 |
| 4.3 | Source Category 2B – Chemical Industry | 198 |
| 4.4 | Source Category 2C – Metal Production | 205 |
| 4.5 | Source Category 2D – Other Production | 211 |
| 4.6 | Source Category 2E – Production of Halocarbons and SF ₆ | 213 |
| 4.7 | Source Category 2F – Consumption of Halocarbons and SF ₆ | 213 |
| 4.8 | Source Category 2G – Other..... | 228 |
| 5 | Solvent and Other Product Use | 231 |
| 5.1 | Overview | 231 |
| 5.2 | Source Category 3A – Paint Application..... | 233 |
| 5.3 | Source Category 3B – Degreasing and Dry Cleaning..... | 235 |
| 5.4 | Source Category 3C – Chemical Products, Manufacture and Processing | 236 |
| 5.5 | Source Category 3D – Other | 239 |
| 6 | Agriculture..... | 243 |
| 6.1 | Overview | 243 |
| 6.2 | Source Category 4A – Enteric Fermentation | 245 |
| 6.3 | Source Category 4B – Manure Management | 255 |
| 6.4 | Source Category 4C – Rice Cultivation | 267 |
| 6.5 | Source Category 4D – Agricultural Soils..... | 267 |
| 6.6 | Source Category 4E – Burning of savannahs..... | 277 |
| 6.7 | Source Category 4F – Field Burning of Agricultural Residues | 277 |
| 7 | LULUCF | 279 |
| 7.1 | Overview of LULUCF | 279 |
| 7.2 | Activity Data – Land Areas | 289 |
| 7.3 | Category 5A – Forest Land | 300 |
| 7.4 | Category 5B – Cropland..... | 335 |
| 7.5 | Category 5C – Grassland..... | 343 |
| 7.6 | Category 5D – Wetlands | 353 |
| 7.7 | Category 5E – Settlements..... | 356 |
| 7.8 | Category 5F – Other Land..... | 359 |
| 8 | Waste..... | 361 |
| 8.1 | Overview | 361 |

| | | |
|---|---|-----|
| 8.2 | Source Category 6A – Solid Waste Disposal on Land | 367 |
| 8.3 | Source Category 6B – Wastewater Handling | 371 |
| 8.4 | Source Category 6C – Waste Incineration..... | 376 |
| 8.5 | Source Category 6D – Other Waste | 380 |
| 9 | Other | 385 |
| 9.1 | Overview | 385 |
| 9.2 | Source Category – Other non-specified..... | 386 |
| 10 | Recalculations | 389 |
| 10.1 | Explanations and Justifications for Recalculation | 389 |
| 10.2 | Implications for Emission Levels 1990 and 2011 | 397 |
| 10.3 | Implications for Emissions Trends, including Time Series Consistency | 398 |
| 10.4 | Recalculations, Including in Response to the Review Process, and Planned Improvements to the Inventory..... | 399 |
| PART 2 | | 401 |
| 11 | KP-LULUCF..... | 401 |
| 11.1 | General Information..... | 407 |
| 11.2 | Land-related Information | 409 |
| 11.3 | Activity-specific Information | 413 |
| 11.4 | Article 3.3. | 421 |
| 11.5 | Article 3.4 | 423 |
| 11.6 | Other Information | 427 |
| 11.7 | Information Relating to Article 6 | 427 |
| 12 | Information on Accounting of Kyoto Units | 429 |
| 12.1 | Background Information | 429 |
| 12.2 | Summary of Information Reported in the SEF Tables..... | 429 |
| 12.3 | Discrepancies and Notifications..... | 430 |
| 12.4 | Publicly Accessible Information | 430 |
| 12.5 | Calculation of the Commitment Period Reserve (CPR) | 430 |
| 12.6 | KP-LULUCF Accounting..... | 431 |
| 13 | Information on Changes in National System | 433 |
| 14 | Information on Changes in National Registry | 435 |
| 15 | Information on Minimization of Adverse Impacts in Accordance with Article 3, Paragraph 14 | 437 |
| 16 | Other Information..... | 441 |
| References | | 455 |
| References to EMIS database comments..... | | 482 |
| Annexes | | 483 |
| Annex 1: Key Category Analysis (KCA) | | 483 |
| Annex 2: Sulphur Dioxide (SO ₂) | | 492 |

| | |
|--|-----|
| Annex 3: Other detailed methodological descriptions for individual source or sink categories | 494 |
| Annex 4: CO ₂ Reference Approach and comparison with Sectoral Approach, and relevant information on the national energy balance | 517 |
| Annex 5: Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded | 520 |
| Annex 6: Additional information to be considered as part of the NIR submission (where relevant) or other useful reference information | 521 |
| Annex 7: Supplementary Information on the Uncertainty Analysis | 527 |
| Annex 8: Supplementary Information under Article 7, paragraph 1 of the Kyoto Protocol ... | 532 |

Glossary

| | |
|--------------------------------------|--|
| AD | Activity data |
| AEF | Area expansion factor |
| AREA1 | Swiss Land Use Statistics 1979/85 (ASCH1 data re-evaluated according to the AREA set of land-use and land-cover categories) |
| AREA2 | Swiss Land Use Statistics 1992/97 (ASCH2 data re-evaluated according to the AREA set of land-use and land-cover categories) |
| AREA3 | Swiss Land Use Statistics, third survey 2004/09 |
| ART | Agroscope Reckenholz-Tänikon Research Station (formerly FAL) since 2014 Agroscope |
| ASCH1 | Swiss Land Use Statistics, first survey 1979/85 |
| ASCH2 | Swiss Land Use Statistics, second survey 1992/97 |
| BEF, BCEF | biomass expansion factor, biomass conversion and expansion factor |
| Carbura | Swiss Central Office for the Import of Liquid Fuels |
| Cemsuisse | Association of the Swiss Cement Industry |
| CC | Combination category |
| CH ₄ | Methane, 1995 IPCC GWP: 21 (UNFCCC 2006b, Table 1) |
| CFC | Chlorofluorocarbon (organic compound: refrigerant, propellant) |
| CHP | Combined heat and power production |
| CO | Carbon monoxide |
| CO ₂ , CO ₂ eq | Carbon dioxide, carbon dioxide equivalent |
| CORINAIR | CORe INventory of AIR emissions (under the European Topic Centre on Air Emissions and under the European Environment Agency) |
| CRF | Common reporting format |
| DBH | Diameter (of trees) at breast height |
| EF | Emission factor |
| EMEP | European Monitoring and Evaluation Programme (under the Convention on Long-range Transboundary Air Pollution) |
| EMIS | Swiss national air pollution database |
| EMPA | Swiss Federal Laboratories for Material Testing and Research |
| EV | Erdöl-Vereinigung (Swiss Petroleum Association) |
| DETEC | Dept. of the Environment, Transport, Energy and Communications |
| FAL | Swiss Federal Research Station for Agroecology and Agriculture (since 2006: ART) |
| FCCC | Framework Convention on Climate Change |
| FiBL | Research Institute of Organic Agriculture |
| FOAG | Federal Office for Agriculture |
| FOCA | Federal Office of Civil Aviation |

| | |
|---------------------|---|
| FOEN | Federal Office for the Environment (former name SAEFL until 2005) |
| FOITT | Federal Office of Information Technology, Systems and Telecommunication |
| Gg | Gigagram (10^9 g = 1'000 tons) |
| GHG | Greenhouse gas |
| GL, GPG | Guidelines, Good Practice Guidance |
| GVS | Swiss Foundry Association |
| GWP | Global Warming Potential |
| ha | hectare |
| HFC | Hydrofluorocarbons (e.g. HFC-32 difluoromethane) |
| HFO | Heavy fuel oil |
| IDM | FOEN Internal Document Management System |
| IDP | Inventory Development Plan |
| IPCC | Intergovernmental Panel on Climate Change |
| KCA | key category analysis |
| kha | kilo hectare |
| LPG | Liquefied Petroleum Gas (Propane/Butane) |
| LTO | Landing-Takeoff-Cycle (Aviation) |
| LULUCF | Land Use, Land-Use Change and Forestry |
| MSW | Municipal solid waste |
| NABO | Swiss Soil Monitoring Network |
| NCV | Net calorific value |
| NFI 1, NFI 2, NFI 3 | First (1983-1985), Second (1993-1995) and Third (2004-2006) National Forest Inventory |
| NIR | National Inventory Report |
| NIS | National Inventory System |
| NMVOC | Non-methane volatile organic compounds |
| N ₂ O | Nitrous oxide; 1995 IPCC GWP: 310 (UNFCCC 2006b, Table 1) |
| NO _x | Nitrogen oxides |
| PCDD/PCDF | Polychlorinated Dibenzodioxins and -furans |
| PFC | Perfluorinated carbon compounds (e.g. Tetrafluoromethane) |
| SAEFL | Swiss Agency for the Environment, Forests and Landscape (since 2006: Federal Office for the Environment FOEN) |
| SF ₆ | Sulphur hexafluoride, 1995 IPCC GWP: 23'900 (UNFCCC 2006b, Table 1) |
| SFOE | Swiss Federal Office of Energy |
| SFSO | Swiss Federal Statistical Office |
| SO ₂ | Sulphur dioxide |
| SOC | Soil organic carbon |

| | |
|----------------|--|
| SVGW/SSIG/SGWA | Schweizerischer Verein des Gas- und Wasserfaches / Société Suisse de l'Industrie du Gaz et des Eaux / Swiss Gas and Water Industry Association |
| SWISSMEM | Swiss Mechanical and Electrical Engineering Industries (Schweizer Maschinen-, Elektro- und Metallindustrie) |
| UNECE | United Nations Economic Commission for Europe |
| UNFCCC | United Nations Framework Convention on Climate Change |
| VOC | Volatile organic compounds |
| VTG | Luftwaffe (Swiss Air Force Administration) |
| WSL | Swiss Federal Institute for Forest, Snow and Landscape Research |

Executive Summary

ES 1 Background Information on Greenhouse Gas Inventories, Climate Change and Supplementary Information Required Under Art. 7.1. KP

ES 1.1 Background Information on Climate Change

Recent data confirms a warming trend in Switzerland with an observed increase in mean annual temperature of 1.75°C between 1864 and 2012 (FOEN 2014d). Over the last 30 years Swiss temperature has increased with an annual average warming rate of 0.35°C/decade (CH2011, 2011). The most visible change in the Alps resulting from global warming is the retreat of glaciers, which showed a volume loss of 12% since 1999 (FOEN 2014d).

The observed trends in precipitation are less distinct than in temperature. They generally show an increase in winter and spring, whereas for summer and autumn no significant trends are detectable. Regional scenarios predict an increase in mean winter precipitation and a decrease in summer, which will have a marked impact on the hydrological cycle. Further, higher intensity of storms and reduced snowfall and snow cover duration are expected, increasing the risk and frequency of floods, landslides and debris flows.

Concerning biodiversity, climate change is expected to affect species composition, distribution, their cycles, synchronicity, the overall genetic diversity and the provision of ecosystem services. It will enhance the vulnerability of forests and potentially impair their protective, productive and social functions.

For agriculture, a moderate warming of 2°C to 3°C might increase productivity; however, if temperature will rise beyond that level, the increase in heat waves and drought periods would prove problematic for the cultivation of land and for livestock husbandry.

Various sectors of the Swiss economy are likely to be adversely affected by progressing climate change: In particular, winter tourism will suffer from increased scarcity of snow, hydroelectric power stations are confronted with altered runoff and sediment transport regimes, and insurance companies may face increased losses due to winter storms and floods. Natural hazards and extreme weather events potentially pose a growing risk to infrastructure and human health. Heat waves and elevated tropospheric ozone levels are cause for serious concern. Finally, it remains to be seen to what extent vector borne diseases spread due to changing climatic conditions. Recently Switzerland has analysed these challenges in detail and developed an effective adaptation strategy in order to hedge against negative effects resulting from climate change in Switzerland (FOEN, 2012b).

ES.1.2 Background Information on Greenhouse Gas Inventories

On 10 December 1993, Switzerland ratified the United Nations Framework Convention on Climate Change (UNFCCC). Since 1996, the submission of its national greenhouse gas inventory has been based on IPCC guidelines. From 1998 onwards, the inventories have been submitted in the Common Reporting Format (CRF). In 2004, Switzerland started submitting a yearly National Inventory Report (NIR) under the UNFCCC.

On 9 July 2003, Switzerland ratified the Kyoto Protocol under the UNFCCC. The Swiss National Inventory System (NIS) according to Article 5.1 of the Kyoto Protocol has been implemented and is fully operational.

The 2014 inventory submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol (FOEN 2014) includes the NIR on hand, the greenhouse gas inventory 1990–2012 and the Kyoto Protocol LULUCF tables 2008–2012 in the common reporting format as well as the SEF tables and the standard independent

assessment report (SIAR) from the National Registry. As a supplement, the update of the Description of the Quality Management System (FOEN 2014a) is provided.

The Federal Office for the Environment (FOEN) is in charge of compiling the emission data and bears overall responsibility for Switzerland's national greenhouse gas inventory and the national registry. In addition to the FOEN, the Swiss Federal Office of Energy (SFOE), the Agroscope Research Station and the Federal Office of Civil Aviation (FOCA) participate directly in the compilation of the inventory. Several other administrative offices and research institutions are involved in inventory preparation.

In preparing the national greenhouse gas inventory, Switzerland took into account the findings of the individual reviews of the inventory by the expert review teams of the UN (UNFCCC 2004, UNFCCC 2006, UNFCCC 2007, UNFCCC 2009, UNFCCC 2010, UNFCCC 2011, UNFCCC 2012, UNFCCC 2013). The recommendations of the Annual Review Report (ARR) for the submission 2013 (UNFCCC 2014) as well as the recommendations of the "Saturday Paper" (UNFCCC 2013a) are included in the present submission (see Chapter 16).

The structure of the NIR corresponds to the UNFCCC annotated outline (UNFCCC 2009a) and it contains three parts: **PART 1** reports the obligations under the UNFCCC, **PART 2** the additional obligations under the Kyoto Protocol and several **Annexes** with detailed information on selected issues of Part 1 and Part 2.

Chapter 1 of the NIR, the introduction, provides an overview of Switzerland's institutional arrangements for producing the inventory, and the process and methodologies used for inventory preparation.

- The data sources used to compile the national inventory and to estimate greenhouse gas emissions and removals are: the Swiss national air pollution database (EMIS), national energy statistics, data from industry associations, as well as further statistics and models for road transportation, off-road vehicles and machinery, agriculture, land use, land-use change and forestry (LULUCF) and waste. Emissions are calculated according to methodologies recommended by the IPCC and contained in the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 1997a, 1997b, 1997c), in the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC 2000), and for LULUCF in the IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC 2003). Furthermore, the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006) have been consulted in a few cases. However, the nomenclature of the Revised 1996 IPCC Guidelines has been used throughout the current NIR. The data in the EMIS database are pre-processed in order to enable transfer to the CRF Reporter required for reporting under the UNFCCC and under the Kyoto Protocol.
- All inventory data are assembled and prepared for input into the CRF Reporter by the GHG Inventory Core Group, which is responsible for ensuring the conformity of the inventory with the Updated UNFCCC Reporting Guidelines on Annual Inventories (UNFCCC 2006b) and the 2008 Kyoto Protocol Reference Manual (UNFCCC 2008). In the preparation of this report, the Inventory Group was supported by consultants. Their mandate included editing of the NIR, and an analysis of the consistency between the emission modelling and the recommendations of the IPCC Good Practice Guidance. Furthermore, the consultants contributed to the key category analyses and carried out the uncertainty analyses. They were also involved in inventory improvement, e.g. by performing tasks contained in the Inventory Development Plan.
- The inventory quality management system is designed to comply with the objectives of good practice guidance, i.e. to ensure and improve transparency, consistency, comparability, completeness, accuracy and confidence in national GHG emission and removal estimates. The QA/QC Officer is responsible for enforcement of the defined quality standards. The National Inventory System complies with the ISO 9001:2008

standard (Quality Management System) and is certified by the Swiss TS Technical Services AG (Swiss-TS 2013).

- A National Inventory System Supervisory Board was established by decision of the FOEN Directorate in summer 2006. The Board oversees activities related to the GHG Inventory and to the National Registry.
- Furthermore, Chapter 1 provides information on key categories and uncertainties.

Chapter 2 contains an analysis of trends in Switzerland's greenhouse gas emissions by sources and removals by sinks for all sectors.

Chapters 3 to 9 provide principal source and sink category estimates.

Chapter 10 justifies, explains and summarises the recalculations and planned improvements. They result in a very small change of -0.30% (without LULUCF) in the base year emissions (1990) compared with the latest submission in 2013 and a small change in the latest year of recalculations 2011 by -0.38%. The chapter also contains an overview of the planned improvements.

In **PART 2**, **Chapter 11** reports KP LULUCF data, **Chapter 12** presents information on accounting of Kyoto Units, **Chapter 13** lists changes in the National System, **Chapter 14** documents changes in the National Registry, **Chapter 15** provides information on the minimization of adverse effects and **Chapter 16** contains other information including the "Saturday Paper" (UNFCCC 2013a) that resulted from the 2013 review, together with the party's responses.

ES.1.3 Background Information on Supplementary Information Required under Article 7.1. of the Kyoto Protocol (KP)

Chapter 11 of PART 2 as mentioned above, provides information on KP-LULUCF. Switzerland has decided to account for Forest Management under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol. In accordance with the Annex to Decision 16/CMP.1, credits from Forest Management are capped in the first commitment period. For Switzerland the cap amounts to 1.83 Mt CO₂ (0.5 Mt C) per year, or 9.167 Mt CO₂ for the whole commitment period.

Switzerland has chosen to account annually for emissions and removals from activities under the Kyoto Protocol. The current submission contains the mandatory inventory years 2008, 2009, 2010, 2011 and 2012. In the NIR, additional information about 1999 to 2007 is included.

ES.2 Summary of National Emission and Removal Related Trends, and Emission and Removals from KP-LULUCF Activities

ES.2.1 GHG Inventory

In 2012, Switzerland emitted 51'449 Gg (kilotonnes) CO₂ equivalent, corresponding to 6.43 tonnes CO₂ equivalent per capita (CO₂: 5.41 tonnes per capita), to the atmosphere, excluding emissions from international bunkers (aviation and marine) and excluding emissions and removals from the sector Land Use, Land-Use Change and Forestry (LULUCF)¹. For the emissions that are relevant under the Kyoto Protocol see chapter ES.3.3.

Several Key Category Analyses (with, without LULUCF and combined) are carried out for 2012 and for the base year 1990.

¹ Inhabitants in Switzerland in 2012: 7.997 million (SFSO 2013a)

- Tier 1 analysis (without LULUCF): For 2012, among a total of 135 categories, 31 have been identified as key categories (level and/or trend) with an aggregated contribution of 97.2% to total national emissions. Of the 31 key categories, 19 are in sector 1 Energy, accounting for 79.5% of total CO₂ equivalent emissions in 2012.
- Tier 2 analysis (without LULUCF): For 2012, among a total of 135 categories, 30 have been identified as key categories (level and/or trend) with an aggregated contribution of 94.0% of the sum of all level assessments weighted with their uncertainty in 2012. Of the 30 key categories, 14 are in sector 1 Energy, accounting for 29.8% of the sum of all level assessments weighted with their uncertainty in 2012. Sector 4 Agriculture accounts for 51.0% of that sum. Tier 2 key category analysis shows that these two sectors have the highest impact on inventory uncertainty.
- A Tier 1 and Tier 2 analysis with LULUCF was conducted as well (see 1.5.1.3 and Annex A1.5).

Table E-1 shows Switzerland's annual GHG emissions by individual GHGs from 1990 (base year) to 2012. Despite clear trends in some GHG emissions (see below), there is no significant trend in the total emissions of the period 1990–2012. Year-to-year variations of total emissions are mainly caused by changing winter temperatures and their effect on CO₂ emissions from fuel combustion (source category 1A4). In 2012, total gross GHG emissions (excluding LULUCF) show a decrease of 2.7% compared to the level recorded for 1990 (see also Table E-2).

Table E-1 Summary of Switzerland's GHG emissions in CO₂ equivalent (Gg), 1990–2012 (from CRF-tables 10s5, 10s5.2 and 10s5.3, upper half). HFCs increased by 5'526'548% compared to 1990 levels (0.02 Gg CO₂ equivalent).

| Greenhouse Gas Emissions | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 42'677 | 44'356 | 44'111 | 40'386 | 40'550 | 40'519 | 41'588 | 40'257 | 41'833 | 42'724 |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 44'639 | 46'369 | 46'258 | 43'718 | 42'981 | 43'683 | 44'340 | 43'514 | 44'812 | 44'894 |
| CH ₄ emissions including CH ₄ from LULUCF | 4'576 | 4'563 | 4'471 | 4'377 | 4'330 | 4'328 | 4'283 | 4'225 | 4'170 | 4'073 |
| CH ₄ emissions excluding CH ₄ from LULUCF | 4'546 | 4'551 | 4'461 | 4'367 | 4'315 | 4'311 | 4'270 | 4'187 | 4'156 | 4'063 |
| N ₂ O emissions including N ₂ O from LULUCF | 3'472 | 3'463 | 3'433 | 3'345 | 3'307 | 3'295 | 3'289 | 3'177 | 3'161 | 3'124 |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'461 | 3'457 | 3'427 | 3'339 | 3'300 | 3'287 | 3'283 | 3'166 | 3'155 | 3'119 |
| HFCs | 0 | 0 | 7 | 15 | 34 | 182 | 228 | 301 | 358 | 421 |
| PFCs | 100 | 85 | 69 | 30 | 18 | 15 | 17 | 20 | 23 | 36 |
| SF ₆ | 144 | 146 | 148 | 126 | 112 | 98 | 94 | 131 | 160 | 147 |
| Total (including LULUCF) | 50'969 | 52'613 | 52'239 | 48'279 | 48'350 | 48'436 | 49'501 | 48'112 | 49'704 | 50'524 |
| Total (excluding LULUCF) | 52'890 | 54'607 | 54'370 | 51'595 | 50'760 | 51'576 | 52'233 | 51'319 | 52'663 | 52'680 |

| Greenhouse Gas Emissions | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 43'949 | 46'171 | 44'805 | 43'584 | 42'967 | 44'337 | 44'238 | 42'145 | 44'658 | 43'453 |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 43'952 | 44'904 | 43'844 | 45'025 | 45'650 | 46'290 | 45'911 | 43'931 | 45'447 | 44'280 |
| CH ₄ emissions including CH ₄ from LULUCF | 3'995 | 3'983 | 3'939 | 3'849 | 3'806 | 3'801 | 3'804 | 3'801 | 3'850 | 3'787 |
| CH ₄ emissions excluding CH ₄ from LULUCF | 3'985 | 3'973 | 3'922 | 3'829 | 3'797 | 3'791 | 3'793 | 3'787 | 3'840 | 3'777 |
| N ₂ O emissions including N ₂ O from LULUCF | 3'114 | 3'133 | 3'106 | 3'047 | 2'997 | 2'981 | 2'977 | 3'006 | 3'043 | 3'008 |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'109 | 3'128 | 3'100 | 3'040 | 2'993 | 2'977 | 2'972 | 3'001 | 3'039 | 3'004 |
| HFCs | 501 | 597 | 635 | 710 | 820 | 905 | 936 | 976 | 1'042 | 1'083 |
| PFCs | 69 | 45 | 40 | 57 | 53 | 33 | 33 | 29 | 39 | 36 |
| SF ₆ | 158 | 157 | 168 | 174 | 190 | 213 | 201 | 186 | 245 | 187 |
| Total (including LULUCF) | 51'787 | 54'085 | 52'694 | 51'421 | 50'834 | 52'271 | 52'189 | 50'144 | 52'878 | 51'554 |
| Total (excluding LULUCF) | 51'775 | 52'805 | 51'710 | 52'835 | 53'503 | 54'209 | 53'846 | 51'910 | 53'653 | 52'366 |

| Greenhouse Gas Emissions | 2010 | 2011 | 2012 | Change baseyear to 2012 (%) |
|---|---------------------------------|---------------|---------------|-----------------------------|
| | CO ₂ equivalent (Gg) | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 44'976 | 39'934 | 42'109 | -1.3% |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 45'923 | 41'848 | 43'251 | -3.1% |
| CH ₄ emissions including CH ₄ from LULUCF | 3'773 | 3'723 | 3'698 | -19.2% |
| CH ₄ emissions excluding CH ₄ from LULUCF | 3'764 | 3'711 | 3'689 | -18.8% |
| N ₂ O emissions including N ₂ O from LULUCF | 3'082 | 3'019 | 3'011 | -13.3% |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'078 | 3'015 | 3'007 | -13.1% |
| HFCs | 1'138 | 1'195 | 1'245 | see caption |
| PFCs | 37 | 40 | 33 | -67.0% |
| SF ₆ | 155 | 164 | 224 | 56.0% |
| Total (including LULUCF) | 53'161 | 48'076 | 50'320 | -1.3% |
| Total (excluding LULUCF) | 54'095 | 49'973 | 51'449 | -2.7% |

With regard to the distribution of emissions by individual greenhouse gases, CO₂ is the largest single contributor to emissions, accounting for 84.1% of total gross GHG emissions (excluding LULUCF) in 2012 (1990: 84.4%). The share of CH₄ decreased from 8.6% (1990) to 7.2% (2012). Over the same period, the share of N₂O decreased from 6.5% to 5.8%, while the share of F-gases increased from 0.5% to 2.9%.

Table E-2 Switzerland's total gross GHG emissions (excluding LULUCF) and the contribution of individual gases in CO₂ equivalent (Gg), selected years.

| Greenhouse Gas Emissions (excluding LULUCF) | 1990 | | 1995 | | 2000 | | 2005 | |
|--|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| CO ₂ | 44'639 | 84.4% | 43'683 | 84.7% | 43'952 | 84.9% | 46'290 | 85.4% |
| CH ₄ | 4'546 | 8.6% | 4'311 | 8.4% | 3'985 | 7.7% | 3'791 | 7.0% |
| N ₂ O | 3'461 | 6.5% | 3'287 | 6.4% | 3'109 | 6.0% | 2'977 | 5.5% |
| HFCs | 0 | 0.0% | 182 | 0.4% | 501 | 1.0% | 905 | 1.7% |
| PFCs | 100 | 0.2% | 15 | 0.0% | 69 | 0.1% | 33 | 0.1% |
| SF ₆ | 144 | 0.3% | 98 | 0.2% | 158 | 0.3% | 213 | 0.4% |
| Total (excluding LULUCF) | 52'890 | 100% | 51'576 | 100% | 51'775 | 100% | 54'209 | 100% |

| Greenhouse Gas Emissions (excluding LULUCF) | 2009 | | 2010 | | 2011 | | 2012 | |
|--|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| CO ₂ | 44'280 | 84.6% | 45'923 | 84.9% | 41'848 | 83.7% | 43'251 | 84.1% |
| CH ₄ | 3'777 | 7.2% | 3'764 | 7.0% | 3'711 | 7.4% | 3'689 | 7.2% |
| N ₂ O | 3'004 | 5.7% | 3'078 | 5.7% | 3'015 | 6.0% | 3'007 | 5.8% |
| HFCs | 1'083 | 2.1% | 1'138 | 2.1% | 1'195 | 2.4% | 1'245 | 2.4% |
| PFCs | 36 | 0.1% | 37 | 0.1% | 40 | 0.1% | 33 | 0.1% |
| SF ₆ | 187 | 0.4% | 155 | 0.3% | 164 | 0.3% | 224 | 0.4% |
| Total (excluding LULUCF) | 52'366 | 100% | 54'095 | 100% | 49'973 | 100% | 51'449 | 100% |

Figure E-1 shows the shares of 2012 emissions contributed by individual greenhouse gases. As the shares of emissions contributed by the individual gases have remained relatively constant, the diagram is also representative of the other years in the period 1990–2012.




Figure E-1 Contribution of individual gases to Switzerland's GHG emissions (excluding LULUCF) in 2012. 100% = 51'449 Gg CO₂ eq. (Numbers may not add to total due to rounding.)

For the emission data of 2012 excluding LULUCF, an uncertainty analysis on Tier 1 level was carried out resulting in a **level uncertainty of 3.65% and a trend uncertainty of 1.87% (1990-2012)**. The analysis was also carried out including the LULUCF sector resulting in increases of the uncertainties to 7.43% (level uncertainty) and 2.46% (trend uncertainty). Tier 2 show somewhat higher uncertainties: **level uncertainty of 3.86% and a trend uncertainty of 3.11%**. Including the LULUCF sector, the level uncertainty is 7.51% and the trend uncertainty 8.82%. In the Tier 2 analyses, asymmetric probability distributions and positive correlations cause higher uncertainties than in the Tier 1 analyses.

Chapter 10 explains and justifies recalculations that have been performed since the previous inventory submission to the UNFCCC secretariat in September 2013 after the centralized review 2013. The recalculations result in a decrease of the total base year (1990) emissions of 0.30% in CO₂ equivalents compared to the previous inventory (without LULUCF). For the year 2011 emissions, the decrease is 0.38% without emissions and removals from LULUCF. If the LULUCF sector is included there is a increase of 2.15% in 1990 and a decrease of 2.83% in 2011.

ES.2.2 KP-LULUCF Activities

Switzerland reports the mandatory LULUCF activities Afforestation and Deforestation (Reforestation is not occurring in Switzerland) under Article 3, paragraph 3 of the Kyoto Protocol, and Forest Management under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol. The total contribution of these activities is shown in Table E-3.

Table E-3 Contribution of activities accounted for under Article 3, paragraph 3 and paragraph 4 (Forest Management) of the Kyoto Protocol, Gg CO₂ eq., 1999-2012.

| Greenhouse gas source and sink activities | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|---|---|---------|--------|--------|----------|----------|----------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| Article 3.3 activities | 216.97 | 216.83 | 215.84 | 214.33 | 212.51 | 210.95 | 174.16 |
| Article 3.4 activities | -3330.01 | -395.31 | 166.35 | 219.39 | -2434.18 | -2810.38 | -2663.60 |

| Greenhouse gas source and sink activities | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|---|---|----------|----------|----------|----------|----------|----------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| Article 3.3 activities | 147.05 | 117.04 | 81.74 | 162.22 | 197.10 | 201.52 | 204.74 |
| Article 3.4 activities | -2804.38 | -1901.42 | -1202.77 | -1419.28 | -2020.23 | -2063.62 | -2236.38 |

ES.3. Overview of Source and Sink Category Estimates and Trends, including KP-LULUCF Activities

ES.3.1 GHG Inventory (Convention on Climate Change)

Table E-4 and Figure E-2 show the GHG emissions and removals by the main source and sink categories. The energy sector is by far the largest source of national emissions, accounting for 80.6% of the total GHG emissions (excluding LULUCF). There are decreasing trends in the source categories 3. Solvent and Other Product Use, 4. Agriculture, and 6. Waste as well as an increasing trend in source category 2 Industrial Processes. However, there is no significant trend in total emissions over the period 1990–2012 due to the dominating emissions of the energy sector with its year-to-year variability caused by changing winter temperatures and their effect on CO₂ emissions from fuel combustion.

Table E-4 Switzerland's GHG emissions and removals by source and sink categories in CO₂ equivalent (Gg), 1990–2012 (from CRF-tables 10s5, 10s5.2 and 10s5.3).

| Source and Sink Categories | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| 1. Energy | 41'989 | 44'121 | 44'191 | 41'915 | 41'029 | 41'916 | 42'787 | 42'147 | 43'474 | 43'533 |
| 1A1 Energy Industries | 2'601 | 2'859 | 2'939 | 2'584 | 2'613 | 2'643 | 2'855 | 2'813 | 3'132 | 3'165 |
| 1A2 Manufacturing Industries and Construction | 6'138 | 6'326 | 5'988 | 5'909 | 5'911 | 6'106 | 5'885 | 5'784 | 5'969 | 5'954 |
| 1A3 Transport | 14'600 | 15'094 | 15'418 | 14'352 | 14'539 | 14'225 | 14'287 | 14'844 | 15'056 | 15'663 |
| 1A4 Other Sectors | 18'095 | 19'273 | 19'276 | 18'500 | 17'407 | 18'393 | 19'207 | 18'138 | 18'745 | 18'227 |
| 1A5 Other (Military) | 206 | 188 | 180 | 171 | 166 | 148 | 137 | 147 | 146 | 132 |
| 1B Fugitive emissions from oil and natural gas | 349 | 381 | 390 | 399 | 393 | 400 | 416 | 420 | 426 | 392 |
| 2. Industrial Processes | 3'320 | 2'957 | 2'793 | 2'484 | 2'649 | 2'626 | 2'498 | 2'431 | 2'517 | 2'580 |
| 3. Solvent and Other Product Use | 470 | 444 | 420 | 392 | 374 | 354 | 331 | 308 | 286 | 273 |
| 4. Agriculture | 6'092 | 6'069 | 5'979 | 5'877 | 5'843 | 5'819 | 5'780 | 5'606 | 5'578 | 5'511 |
| 6. Waste | 1'007 | 1'004 | 976 | 914 | 852 | 847 | 823 | 814 | 795 | 768 |
| 7. Other | 12 | 12 | 13 | 13 | 13 | 13 | 13 | 13 | 14 | 14 |
| Total (excluding LULUCF) | 52'890 | 54'607 | 54'370 | 51'595 | 50'760 | 51'576 | 52'233 | 51'319 | 52'663 | 52'680 |
| 5. Land Use, Land-Use Change and Forestry | -1'921 | -1'995 | -2'131 | -3'316 | -2'410 | -3'140 | -2'732 | -3'207 | -2'959 | -2'155 |
| Total (including LULUCF) | 50'969 | 52'613 | 52'239 | 48'279 | 48'350 | 48'436 | 49'501 | 48'112 | 49'704 | 50'524 |

| Source and Sink Categories | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| 1. Energy | 42'429 | 43'330 | 42'297 | 43'497 | 43'922 | 44'432 | 44'084 | 42'078 | 43'628 | 42'506 |
| 1A1 Energy Industries | 3'064 | 3'187 | 3'268 | 3'295 | 3'617 | 3'804 | 4'078 | 3'835 | 4'025 | 3'949 |
| 1A2 Manufacturing Industries and Construction | 5'839 | 6'125 | 5'865 | 6'001 | 6'120 | 6'173 | 6'320 | 6'159 | 6'173 | 5'798 |
| 1A3 Transport | 15'896 | 15'597 | 15'522 | 15'689 | 15'767 | 15'827 | 15'939 | 16'257 | 16'624 | 16'427 |
| 1A4 Other Sectors | 17'133 | 17'946 | 17'186 | 18'092 | 18'021 | 18'239 | 17'360 | 15'462 | 16'450 | 15'983 |
| 1A5 Other (Military) | 136 | 134 | 140 | 125 | 114 | 124 | 127 | 120 | 115 | 116 |
| 1B Fugitive emissions from oil and natural gas | 362 | 341 | 317 | 295 | 283 | 265 | 260 | 245 | 242 | 232 |
| 2. Industrial Processes | 2'836 | 2'938 | 2'922 | 2'970 | 3'235 | 3'425 | 3'397 | 3'421 | 3'535 | 3'446 |
| 3. Solvent and Other Product Use | 259 | 245 | 234 | 225 | 212 | 211 | 206 | 205 | 202 | 201 |
| 4. Agriculture | 5'496 | 5'561 | 5'536 | 5'461 | 5'447 | 5'474 | 5'493 | 5'556 | 5'645 | 5'587 |
| 6. Waste | 741 | 717 | 708 | 669 | 673 | 654 | 653 | 637 | 629 | 612 |
| 7. Other | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 |
| Total (excluding LULUCF) | 51'775 | 52'805 | 51'710 | 52'835 | 53'503 | 54'209 | 53'846 | 51'910 | 53'653 | 52'366 |
| 5. Land Use, Land-Use Change and Forestry | 12 | 1'281 | 984 | -1'414 | -2'669 | -1'939 | -1'657 | -1'766 | -775 | -813 |
| Total (including LULUCF) | 51'787 | 54'085 | 52'694 | 51'421 | 50'834 | 52'271 | 52'189 | 50'144 | 52'878 | 51'554 |

| Source and Sink Categories | 2010 | 2011 | 2012 | 2012/1990 |
|--|---------------------------------|--------|--------|-----------|
| | CO ₂ equivalent (Gg) | | | % |
| 1. Energy | 44'004 | 39'945 | 41'477 | -1.2% |
| 1A1 Energy Industries | 4'180 | 3'968 | 4'064 | 56.3% |
| 1A2 Manufacturing Industries and Construction | 5'954 | 5'449 | 5'515 | -10.1% |
| 1A3 Transport | 16'321 | 16'205 | 16'331 | 11.9% |
| 1A4 Other Sectors | 17'193 | 13'994 | 15'240 | -15.8% |
| 1A5 Other (Military) | 121 | 108 | 116 | -43.6% |
| 1B Fugitive emissions from oil and natural gas | 234 | 220 | 209 | -40.0% |
| 2. Industrial Processes | 3'634 | 3'642 | 3'628 | 9.3% |
| 3. Solvent and Other Product Use | 199 | 202 | 200 | -57.5% |
| 4. Agriculture | 5'637 | 5'572 | 5'539 | -9.1% |
| 6. Waste | 608 | 598 | 591 | -41.3% |
| 7. Other | 14 | 14 | 14 | 16.2% |
| Total (excluding LULUCF) | 54'095 | 49'973 | 51'449 | -2.7% |
| 5. Land Use, Land-Use Change and Forestry | -934 | -1'897 | -1'129 | -41.2% |
| Total (including LULUCF) | 53'161 | 48'076 | 50'320 | -1.3% |




Figure E-2 Switzerland's greenhouse gas emissions in CO₂ equivalent (Gg) by main source categories, 1990–2012 (excluding LULUCF).

Table E-5 shows the contributions of individual sectors to total emissions excl. LULUCF for selected years in more detail. Between 1990 and 2012, the relative contribution of sector 1 Energy increased marginally from 79.4% to 80.6%, whereas emissions from sector 4 Agriculture decreased from 11.5% to 10.8% and those from sector 6 Waste changed from 1.9% to 1.1%. Sector 2 Industrial Processes contributed 6.3% to total emissions in 1990 and 7.1 % in 2012, but with lower values in between (1995, 2000).

Table E-5 Switzerland's total gross GHG emissions (excluding LULUCF) in CO₂ equivalent (Gg) and the contribution of individual source categories, selected years.

| Source and Sink Categories | 1990 | | 1995 | | 2000 | | 2005 | | 2007 | |
|--|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| 1. Energy | 41'989 | 79.4% | 41'916 | 81.3% | 42'429 | 81.9% | 44'432 | 82.0% | 42'078 | 81.1% |
| 1A1 Energy Industries | 2'601 | 4.9% | 2'643 | 5.1% | 3'064 | 5.9% | 3'804 | 7.0% | 3'835 | 7.4% |
| 1A2 Manufacturing Industries and Construction | 6'138 | 11.6% | 6'106 | 11.8% | 5'839 | 11.3% | 6'173 | 11.4% | 6'159 | 11.9% |
| 1A3 Transport | 14'600 | 27.6% | 14'225 | 27.6% | 15'896 | 30.7% | 15'827 | 29.2% | 16'257 | 31.3% |
| 1A4 Other Sectors | 18'095 | 34.2% | 18'393 | 35.7% | 17'133 | 33.1% | 18'239 | 33.6% | 15'462 | 29.8% |
| 1A5 Other (Military) | 206 | 0.4% | 148 | 0.3% | 136 | 0.3% | 124 | 0.2% | 120 | 0.2% |
| 1B Fugitive emissions from oil and natural gas | 349 | 0.7% | 400 | 0.8% | 362 | 0.7% | 265 | 0.5% | 245 | 0.5% |
| 2. Industrial Processes | 3'320 | 6.3% | 2'626 | 5.1% | 2'836 | 5.5% | 3'425 | 6.3% | 3'421 | 6.6% |
| 3. Solvent and Other Product Use | 470 | 0.9% | 354 | 0.7% | 259 | 0.5% | 211 | 0.4% | 205 | 0.4% |
| 4. Agriculture | 6'092 | 11.5% | 5'819 | 11.3% | 5'496 | 10.6% | 5'474 | 10.1% | 5'556 | 10.7% |
| 6. Waste | 1'007 | 1.9% | 847 | 1.6% | 741 | 1.4% | 654 | 1.2% | 637 | 1.2% |
| 7. Other | 12 | 0.0% | 13 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% |
| Total (excluding LULUCF) | 52'890 | 100.0% | 51'576 | 100.0% | 51'775 | 100.0% | 54'209 | 100.0% | 51'910 | 100.0% |

| Source and Sink Categories | 2008 | | 2009 | | 2010 | | 2011 | | 2012 | |
|--|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| 1. Energy | 43'628 | 81.3% | 42'506 | 81.2% | 44'004 | 81.3% | 39'945 | 79.9% | 41'477 | 80.6% |
| 1A1 Energy Industries | 4'025 | 7.5% | 3'949 | 7.5% | 4'180 | 7.7% | 3'968 | 7.9% | 4'064 | 7.9% |
| 1A2 Manufacturing Industries and Construction | 6'173 | 11.5% | 5'798 | 11.1% | 5'954 | 11.0% | 5'449 | 10.9% | 5'515 | 10.7% |
| 1A3 Transport | 16'624 | 31.0% | 16'427 | 31.4% | 16'321 | 30.2% | 16'205 | 32.4% | 16'331 | 31.7% |
| 1A4 Other Sectors | 16'450 | 30.7% | 15'983 | 30.5% | 17'193 | 31.8% | 13'994 | 28.0% | 15'240 | 29.6% |
| 1A5 Other (Military) | 115 | 0.2% | 116 | 0.2% | 121 | 0.2% | 108 | 0.2% | 116 | 0.2% |
| 1B Fugitive emissions from oil and natural gas | 242 | 0.5% | 232 | 0.4% | 234 | 0.4% | 220 | 0.4% | 209 | 0.4% |
| 2. Industrial Processes | 3'535 | 6.6% | 3'446 | 6.6% | 3'634 | 6.7% | 3'642 | 7.3% | 3'628 | 7.1% |
| 3. Solvent and Other Product Use | 202 | 0.4% | 201 | 0.4% | 199 | 0.4% | 202 | 0.4% | 200 | 0.4% |
| 4. Agriculture | 5'645 | 10.5% | 5'587 | 10.7% | 5'637 | 10.4% | 5'572 | 11.1% | 5'539 | 10.8% |
| 6. Waste | 629 | 1.2% | 612 | 1.2% | 608 | 1.1% | 598 | 1.2% | 591 | 1.1% |
| 7. Other | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% |
| Total (excluding LULUCF) | 53'653 | 100.0% | 52'366 | 100.0% | 54'095 | 100.0% | 49'973 | 100.0% | 51'449 | 100.0% |

ES.3.2 KP-LULUCF Activities

An overview of net CO₂ equivalent emissions and removals of activities under Article 3, paragraph 3 and Forest Management under paragraph 4 of the Kyoto Protocol is shown in Table E-6. In 2012, Deforestations were responsible for an emission of 221.87 Gg CO₂ equivalent, whereas Afforestations stored -17.13 Gg CO₂ equivalent and Forest Management -2236.38 Gg CO₂ equivalent.

Detailed quantitative information of the inventory years 2008, 2009, 2010, 2011 and 2012 as well as data for the previous years 1999–2007 are reported in Chapter 11.4, Chapter 11.5 and displayed in Table 11-4. Annual changes in the emissions from Deforestation can directly be attributed to the changes in the area of Deforestations. Year-to-year fluctuations in removals from Afforestations are mainly due to changes in the yearly afforested area and the application of a logistical growth curve for Afforestations. Fluctuations in the contribution of Forest Management can mainly be explained by differences in the losses of living (cut and mortality) and dead biomass (dead wood and litter), whereas changes in the area of managed forest are relatively small.

Table E-6 Contribution of the carbon pools under Activities under Article 3, paragraph 3 and paragraph 4 (Forest Management) of the Kyoto Protocol, Gg CO₂ eq., 1999-2012.

| Greenhouse gas source and sink activities | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|---|---|----------------|---------------|---------------|-----------------|-----------------|-----------------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| A. Article 3.3 activities | 216.97 | 216.83 | 215.84 | 214.33 | 212.51 | 210.95 | 174.16 |
| A.1. Afforestation and Reforestation | -6.09 | -6.82 | -7.80 | -9.17 | -10.83 | -12.60 | -14.86 |
| A.1.1. Units of land not harvested since the beginning of the commitment period | -6.09 | -6.82 | -7.80 | -9.17 | -10.83 | -12.60 | -14.86 |
| A.1.2. Units of land harvested since the beginning of the commitment period | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| A.2. Deforestation | 223.06 | 223.66 | 223.63 | 223.50 | 223.34 | 223.55 | 189.02 |
| B. Article 3.4 activities | -3330.01 | -395.31 | 166.35 | 219.39 | -2434.18 | -2810.38 | -2663.60 |
| B.1. Forest Management incl. biomass burning | -3330.01 | -395.31 | 166.35 | 219.39 | -2434.18 | -2810.38 | -2663.60 |
| gains above ground living biomass | -9617.06 | -9624.02 | -9630.98 | -9637.94 | -9644.90 | -9651.86 | -9662.29 |
| gains below ground living biomass | -2839.47 | -2841.80 | -2844.13 | -2846.46 | -2848.79 | -2851.12 | -2856.62 |
| losses above ground living biomass | 7758.43 | 10051.40 | 10371.01 | 10243.29 | 8273.08 | 8004.19 | 8365.83 |
| losses below ground living biomass | 2252.11 | 2847.55 | 2925.45 | 2894.29 | 2392.50 | 2319.62 | 2410.48 |
| changes litter | -335.15 | -271.92 | -166.56 | -34.20 | -201.14 | -218.05 | -467.97 |
| changes dead wood | -555.68 | -563.91 | -495.37 | -423.47 | -435.68 | -418.28 | -458.88 |
| changes soil C min. soils | -2.28 | -2.95 | -3.46 | -3.79 | -4.07 | -4.50 | -5.04 |
| changes soil C org. soils | 8.68 | 8.68 | 8.68 | 8.69 | 8.69 | 8.69 | 8.70 |
| sum forest management excl. biomass burning | -3330.43 | -396.98 | 164.63 | 200.40 | -2460.30 | -2811.31 | -2665.78 |
| biomass burning | 0.42 | 1.67 | 1.71 | 18.99 | 26.12 | 0.93 | 2.18 |
| B.2. Cropland Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.3. Grazing Land Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.4. Revegetation (if elected) | NA | NA | NA | NA | NA | NA | NA |

| Greenhouse gas source and sink activities | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|---|---|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| A. Article 3.3 activities | 147.05 | 117.04 | 81.74 | 162.22 | 197.10 | 201.52 | 204.74 |
| A.1. Afforestation and Reforestation | -17.18 | -20.14 | -22.17 | -24.33 | -23.34 | -19.62 | -17.13 |
| A.1.1. Units of land not harvested since the beginning of the commitment period | -17.18 | -20.14 | -22.17 | -24.33 | -23.00 | -18.89 | -15.89 |
| A.1.2. Units of land harvested since the beginning of the commitment period | 0.00 | 0.00 | 0.00 | 0.00 | -0.34 | -0.73 | -1.25 |
| A.2. Deforestation | 164.23 | 137.19 | 103.91 | 186.56 | 220.45 | 221.14 | 221.87 |
| B. Article 3.4 activities | -2804.38 | -1901.42 | -1202.77 | -1419.28 | -2020.23 | -2063.62 | -2236.38 |
| B.1. Forest Management incl. biomass burning | -2804.38 | -1901.42 | -1202.77 | -1419.28 | -2020.23 | -2063.62 | -2236.38 |
| gains above ground living biomass | -9844.35 | -10013.44 | -10184.68 | -10196.75 | -10201.54 | -10206.14 | -10210.74 |
| gains below ground living biomass | -2925.08 | -2989.60 | -3055.14 | -3060.06 | -3062.02 | -3063.86 | -3065.70 |
| losses above ground living biomass | 8106.69 | 8426.45 | 8467.86 | 8105.81 | 7882.81 | 7839.72 | 7757.11 |
| losses below ground living biomass | 2288.18 | 2387.54 | 2417.30 | 2332.60 | 2285.48 | 2283.19 | 2264.70 |
| changes litter | -222.74 | 71.47 | 383.78 | 354.85 | 29.07 | 115.00 | 134.28 |
| changes dead wood | -214.87 | 202.23 | 761.98 | 1037.66 | 1039.35 | 954.69 | 876.33 |
| changes soil C min. soils | -5.60 | -5.63 | -5.05 | -4.07 | -3.27 | -2.69 | -2.12 |
| changes soil C org. soils | 8.71 | 8.72 | 8.73 | 8.74 | 8.74 | 8.74 | 8.74 |
| sum forest management excl. Biomass burning | -2809.05 | -1912.26 | -1205.22 | -1421.23 | -2021.39 | -2071.35 | -2237.40 |
| biomass burning | 4.68 | 10.84 | 2.45 | 1.95 | 1.16 | 7.73 | 1.02 |
| B.2. Cropland Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.3. Grazing Land Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.4. Revegetation (if elected) | NA | NA | NA | NA | NA | NA | NA |




Figure E-3: Emissions (positive sign) and removals (negative sign) of CO₂ eq from Afforestation and Deforestation under Article 3, paragraph 3 and Forest Management under Article 3, paragraph 4, 1999-2012.

ES.3.3 GHG Inventory (Kyoto Protocol)

Relevant emissions and removals under the Kyoto Protocol are shown in table E-7 and E-8, sorted by sectors and gases respectively. The reported total emissions differ from those reported under the UNFCCC, as sector 7 Other – in addition to LULUCF and international bunkers – is not accounted for under the Kyoto Protocol. On the other hand, activities under article 3.3 (Afforestation, Reforestation and Deforestation) and 3.4 (forest, cropland and grazing management and revegetation) are taken into account over the commitment period 2008-2012. Under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol, Switzerland only accounts for Forest Management. Base year emissions (as shown in tables E-7 and E-8) for the first commitment period are fixed at the value reported in the Initial Report 2006 (FOEN 2006h, UNFCCC 2007a).

Table E-7 Summary of Switzerland's GHG emissions in CO₂ equivalent (Gg), 1990–2012 excluding emissions from sectors LULUCF, Other and International Bunkers.

| Annex A sources | Sector | Base year initial report | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| Annex A sources | 1 Energy | 42'134 | 41'989 | 44'121 | 44'191 | 41'915 | 41'029 | 41'916 | 42'787 | 42'147 | 43'474 |
| | 2 Industrial Processes | 3'258 | 3'320 | 2'957 | 2'793 | 2'484 | 2'649 | 2'626 | 2'498 | 2'431 | 2'517 |
| | 3 Solvent and Other Product Use | 466 | 470 | 444 | 420 | 392 | 374 | 354 | 331 | 308 | 286 |
| | 4 Agriculture | 5'903 | 6'092 | 6'069 | 5'979 | 5'877 | 5'843 | 5'819 | 5'780 | 5'606 | 5'578 |
| | 6 Waste | 1'030 | 1'007 | 1'004 | 976 | 914 | 852 | 847 | 823 | 814 | 795 |
| | Total (Annex A sources) | 52'791 | 52'878 | 54'595 | 54'358 | 51'582 | 50'747 | 51'563 | 52'220 | 51'305 | 52'650 |

| Annex A sources | Sector | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| Annex A sources | 1 Energy | 43'533 | 42'429 | 43'330 | 42'297 | 43'497 | 43'922 | 44'432 | 44'084 | 42'078 | 43'628 |
| | 2 Industrial Processes | 2'580 | 2'836 | 2'938 | 2'922 | 2'970 | 3'235 | 3'425 | 3'397 | 3'421 | 3'535 |
| | 3 Solvent and Other Product Use | 273 | 259 | 245 | 234 | 225 | 212 | 211 | 206 | 205 | 202 |
| | 4 Agriculture | 5'511 | 5'496 | 5'561 | 5'536 | 5'461 | 5'447 | 5'474 | 5'493 | 5'556 | 5'645 |
| | 6 Waste | 768 | 741 | 717 | 708 | 669 | 673 | 654 | 653 | 637 | 629 |
| | Total (Annex A sources) | 52'666 | 51'761 | 52'790 | 51'696 | 52'821 | 53'489 | 54'195 | 53'832 | 51'896 | 53'639 |

| KP-LULUCF | Art. 3.3 | Sector | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|-----------|----------|-------------------------------|---------------------------------|------|------|------|------|------|------|------|------|---------------|
| | | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | | |
| KP-LULUCF | Art. 3.3 | Afforestation & reforestation | | | | | | | | | | -22 |
| | | Deforestation | | | | | | | | | | 104 |
| | | Forest management | | | | | | | | | | -1'203 |
| | Art. 3.4 | Cropland management | | | | | | | | | | NA |
| | | Grazing land management | | | | | | | | | | NA |
| | | Revegetation | | | | | | | | | | NA |
| | | Total (Art. 3.3 + 3.4) | | | | | | | | | | -1'121 |

| Annex A sources | Sector | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|---------------------|
| | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | |
| Annex A sources | 1 Energy | 42'506 | 44'004 | 39'945 | 41'477 | -1.6% |
| | 2 Industrial Processes | 3'446 | 3'634 | 3'642 | 3'628 | 11.4% |
| | 3 Solvent and Other Product Use | 201 | 199 | 202 | 200 | -57.1% |
| | 4 Agriculture | 5'587 | 5'637 | 5'572 | 5'539 | -6.2% |
| | 6 Waste | 612 | 608 | 598 | 591 | -42.5% |
| | Total (Annex A sources) | 52'352 | 54'081 | 49'959 | 51'435 | -2.6% |

| KP-LULUCF | Art. 3.3 | Sector | 2009 | 2010 | 2011 | 2012 |
|-----------|----------|-------------------------------|---------------------------------|---------------|---------------|---------------|
| | | | CO ₂ equivalent (Gg) | | | |
| | | | | | | |
| KP-LULUCF | Art. 3.3 | Afforestation & reforestation | -24 | -23 | -20 | -17 |
| | | Deforestation | 187 | 220 | 221 | 222 |
| | | Forest management | -1'419 | -2'020 | -2'064 | -2'236 |
| | Art. 3.4 | Cropland management | NA | NA | NA | NA |
| | | Grazing land management | NA | NA | NA | NA |
| | | Revegetation | NA | NA | NA | NA |
| | | Total (Art. 3.3 + 3.4) | -1'257 | -1'823 | -1'862 | -2'032 |

Table E-8 Switzerland's total GHG emissions (excluding LULUCF, Other and International Bunkers) and the contribution of individual gases in CO₂ equivalent (Gg), 1990-2012.

| Annex A sources | GHG | Base year initial report | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|-----------------|--------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| | CO ₂ | 44'553 | 44'628 | 46'357 | 46'247 | 43'706 | 42'969 | 43'672 | 44'328 | 43'502 | 44'799 |
| | CH ₄ | 4'370 | 4'545 | 4'551 | 4'460 | 4'367 | 4'315 | 4'310 | 4'269 | 4'186 | 4'155 |
| | N ₂ O | 3'623 | 3'461 | 3'456 | 3'426 | 3'339 | 3'299 | 3'287 | 3'283 | 3'165 | 3'154 |
| | HFCs | 0.0 | 0.0 | 0.2 | 7 | 15 | 34 | 182 | 228 | 301 | 358 |
| | PFCs | 100 | 100 | 85 | 69 | 30 | 18 | 15 | 17 | 20 | 23 |
| | SF ₆ | 144 | 144 | 146 | 148 | 126 | 112 | 98 | 94 | 131 | 160 |
| | Total (Annex A sources) | 52'791 | 52'878 | 54'595 | 54'358 | 51'582 | 50'747 | 51'563 | 52'220 | 51'305 | 52'650 |

| Annex A sources | GHG | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|-----------------|--------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| | CO ₂ | 44'881 | 43'939 | 44'891 | 43'831 | 45'012 | 45'637 | 46'277 | 45'898 | 43'918 | 45'434 |
| | CH ₄ | 4'063 | 3'985 | 3'972 | 3'922 | 3'828 | 3'797 | 3'791 | 3'793 | 3'786 | 3'840 |
| | N ₂ O | 3'118 | 3'109 | 3'127 | 3'099 | 3'039 | 2'992 | 2'976 | 2'971 | 3'001 | 3'039 |
| | HFCs | 421 | 501 | 597 | 635 | 710 | 820 | 905 | 936 | 976 | 1'042 |
| | PFCs | 36 | 69 | 45 | 40 | 57 | 53 | 33 | 33 | 29 | 39 |
| | SF ₆ | 147 | 158 | 157 | 168 | 174 | 190 | 213 | 201 | 186 | 245 |
| | Total (Annex A sources) | 52'666 | 51'761 | 52'790 | 51'696 | 52'821 | 53'489 | 54'195 | 53'832 | 51'896 | 53'639 |

| KP-LULUCF | Art.3.3 | GHG | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------|---------|--------------------------------|---------------------------------|---------------|---------------|---------------|------------------|
| | | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | | |
| | | CO ₂ | 44'267 | 45'910 | 41'835 | 43'238 | -3.0% |
| | | CH ₄ | 3'776 | 3'763 | 3'710 | 3'688 | -15.6% |
| | | N ₂ O | 3'003 | 3'077 | 3'014 | 3'007 | -17.0% |
| | | HFCs | 1'083 | 1'138 | 1'195 | 1'245 | NA |
| | | PFCs | 36 | 37 | 40 | 33 | -67.0% |
| | | SF ₆ | 187 | 155 | 164 | 224 | 56.0% |
| | | Total (Annex A sources) | 52'352 | 54'081 | 49'959 | 51'435 | -2.6% |

| KP-LULUCF | Art.3.4 | GHG | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------|---------|-------------------------------|---------------------------------|---------------|---------------|---------------|------------------|
| | | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | | |
| | | CO ₂ | 162 | 197 | 202 | 205 | |
| | | CH ₄ | NO | NO | NO | NO | |
| | | N ₂ O | 0.0 | 0.0 | 0.0 | 0.0 | |
| | | CO ₂ | -1'420 | -2'021 | -2'068 | -2'237 | |
| | | CH ₄ | 0.8 | 0.5 | 3.2 | 0.4 | |
| | | N ₂ O | 0.2 | 0.1 | 0.7 | 0.1 | |
| | | Total (Art. 3.3 + 3.4) | -1'257 | -1'823 | -1'862 | -2'032 | |

ES.4. Other information

Emission trends for indirect greenhouse gases show a very pronounced decline (see Table 2-6 and Figure 2-9). A strict air pollution control policy and the implementation of a large number of emission reduction measures led to a decrease of 49% to 74% in the period 1990-2012 in emissions of air pollutants. The main reduction measures were abatement of exhaust emissions from road vehicles and stationary combustion equipment, taxation of solvents and sulphured fuels, and voluntary agreements with industry sectors (FOEN 2010i, Swiss Confederation 1985, 1997).

Acknowledgements

The GHG inventory preparation is a joint effort which is based on input from many federal agencies, institutions, associations, companies and individuals. Their effort was essential for the successful completion of the present inventory report.

The Federal Office for the Environment would like to acknowledge the valuable support it has received from the many contributors to this document. In particular, it would like to thank all the data suppliers, including the Office of Environmental Protection of the Principality of Liechtenstein for providing its fossil fuel consumption data, as well as experts, authors and both national and international reviewers.

PART 1

1 Introduction

1.1 *Background Information on Swiss Greenhouse Gas Inventories, Climate Change and Supplementary Information of the Kyoto Protocol (KP)*

1.1.1 Information on Climate Change

The report of the Swiss Advisory Body on Climate Change (OcCC) provides an assessment of the observed and expected impacts of climate change on Switzerland and the vulnerability of various ecological and socio-economic systems (OcCC, 2008). In the course of the 21st century, Swiss climate is projected to depart significantly from present and past conditions (CH2011 2011). Recent data confirms a warming trend with an observed increase in mean annual temperature of 1.75°C between 1864 and 2012 (FOEN 2014d). Over the last 100 years, mean annual temperatures increased by 0.13-0.20°C per decade, with a substantially accelerated warming in recent decades. According to the non-intervention scenarios (A2, A1B), seasonal mean temperatures will rise by another 2.7 °C to 4.8 °C by the end of this century compared to the period 1980-2009. Under the climate stabilization scenario (RCP3PD), Swiss climate would still change over the next decades, but is projected to stabilize at a mean warming of 1.2-1.8°C (FOEN 2014d).

The most visible change in the Alps resulting from global warming is the retreat of glaciers, which showed a volume loss of 12% since 1999 (FOEN 2014d). The area covered by alpine glaciers continuously diminishes. From about 2'900 km² of Alpine glacier area in the mid-1970s, only 2'100 km² remained in 2003 and an estimated 1'900 km² in 2013. A dramatic future loss in glacier covered area of 50-90% by 2100 has recently been modelled for a temperature increase between 2°C and 6°C for Switzerland (FOEN 2014d).

The observed trends in precipitation are less distinct than in temperature. Compared to the last 30 years, and depending on the scenario considered, the best estimates of summer mean precipitation for all Swiss regions is projected to decrease by 8-28% over the 21st century. Uncertainties due to climate model imperfections and natural variability typically amount to 15% in precipitation (CH 2011 2011). This change in summer mean precipitation will have a marked impact on the hydrological cycle: on the Central Plateau and in the very south of Switzerland, small and medium watercourses will dry up more frequently and natural replenishment of groundwater will decrease accordingly. Apart from changes to the mean temperature and precipitation, the nature of extreme events is also expected to change (CH2011 2011). More frequent, intense and longer-lasting summer warm spells and heat waves are expected, while the number of cold winter days and nights decrease in the projections for future climate in Switzerland. This is particularly relevant for alpine areas, tourism and forestry due to the risk of more frequent floods, landslides and debris flows.

The warming trend and changing precipitation patterns are expected to have significant effects on ecosystems. The Biodiversity Monitoring Switzerland reports that impacts of climate change are already being observed. A report about climate change in Switzerland summarizes several climate change affected indicators such as the phenological spring phases, flowering indices and animal specific indices (FOEN 2014d). The indicators shows significant changes in a wide range of ecosystems during the last decades. The report also emphasizes that typical alpine vascular plants have shifted uphill over the past century. Generally, climate change is expected to affect species composition, distribution, their cycles, synchronicity, the overall genetic diversity and the provision of ecosystem services. It will enhance the vulnerability of forests and impair their protective, productive and social functions. For agriculture, a moderate warming of 2°C to 3°C might increase productivity,

however, if temperature will rise beyond that level, the increase in heat waves and drought periods would prove problematic for the cultivation of land and for livestock husbandry.

Various sectors of the Swiss economy are likely to be affected by progressing climate change. In particular, the tourism industry will be hit, as the potentially beneficial effects for summer tourism will not compensate for the loss of income in mountain resorts during winter due to scarcity of snow. Cable car stations may lose their stability due to instabilities of permafrost soils. Hydroelectric power stations may be affected by altered runoff and sediment transport regimes, and insurance companies may face increased losses due to winter storms and floods. Natural hazards and extreme weather events potentially pose a growing risk to infrastructure and human health. Heat waves and elevated tropospheric ozone levels are cause for serious concern, as evidenced by the impacts of the heat wave in 2003. Finally, it remains to be seen to what extent vector borne diseases spread due to changing climatic conditions. Recently Switzerland has analysed these challenges in detail and developed an effective adaptation strategy in order to hedge against negative effects resulting from climate change in Switzerland (FOEN, 2012b).

1.1.2 Information on the Greenhouse Gas Inventory

On 10 December 1993, Switzerland ratified the United Nations Framework Convention on Climate Change (UNFCCC). Since 1996, the submission of its national greenhouse gas inventory has been based on IPCC guidelines. From 1998 onwards, the inventories have been submitted in the Common Reporting Format (CRF). In 2004, Switzerland started submitting an annual National Inventory Report (NIR) under the UNFCCC.

On 9 July 2003, Switzerland ratified the Kyoto Protocol under the UNFCCC. In November 2006 Switzerland submitted its Initial Report under Article 7, paragraph 4 of the Kyoto Protocol (FOEN 2006h). The Swiss National Inventory System (NIS) according to Article 5.1 of the Kyoto Protocol has been implemented and is fully operational. On 6 December 2007, the NIS quality management system was certified to comply with ISO 9001:2000 requirements (SQS 2008); it has been audited and recertified in November 2010 and 2013. It includes the accounting and reporting of the national registry as well (ISO 9001:2008, SQS 2010, Swiss TS 2013). The April 2008 submission of the Swiss GHG inventory (FOEN 2008) has been Switzerland's first submission under both the UNFCCC and the Kyoto Protocol.

For the submission in 2010, the NIR has been restructured according to the new outline (UNFCCC 2009a), which includes extended reporting under the Kyoto Protocol.

The 2014 inventory submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol (FOEN 2014) includes the NIR on hand, the greenhouse gas inventory 1990–2012 and the Kyoto Protocol LULUCF tables 2008–2012 in the common reporting format as well as the SEF tables and the standard independent assessment report (SIAR) from the National Registry. As a supplement, the update of the Description of the Quality Management System (FOEN 2014a) is provided.

1.1.3 Supplementary Information Required under Art. 7.1. KP

Switzerland has decided to account for Forest Management under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol. In accordance with the Annex to Decision 16/CMP.1, credits from Forest Management are capped in the first commitment period. For Switzerland, the cap amounts to 1.83 Mt CO₂ (0.5 Mt C) per year, or 9.167 Mt CO₂ for the whole commitment period.

Switzerland has chosen to account annually for emissions and removals from the LULUCF sector. The current submission contains the mandatory inventory years 2008–2012 in the

Common Reporting Format. In addition, Switzerland includes KP-LULUCF information for the years 1999-2007 on a voluntary basis in the NIR.

1.2 Institutional Arrangements for Inventory Preparation

1.2.1 Overview of Institutional, Legal and Procedural Arrangements for Compiling GHG Inventory and Supplementary Information for KP

The Swiss National Inventory System (NIS) is developed and managed under the auspices of the Federal Department of the Environment, Transport, Energy and Communications (DETEC). It is hosted by a DETEC agency, the Federal Office for the Environment (FOEN). As stipulated in the Ordinance on the Internal Organization of DETEC of 13 December 2005, this agency has the lead within the federal administration regarding climate policy and its implementation.

As part of a comprehensive project (Swiss Climate Reporting Project), the FOEN directorate mandated its Economics, Research and Environmental Observation Division in early 2004 to design and establish the NIS in order to ensure full compliance with the reporting requirements of the UNFCCC and the Kyoto Protocol by 2006. Today, the NIS is fully operational. The responsibility lies within the Climate Division of the FOEN which was established on 1st January 2010. Having regard to the provisions of Art. 5, paragraph 1 of the Kyoto Protocol, the NIS covers the following elements:

- arrangements with partner institutions, relating to roles and responsibilities, participation in the inventory development process, data use, communication and publication,
- inventory development plan,
- QA/QC system,
- official consideration and approval of data,
- upgrading and updating of the national air pollution database (EMIS),
- data documentation and storage.

With the formal approval of Switzerland's initial report under article 7, paragraph 4 of the Kyoto Protocol (FOEN 2006h) by the Federal Council on 8 November 2006 the Swiss NIS became operative. By providing for structures and in defining tasks and responsibilities of institutions, organisations and consultants involved, the NIS itself is a key tool in ensuring and improving the quality as well as the process management of inventory preparation. Figure 1-1 gives a schematic overview of the institutional setting of the NIS.




Figure 1-1 Institutional setting of the National Inventory System. The red frame marks the institutions that are included in the ISO 9001:2008 certification.

The **national inventory system supervisory board (NISSB)** was established by decision of the FOEN directorate in summer 2006. The board oversees activities related to the GHG inventory and to the national registry. It is independent of the inventory preparation and the registry administration and, by its composition, combines technical expertise and political authority. In order to put more emphasis on operational and security issues of the national registry, the national supervisory board has updated its formal mandate in 2011 to explicitly cover registry specific issues and assign the corresponding responsibilities.

The main tasks of the national supervisory board are:

- official consideration of the annual inventory submission and recommendation of the inventory for official approval by the FOEN directorate;
- assessment and approval of the recalculation of inventory data;
- handling of any issues arising from the UNFCCC review process that cannot be resolved at the level of the inventory project management or the registry administration;
- facilitation of any non-technical negotiation, consideration or approval processes involving other institutions within the federal administration;
- support of the registry administration in maintaining a secure and reliable registry environment.

The **national registry** is largely run independently of the national greenhouse gas inventory. Its operation is coordinated by the **registry administrator**, whose work is overseen by the registry **quality supervisor**.

The GHG **QA/QC officer** is responsible for enforcement of the defined quality standards of the national inventory. The officer also advises the national supervisory board on matters

relating to the conformity of the inventory with reporting requirements. Tasks and competencies are described in detail in the Description of the Quality Management System (FOEN 2014a), annexed to this report.

The **GHG inventory working group** encompasses all technical personnel involved in the inventory preparation process or representing institutions that play a significant role as suppliers of data. The group as a whole meets at least once per year to take stock of the state of the inventory, discuss priorities in the inventory development process, and to address specific issues of general interest that arise, e.g., from domestic or international reviews.

The **GHG inventory core group** comprises the inventory experts employed at the FOEN or mandated on a regular basis, which are entrusted with specific, major responsibilities for inventory planning, preparation and/or management. All inventory data are assembled and prepared for input into the CRF Reporter by the GHG inventory core group, which is responsible for ensuring the conformity of the inventory with the updated UNFCCC Reporting Guidelines on Annual Inventories (UNFCCC 2006b) and the 2008 Kyoto Protocol Reference Manual (UNFCCC 2008). Further details of the function of the core group and the roles and responsibilities of its members are given in the Description of the Quality Management System (FOEN 2014a).

The core group consists of

- the inventory project management (with overall responsibility for the integrity of the inventory, communication of data, and information exchange with the UNFCCC secretariat);
- the national inventory compiler (responsible for the EMIS inventory data base, key category analyses, and for the CRF-tables);
- the NIR lead authors (responsible for the inventory report and carrying out centralized data assessments such as uncertainty analysis);
- selected sectoral experts.

The QA/QC officer, albeit no formal member, attends the meetings of the core group.

The GHG inventory core group coordinates and integrates the activities of data suppliers within and outside the FOEN as well as those of mandated experts. Further data suppliers contributing to the inventory are research institutions and industry associations (Table 1-1). The latter are obliged by Art. 46 of the Environmental Protection Act (Swiss Confederation 1983) to provide the authorities with the information needed to enforce the law and, where necessary, to carry out inquiries.

The formal arrangements (agreements, contracts, and documentations of roles and responsibilities) that have been established to consolidate and formalize cooperation between the relevant partners contributing to, or involved in, the GHG inventory preparation process are described in Chapter H.1.1 of Switzerland's Initial Report under Article 7, paragraph 4 of the Kyoto Protocol (FOEN 2006h). Changes to the national system are reported in chapter 13 of the NIR.

Information relating to the Swiss GHG inventory is made publicly accessible through a website hosted by FOEN (www.climatereporting.ch), where detailed contact information is also available.

Table 1-1 Suppliers of raw and processed data: 1–15 provide annual updates; 16–20 provide sporadic updates. The IPCC nomenclature (IPCC 1997a) is used for the inventory categories (1A1 = Energy Industries, 1A2 = Manufacturing Industries and Construction etc.). RA = Reference Approach. For further abbreviations and acronyms see the glossary. Coloured boxes mark those sectors to which each data supplier contributes.

| | Institution | Subject | Data supplied for inventory category | | | | | | | | | | | | | |
|----|--------------------------------------|--|--------------------------------------|-----|-----|-----|-----|----|----|---|---|---|------|---|---|--|
| | | | 1A1 | 1A2 | 1A3 | 1A4 | 1A5 | 1B | RA | 2 | 3 | 4 | 5/KP | 6 | 7 | |
| | Data suppliers (annual updates) | | | | | | | | | | | | | | | |
| 1 | FOEN, Air Pollution Control | EMIS Database | | | | | | | | | | | | | | |
| 2 | FOEN, Waste and Raw Materials | Waste Statistics | | | | | | | | | | | | | | |
| 3 | FOEN, Forest Div. | Forest Statistics | | | | | | | | | | | | | | |
| 4 | SFOE | Swiss overall energy statistics | | | | | | | | | | | | | | |
| 5 | SFOE | Swiss wood energy statistics | | | | | | | | | | | | | | |
| 6 | FOCA | Civil Aviation | | | | | | | | | | | | | | |
| 7 | Swiss Air Force Administration | Military Aviation | | | | | | | | | | | | | | |
| 8 | SFSO | Agriculture, LULUCF | | | | | | | | | | | | | | |
| 9 | Agroscope | Agriculture, LULUCF | | | | | | | | | | | | | | |
| 10 | WSL | National Forest Inventory | | | | | | | | | | | | | | |
| 11 | Prognos | Energy Consum-ption | | | | | | | | | | | | | | |
| 12 | Carbotech | F-gases | | | | | | | | | | | | | | |
| 13 | Industry Assoc.: Swissmem, VSAI etc. | Ind. Processes, Solvents and Other Prod. Use | | | | | | | | | | | | | | |
| 14 | Swiss Petroleum Association | Oil Statistics | | | | | | | | | | | | | | |
| 15 | Sigmaplan, Meteotest | LULUCF | | | | | | | | | | | | | | |
| | Data suppliers (sporadic updates) | | | | | | | | | | | | | | | |
| 16 | FOEN, Air Pollution Control | Off-road Data-base, NMVOC | | | | | | | | | | | | | | |
| 17 | SGWA | Gas Distribution Losses | | | | | | | | | | | | | | |
| 18 | EMPA/Intertek | Various Em Fact. | | | | | | | | | | | | | | |
| 19 | INFRAS | On-road Emission Model | | | | | | | | | | | | | | |
| 20 | INFRAS | Off-road Emission Model | | | | | | | | | | | | | | |

1.2.2 Overview of Inventory Planning

Inventory planning, preparation, and management follow an annual cycle that is documented in Table 1 of the QMS (FOEN, 2014a). It marks milestones in the planning and preparation process in relation to QA/QC activities as specified in the quality manual. Key elements of the cycle contain:

- meetings of the supervisory board, the core group and the working group
- modelling of emissions / removals and implementation in the CRF reporter
- QA/QC activities including checklists and reviews and their inclusion in the inventory development plan
- key category and uncertainty analyses
- official consideration, approval, and submission
- publication and archiving

1.2.3 Overview of Inventory Preparation and Management, Including for Supplementary Information for Kyoto Protocol

The overall responsibility of the inventory preparation is held by the Climate Division at FOEN. The project leader coordinates the activities and oversees the compilation of the inventory and related documentation. QA/QC procedures are also coordinated by the Climate Division, and the QA/QC officer ensures archiving of all relevant data and documentation on the internal document management system of the FOEN. Details regarding the inventory preparation are given in section 1.3, while the QA/QC system is described briefly in section 1.6 and more comprehensively in the QMS supplement (FOEN 2014a).

1.3 Process for Inventory Preparation

1.3.1 GHG Inventory and KP-LULUCF Inventory

All inventory data, including activity data and emission factors for both inventories are compiled centrally by the FOEN. While emissions and removals from sector 5 LULUCF and KP-LULUCF are calculated by the Forest Division, all other sectors are calculated or compiled by the Air Pollution Control and Chemicals Division. Activity data are provided by the data suppliers (Table 1-1), while emission factors are partly updated by the data suppliers and partly by the Air Pollution Control Division.

1.3.2 Data Collection, Processing and Storage, Including for KP-LULUCF Inventory

The data needed to prepare the UNFCCC greenhouse gas inventory in the CRF is collected by the various data suppliers (Table 1-1). Since the individual data suppliers bear the main responsibility for the quality of data provided, they are also responsible for the collection of activity data, emission factors, and for the selection of methods compliant with the relevant guidelines (IPCC 1997a, 1997b, 1997c, 2000, 2003). Some data suppliers have further started to adopt the good practice guidance presented in the 2006 IPCC guidelines (IPCC 2006). Several QA/QC activities (see Chapter 1.6.1 and FOEN 2014a) ensure and continuously improve the quality of inventory data.

The Air Pollution Control and Chemical Division (formerly Air Pollution Control and Non-Ionizing Radiation Division) at the FOEN maintains the EMIS database, which contains all the basic data needed to prepare the GHG inventory in the CRF. At the same time, background information on data sources, activity data, emission factors and methods used for emission estimation is documented in the database and/or the NIR.

Figure 1-2 illustrates in a simplified manner the data collection and processing steps leading to the CRF-tables required for reporting under the UNFCCC and under the Kyoto Protocol. From EMIS, an interface transfers the data to the CRF Reporter (Version 3.7) that generates the CRF-tables that are to be submitted using the UNFCCC submission portal released in February 2009. Representative data from the CRF-tables are shown in the NIR. The NIR authors and the reviewers control the correctness of the data transferred from EMIS into the NIR. Figures and tables shown in the NIR are exported directly from EMIS. The NIR authors check the correspondence between the exports and the CRF-tables. A detailed illustration of the sectoral steps of inventory processing is given in the monitoring protocols of NIS core processes and sub-processes, as shown in a couple of examples in FOEN (2014a).




Figure 1-2 Schematic overview: Data collection for EMIS database, CRF Reporter and National Inventory Report (NIR).

1.3.3 QA /QC procedures and extensive review of GHG Inventory and KP-LULUCF Inventory

The national inventory system has an established quality management system (QMS) that complies with the requirements of ISO 9001:2008. Certification has been obtained in 2007 and upheld since through annual audits. An overview over QA/QC procedures and review activities is given in section 1.6.1, a full description of the QMS is provided as a supplement (FOEN 2014a) to the national inventory report.

1.4 Methodologies and Data Sources

1.4.1 GHG Inventory

1.4.1.1 General Description

Emissions are calculated on the basis of the standard methods and procedures published in the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 1997a, 1997b, 1997c), in IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC 2000), and in IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC 2003). Under the UNFCCC, these guidelines have been adopted for mandatory use in reporting on GHG inventories. The 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006), adopted in April 2006 by the IPCC, have been consulted in a few cases.

The national approach for sector 1 Energy is based on import and fuel consumption statistics (fuel sales in the transport sector) in Switzerland (see Chapter 1.4.1.2). The other sectors rely on national statistics and data surveys. For the various sectors, Tier 1, Tier 2 and Tier 3 methodologies according to IPCC Guidelines (IPCC 1997b) and Good Practice Guidance (IPCC 2000) are used. GHG emissions by sources and removals by sinks due to land use, land-use change and forestry (LULUCF sector) are calculated according to IPCC 2003. The following list (Table 1-2) indicates the approaches adopted.

Table 1-2 Summary table for emission factors and methods used (from CRF-tables Summary3) in 2012.

| GREENHOUSE GAS SOURCE AND SINK CATEGORIES | CO ₂ | | CH ₄ | | N ₂ O | |
|---|-----------------|-----------------|-----------------|-----------------|------------------|-----------------|
| | Method applied | Emission factor | Method applied | Emission factor | Method applied | Emission factor |
| 1. Energy | CS,D,T2,T3 | CS,D | CS,D,T2,T3 | CR,CS,D | CS,D,T2,T3 | CS,D |
| A. Fuel Combustion | CS,T2,T3 | CS | CS,T2,T3 | CR,CS | CS,D,T2,T3 | CS,D |
| 1. Energy Industries | CS,T2 | CS | CS,T2 | CS | CS,D | CS,D |
| 2. Manufacturing Industries and Constr. | CS,T2 | CS | CS,T2,T3 | CS | D | D |
| 3. Transport | T2,T3 | CS | T2,T3 | CR,CS | CS,D,T2,T3 | CS,D |
| 4. Other Sectors | CS,T2 | CS | CS,T2 | CS | D | D |
| 5. Other | T2 | CS | T2 | CS | T2 | CS |
| B. Fugitive Emissions from Fuels | CS,D | CS,D | CS,D | CS,D | D | D |
| 1. Solid Fuels | NA | NA | NA | NA | NA | NA |
| 2. Oil and Natural Gas | CS,D | CS,D | CS,D | CS,D | D | D |
| 2. Industrial Processes | CS,T2 | CS,D,PS | CS,T2 | D,PS | CS,T2 | PS |
| A. Mineral Products | CS,T2 | CS,D,PS | NA | NA | NA | NA |
| B. Chemical Industry | CS,T2 | PS | CS,T2 | D,PS | CS,T2 | PS |
| C. Metal Production | CS | CS | NA | NA | NA | NA |
| D. Other Production | NA | NA | | | | |
| E. Production of Halocarbons and SF ₆ | | | | | | |
| F. Consumption of Halocarbons and SF ₆ | | | | | | |
| G. Other | CS | CS | NA | NA | NA | NA |
| 3. Solvent and Other Product Use | CS | CS | | | CS | CS |
| 4. Agriculture | | | T2 | CS,D | CS,T1b | D |
| A. Enteric Fermentation | | | T2 | CS | | |
| B. Manure Management | | | T2 | CS,D | CS | D |
| C. Rice Cultivation | | | NA | NA | | |
| D. Agricultural Soils | | | NA | NA | CS,T1b | D |
| E. Prescribed Burning of Savannas | | | NA | NA | NA | NA |
| F. Field Burning of Agricultural Residues | | | NA | NA | NA | NA |
| G. Other | | | NA | NA | NA | NA |
| 5. Land Use, Land-Use Change and Forestry | T2 | CS | T1 | CS | T1 | D |
| A. Forest Land | T2 | CS | T1 | CS | T1 | D |
| B. Cropland | T2 | CS | NA | NA | T1 | D |
| C. Grassland | T2 | CS | T1 | CS | T1 | D |
| D. Wetlands | T2 | CS | T1 | CS | NA | NA |
| E. Settlements | T2 | CS | NA | NA | NA | NA |
| F. Other Land | T2 | CS | NA | NA | NA | NA |
| G. Other | NA | NA | NA | NA | NA | NA |
| 6. Waste | CS | CS | CS,D | CS,D | CS | CS |
| A. Solid Waste Disposal on Land | NA | NA | CS,D | CS,D | | |
| B. Waste-water Handling | | | CS,D | CS,D | CS | CS |
| C. Waste Incineration | CS | CS | CS | CS | CS | CS |
| D. Other | NA | NA | CS | CS | CS | CS |
| 7. Other (as specified in Summary 1.A) | T1 | CS | T1 | CS | CS,T1b | CS,D |

| GREENHOUSE GAS SOURCE AND SINK | HFCs | | PFCs | | SF ₆ | |
|---|-------|------|-------|------|-----------------|------|
| 2. Industrial Processes | T1,T2 | CS,D | T1,T2 | CS,D | T1,T2,T3 | CS,D |
| A. Mineral Products | | | | | | |
| B. Chemical Industry | NA | NA | NA | NA | NA | NA |
| C. Metal Production | NA | NA | NA | NA | T1,T2,T3 | D |
| D. Other Production | | | | | | |
| E. Production of Halocarbons and SF ₆ | NA | NA | NA | NA | NA | NA |
| F. Consumption of Halocarbons and SF ₆ | T1,T2 | CS,D | T1,T2 | CS,D | T1,T2,T3 | CS,D |
| G. Other | NA | NA | NA | NA | NA | NA |

1.4.1.2 National and Reference Approach for Sector 1 Energy

The Reference Approach is used as a check for (i) overall energy consumption and (ii) the resulting CO₂ emissions reported in source category 1 Energy. In Switzerland, it is applied on the basis of data published in the Swiss overall energy statistics (SFOE 2013). The results of the Reference Approach are compared with the results of the sectoral approach for sector 1 Energy in order to test the quality and completeness of the inventory. For the current

inventory, the two approaches show a good correspondence; with CO₂ emissions differing by 0.89% and energy consumption by 0.62% in 2012 (see chapter 3.2.1).

1.4.1.3 National Air Pollution Database EMIS

A large body of emission data is adopted from Switzerland's national air pollution database EMIS, which is operated by FOEN (FOEN 2006c). EMIS was established at SAEFL (former name of FOEN) in the late 1980s. Its initial purpose was to record and monitor emissions of air pollutants. It has since been extended to cover greenhouse gases, too. Its structure corresponds to the EMEP/CORINAIR system for classifying emission-generating activities. EMEP/CORINAIR uses the Nomenclature for Reporting ("NFR code", UNECE 2003). The Revised 1996 IPCC Guidelines provide a correspondence key between IPCC and EMEP/CORINAIR source categories (IPCC 1997a: Annex 2). EMIS thus contains cross-references to IPCC/UNFCCC coding formats.

EMIS calculates emissions for various pollutants using emission factors and activity data according to the EMEP/CORINAIR methodology. Pollutants in EMIS include sulphur dioxide (SO₂), nitrogen oxides (NO_x), nitrous oxide (N₂O), ammonia (NH₃), non-methane volatile organic compounds (NMVOC), carbon monoxide (CO), black carbon (BC), hydrochloric acid (HCl), particulate matter, heavy metals (lead, zinc, cadmium, mercury), polychlorinated dibenzodioxins and -furans (PCDD/PCDF), hexachlorobenzene (HCB), hydrogen fluoride (HF), hydrofluorocarbons (HFC), perfluorinated carbon compounds (PFC), sulphur hexafluoride (SF₆), methane (CH₄), carbon dioxide CO₂ (fossil/geological origin) and CO₂ (biogenic). The input data originate from a variety of sources, such as production data and emission factors from industry, industry associations and research institutions, as well as population, employment, waste and agriculture statistics. EMIS is documented in an internal FOEN manual for the database (FOEN 2006c).

The original EMIS database underwent a full redesign in 2005/2006. It was extended to incorporate more data sources, updated, and migrated to a new software platform. At the same time, activity data and emission factors were being checked and updated. Emission data from EMIS that are relevant for the GHG inventory are exported to the CRF reporter.

Input data for the EMIS database comprise the SFOE Swiss overall energy statistics, the SFOE Swiss wood energy statistics, FOEN statistics and models for emissions from road transportation, statistics and models of off-road activities, modelled emissions based on the import statistics for F-gases, waste and agricultural statistics, extracts from the National Forest Inventory and the National Forest Statistics (see Figure 1-2).

1.4.2 KP- LULUCF Inventory

Emission factors for parts of sector 5 LULUCF (forest land) and the KP-LULUCF tables are calculated by the Forest Division of the FOEN. A detailed description of the calculation of these emission factors can be found in Chapter 7.3 and Chapter 11.3. Both data sets are imported in the EMIS database (FOEN 2006c).

1.5 Description of Key Categories

1.5.1 GHG Inventory

1.5.1.1 Methodology

The key category analyses are performed according to the IPCC Good Practice Guidance (IPCC 2000, chapter 7) for 1990 and the latest reported year. A Tier 1 level and trend assessment is applied with the proposed threshold of 95%. A Tier 2 key category analyses has also been carried out for this submission with the proposed threshold of 90% of the sum

of all level assessments weighted with their relative source uncertainty. The uncertainty used for the calculations is the Tier 1 uncertainty (see chapter 1.7).

According to good practice guidance (IPCC 2000), the result of Tier 2 key category analysis should be used when results between Tier 1 and Tier 2 differ. However, it would also be possible to keep Tier 1 key categories as key categories based on qualitative criteria. The GHG inventory core group has agreed to keep Tier 1 key categories in this submission as key categories, even if they are not key in Tier 2 (and vice versa). This procedure would also be compatible with the 2006 IPCC Guidelines (IPCC 2006), which recommend exactly such a procedure of combining results from Tier 1 and Tier 2 categories if results from the two approaches differ. When combining Tier 1 and Tier 2 key category analysis results, we consider a category to be key because of level, if the category is key due to level according to Tier 1 or Tier 2, and a category is considered to be key because of trend, if the category is key due to trend according to Tier 1 or Tier 2.

1.5.1.2 KCA without LULUCF categories

Tier 1

For 2012, among a total of 135 categories, 31 have been identified as key categories with an aggregated contribution of 97.2% to total national emissions. 23 categories are key due to the level assessment, 27 due to the trend assessment.

Of the 31 key categories, 19 are in sector 1 Energy, accounting for 79.5% of total CO₂ equivalent emissions in 2012. The other key categories are from sectors 2 Industrial Processes (5.7%), 3 Solvent and Other Product Use (0.3%), 4 Agriculture (10.7%) and 6 Waste (1.0%).

There are three major key sources each contributing more than 10 % to the level assessment:

- 1A3b Energy, Fuel Combustion, Road Transportation, Gasoline, CO₂, level contribution 17.5%
- 1A4b Energy, Fuel Combustion, Other Sectors, Residential, Liquid Fuels, CO₂, level contribution 14.3%
- 1A3b Energy, Fuel Combustion, Road Transportation, Diesel, CO₂, level contribution 13.2%

Compared to the key category analysis in the previous inventory report of April 2013 (FOEN 2013), the following category is a new key category:

- CO₂ emissions from 1A3b Road Transportation, Gaseous Fuels: the reason for being a new key category is the significant higher activity data based on recalculations in in this source category (see chapter 3.2.8.5)

The following category is no longer key category in Tier 1 compared to the previous submission of April 2013:

- CO₂ emissions from 2C1 Industrial Processes, Steel Production: The reason for not being key category anymore is the significant reduction of emissions based on recalculations in the implied emission factor in this sector (see chapter 4.4.5)

The following table shows the contributions of the individual key categories. The complete results of the key category analysis for 2012 are given in Annex A1.2.

Table 1-3 List of Switzerland's Tier 1 key categories 2012 without LULUCF categories, sorted by category code.

| Tier 1 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | | |
|---|--|----------------------------------|--|-----------------------------------|---------------|------------|---|--|---------------|---------------|--------------------|----------------------|----------------------|
| A | | | | | | B | C | D | E-L | E-T | F-T | M | N |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | Direct GHG | Base Year 1990 Estimate [Gg CO ₂ eq] | Year 2012 Estimate [Gg CO ₂ eq] | Level Assessm | Trend Assessm | % Contrib in Trend | Result level assessm | Result trend assessm |
| 1 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Gaseous Fuels | CO2 | 289.73 | 498.74 | 0.97% | 0.00434 | 1.0% | KC level | KC trend |
| 2 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 1.56% | 0.00261 | 0.6% | KC level | KC trend |
| 3 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 5.28% | 0.02471 | 5.8% | KC level | KC trend |
| 4 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 4.07% | 0.02102 | 5.0% | KC level | KC trend |
| 5 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 5.13% | 0.01900 | 4.5% | KC level | KC trend |
| 6 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Other Fuels | CO2 | 134.15 | 288.60 | 0.56% | 0.00316 | 0.7% | KC level | KC trend |
| 7 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.88% | 0.01432 | 3.4% | KC level | KC trend |
| 8 | 1A3a | 1. Energy | A. Fuel Comb. | 3. Transport; Civil Aviation | | CO2 | 252.55 | 136.65 | 0.27% | 0.00218 | 0.5% | | KC trend |
| 9 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Diesel | CO2 | 2587.68 | 6767.05 | 13.15% | 0.08494 | 20.1% | KC level | KC trend |
| 10 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | CO2 | 11335.27 | 9016.58 | 17.53% | 0.04013 | 9.5% | KC level | KC trend |
| 11 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | N2O | 142.38 | 27.52 | 0.05% | 0.00222 | 0.5% | - | KC trend |
| 12 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gaseous Fuels | CO2 | 0.00 | 83.59 | 0.16% | 0.00167 | 0.4% | - | KC trend |
| 13 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 2.88% | 0.01044 | 2.5% | KC level | KC trend |
| 14 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 5.91% | 0.02881 | 6.8% | KC level | KC trend |
| 15 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 5.15% | 0.02527 | 6.0% | KC level | KC trend |
| 16 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 14.33% | 0.05183 | 12.3% | KC level | KC trend |
| 17 | 1A4c | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Agric./Forestry | Liquid Fuels | CO2 | 547.34 | 540.01 | 1.05% | 0.00015 | 0.0% | KC level | - |
| 18 | 1A5 | 1. Energy | A. Fuel Comb. | 5. Other | Liquid Fuels | CO2 | 203.58 | 114.80 | 0.22% | 0.00166 | 0.4% | - | KC trend |
| 19 | 1B2 | 1. Energy | B. Fugitive Emissi | 2. Oil and Natural Gas | | CH4 | 263.72 | 169.45 | 0.33% | 0.00174 | 0.4% | - | KC trend |
| 20 | 2A1 | 2. Ind. Prod | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 1787.11 | 3.47% | 0.01336 | 3.2% | KC level | KC trend |
| 21 | 2F1 | 2. Ind. Prod | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | | HFC | 0.02 | 1137.81 | 2.21% | 0.02274 | 5.4% | KC level | KC trend |
| 22 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 155.28 | 0.30% | 0.00389 | 0.9% | - | KC trend |
| 23 | 4A | 4. Agric. | A. Enteric Fermentation | | | CH4 | 2635.45 | 2496.98 | 4.85% | 0.00132 | 0.3% | KC level | - |
| 24 | 4B | 4. Agric. | B. Manure Management | | | CH4 | 671.61 | 646.11 | 1.26% | 0.00014 | 0.0% | KC level | - |
| 25 | 4B | 4. Agric. | B. Manure Management | | | N2O | 454.68 | 335.81 | 0.65% | 0.00213 | 0.5% | KC level | KC trend |
| 26 | 4D1 | 4. Agric. | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 1143.10 | 2.22% | 0.00342 | 0.8% | KC level | KC trend |
| 27 | 4D2 | 4. Agric. | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 220.79 | 0.43% | 0.00192 | 0.5% | KC level | KC trend |
| 28 | 4D3 | 4. Agric. | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 674.93 | 1.31% | 0.00250 | 0.6% | KC level | KC trend |
| 29 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 158.26 | 0.31% | 0.01021 | 2.4% | - | KC trend |
| 30 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 240.28 | 0.47% | 0.00121 | 0.3% | KC level | - |
| 31 | 6D | 6. Waste | D. Other | | | CH4 | 29.94 | 113.76 | 0.22% | 0.00169 | 0.4% | - | KC trend |

Table 1-4 List of Switzerland's Tier 1 key categories for the base year 1990 without LULUCF categories, sorted by category code.

| Tier 1 Key category analysis for the base year 1990 without LULUCF categories | | | | | | | | | |
|---|---|----------------------------------|--|--|---------------|------------------|------------------------------|-----------------------|----------------------------|
| No. | A IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | B Direct GHG | C Base Year 1990 Estimate | E-L Level Assessm. | M Result level assessm. |
| 1 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CO ₂ | 289.73 | 0.55% | KC level |
| 2 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO ₂ | 693.69 | 1.31% | KC level |
| 3 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO ₂ | 1519.73 | 2.87% | KC level |
| 4 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Gaseous Fuels | CO ₂ | 1074.09 | 2.03% | KC level |
| 5 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | CO ₂ | 3692.22 | 6.98% | KC level |
| 6 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Solid Fuels | CO ₂ | 1204.47 | 2.28% | KC level |
| 7 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CO ₂ | 252.55 | 0.48% | KC level |
| 8 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CO ₂ | 2587.68 | 4.89% | KC level |
| 9 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO ₂ | 11335.27 | 21.43% | KC level |
| 10 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Gaseous Fuels | CO ₂ | 987.24 | 1.87% | KC level |
| 11 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | CO ₂ | 4606.43 | 8.71% | KC level |
| 12 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO ₂ | 1424.38 | 2.69% | KC level |
| 13 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO ₂ | 10248.79 | 19.38% | KC level |
| 14 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CO ₂ | 547.34 | 1.03% | KC level |
| 15 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO ₂ | 203.58 | 0.38% | KC level |
| 16 | 1B2 | 1. Energy | B. Fugitive Emissions from | 2. Oil and Natural Gas | | CH ₄ | 263.72 | 0.50% | KC level |
| 17 | 2A1 | 2. Industrial P | A. Mineral Products; Cement Production-CO ₂ | | | CO ₂ | 2524.68 | 4.77% | KC level |
| 18 | 3 | 3. Solvent and Other Product Use | | | | CO ₂ | 360.04 | 0.68% | KC level |
| 19 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH ₄ | 2635.45 | 4.98% | KC level |
| 20 | 4B | 4. Agriculture | B. Manure Management | | | CH ₄ | 671.61 | 1.27% | KC level |
| 21 | 4B | 4. Agriculture | B. Manure Management | | | N ₂ O | 454.68 | 0.86% | KC level |
| 22 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N ₂ O | 1351.48 | 2.56% | KC level |
| 23 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N ₂ O | 822.48 | 1.56% | KC level |
| 24 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH ₄ | 688.16 | 1.30% | KC level |

There are 24 level key categories in the base year 1990 (see Table 1-4). All of them are also key categories in 2012.

Compared to the key category analysis in the previous inventory report of April 2013, the key categories in the base year 1990 are the same.

Tier 2

For 2012, among a total of 135 categories, 30 have been identified as key categories with an aggregated contribution of 94.0% of the sum of all level assessments weighted with their

uncertainty in 2012. 23 categories are key due to the level assessment, 27 due to the trend assessment.

Of the 30 key categories, 14 are in sector 1 Energy, accounting for 29.8% of the sum of all level assessments weighted with their uncertainty in 2012 (12.5%, see Table A - 4). Sector 4 Agriculture accounts for 51.0% of that sum. The other key categories are from sectors 2 Industrial Processes (6.3%), 3 Solvent and Other Product Use (1.8%) and 6 Waste (5.1%). There are three major key categories contributing more than 10% to the level assessment weighted with their uncertainty:

- 4D3, Agricultural Soils; Indirect Emissions, N₂O, contribution of 17.4% to the sum of all level assessments weighted with their uncertainty.
- 4D1, Agricultural Soils; Direct Soil Emissions, N₂O, contribution of 14.8% to the sum of all level assessments weighted with their uncertainty.
- 1A1 Energy, Fuel Combustion, Energy Industries, Other Fuels, CO₂, contribution of 13.3% to the sum of all level assessments weighted with their uncertainty.

Table 1-5 shows the contributions of the individual key categories. The complete results of the key category analysis for 2012 are given in Annex A1.4.

Compared to the submission of April 2013, the following categories are new key category in Tier 2:

- CO₂ emissions from 2A3 Industrial Processes, Limestone and Dolomite use: the reason for being new key category is an increase in emissions based on recalculations in the implied emission factor in this sector (see chapter 4.2.5)
- SF₆ emissions from 2F9 Industrial Processes, consumption of Halocarbons and SF₆: the reason for being new key category is an increase in the emissions in 2012 compared to 2011.

No longer key in Tier 2 is the following category:

- CO₂ emissions from 2C1 Industrial Processes, Steel Production: the reason for not being key category anymore is the significant reduction of emissions based on recalculations in the implied emission factor in this sector (see chapter 4.4.5)

Table 1-5 List of Switzerland's Tier 2 key categories 2012 without LULUCF categories, sorted by category code.

| Tier 2 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | | |
|---|--|----------------------------------|---|--------------------------------|---------------|------------|--|-----------------------------------|---------------------------------|---------------------------------|---------------------|----------------------|----------------------|
| A | | | | | | B | C | D | E-L | E-T | F-T | M | N |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | Level Assessm. with Uncertainty | Trend Assessm. with Uncertainty | % Contrib. in Trend | Result level assessm | Result trend assessm |
| 1 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 1.67% | 0.00781 | 15.3% | KC level | KC trend |
| 2 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 0.20% | 0.00105 | 2.1% | KC level | KC trend |
| 3 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Other Fuels | CO2 | 134.15 | 288.60 | 0.18% | 0.00100 | 2.0% | KC level | KC trend |
| 4 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.07% | 0.00111 | 2.2% | - | KC trend |
| 5 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Diesel | CO2 | 2587.68 | 6767.05 | 0.29% | 0.00190 | 3.7% | KC level | KC trend |
| 6 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | CH4 | 97.47 | 18.82 | 0.01% | 0.00056 | 1.1% | - | KC trend |
| 7 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | CO2 | 11335.27 | 9016.58 | 0.45% | 0.00103 | 2.0% | KC level | KC trend |
| 8 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | N2O | 142.38 | 27.52 | 0.03% | 0.00111 | 2.2% | - | KC trend |
| 9 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 0.14% | 0.00052 | 1.0% | KC level | KC trend |
| 10 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 0.08% | 0.00040 | 0.8% | - | KC trend |
| 11 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 33.78 | 0.04% | 0.00078 | 1.5% | - | KC trend |
| 12 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 0.26% | 0.00127 | 2.5% | KC level | KC trend |
| 13 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 0.20% | 0.00072 | 1.4% | KC level | KC trend |
| 14 | 1B2 | 1. Energy | B. Fugitive Emis | 2. Oil and Natural Gas | | CH4 | 263.72 | 169.45 | 0.10% | 0.00052 | 1.0% | KC level | KC trend |
| 15 | 2A1 | 2. Ind. Proc | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 1787.11 | 0.10% | 0.00038 | 0.7% | KC level | - |
| 16 | 2A3 | 2. Ind. Proc | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | | CO2 | 150.39 | 98.48 | 0.10% | 0.00049 | 1.0% | - | KC trend |
| 17 | 2F1 | 2. Ind. Proc | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | | HFC | 0.02 | 1137.81 | 0.27% | 0.00273 | 5.3% | KC level | KC trend |
| 18 | 2F9 | 2. Ind. Proc | F. Consumption of Halocarbons and SF6; Other | | | HFC | 0.00 | 76.14 | 0.12% | 0.00122 | 2.4% | KC level | KC trend |
| 19 | 2F9 | 2. Ind. Proc | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 135.91 | 0.21% | 0.00094 | 1.8% | KC level | KC trend |
| 20 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 155.28 | 0.15% | 0.00195 | 3.8% | KC level | KC trend |
| 21 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 44.62 | 0.07% | 0.00100 | 2.0% | - | KC trend |
| 22 | 4A | 4. Agric. | A. Enteric Fermentation | | | CH4 | 2635.45 | 2496.98 | 0.89% | 0.00024 | 0.5% | KC level | - |
| 23 | 4B | 4. Agric. | B. Manure Management | | | CH4 | 671.61 | 646.11 | 0.68% | 0.00008 | 0.2% | KC level | - |
| 24 | 4B | 4. Agric. | B. Manure Management | | | N2O | 454.68 | 335.81 | 0.41% | 0.00135 | 2.6% | KC level | KC trend |
| 25 | 4D1 | 4. Agric. | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 1143.10 | 1.85% | 0.00285 | 5.6% | KC level | KC trend |
| 26 | 4D2 | 4. Agric. | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 220.79 | 0.36% | 0.00163 | 3.2% | KC level | KC trend |
| 27 | 4D3 | 4. Agric. | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 674.93 | 2.18% | 0.00415 | 8.1% | KC level | KC trend |
| 28 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 158.26 | 0.18% | 0.00596 | 11.7% | KC level | KC trend |
| 29 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 240.28 | 0.23% | 0.00061 | 1.2% | KC level | KC trend |
| 30 | 6D | 6. Waste | D. Other | | | CH4 | 29.94 | 113.76 | 0.22% | 0.00170 | 3.3% | KC level | KC trend |

Table 1-6 List of Switzerland's Tier 2 key categories for the base year 1990 without LULUCF categories, sorted by category code.

| Tier 2 Key category analysis for the base year 1990 without LULUCF categories | | | | | | | | | |
|---|---|----------------------------------|---|-----------------------------------|---------------|------------|-------------------------------------|---------------------------------|-----------------------|
| No. | A | | | | | B | C | E-L | M |
| | IPCC Source Categories and fuels if applicable (combined without LULUCF categories) | | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Level Assessm. with Uncertainty | Result level assessm. |
| 1 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 0.91% | KC level |
| 2 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 0.18% | KC level |
| 3 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO2 | 11335.27 | 0.55% | KC level |
| 4 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | N2O | 142.38 | 0.13% | KC level |
| 5 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Com./Instit. | Liquid Fuels | CO2 | 4606.43 | 0.12% | KC level |
| 6 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 0.12% | KC level |
| 7 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 0.14% | KC level |
| 8 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 0.27% | KC level |
| 9 | 1B2 | 1. Energy | B. Fugitive Emissions from | 2. Oil and Natural Gas | | CH4 | 263.72 | 0.15% | KC level |
| 10 | 2A1 | 2. Industrial Proc. | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 0.14% | KC level |
| 11 | 2A3 | 2. Industrial Proc. | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | | CO2 | 150.39 | 0.15% | KC level |
| 12 | 2F9 | 2. Industrial Proc. | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 0.12% | KC level |
| 13 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 0.34% | KC level |
| 14 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 0.17% | KC level |
| 15 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH4 | 2635.45 | 0.91% | KC level |
| 16 | 4B | 4. Agriculture | B. Manure Management | | | CH4 | 671.61 | 0.69% | KC level |
| 17 | 4B | 4. Agriculture | B. Manure Management | | | N2O | 454.68 | 0.55% | KC level |
| 18 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 2.12% | KC level |
| 19 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 0.21% | KC level |
| 20 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 2.58% | KC level |
| 21 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 0.76% | KC level |
| 22 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 0.17% | KC level |

There are 22 level Tier 2 key categories in the base year 1990 (see Table 1-6). All of them are also key categories in 2012.

Compared to the key category analysis in the previous inventory report of April 2013, there is one new key category in 2012 for the base year 1990:

CO₂ emissions from 2A3 Industrial Processes, Limestone and Dolomite use: The reason for being new key category is an increase in emissions based on recalculations in the implied emission factor in this sector (see chapter 4.2.5).

1.5.1.3 Combined KCA without and with LULUCF categories

The key category analysis including LULUCF categories has also been carried out for 2012 and 1990. The complete results of the key category analysis for 2012 are shown in Annex

A1. According to IPCC Good Practice Guidance for LULUCF (IPCC 2003, Section 5.4.2), the set of key categories consists of all non-LULUCF key categories that result from the KCA without LULUCF combined with all LULUCF key categories that result from the KCA including LULUCF.

Tier 1

In the Tier 1 KCA for the year 2012 including LULUCF categories there are five additional categories out of the LULUCF sector:

- CO₂ emissions from 5A1 Forest Land remaining Forest Land (level and trend key category)
- CO₂ emissions from 5A2 Land converted to Forest Land (level and trend key category)
- CO₂ emissions from 5B1 Cropland remaining Cropland (level and trend key category)
- CO₂ emissions from 5C2 Land converted to Grassland (trend key category)
- CO₂ emissions from 5E2 Land converted to Settlements (level key category)

The categories 5A1 Forest Land remaining Forest Land and 5B1 Cropland remaining Cropland are large categories, contributing for 3.8% and 1.3% to the level assessment. Categories 5A2, 5C2 and 5E2 contribute less to the level assessment with 0.9%, 0.3% and 0.5%, respectively.

The five LULUCF key categories 5A1, 5A2, 5B1, 5C2 and 5E2 were also key in the analysis for 2011 as contained in the previous inventory report of April 2013 (FOEN 2013).

For the combined KCA without and with LULUCF, these categories are added to the other 31 key categories from the KCA without LULUCF.

In the KCA for the year 1990, four of these five LULUCF categories are also key categories. Categories 5C2 is not key category for the year 1990.

For the combined KCA without and with LULUCF categories, these categories are added to the other 24 key categories from the KCA without LULUCF. The results of the combined Tier 1 KCA are summarised in Table 1-7 (year 2012) and Table 1-8 (1990).

Tier 2

In the Tier 2 KCA for 2012 including LULUCF categories, there are seven additional categories out of the LULUCF sector: CO₂ emissions from 5A1, 5A2, 5B1, 5B2, 5C1, 5C2 and 5E2. Five of these categories are also key categories out of the LULUCF sector as in Tier 1. Additionally, CO₂ emissions from 5B2 and 5C1 are key in Tier 2.

The categories 5C1 Grassland remaining Grassland, 5A1 Forest Land remaining Forest Land and 5B1 Cropland remaining Cropland are large categories, contributing for 23.2%, 11.1% and 6.4% of the sum of all level assessments weighted with their uncertainty. Source categories 5A2, 5B2, 5C2 and 5E2 contribute less, with 2.7%, 0.3%, 0.9% and 1.3%, respectively.

The six of the seven LULUCF key categories were also key in the analysis for 2011 as contained in the previous inventory report of April 2013 (FOEN 2013). Source category 5B2 land converted to Cropland is new key category for 2012.

For the combined KCA without and with LULUCF categories, these categories are added to the other 30 key categories from the KCA without LULUCF.

In the KCA for the year 1990, five of these seven LULUCF categories are also key categories. Source categories 5B2 and 5C2 are not key categories for the year 1990.

Compared to the previous submission of April 2013 (FOEN 2013), 5C1 is new key category.

For the combined KCA without and with LULUCF categories, these categories are added to the other 22 key categories from the KCA without LULUCF. The results of the combined Tier 2 KCA are summarised in Table 1-9 (year 2012) and Table 1-10 (year 1990).

Table 1-7 List of Switzerland's Tier 1 key categories, combined KCA without and with LULUCF (in italic) categories 2012, sorted by category code.

| Combined Tier 1 Key category analysis 2012 without and with LULUCF categories | | | | | | | | | | | | | |
|---|--|----------------------------------|--|--------------------------------------|---------------|-------------------------------------|--------------------------------|---------------|---------------|--------------------|----------------------|----------------------|----------|
| A | | | | | B | C | D | E-L | E-T | F-T | M | N | |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | Level Assessm | Trend Assessm | % Contrib in Trend | Result level assessm | Result trend assessm | |
| 1 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Gaseous Fuels | CO2 | 289.73 | 498.74 | 0.97% | 0.00434 | 1.0% | KC level | KC trend |
| 2 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 1.56% | 0.00261 | 0.6% | KC level | KC trend |
| 3 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 5.28% | 0.02471 | 5.8% | KC level | KC trend |
| 4 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 4.07% | 0.02102 | 5.0% | KC level | KC trend |
| 5 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 5.13% | 0.01900 | 4.5% | KC level | KC trend |
| 6 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Other Fuels | CO2 | 134.15 | 288.60 | 0.56% | 0.00316 | 0.7% | KC level | KC trend |
| 7 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.88% | 0.01432 | 3.4% | KC level | KC trend |
| 8 | 1A3a | 1. Energy | A. Fuel Comb. | 3. Transport: Civil Aviation | | CO2 | 252.55 | 136.65 | 0.27% | 0.00218 | 0.5% | - | KC trend |
| 9 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport: Road Transp. | Diesel | CO2 | 2587.68 | 6767.05 | 13.15% | 0.08494 | 20.1% | KC level | KC trend |
| 10 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport: Road Transp. | Gasoline | CO2 | 11335.27 | 9016.58 | 17.53% | 0.04013 | 9.5% | KC level | KC trend |
| 11 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport: Road Transp. | Gasoline | N2O | 142.38 | 27.52 | 0.05% | 0.00222 | 0.5% | - | KC trend |
| 12 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport: Road Transp. | Gaseous Fuels | CO2 | 0.00 | 83.59 | 0.16% | 0.00167 | 0.4% | - | KC trend |
| 13 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 2.88% | 0.01044 | 2.5% | KC level | KC trend |
| 14 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Instit. | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 5.91% | 0.02881 | 6.8% | KC level | KC trend |
| 15 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 5.15% | 0.02527 | 6.0% | KC level | KC trend |
| 16 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 14.33% | 0.05183 | 12.3% | KC level | KC trend |
| 17 | 1A4c | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Agric./Forestry | Liquid Fuels | CO2 | 547.34 | 540.01 | 1.05% | 0.00015 | 0.0% | KC level | - |
| 18 | 1A5 | 1. Energy | A. Fuel Comb. | 5. Other | Liquid Fuels | CO2 | 203.58 | 114.80 | 0.22% | 0.00166 | 0.4% | - | KC trend |
| 19 | 1B2 | 1. Energy | B. Fugitive Emis | 2. Oil and Natural Gas | | CH4 | 263.72 | 169.45 | 0.33% | 0.00174 | 0.4% | - | KC trend |
| 20 | 2A1 | 2. Ind. Proc. | A. Mineral Products; Cement Production-CO2 | | CO2 | 2524.68 | 1787.11 | 3.47% | 0.01336 | 3.2% | KC level | KC trend | |
| 21 | 2F1 | 2. Ind. Proc. | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | HFC | 0.02 | 1137.81 | 2.21% | 0.02274 | 5.4% | KC level | KC trend | |
| 22 | 3 | 3. Solvent and Other Product Use | | | CO2 | 360.04 | 155.28 | 0.30% | 0.00389 | 0.9% | - | KC trend | |
| 23 | 4A | 4. Agric. | A. Enteric Fermentation | | CH4 | 2635.45 | 2496.98 | 4.85% | 0.00132 | 0.3% | KC level | - | |
| 24 | 4B | 4. Agric. | B. Manure Management | | CH4 | 671.61 | 646.11 | 1.26% | 0.00014 | 0.0% | KC level | - | |
| 25 | 4B | 4. Agric. | B. Manure Management | | N2O | 454.68 | 335.81 | 0.65% | 0.00213 | 0.5% | KC level | KC trend | |
| 26 | 4D1 | 4. Agric. | D. Agricultural Soils; Direct Soil Emissions | | N2O | 1351.48 | 1143.10 | 2.22% | 0.00342 | 0.8% | KC level | KC trend | |
| 27 | 4D2 | 4. Agric. | D. Agricultural Soils; Pasture, Range and Paddock Manure | | N2O | 128.10 | 220.79 | 0.43% | 0.00192 | 0.5% | KC level | KC trend | |
| 28 | 4D3 | 4. Agric. | D. Agricultural Soils; Indirect Emissions | | N2O | 822.48 | 674.93 | 1.31% | 0.00250 | 0.6% | KC level | KC trend | |
| 29 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO2 | -2416.89 | -2134.56 | 3.84% | 0.00409 | 1.0% | KC level | KC trend |
| 30 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO2 | -621.57 | -518.61 | 0.93% | 0.00161 | 0.4% | KC level | KC trend |
| 31 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO2 | 345.17 | 707.27 | 1.27% | 0.00683 | 1.7% | KC level | KC trend |
| 32 | 5C2 | 5. LULUCF | C. Grassland | 2. Land converted to Grassland | | CO2 | 59.85 | 169.07 | 0.30% | 0.00204 | 0.5% | - | KC trend |
| 33 | 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | | CO2 | 382.71 | 302.93 | 0.54% | 0.00129 | 0.3% | KC level | - |
| 34 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CH4 | 688.16 | 158.26 | 0.31% | 0.01021 | 2.4% | - | KC trend | |
| 35 | 6B | 6. Waste | B. Wastewater Handling | | N2O | 184.72 | 240.28 | 0.47% | 0.00121 | 0.3% | KC level | - | |
| 36 | 6D | 6. Waste | D. Other | | CH4 | 29.94 | 113.76 | 0.22% | 0.00169 | 0.4% | - | KC trend | |

Table 1-8 List of Switzerland's Tier 1 key categories for the base year 1990, combined KCA without and with LULUCF (*in italic*) categories, sorted by category code.

| Combined Tier 1 Key category analysis for the base year 1990 without and with LULUCF categories | | | | | | | | | |
|---|---|----------------------------------|--|--------------------------------------|---------------|------------------|--|-----------------------|----------------------------|
| No. | A IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | B Direct GHG | C Base Year 1990 Estimate [Gg CO ₂ eq] | E-L Level Assessm. | M Result level assessm. |
| 1 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CO ₂ | 289.73 | 0.55% | KC level |
| 2 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO ₂ | 693.69 | 1.31% | KC level |
| 3 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO ₂ | 1519.73 | 2.87% | KC level |
| 4 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and C | Gaseous Fuels | CO ₂ | 1074.09 | 2.03% | KC level |
| 5 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and C | Liquid Fuels | CO ₂ | 3692.22 | 6.98% | KC level |
| 6 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and C | Solid Fuels | CO ₂ | 1204.47 | 2.28% | KC level |
| 7 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CO ₂ | 252.55 | 0.48% | KC level |
| 8 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CO ₂ | 2587.68 | 4.89% | KC level |
| 9 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO ₂ | 11335.27 | 21.43% | KC level |
| 10 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Ins | Gaseous Fuels | CO ₂ | 987.24 | 1.87% | KC level |
| 11 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Ins | Liquid Fuels | CO ₂ | 4606.43 | 8.71% | KC level |
| 12 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO ₂ | 1424.38 | 2.69% | KC level |
| 13 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO ₂ | 10248.79 | 19.38% | KC level |
| 14 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Fore | Liquid Fuels | CO ₂ | 547.34 | 1.03% | KC level |
| 15 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO ₂ | 203.58 | 0.38% | KC level |
| 16 | 1B2 | 1. Energy | B. Fugitive Emissions | 2. Oil and Natural Gas | | CH ₄ | 263.72 | 0.50% | KC level |
| 17 | 2A1 | 2. Industrial | A. Mineral Products; Cement Production | CO ₂ | | CO ₂ | 2524.68 | 4.77% | KC level |
| 18 | 3 | 3. Solvent and Other Product Use | | | | CO ₂ | 360.04 | 0.68% | KC level |
| 19 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH ₄ | 2635.45 | 4.98% | KC level |
| 20 | 4B | 4. Agriculture | B. Manure Management | | | CH ₄ | 671.61 | 1.27% | KC level |
| 21 | 4B | 4. Agriculture | B. Manure Management | | | N ₂ O | 454.68 | 0.86% | KC level |
| 22 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N ₂ O | 1351.48 | 2.56% | KC level |
| 23 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N ₂ O | 822.48 | 1.56% | KC level |
| 24 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO ₂ | -2416.89 | 4.24% | KC level |
| 25 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO ₂ | -621.57 | 1.09% | KC level |
| 26 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO ₂ | 345.17 | 0.61% | KC level |
| 27 | 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | | CO ₂ | 382.71 | 0.67% | KC level |
| 28 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH ₄ | 688.16 | 1.30% | KC level |

Table 1-9 List of Switzerland's Tier 2 key categories, combined KCA without and with LULUCF (*in italic*) categories 2012, sorted by category code.

| Combined Tier 2 Key category analysis 2012 without and with LULUCF categories | | | | | | | | | | | | | |
|---|---|----------------------------------|---|--------------------------------------|---------------|-----------------|--|-------------------------------------|--|--|----------------------------|---------------------------|---------------------------|
| No. | A IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | B Direct GHG | C Base Year 1990 Estimate [Gg CO2 eq] | D Year 2012 Estimate [Gg CO2 eq] | E-L Level Assessm. with Uncertainty | E-T Trend Assessm. with Uncertainty | F-T % Contrib. in Trend | M Result level assessm | N Result trend assessm |
| 1 | 1A1 | 1. Energy | A. Fuel Comb. | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 1.67% | 0.00781 | 15.3% | KC level | KC trend |
| 2 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 0.20% | 0.00105 | 2.1% | KC level | KC trend |
| 3 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Other Fuels | CO2 | 134.15 | 288.60 | 0.18% | 0.00100 | 2.0% | KC level | KC trend |
| 4 | 1A2 | 1. Energy | A. Fuel Comb. | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.07% | 0.00111 | 2.2% | - | KC trend |
| 5 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Diesel | CO2 | 2587.68 | 6767.05 | 0.29% | 0.00190 | 3.7% | KC level | KC trend |
| 6 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | CH4 | 97.47 | 18.82 | 0.01% | 0.00056 | 1.1% | - | KC trend |
| 7 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | CO2 | 11335.27 | 9016.58 | 0.45% | 0.00103 | 2.0% | KC level | KC trend |
| 8 | 1A3b | 1. Energy | A. Fuel Comb. | 3. Transport; Road Transp. | Gasoline | N2O | 142.38 | 27.52 | 0.03% | 0.00111 | 2.2% | - | KC trend |
| 9 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Inst. | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 0.14% | 0.00052 | 1.0% | KC level | KC trend |
| 10 | 1A4a | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Com./Inst. | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 0.08% | 0.00040 | 0.8% | - | KC trend |
| 11 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 33.78 | 0.04% | 0.00078 | 1.5% | - | KC trend |
| 12 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 0.26% | 0.00127 | 2.5% | KC level | KC trend |
| 13 | 1A4b | 1. Energy | A. Fuel Comb. | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 0.20% | 0.00072 | 1.4% | KC level | KC trend |
| 14 | 1B2 | 1. Energy | B. Fugitive Emis | 2. Oil and Natural Gas | | CH4 | 263.72 | 169.45 | 0.10% | 0.00052 | 1.0% | KC level | KC trend |
| 15 | 2A1 | 2. Ind. Proc. | A. Mineral Products; Cement Production | CO2 | | CO2 | 2524.68 | 1787.11 | 0.10% | 0.00038 | 0.7% | KC level | - |
| 16 | 2A3 | 2. Ind. Proc. | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | | CO2 | 150.39 | 98.48 | 0.10% | 0.00049 | 1.0% | - | KC trend |
| 17 | 2F1 | 2. Ind. Proc. | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | | HFC | 0.02 | 1137.81 | 0.27% | 0.00273 | 5.3% | KC level | KC trend |
| 18 | 2F9 | 2. Ind. Proc. | F. Consumption of Halocarbons and SF6; Other | | | HFC | 0.00 | 76.14 | 0.12% | 0.00122 | 2.4% | KC level | KC trend |
| 19 | 2F9 | 2. Ind. Proc. | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 135.91 | 0.21% | 0.00094 | 1.8% | KC level | KC trend |
| 20 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 155.28 | 0.15% | 0.00195 | 3.8% | KC level | KC trend |
| 21 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 44.62 | 0.07% | 0.00100 | 2.0% | - | KC trend |
| 22 | 4A | 4. Agric. | A. Enteric Fermentation | | | CH4 | 2635.45 | 2496.98 | 0.89% | 0.00024 | 0.5% | KC level | - |
| 23 | 4B | 4. Agric. | B. Manure Management | | | CH4 | 671.61 | 646.11 | 0.68% | 0.00008 | 0.2% | KC level | - |
| 24 | 4B | 4. Agric. | B. Manure Management | | | N2O | 454.68 | 335.81 | 0.41% | 0.00135 | 2.6% | KC level | KC trend |
| 25 | 4D1 | 4. Agric. | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 1143.10 | 1.85% | 0.00285 | 5.6% | KC level | KC trend |
| 26 | 4D2 | 4. Agric. | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 220.79 | 0.36% | 0.00163 | 3.2% | KC level | KC trend |
| 27 | 4D3 | 4. Agric. | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 674.93 | 2.18% | 0.00415 | 8.1% | KC level | KC trend |
| 28 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO2 | -2416.89 | -2134.56 | 2.41% | 0.00257 | 3.5% | KC level | KC trend |
| 29 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO2 | -621.57 | -518.61 | 0.59% | 0.00101 | 1.4% | KC level | KC trend |
| 30 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO2 | 345.17 | 707.27 | 1.40% | 0.00751 | 10.2% | KC level | KC trend |
| 31 | 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | | CO2 | 43.33 | 22.26 | 0.06% | 0.00053 | 0.7% | - | KC trend |
| 32 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CO2 | 107.09 | 134.74 | 5.05% | 0.01165 | 15.8% | KC level | KC trend |
| 33 | 5C2 | 5. LULUCF | C. Grassland | 2. Land converted to Grassland | | CO2 | 59.85 | 169.07 | 0.21% | 0.00138 | 1.9% | KC level | KC trend |
| 34 | 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | | CO2 | 382.71 | 302.93 | 0.27% | 0.00065 | 0.9% | KC level | KC trend |
| 35 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 158.26 | 0.18% | 0.00596 | 11.7% | KC level | KC trend |
| 36 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 240.28 | 0.23% | 0.00061 | 1.2% | KC level | KC trend |
| 37 | 6D | 6. Waste | D. Other | | | CH4 | 29.94 | 113.76 | 0.22% | 0.00170 | 3.3% | KC level | KC trend |

Table 1-10 List of Switzerland's Tier 2 key categories for the base year 1990, combined KCA without and with LULUCF (in italic) categories, sorted by category code

| Combined Tier 2 Key category analysis for the base year 1990 without and with LULUCF categories | | | | | | | | | |
|---|---|----------------------------------|---|--------------------------------------|---------------|--|---------------------------------|-----------------------|----------|
| A | | | | | B | C | E-L | M | |
| No. | IPCC Source Categories and fuels if applicable (combined without LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Level Assessm. with Uncertainty | Result level assessm. | |
| 1 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 0.91% | KC level |
| 2 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manuf. Ind. and Constr. | Solid Fuels | CO2 | 1204.47 | 0.18% | KC level |
| 3 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO2 | 11335.27 | 0.55% | KC level |
| 4 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | N2O | 142.38 | 0.13% | KC level |
| 5 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Com./Instit. | Liquid Fuels | CO2 | 4606.43 | 0.12% | KC level |
| 6 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 0.12% | KC level |
| 7 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 0.14% | KC level |
| 8 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 0.27% | KC level |
| 9 | 1B2 | 1. Energy | B. Fugitive Emissions from F2. Oil and Natural Gas | | | CH4 | 263.72 | 0.15% | KC level |
| 10 | 2A1 | 2. Ind Proc. | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 0.14% | KC level |
| 11 | 2A3 | 2. Ind Proc. | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | | CO2 | 150.39 | 0.15% | KC level |
| 12 | 2F9 | 2. Ind Proc. | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 0.12% | KC level |
| 13 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 0.34% | KC level |
| 14 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 0.17% | KC level |
| 15 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH4 | 2635.45 | 0.91% | KC level |
| 16 | 4B | 4. Agriculture | B. Manure Management | | | CH4 | 671.61 | 0.69% | KC level |
| 17 | 4B | 4. Agriculture | B. Manure Management | | | N2O | 454.68 | 0.55% | KC level |
| 18 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 2.12% | KC level |
| 19 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 0.21% | KC level |
| 20 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 2.58% | KC level |
| 21 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO2 | -2416.89 | 2.66% | KC level |
| 22 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO2 | -621.57 | 0.68% | KC level |
| 23 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO2 | 345.17 | 0.66% | KC level |
| 24 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CO2 | 107.09 | 3.91% | KC level |
| 25 | 5E2 | 5. LULUCF | E. Settlements | | | CO2 | 382.71 | 0.34% | KC level |
| 26 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 0.76% | KC level |
| 27 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 0.17% | KC level |

Overview of KCA for Tier 1 and Tier 2

Table 1-11 presents an overview on key categories according to the combined KCA without and with LULUCF categories for both Tier 1 and Tier 2, and for both 1990 and 2012.

Table 1-11 Overview on key categories according to the combined KCA without and with LULUCF categories for both Tier 1 and Tier 2, and for both 2012 and 1990, sorted by category code.

| Overview on key categories according to the combined KCA without and with LULUCF categories for both Tier 1 and 2, and for both the submission and the base year | | | | | | | | | | | |
|--|----------------------------------|--|--|---------------|-----|------------|----------|----------|----------|----------|----------|
| IPCC Source Categories and fuels if applicable (with LULUCF categories) | | | | | | 2012 | | 1990 | 2012 | | 1990 |
| | | | | | | Direct GHG | Tier 1 | Tier 1 | Tier 1 | Tier 2 | Tier 2 |
| 1A1 | 1. Energy | A. Fuel Comb | 1. Energy Industries | Gaseous Fuels | CO2 | | KC level | KC trend | KC level | - | - |
| 1A1 | 1. Energy | A. Fuel Comb | 1. Energy Industries | Liquid Fuels | CO2 | | KC level | KC trend | KC level | - | - |
| 1A1 | 1. Energy | A. Fuel Comb | 1. Energy Industries | Other Fuels | CO2 | | KC level | KC trend | KC level | KC trend | KC level |
| 1A2 | 1. Energy | A. Fuel Comb | 2. Manuf Ind and Constr | Gaseous Fuels | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A2 | 1. Energy | A. Fuel Comb | 2. Manuf Ind and Constr | Liquid Fuels | CO2 | | KC level | KC trend | KC level | - | - |
| 1A2 | 1. Energy | A. Fuel Comb | 2. Manuf Ind and Constr | Other Fuels | CO2 | | KC level | KC trend | - | KC level | KC trend |
| 1A2 | 1. Energy | A. Fuel Comb | 2. Manuf Ind and Constr | Solid Fuels | CO2 | | KC level | KC trend | KC level | - | KC trend |
| 1A3a | 1. Energy | A. Fuel Comb | 3. Transport; Civil Aviation | | CO2 | | - | KC trend | KC level | - | - |
| 1A3b | 1. Energy | A. Fuel Comb | 3. Transport; Road Transp | Diesel | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A3b | 1. Energy | A. Fuel Comb | 3. Transport; Road Transp | Gasoline | CH4 | | - | - | - | - | KC trend |
| 1A3b | 1. Energy | A. Fuel Comb | 3. Transport; Road Transp | Gasoline | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A3b | 1. Energy | A. Fuel Comb | 3. Transport; Road Transp | Gasoline | N2O | | - | KC trend | - | - | KC trend |
| 1A3b | 1. Energy | A. Fuel Comb | 3. Transport; Road Transp | Natural Gas | CO2 | | - | KC trend | - | - | - |
| 1A4a | 1. Energy | A. Fuel Comb | 4. Other Sectors; Com/Instit | Gaseous Fuels | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A4a | 1. Energy | A. Fuel Comb | 4. Other Sectors; Com/Instit | Liquid Fuels | CO2 | | KC level | KC trend | KC level | - | KC trend |
| 1A4b | 1. Energy | A. Fuel Comb | 4. Other Sectors; Residential | Biomass | CH4 | | - | - | - | - | KC trend |
| 1A4b | 1. Energy | A. Fuel Comb | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A4b | 1. Energy | A. Fuel Comb | 4. Other Sectors; Residential | Liquid Fuels | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 1A4c | 1. Energy | A. Fuel Comb | 4. Other Sectors; Agric/Forestry | Liquid Fuels | CO2 | | KC level | - | KC level | - | - |
| 1A5 | 1. Energy | A. Fuel Comb | 5. Other | Liquid Fuels | CO2 | | - | KC trend | KC level | - | - |
| 1B2 | 1. Energy | B. Fugitive Emiss | 2. Oil and Natural Gas | | CH4 | | - | KC trend | KC level | KC level | KC trend |
| 2A1 | 2. Ind Proc. | A. Mineral Products | Cement Production-CO2 | | CO2 | | KC level | KC trend | KC level | KC level | - |
| 2A3 | 2. Ind Proc. | A. Mineral Products | Limestone and Dolomite Use, Emissions, CO2 | | CO2 | | - | - | - | - | KC trend |
| 2F1 | 2. Ind Proc. | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | HFC | | | KC level | KC trend | - | KC level | KC trend |
| 2F9 | 2. Ind Proc. | F. Consumption of Halocarbons and SF6; Other | | HFC | | | - | - | - | KC level | KC trend |
| 2F9 | 2. Ind Proc. | F. Consumption of Halocarbons and SF6; Other | | SF6 | | | - | - | - | KC level | KC trend |
| 3 | 3. Solvent and Other Product Use | | | | CO2 | | - | KC trend | KC level | KC level | KC trend |
| 3 | 3. Solvent and Other Product Use | | | | N2O | | - | - | - | - | KC trend |
| 4A | 4. Agric | A. Enteric Fermentation | | | CH4 | | KC level | - | KC level | KC level | - |
| 4B | 4. Agric | B. Manure Management | | | CH4 | | KC level | - | KC level | KC level | - |
| 4B | 4. Agric | B. Manure Management | | | N2O | | KC level | KC trend | KC level | KC level | KC trend |
| 4D1 | 4. Agric | D. Agricultural Soils; Direct Soil Emissions | | | N2O | | KC level | KC trend | KC level | KC level | KC trend |
| 4D2 | 4. Agric | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | | KC level | KC trend | - | KC level | KC trend |
| 4D3 | 4. Agric | D. Agricultural Soils; Indirect Emissions | | | N2O | | KC level | KC trend | KC level | KC level | KC trend |
| 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO2 | | KC level | KC trend | KC level | KC level | KC trend |
| 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | | CO2 | | - | - | - | - | KC trend |
| 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CO2 | | - | - | - | KC level | KC trend |
| 5C2 | 5. LULUCF | C. Grassland | 2. Land converted to Grassland | | CO2 | | - | KC trend | - | KC level | KC trend |
| 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | | CO2 | | KC level | - | KC level | KC level | KC trend |
| 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | | - | KC trend | KC level | KC level | KC trend |
| 6B | 6. Waste | B. Wastewater Handling | | | N2O | | KC level | - | - | KC level | KC trend |
| 6D | 6. Waste | D. Other | | | CH4 | | - | KC trend | - | KC level | KC trend |

1.5.2 KP-LULUCF Inventory

Switzerland identified three key categories for activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol (Forest Management, Afforestation and Reforestation, Deforestation). The approach relies on full inventory KCA (with LULUCF), KP - CRF association and qualitative assessment. A detailed description is presented in chapter 11.6.1 and in Table 11-3.

1.6 Quality Assurance and Quality Control (QA/QC)

1.6.1 QA / QC Procedures

In 2002, a total quality management system was introduced within the Federal Office for the Environment (FOEN), within which the GHG inventory was registered as a process. Subsequent to an audit in 2004, an inventory-specific quality management system (QMS) was developed. This QMS is designed to comply with the quality objectives of Good Practice

Guidance of IPCC (2000), to ensure and continuously improve transparency, consistency, comparability, completeness, accuracy, and confidence in national GHG emission and removal estimates. Furthermore, Switzerland adopted timeliness as a quality criterion. Switzerland's inventory system is designed to produce a high quality inventory that ensures full compliance with the reporting requirements of the UNFCCC and the Kyoto Protocol.

The quality management system is designed according to a plan-do-check-act cycle (PDCA cycle), which is a generally accepted model according to international standards. Key findings from QA/QC procedures are included in the inventory development plan (IDP), which represents the main instrument for continuous improvement in subsequent inventory cycles. This approach is in accordance with procedures described in decision 19/CMP.1 (UNFCCC 2006a) and in the IPCC Good Practice Guidance (IPCC 2000, chapter 8). The QMS complies with the ISO 9001:2008 standard and has been certified by the Swiss association for quality and management systems (SQS) in December 2007 (SQS, 2008), re-certified in 2010 (SQS, 2010) and in 2013 (Swiss-TS, 2013). Certification is upheld since through annual audits. Annual audits are part of the recertification procedure.

The main QMS elements are summarized below. The detailed state of its implementation is documented in the Description of the Quality Management System (FOEN 2014a), submitted alongside this report. All activities are embedded in an annual cycle of inventory planning, preparation, and management (see Table 1 in FOEN 2014a).

1.6.1.1 Responsibilities for QA/QC activities

The national inventory system has a dedicated QA/QC officer who is responsible for coordinating and ensuring compliance with procedures related to quality control and quality assurance. QA/QC activities are carried out by everyone involved in inventory preparation, and various cross-checks are set up to minimise inconsistencies and errors in the inventory. Individual responsibilities are described in detail in sections 2.1 and 5.1 of FOEN (2014a). Results from QA/QC activities are documented and reviewed by the QA/QC officer. Based on these feedbacks, suggestions for further improvements of the inventory are developed by the QA/QC officer, which are then discussed in the GHG inventory core group, added to the inventory development plan and assigned to the relevant expert.

1.6.1.2 QA/QC plan

The QA/QC plan is represented by a quality manual as required by the ISO 9001:2008 standard. This quality manual constitutes the core of the quality management system. It consists of a systematic compilation of all documents relevant to quality issues on the FOEN internal document management system. The quality manual contains information regarding requirements, core processes and results of the inventory process and the national registry, as well as QA/QC activities, management and supporting documents (Figure 1-3). The core processes are represented by detailed flowcharts that specify tasks and responsibilities, data sources and collection processes, reference material and guidelines, and archived documents.

The quality manual is reviewed annually by the QA/QC officer and modified after consultation with the project management if necessary. Since 2007, most contributors to the GHG inventory are authorised to access the FOEN-based inventory files by means of a SSL connection to a web platform, including the quality manual with the underlying documents.

Quality Manual – Swiss National System




Figure 1-3: Overview of the quality manual of the national inventory system

1.6.1.3 QC procedures

All contributors to the inventory complete checklists that have been designed according to table 8.1 of the Good Practice Guidance (IPCC 2000). During the period of data collection, the data suppliers fill in the checklists. Once completed, the checklists are returned to FOEN. Simultaneously to GHG inventory preparation, the suppliers of emission data, the national inventory compiler, the NIR lead authors, and the project management complete the respective checklists. The QA/QC officer reviews and archives the checklists and contacts the suppliers if concerns about data integrity and/or the performance of quality control procedures arise and arranges for necessary measures to be taken.

In addition to general QC, the inventory project management promotes specific Tier 2 QC procedures both by providing for a FOEN (co-)funding of selected research projects and by initiating internal studies, where appropriate (see FOEN 2014a, Annex D for a list of past and current activities). Significant outcomes are fed into the inventory development plan (IDP; FOEN 2014a, chapter 3, Annex E) in order to be considered in future inventory submissions.

1.6.1.4 QA review procedures

Apart from the **UNFCCC reviews** of the Swiss inventory, various other efforts are made to assure the high quality standards set out in the quality objectives:

Expert peer reviews are commissioned periodically to provide in-depth analysis of specific sectors. In 2006, energy and industrial processes have been scrutinized, as well as methane emissions from agriculture. In 2009, the waste sector was subject to a domestic expert review. At the end of 2010, a thorough review of the LULUCF sector has taken place. Furthermore the industrial processes sector has undergone a substantial external review in 2013 (CSD, 2013). Further reviews are planned after the implementation of the revised reporting guidelines.

Internal reviews of the NIR, GHG inventory CRF-tables, Kyoto Protocol LULUCF CRF-tables, and the QA/QC supplement are made prior to each submission. They are performed by members of the GHG inventory core group as well as by the staff of the consultancies involved in inventory compilation.

The outcomes of all those reviewing activities are evaluated by the project management and the QA/QC officer, resulting in suggestions for amendments and improvements. The core group decides which items are to be followed up and who will take on the responsibility for implementation of the changes in future submissions (see inventory development plan).

FOEN operates a homepage (www.climate reporting.ch) where the Swiss GHG inventories (NIR, CRF-tables, QA/QC supplement, UNFCCC review reports); the Swiss national communications and other reports submitted to the UNFCCC and the Kyoto Protocol may be downloaded. On this web site, most papers, internal reports, domestic reviews, Excel calculation sheets, and other difficult-to-access materials ('grey literature') quoted in the Swiss GHG inventory are provided online. The climate reporting homepage thus provides the option for public review.

1.6.1.5 Implementation of the recommendations of the ERT

All issues raised by the ERT are added to the inventory development plan (see also chapter 1.6.1.2 and FOEN2014a). The tables below give an overview over how and where recommendations (Table 1-12) and encouragements (Table 1-13) of the ERT have been implemented in the current submission. Outstanding issues, which could not be fully implemented in the current submission remain in the IDP and will be followed-up for future submissions (see Annex E in FOEN 2014a).

The first column in table (Table 1-12) refers to the relevant paragraph in the review report. The second column gives a brief description of the issue; the third column shows what has been improved and where the changes have been implemented. The fourth column refers to the corresponding entry in the inventory development plan (see Annex E in FOEN 2014a).

Table 1-12 Implementation of recommendations of the Expert Review Team.

| ARR (para.) | Recommendation | Improvement – NIR chapter | IDP |
|--------------------|---|---|--------|
| Energy | | | |
| 22 | Adhere to QA/QC procedures | In order to identify any mistakes or errors, QA/QC procedures were conscientiously implemented. However, even with rigorous QA/QC procedures, minor errors may pass unnoticed or may be identified too late to be corrected for the current submission. The QA/QC system in place strives to continuously improve the inventory and will ensure progress. | - |
| 23 | Provide a better description of the fuel allocation in the inventory in comparison to the national energy statistics | The entire chapter describing emissions in the energy sector has been overhauled. The allocation of energy use to the different categories is described in section 3.2.5 Country-specific Issues. | 10, 17 |
| 24 | Implementation of the results of the ongoing study on CO ₂ EFs and NCVs from liquid fuels in the 2015 submission | Study is on-going. Results are expected for mid-2014. Implementation is planned for the 2015 submission. | 19 |
| 28 | Disaggregation of feedstocks and non-energy use of fuels | Naphtha and LPG are reported separately (see chapter 3.2.3). A follow-up with the industry is planned in the coming year. Data for naphtha and LPG use are confidential and can be provided on demand to the reviewers. | 8 |
| 29 | Review of the CO ₂ EF of refinery gas | The emission factor has been reassessed. It resulted in a recalculation of the entire time series. See section 3.2.6.2 for a detailed description of the revised EF. | 16 |
| 30 | Provide documentation of CO ₂ EFs of solid fuels and use correct values | The CO ₂ EFs are documented in section 3.2.5.2. The error with regard to the CO ₂ EF of coal that occurred in the last submission has been corrected. | 9 |
| 37 | Communicate correct charcoal production data to FAO | The correct numbers were sent to FAO. | 18 |
| 40 | Reporting and documentation of emissions from oil pipelines | Emissions from oil pipelines are reported using the IPCC tier 1 default methodology and documented in section 3.3.2.2 | 12 |
| 42 | Estimate emissions from gas production from 1990-1994 | Emissions from gas production are reported using the IPCC tier 1 default methodology and documented in section 3.3.3.2 | 14 |
| 43 | Review of emissions from the gas industry and improve documentation of fugitive emissions | The emissions from the Swiss gas industry were reassessed and led to a recalculation of the entire time series. Documentation can be found in section 3.3.3.2 | 14 |
| 44 | Expand the description of the methodology used to estimate fugitive emissions from venting and flaring | The description of the entire source category 1B has been overhauled (see section 3.3.3.2) | 13 |
| IP | | | |
| 47 | Recalculation of emissions from blasting using the correct EF | The correct emission factor is used (see 4.2.2.1) | 6 |
| 50/51 | Provide improved estimate and documentation of the CO ₂ EF for brick and tile production | New data for brick and tile production could be obtained. The procedure to derive the CO ₂ EF for brick and tile production is described in section 4.2.2.3. | 8 |
| 52 | Include information regarding the justification of the N ₂ O EF for nitric acid production | Section 4.3.2.2 describes how the N ₂ O EF for nitric acid production was derived. | 9 |
| Agriculture | | | |
| 58 | Include information on conversion factors used for energy requirements of dairy cattle | All energy conversion factors are included in Table 6-3 (see section 6.2.2). | 10 |
| 60 | Include information and references regarding the choice of MCF of 10.0 per cent | Studies supporting the MCF of 10% for deep litter are included now in section 6.3.2. | 5 |

| | | | |
|------------------|---|--|---------|
| LULUCF | | | |
| 67 | Provide full coverage of the land use statistics AREA | Full coverage of the land use statistics AREA is provided (see Chapter 7.2.2.1) | 2 |
| 70 | Improve presentation of the methods and include criteria for the use of each method, the reasoning behind it and the relevant references | The methodology of calculating carbon stock changes was changed, simplified and harmonized with the reporting of carbon stock changes under UNFCCC and under the Kyoto Protocol. Applied methods and the reasoning behind it are described in Chapter 7.1.3.2, Chapter 11.3.1.1 and in Chapter 11.3.1.4. All information provided in these Chapters is referenced, including specifications for AD, factors and parameters used. | 3 |
| 71 | Provide references and improve documentation to justify not reporting certain carbon pools under afforestation | Requested references and supplementary information were inserted in Chapter 11.3.1.1 and in Chapter 11.3.1.2. | 13 |
| 72 | Report above-ground and below-ground carbon pools separately | Above- and belowground living biomass is reported separately under the Kyoto Protocol. The approach is described in Chapter 7.3.4.6. | 3 |
| 74 | Improve documentation to justify not reporting soil organic carbon changes in mineral soils for unproductive forests | The description of unproductive forests has been extended. The reasoning why the carbon pools of unproductive forests are not a net source of emissions was supported with references (see section 7.3.4.9). | 12 |
| Waste | | | |
| 77 | Improve documentation of the methodology used to estimate waste water emissions | See improved description in Chapter 8.3.2. | 9 |
| 78 | Further expand on the amounts of waste that are reported in the energy sector | See new Figure 8-4 and respective description. | 6 |
| 79 | Provide information regarding the composition of wastes in SWDS used to derive the degradable organic fraction | See new table 8-3. | 11 |
| 81 | Disaggregate the emissions of subcategory 6D. | Emissions from subcategory 6D are listed separately for composting, fermentation, shredder and other non-specified sources in CRF table 6. | 8 |
| KP-LULUCF | | | |
| 83 | Improve comparability of reporting of land converted to forest land under the convention and afforestation and reforestation under the Kyoto Protocol | An area budget for LULUCF and KP-LULUCF reporting was introduced in Chapter 11.2.3. Further, see comment on para 70. | 3, 10 |
| 86 | Use notation key "IE" for losses in living biomass in KP-LULUCF table 5 (KP-I)A.1.2 | The methodology was changed. For afforestations older than 20 years, emission factors of productive forests were applied (see Chapter 7.1.3.2 and Chapter 11.3.1.1). The losses in living biomass were noted as "R" (reported). | NA, (3) |
| 87 | Improve documentation regarding emissions from mineral soils in afforested units harvested and of dead wood and litter in units harvested | Requested references and supplementary information were inserted in Chapter 11.3.1.1 and in Chapter 11.3.1.2. | 13 |
| 88 | Improve documentation of the methodology used to calculate carbon stock changes under forest management | Supplementary documentation is provided by Didion (2014). A summary of this report was inserted in Chapter 11.3.2. | 14 |

Table 1-13 Implementation of encouragements of the Expert Review Team

| ARR (para.) | Encouragement | Improvement – NIR chapter | IDP |
|--------------------|--|---|-----|
| General | | | |
| Table 4 | Highlight planned improvements in key categories | The revised reporting guidelines will give rise to a considerable workload. Therefore, major improvements linked to the revised guidelines will be given priority in the next submission. This is stated accordingly in the relevant sections of the NIR ("Planned improvements"). | 5 |
| 14 | Explain changes in uncertainty estimates between submissions | Changes in uncertainty estimates is included in chapter 1.7.1.3 | |
| Energy | | | |
| 44 | Reassess emission factors used for venting and flaring | Emissions from venting and flaring are of minor importance in Switzerland, as there is no gas and oil production. Therefore, the reassessment of emission factors for venting and flaring is assigned a low priority. It will not be possible to implement the encouragement for the next submission. | - |
| Agriculture | | | |
| 56 | Improve the reporting of agricultural residues in biogas digesters and the split in the energy, agriculture and waste sector | The reporting of manure used for biogas production could not yet be improved and harmonized due to delays in the regulation for national compensation projects. The planned improvement has been postponed to the submission 2015 (see also section 6.3.6). | 3 |
| 57 | Implement a comparison between Swiss estimates and the IPCC tier 2 default to calculate the CH ₄ emissions for all animal categories | Country specific emission factor for enteric fermentation of cattle have been compared to IPCC Tier 2 emission factor. Results are presented in ART 2013a (see also section 6.2.4). | 9 |
| 59 | Split the sheep population to use the corresponding CH ₄ conversion rates for mature sheep and other categories provided by IPCC | Postponed to submission 2015. | 11 |
| 62 | Provide information regarding the causes for the trend of percentage of dairy cattle on pasture | Information provided in section 6.3.2. | 12 |
| 64 | Explain the differences in N ₂ O emissions from agricultural soils calculated using the country-specific method and the IPCC default method | Differences between the country specific and the IPCC default model are explained in section 6.5.2 and in ART 2013a. A more extensive comparison will not be conducted earliest with the implementation of the new 2006 IPCC Guidelines. | 7 |
| LULUCF | | | |
| 67 | Report CH ₄ and N ₂ O emissions from the non-mandatory categories that are currently not reported | The implementation of this encouragement was evaluated. The party decided not to report these emissions. | 4 |
| 74 | Use a coherent approach for LULUCF reporting under the Convention and under the Kyoto | See comments on para 70 and on para 83. | 3 |
| Waste | | | |
| 80 | Enhance investigations regarding N ₂ O emissions from waste water handling | Emissions from waste water are on the list of potential areas for improvements in future emissions. The encouragement will be taken into consideration when the sector waste water handling will be fully reassessed. | 12 |
| KP-LULUCF | | | |
| 85 | Clarify in the NIR that afforested and reforested areas are not reclassified to forest management | Supplementary text was inserted in Chapter 11.2.3, Chapter 11.3.1.1 and in the new Chapter 11.5.4. | NA |

1.6.1.6 Documentation and archiving procedures

Inventory data as well as background information on activity data and emission factors are archived by the national inventory compiler in the EMIS data base. EMIS allows to file background information (e.g. interim worksheets; references; rationale for choice of methods) for any subset of inventory-related data. Whenever such documents are used they are labeled as "EMIS 2014/(NFR-Code)" in this report.

Information on the QMS, all QA/QC activities performed, decisions reached by the experts (minutes), results of key category analyses and uncertainty analyses as well as inventory development (IDP) is documented and archived in the FOEN IDM system and accessible to authorised collaborators via the GHG inventory web platform. All inventory information, as far as needed to reconstruct and interpret inventory data and to describe the inventory system and its functions, is archived after each submission. It is accessible at a single location at the FOEN in Ittigen near Bern.

Data backup is managed by the Federal Office of Information Technology, Systems and Telecommunication (FOITT) using a Storage Area Network. FOITT runs backup facilities at two distinct locations on a daily as well as on a weekly basis.

1.6.2 Verification Activities

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

For each check, the CRF-table cells are marked in green if values are identical, in grey if they differ by no more than 20%, in orange if they differ by 20% to 50%, and in red if they differ by more than 50%. The findings are discussed among the core group members and the modelling specialists. All differences are investigated and the reasons for the differences sought. This procedure has already led to the identification of several mistakes, which were subsequently corrected before submission.

The current submission has been reviewed by personnel not directly involved in the preparation of a particular section of the inventory and revised accordingly.

The FOEN supports a monitoring campaign at the high altitude research station Jungfraujoch, where various greenhouse gases are measured continuously. The location of the research station normally provides for analysis of tropospheric background concentrations. However, under special meteorological conditions, an estimate of Swiss emissions can be derived from the measurements. For a couple of F-gases, a comparison of the inventory data with the inferred emissions is presented in Annex A6.1. Further research is needed to refine the approach and apply it to other greenhouse gases.

As an additional activity, the emission factor of all source categories used in the Swiss Inventory have been compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (INFRAS 2012). If respective Swiss values deviate more than $\pm 10\%$ from other countries' average or from the IPCC default value, explanations for the divergence are provided.

1.6.3 Treatment of Confidentiality Issues

Nearly all of the data necessary to compile the Swiss GHG inventory are publicly available. There are, however, a few exceptions:

- (i) Emission data that refers to a single enterprise is in general confidential.
- (ii) The reporting of disaggregated emissions from F-gases is confidential (not confidential as aggregated data).
- (iii) In the civil aviation sub-sector one data source (FOCA 1991) has been marked confidential by the Federal Office of Civil Aviation (FOCA).
- (iv) Unpublished AREA land use statistics raw data have been temporarily classified confidential by the Swiss Federal Statistical Office (SFSO).

The FOEN collects the data needed for calculating emissions of HFCs, PFCs and SF₆ from private companies or industry associations. In the National Inventory Report, the activity data underlying emission estimates of HFCs, PFCs and SF₆ are only partly presented at the most

disaggregated level for reasons of confidentiality. However, complete emissions are reported in aggregated tables.

Confidential data will be made available by the FOEN in line with the procedures agreed under the UNFCCC for the technical review of GHG inventories (UNFCCC 2003).

1.7 Uncertainty Evaluation

1.7.1 GHG Inventory

1.7.1.1 Tier 1 and Tier 2 analysis

This chapter presents the main results of the uncertainty evaluation Tier 1 and Tier 2 in accordance with the IPCC Good Practice Guidance:

- Tier 1 methodology (IPCC 2000: p. 6.13ff.)
- Tier 2 methodology, Monte Carlo simulation (IPCC 2000: p. 6.18ff.)

All uncertainties are given as half of the 95% confidence interval divided by the mean and expressed as a percentage (approximately two standard deviations) as suggested by the IPCC Guidelines (IPCC 1997a).

The uncertainty analysis Tier 1 is updated yearly and the uncertainty analysis Tier 2 (Monte Carlo simulation) is carried out every two years. Within this submission, the Tier 1 and Tier 2 analysis have been carried out.

The following chapters present the overall results of the uncertainty evaluation. Specific information about the uncertainty estimation for activity data, emission factors or emissions of each source category is included in the respective sectoral chapters (3–9) below.

1.7.1.2 Data Used

The evaluation includes uncertainties regarding activity data and emission factors. Uncertainties in the GWP values are not taken into account.

Uncertainty distributions are assumed to be symmetric for the Tier 1 method. For the Monte Carlo simulation, asymmetric distributions (triangle) were also adopted.

For source categories with quantitative uncertainty data available, the input information from studies or from the data suppliers is used for the uncertainty evaluation. This is mainly the case for key categories. For several key categories, no explicit information on uncertainties is available. For these cases, authors of the NIR chapters, FOEN experts involved and several data suppliers derived estimates of uncertainties based on the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC 2000) default values and on information concerning the process of data collection for activity data and emission factors (import or sales statistics, surveys or modelling). Several experts from data suppliers were contacted for further information on some of the uncertainties. Some industry associations/sources also provided published or unpublished uncertainty estimates for their data. The data sources can be found in the relevant sub-sections on “Uncertainties and Time-Series Consistency” in each of the sectoral chapters (3–9) below.

For categories with no quantitative uncertainty data available, the NIR provides qualitative estimates of uncertainties. The elaboration of a quantitative uncertainty assessment for these categories would present a large effort with only limited effect on the overall uncertainty and therefore it has been decided to realize a semi-quantitative assessment. This includes the definition of a list of the combined uncertainties for all gases and three uncertainty levels: low, medium and high (see Table 1-14). These values are motivated by the comparison of uncertainty analyses of several countries carried out by de Keizer et al. (2007), as presented at the 2nd Internat. Workshop on Uncertainty in Greenhouse Gas Inventories (Vienna 27-28

September 2007), and by Table A1-1 of IPCC Guidelines, Vol. 1, Annex 1, Managing uncertainties (IPCC 1997a).

Table 1-14 Semi-quantitative (combined) uncertainties (U) for the emission of categories with no quantitative uncertainty data available.

| Gas | Uncertainty Category | Combined Uncertainty |
|------------------|----------------------|----------------------|
| CO ₂ | low | 2% |
| | medium | 10% |
| | high | 40% |
| CH ₄ | low | 15% |
| | medium | 30% |
| | high | 60% |
| N ₂ O | low | 40% |
| | medium | 80% |
| | high | 150% |
| HFC | medium | 20% |
| PFC | medium | 20% |
| SF ₆ | medium | 20% |

Despite the investigation carried out for the current uncertainty analyses it will be necessary to further motivate institutions to supply not only average data but also estimates of associated uncertainties.

1.7.1.3 Results of Tier 1 Uncertainty Evaluation

With this submission, updated results of the uncertainty evaluation are presented. There is a calculation of the uncertainty excluding the LULUCF sector and an uncertainty evaluation including LULUCF. As described in IPCC (2000) and IPCC (2003), the uncertainty estimates of the LULUCF sector were combined with the uncertainty estimates of the non-LULUCF sector to obtain the total inventory uncertainty.

The resulting **Tier 1 level uncertainty in the national total annual CO₂ equivalent emissions without LULUCF is estimated to be 3.65%. Tier 1 trend uncertainty is 1.87%** meaning that the change of the base year (1990) to 2012, reported as -2.7%, lies with a probability of 95% between -4.6% and -0.8%.

Compared to the results of the previous inventory 2012 (level 3.55%, trend 1.89%; FOEN 2013), the level and the trend uncertainties for 2012 for the emissions without the LULUCF sector are slightly higher. This is mainly based on higher activity data and emission factor uncertainty in the sectors Energy - Fugitive CO₂-Emissions (1B2), N₂O and CH₄ emissions in Agriculture (4A, 4B and 4D), Industrial Processes (CO₂ emissions in 2A3, HFC emissions in 2F4 and PFC emissions in 2F9), and N₂O and CH₄ emissions in Others (7). This is partially compensated by the lower activity data uncertainty in the Energy sector and the correction of the emission factor uncertainty of the cement production (2A1).

The resulting **Tier 1 level uncertainty in the national total annual CO₂ equivalent emissions including LULUCF sector is estimated to be 7.43%. Tier 1 trend uncertainty is 2.46%.**

Compared to the results of the previous inventory 2012 (level 4.79%, trend 1.99%; FOEN 2013), the level and the trend uncertainties for 2012 for the emissions including LULUCF sector are higher. This is based on the significant higher emission factor uncertainty of CO₂-Emissions of 5C1 Grassland remaining Grassland (further information is included in chapter 7.5.5).

The results of the Tier 1 uncertainty analysis for GHG emissions 2012 are summarized in Table 1-16. Details of the uncertainty estimates for specific sources are provided in the sub-sections on “Uncertainties and Time-Series Consistency” in each of the chapters on source categories below.

It should be noted that the present results of the Tier 1 uncertainty analysis for GHG emissions do not, or not fully, take into account the following factors that may further increase uncertainties:

- correlations existing between source categories that have not been considered by the Tier 1 approach (e.g. production data used for industry emissions in both categories 1A2 Manufacturing Industries and 2 Industrial Processes, or cattle numbers used for emissions related to enteric fermentation and animal manure production);
- errors due to the assumption of constant parameters;
- errors due to non-normal, asymmetric distribution of the uncertainties;
- errors due to methodological shortcomings;
- errors due to sources not reported (these are assumed to be very small).

On the other hand, the Tier 2 uncertainty evaluation described below explicitly takes into account correlations between sources and asymmetric distributions.

Ranked by their contribution to uncertainty in the total national emissions level, the following categories are the top contributors to uncertainty with a contribution of over 97% to level and trend uncertainty (cf. Column H, Table 1-15):

- CO₂ from 5C1 Grassland remaining Grassland (combined uncertainty 5.6%)
- CO₂ from 5A1 Forest Land remaining Forest Land (2.7%)
- Indirect (4D3) and direct (4D1) emissions of N₂O from Agricultural Soils (2.3%)
- CO₂ from 1A1 Energy Industries (Other fuels) (1.9%)
- CO₂ from 5B1 Cropland remaining Cropland (1.7%)

These six source categories contribute with 7.2% to the level (root of sumsquare of the individual combined uncertainty) and with 2.1% to the trend uncertainty including LULUCF (root of sumsquare of the individual uncertainty introduced into the trend in total national emissions).

This allows for the identification of future areas of improvement in the context of the Inventory Development Plan (IDP).

Table 1-15 Ranked combined level uncertainties for sources in Switzerland.

Tier 1 Uncertainty Calculation and Reporting

| A | B | C | D | E | F | G | H |
|--|-----------------------|--------------------------|---------------------|---------------------------|-----------------------------|----------------------|--|
| IPCC Source category | Gas | Base year emissions 1990 | Year 2012 emissions | Activity data uncertainty | Emission factor uncertainty | Combined uncertainty | Combined uncertainty as % of total national emission in year t |
| | Input data | Input data | Input data | Input data | Calc/Input | | |
| | Gg CO ₂ eq | Gg CO ₂ eq | % | % | % | % | % |
| 5C15. LULUCFC. Grassland1. Grassland remaining Grassland | CO ₂ | 107.09 | 134.74 | 6.0 | 2084.30 | 2084.3 | 5.581 |
| 5A15. LULUCFA. Forest Land1. Forest Land remaining Forest Land | CO ₂ | -2'416.89 | -2'134.56 | 2.0 | 62.80 | 62.8 | 2.665 |
| 4D34. AgricultureD. Agricultural Soils; Indirect Emissions | N ₂ O | 822.48 | 674.93 | 31.8 | 163.00 | 166.1 | 2.227 |
| 4D14. AgricultureD. Agricultural Soils; Direct Soil Emissions | N ₂ O | 1'351.48 | 1'143.10 | 20.4 | 80.63 | 83.2 | 1.889 |
| 1A11. Energy A. Fuel Combustion 1. Energy IndustriesOther Fuels | CO ₂ | 1'519.73 | 2'714.50 | 10.0 | 30.00 | 31.6 | 1.706 |
| 5B15. LULUCFB. Cropland1. Cropland remaining Cropland | CO ₂ | 345.17 | 707.27 | 5.0 | 109.79 | 109.9 | 1.545 |
| 4A4. AgricultureA. Enteric Fermentation | CH ₄ | 2'635.45 | 2'496.98 | 6.4 | 17.17 | 18.3 | 0.910 |
| 4B4. AgricultureB. Manure Management | CH ₄ | 671.61 | 646.11 | 6.4 | 54.13 | 54.5 | 0.700 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationGasoline | CO ₂ | 11'335.27 | 9'016.58 | 2.2 | 1.36 | 2.6 | 0.462 |
| 4B4. AgricultureB. Manure Management | N ₂ O | 454.68 | 335.81 | 29.5 | 56.25 | 63.5 | 0.424 |
| 4D24. AgricultureD. Agricultural Soils; Pasture, Range and Paddock Manure | N ₂ O | 128.10 | 220.79 | 57.3 | 62.50 | 84.8 | 0.372 |
| 5E25. LULUCFE. Settlements2. Land converted to Settlements | CO ₂ | 382.71 | 302.93 | 5.0 | 50.00 | 50.2 | 0.303 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationDiesel | CO ₂ | 2'587.68 | 6'767.05 | 2.2 | 0.47 | 2.2 | 0.301 |
| 2F12. Industrial Proc.F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | HFC | 0.02 | 1'137.81 | 8.5 | 8.49 | 12.0 | 0.271 |
| 1A4b1. Energy A. Fuel Combustion 4. Other Sectors; ResidentialGaseous Fuels | CO ₂ | 1'424.38 | 2'649.60 | 2.0 | 4.60 | 5.0 | 0.264 |
| 6B6. Waste B. Wastewater Handling | N ₂ O | 184.72 | 240.28 | 1.3 | 50.00 | 50.0 | 0.239 |
| 6D6. Waste D. Other | CH ₄ | 29.94 | 113.76 | 10.0 | 100.00 | 100.5 | 0.227 |
| 5C25. LULUCFC. Grassland2. Land converted to Grassland | CO ₂ | 59.85 | 169.07 | 6.0 | 67.34 | 67.6 | 0.227 |
| 2F92. Industrial Proc.F. Consumption of Halocarbons and SF6; Other | SF ₆ | 79.58 | 135.91 | 56.6 | 56.57 | 80.0 | 0.216 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionGaseous Fuels | CO ₂ | 1'074.09 | 2'096.41 | 2.0 | 4.60 | 5.0 | 0.209 |
| 1A4b1. Energy A. Fuel Combustion 4. Other Sectors; ResidentialLiquid Fuels | CO ₂ | 10'248.79 | 7'374.50 | 1.3 | 0.53 | 1.4 | 0.204 |
| 6A6. Waste A. Solid Waste Disposal on Land | CH ₄ | 688.16 | 158.26 | 30.0 | 50.00 | 58.3 | 0.183 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionOther Fuels | CO ₂ | 134.15 | 288.60 | 10.0 | 30.00 | 31.6 | 0.181 |
| 3 3. Solvent and Other Product Use | CO ₂ | 360.04 | 155.28 | 35.4 | 35.36 | 50.0 | 0.154 |
| 1A4a1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutionalGaseous Fuels | CO ₂ | 987.24 | 1'482.76 | 2.0 | 4.60 | 5.0 | 0.148 |
| 2F92. Industrial Proc.F. Consumption of Halocarbons and SF6; Other | HFC | 0.00 | 76.14 | 56.6 | 56.57 | 80.0 | 0.121 |
| 5F25. LULUCFF. Other Land2. Land converted to Other Land | CO ₂ | 91.98 | 112.28 | 4.0 | 50.00 | 50.2 | 0.112 |
| 1B21. Energy B. Fugitive Emissions from Fuels2. Oil and Natural Gas | CH ₄ | 263.72 | 169.45 | 21.2 | 21.21 | 30.0 | 0.101 |
| 2A12. Industrial Proc.A. Mineral Products; Cement Production-CO ₂ | CO ₂ | 2'524.68 | 1'787.11 | 2.0 | 2.00 | 2.8 | 0.100 |
| 2A32. Industrial Proc.A. Mineral Products; Limestone and Dolomite Use, Emissions, CO ₂ | CO ₂ | 150.39 | 98.48 | 1.4 | 50.98 | 51.0 | 0.100 |
| 1A4a1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutionalLiquid Fuels | CO ₂ | 4'606.43 | 3'038.51 | 1.3 | 0.53 | 1.4 | 0.084 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionLiquid Fuels | CO ₂ | 3'692.22 | 2'640.18 | 1.3 | 0.53 | 1.4 | 0.073 |
| 3 3. Solvent and Other Product Use | N ₂ O | 110.14 | 44.62 | 35.4 | 71.76 | 80.0 | 0.071 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionSolid Fuels | CO ₂ | 1'204.47 | 454.87 | 5.9 | 5.00 | 7.7 | 0.070 |
| 5B25. LULUCFB. Cropland2. Land converted to Cropland | CO ₂ | 43.33 | 22.26 | 6.0 | 143.39 | 143.5 | 0.063 |
| 1A11. Energy A. Fuel Combustion 1. Energy IndustriesGaseous Fuels | CO ₂ | 289.73 | 498.74 | 2.0 | 4.60 | 5.0 | 0.050 |
| 2B2. Industrial Proc.B. Chemical Industry | N ₂ O | 68.13 | 53.57 | 7.1 | 40.39 | 41.0 | 0.044 |
| 1A4b1. Energy A. Fuel Combustion 4. Other Sectors; ResidentialBiomass | CH ₄ | 97.87 | 33.78 | 21.2 | 60.00 | 63.6 | 0.043 |
| 6D6. Waste D. Other | N ₂ O | 5.82 | 25.55 | 10.0 | 79.37 | 80.0 | 0.041 |
| 5E15. LULUCFE. Settlements1. Settlements remaining Settlements | CO ₂ | 3.60 | 33.69 | 5.0 | 50.00 | 50.2 | 0.034 |
| 1A11. Energy A. Fuel Combustion 1. Energy IndustriesOther Fuels | N ₂ O | 20.85 | 20.96 | 10.0 | 79.37 | 80.0 | 0.033 |
| 4D44. AgricultureD. Agricultural Soils; Use of sewage sludge as fertilizers | N ₂ O | 28.30 | 20.85 | 8.1 | 80.00 | 80.4 | 0.033 |
| 1A11. Energy A. Fuel Combustion 1. Energy IndustriesBiomass | N ₂ O | 27.72 | 19.75 | 21.2 | 77.14 | 80.0 | 0.031 |
| 1A4b1. Energy A. Fuel Combustion 4. Other Sectors; ResidentialLiquid Fuels | N ₂ O | 25.94 | 18.69 | 1.3 | 79.99 | 80.0 | 0.030 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationDiesel | N ₂ O | 5.74 | 66.06 | 2.2 | 22.00 | 22.1 | 0.029 |
| 5D25. LULUCFD. Wetlands2. Land converted to Wetlands | CO ₂ | 20.06 | 28.95 | 5.0 | 50.00 | 50.2 | 0.029 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationGasoline | N ₂ O | 142.38 | 27.52 | 2.2 | 50.00 | 50.0 | 0.027 |
| 1B21. Energy B. Fugitive Emissions from Fuels2. Oil and Natural Gas | CO ₂ | 84.62 | 39.29 | 21.2 | 21.21 | 30.0 | 0.023 |
| 1A11. Energy A. Fuel Combustion 1. Energy IndustriesLiquid Fuels | CO ₂ | 693.69 | 805.16 | 1.3 | 0.53 | 1.4 | 0.022 |
| 2B2. Industrial Proc.B. Chemical Industry | CO ₂ | 111.22 | 110.47 | 7.1 | 7.07 | 10.0 | 0.022 |
| 2F92. Industrial Proc.F. Consumption of Halocarbons and SF6; Other | PFC | 0.00 | 13.10 | 56.6 | 56.57 | 80.0 | 0.021 |
| 6C6. Waste C. Waste Incineration | N ₂ O | 19.06 | 25.86 | 28.3 | 28.28 | 40.0 | 0.021 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionLiquid Fuels | N ₂ O | 13.92 | 11.76 | 1.3 | 79.99 | 80.0 | 0.019 |
| 2F72. Industrial Proc.F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | SF ₆ | 0.00 | 19.55 | 28.3 | 28.28 | 40.0 | 0.016 |
| 1A4c1. Energy A. Fuel Combustion 4. Other Sectors; Agriculture/ForestryLiquid Fuels | CO ₂ | 547.34 | 540.01 | 1.3 | 0.53 | 1.4 | 0.015 |
| 1A4b1. Energy A. Fuel Combustion 4. Other Sectors; ResidentialBiomass | N ₂ O | 10.79 | 9.07 | 21.2 | 77.14 | 80.0 | 0.014 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationGasoline | CH ₄ | 97.47 | 18.82 | 2.2 | 37.00 | 37.1 | 0.014 |
| 2F22. Industrial Proc.F. Consumption of Halocarbons and SF6; Hard Foam | HFC | 0.00 | 13.27 | 34.6 | 34.65 | 49.0 | 0.013 |
| 5D25. LULUCFD. Land converted to Wetlands5(II) Non-CO ₂ emissions from drainage of soils at | CH ₄ | 9.03 | 9.03 | 10.0 | 70.00 | 70.7 | 0.013 |
| 1A4a1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutionalLiquid Fuels | N ₂ O | 11.70 | 7.80 | 1.3 | 79.99 | 80.0 | 0.012 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionBiomass | N ₂ O | 1.68 | 6.85 | 21.2 | 77.14 | 80.0 | 0.011 |
| 2F42. Industrial Proc.F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and Other | HFC | 0.00 | 13.22 | 28.3 | 28.28 | 40.0 | 0.011 |
| 77. Other | CO ₂ | 10.96 | 12.92 | 28.3 | 28.28 | 40.0 | 0.010 |
| 6C6. Waste C. Waste Incineration | CO ₂ | 54.10 | 12.34 | 28.3 | 28.28 | 40.0 | 0.010 |
| 2C2. Industrial Proc.C. Metal Production; Magnesium Foundries | SF ₆ | 0.00 | 31.72 | 10.6 | 10.61 | 15.0 | 0.009 |
| 1A4c1. Energy A. Fuel Combustion 4. Other Sectors; Agriculture/ForestryLiquid Fuels | N ₂ O | 4.96 | 5.54 | 1.3 | 79.99 | 80.0 | 0.009 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationNatural Gas | CO ₂ | 0.00 | 83.59 | 3.5 | 3.55 | 5.0 | 0.008 |
| 1A21. Energy A. Fuel Combustion 2. Manufacturing Industries and ConstructionOther Fuels | N ₂ O | 2.28 | 5.24 | 10.0 | 79.37 | 80.0 | 0.008 |
| 2C12. Industrial Proc.C. Metal Production; Steel Production | CO ₂ | 9.20 | 9.89 | 5.0 | 40.00 | 40.3 | 0.008 |
| 6C6. Waste C. Waste Incineration | CH ₄ | 11.58 | 6.14 | 28.3 | 52.92 | 60.0 | 0.007 |
| 2F82. Industrial Proc.F. Consumption of Halocarbons and SF6; Electrical Eq. | SF ₆ | 64.04 | 36.81 | 7.1 | 7.07 | 10.0 | 0.007 |
| 1A3a1. EnergyA. Fuel Combustion 3. Transport; Civil Aviation | CO ₂ | 252.55 | 136.65 | 2.2 | 1.16 | 2.5 | 0.007 |
| 5B25. LULUCFB. Cropland2. Land converted to Cropland | N ₂ O | 5.58 | 3.58 | 6.0 | 90.00 | 90.2 | 0.006 |
| 6B6. Waste B. Wastewater Handling | CH ₄ | 4.65 | 9.28 | 1.3 | 29.97 | 30.0 | 0.006 |
| 1A3b1. EnergyA. Fuel Combustion 3. Transport; Road TransportationNatural Gas | N ₂ O | 0.00 | 3.47 | 3.5 | 79.92 | 80.0 | 0.006 |

Table 1-16 Tier 1 uncertainty results for sources in Switzerland 2012 (IPCC 2000, Table 6.1, IPCC 2003).

| IPCC GPG Table 6.1 Tier 1 Uncertainty Calculation and Reporting | | | | | | | | | | | | | | |
|--|------------|--------------------------|---------------------|---------------------------|-----------------------------|----------------------|--|--------------------|--------------------|--|--|---|------|--|
| A | B | C | D | E | F | G | H | I | J | K | L | M | | |
| IPCC Source category | Gas | Base year emissions 1990 | Year 2012 emissions | Activity data uncertainty | Emission factor uncertainty | Combined uncertainty | Combined uncertainty as % of total national emission in year t | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions | | |
| | Input data | Input data | Input data | % | % | Calc/Input | % | % | % | % | % | % | | |
| | Gg CO2 eq | Gg CO2 eq | | | | | | | | | | | | |
| Total Uncertainty including LULUCF | | | | | | | | | | | | | | |
| | 50'968.63 | 50'320.09 | | | | | 7.43 | | | | | | 2.46 | |
| Emissions without LULUCF | | | | | | | | | | | | | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Gaseous Fuels | CH4 | 0.65 | 1.12 | 2.0 | 29.9 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Liquid Fuels | CH4 | 0.49 | 0.67 | 1.3 | 30.0 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Solid Fuels | CH4 | 0.10 | 0.00 | 5.9 | 29.4 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Biomass | CH4 | 0.33 | 0.43 | 21.2 | 21.2 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Gaseous Fuels | CO2 | 289.73 | 498.74 | 2.0 | 4.6 | 5.0 | 0.050 | 0.0042 | 0.0098 | 0.02 | 0.03 | 0.03 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Liquid Fuels | CO2 | 693.69 | 805.16 | 1.3 | 0.5 | 1.4 | 0.022 | 0.0024 | 0.0158 | 0.00 | 0.03 | 0.03 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Solid Fuels | CO2 | 44.84 | 0.00 | 5.9 | 5.0 | 7.7 | 0.000 | -0.0009 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Other Fuels | CO2 | 1519.73 | 2714.50 | 10.0 | 30.0 | 31.6 | 1.706 | 0.0238 | 0.0533 | 0.71 | 0.75 | 1.04 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Gaseous Fuels | N2O | 0.16 | 0.28 | 2.0 | 80.0 | 80.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Liquid Fuels | N2O | 2.15 | 2.86 | 1.3 | 80.0 | 80.0 | 0.005 | 0.0000 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Solid Fuels | N2O | 0.24 | 0.00 | 5.9 | 79.8 | 80.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Biomass | N2O | 27.72 | 19.75 | 21.2 | 77.1 | 80.0 | 0.031 | -0.0001 | 0.0004 | -0.01 | 0.01 | 0.02 | | |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Other Fuels | N2O | 20.85 | 20.96 | 10.0 | 79.4 | 80.0 | 0.033 | 0.0000 | 0.0004 | 0.00 | 0.01 | 0.01 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsGaseous Fuels | CH4 | 2.66 | 4.74 | 2.0 | 29.9 | 30.0 | 0.003 | 0.0000 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsLiquid Fuels | CH4 | 2.32 | 1.04 | 1.3 | 30.0 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsSolid Fuels | CH4 | 0.40 | 0.14 | 5.9 | 29.4 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsBiomass | CH4 | 2.46 | 1.56 | 21.2 | 21.2 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsOther Fuels | CH4 | 0.54 | 0.37 | 10.0 | 28.3 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsGaseous Fuels | CO2 | 1074.09 | 2096.41 | 2.0 | 4.6 | 5.0 | 0.209 | 0.0203 | 0.0411 | 0.09 | 0.12 | 0.15 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsLiquid Fuels | CO2 | 3692.22 | 2640.18 | 1.3 | 0.5 | 1.4 | 0.073 | -0.0197 | 0.0518 | -0.01 | 0.09 | 0.09 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsSolid Fuels | CO2 | 1204.47 | 454.87 | 5.9 | 5.0 | 7.7 | 0.070 | -0.0144 | 0.0089 | -0.07 | 0.07 | 0.10 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsOther Fuels | CO2 | 134.15 | 288.60 | 10.0 | 30.0 | 31.6 | 0.181 | 0.0031 | 0.0057 | 0.09 | 0.08 | 0.12 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsGaseous Fuels | N2O | 0.59 | 1.15 | 2.0 | 80.0 | 80.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsLiquid Fuels | N2O | 13.92 | 11.76 | 1.3 | 80.0 | 80.0 | 0.019 | 0.0000 | 0.0002 | 0.00 | 0.00 | 0.00 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsSolid Fuels | N2O | 6.44 | 2.41 | 5.9 | 79.8 | 80.0 | 0.004 | -0.0001 | 0.0000 | -0.01 | 0.00 | 0.01 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsBiomass | N2O | 1.68 | 6.85 | 21.2 | 77.1 | 80.0 | 0.011 | 0.0001 | 0.0001 | 0.01 | 0.00 | 0.01 | | |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and ConsOther Fuels | N2O | 2.28 | 5.24 | 10.0 | 79.4 | 80.0 | 0.008 | 0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A3a 1. Energy A. Fuel Combustion 3. Transport; Civil Aviation | CH4 | 0.24 | 0.25 | 2.2 | 60.0 | 60.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3a 1. Energy A. Fuel Combustion 3. Transport; Civil Aviation | CO2 | 252.55 | 136.65 | 2.2 | 1.2 | 2.5 | 0.007 | -0.0022 | 0.0027 | 0.00 | 0.01 | 0.01 | | |
| 1A3a 1. Energy A. Fuel Combustion 3. Transport; Civil Aviation | N2O | 2.49 | 1.35 | 2.2 | 150.0 | 150.0 | 0.004 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Natural Gas | CH4 | 0.00 | 0.10 | 3.5 | 29.8 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Diesel | CH4 | 1.38 | 0.55 | 2.2 | 20.0 | 20.1 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Gasoline | CH4 | 97.47 | 18.82 | 2.2 | 37.0 | 37.1 | 0.014 | -0.0015 | 0.0004 | -0.06 | 0.00 | 0.06 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Biomass | CH4 | 0.00 | -0.02 | 21.2 | 56.1 | 60.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Diesel | CO2 | 2587.68 | 6767.05 | 2.2 | 0.5 | 2.2 | 0.301 | 0.0826 | 0.1328 | 0.04 | 0.41 | 0.41 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Gasoline | CO2 | 11335.27 | 9018.58 | 2.2 | 1.4 | 2.6 | 0.462 | -0.0426 | 0.1769 | -0.06 | 0.55 | 0.55 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Natural Gas | CO2 | 0.00 | 83.59 | 3.5 | 3.5 | 5.0 | 0.008 | 0.0016 | 0.0016 | 0.01 | 0.01 | 0.01 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Diesel | N2O | 5.74 | 66.06 | 2.2 | 22.0 | 22.1 | 0.029 | 0.0012 | 0.0013 | 0.03 | 0.00 | 0.03 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Gasoline | N2O | 142.38 | 27.52 | 2.2 | 50.0 | 50.0 | 0.027 | -0.0022 | 0.0005 | -0.11 | 0.00 | 0.11 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Natural Gas | N2O | 0.00 | 3.47 | 3.5 | 79.9 | 80.0 | 0.006 | 0.0001 | 0.0001 | 0.01 | 0.00 | 0.01 | | |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Biomass | N2O | 0.00 | 0.70 | 21.2 | 148.5 | 150.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3c 1. Energy A. Fuel Combustion 3. Transport; Railways Liquid Fuels | CH4 | 0.01 | 0.01 | 1.3 | 30.0 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3c 1. Energy A. Fuel Combustion 3. Transport; Railways Liquid Fuels | CO2 | 28.69 | 39.69 | 1.3 | 0.5 | 1.4 | 0.001 | 0.0002 | 0.0008 | 0.00 | 0.00 | 0.00 | | |
| 1A3c 1. Energy A. Fuel Combustion 3. Transport; Railways Liquid Fuels | N2O | 0.38 | 0.52 | 1.3 | 150.0 | 150.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3d 1. Energy A. Fuel Combustion 3. Transport; Navigation Gas/Diesel Oil | CH4 | 0.01 | 0.02 | 2.2 | 29.9 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3d 1. Energy A. Fuel Combustion 3. Transport; Navigation Gasoline | CH4 | 0.58 | 0.54 | 2.2 | 29.9 | 30.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3d 1. Energy A. Fuel Combustion 3. Transport; Navigation Gas/Diesel Oil | CO2 | 111.93 | 121.14 | 2.2 | 0.5 | 2.3 | 0.005 | 0.0002 | 0.0024 | 0.00 | 0.01 | 0.01 | | |
| 1A3d 1. Energy A. Fuel Combustion 3. Transport; Navigation Gasoline | N2O | 0.66 | 0.82 | 2.2 | 150.0 | 150.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3e 1. Energy A. Fuel Combustion 3. Transport; Other non-specified | CH4 | 0.06 | 0.03 | 2.2 | 35.4 | 35.4 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A3e 1. Energy A. Fuel Combustion 3. Transport; Other non-specified | CO2 | 31.42 | 45.44 | 2.2 | 4.5 | 5.0 | 0.005 | 0.0003 | 0.0009 | 0.00 | 0.00 | 0.00 | | |
| 1A3e 1. Energy A. Fuel Combustion 3. Transport; Other non-specified | N2O | 0.02 | 0.03 | 2.2 | 80.0 | 80.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutGaseous Fuels | CH4 | 2.41 | 3.83 | 2.0 | 29.9 | 30.0 | 0.002 | 0.0000 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutLiquid Fuels | CH4 | 3.06 | 1.40 | 1.3 | 30.0 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutBiomass | CH4 | 9.74 | 4.13 | 21.2 | 21.2 | 30.0 | 0.002 | -0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutGaseous Fuels | CO2 | 987.24 | 1482.76 | 2.0 | 4.6 | 5.0 | 0.148 | 0.0100 | 0.0291 | 0.05 | 0.08 | 0.09 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutLiquid Fuels | CO2 | 4806.43 | 3038.51 | 1.3 | 0.5 | 1.4 | 0.084 | -0.0296 | 0.0596 | -0.02 | 0.11 | 0.11 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutGaseous Fuels | N2O | 0.55 | 0.82 | 2.0 | 80.0 | 80.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutLiquid Fuels | N2O | 11.70 | 7.80 | 1.3 | 80.0 | 80.0 | 0.000 | 0.0012 | 0.0001 | 0.0002 | -0.01 | 0.00 | 0.01 | |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/InstitutBiomass | N2O | 1.45 | 3.23 | 21.2 | 77.1 | 80.0 | 0.005 | 0.0000 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Gaseous Fuels | CH4 | 3.24 | 6.10 | 2.0 | 29.9 | 30.0 | 0.004 | 0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Liquid Fuels | CH4 | 6.00 | 2.26 | 1.3 | 30.0 | 30.0 | 0.001 | -0.0001 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Solid Fuels | CH4 | 3.71 | 2.28 | 5.9 | 29.4 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Biomass | CH4 | 97.87 | 33.78 | 21.2 | 60.0 | 63.6 | 0.043 | -0.0012 | 0.0007 | -0.07 | 0.02 | 0.08 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 2.0 | 4.6 | 5.0 | 0.264 | 0.0244 | 0.0424 | 0.520 | 0.11 | 0.15 | 0.18 | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Liquid Fuels | CO2 | 10248.79 | 7374.50 | 1.3 | 0.5 | 1.4 | 0.204 | -0.0537 | 0.1447 | -0.03 | 0.26 | 0.27 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Solid Fuels | CO2 | 54.59 | 33.60 | 5.9 | 5.0 | 7.7 | 0.005 | -0.0004 | 0.0007 | 0.00 | 0.01 | 0.01 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Gaseous Fuels | N2O | 0.79 | 1.46 | 2.0 | 80.0 | 80.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Liquid Fuels | N2O | 25.94 | 18.69 | 1.3 | 80.0 | 80.0 | 0.030 | -0.0001 | 0.0004 | -0.01 | 0.00 | 0.01 | | |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Solid Fuels | N2O | 0.29 | 0.18 | 5.9 | 79.8 | 80.0 | 0.000 | 0. | | | | | | |

Table 1-16 continued. Tier 1 uncertainty results for sources in Switzerland 2012 (IPCC 2000, Table 6.1, IPCC 2003).

IPCC GPG Table 6.1
Tier 1 Uncertainty Calculation and Reporting

| A | B | C | D | E | F | G | H | I | J | K | L | M |
|--|-----|--------------------------|-----------------------|---------------------------|-----------------------------|----------------------|--|--------------------|--------------------|--|--|---|
| IPCC Source category | Gas | Base year emissions 1990 | Year 2012 emissions | Activity data uncertainty | Emission factor uncertainty | Combined uncertainty | Combined uncertainty as % of total national emission in year t | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
| | | Input data | Input data | Input data | Input data | Calc/Input | % | % | % | % | % | % |
| | | Gg CO ₂ eq | Gg CO ₂ eq | % | % | % | % | % | % | % | % | % |
| Total Uncertainty including LULUCF | | 50'968.63 | 50'320.09 | | | | 7.43 | | | | | 2.46 |
| Emissions without LULUCF | | | | | | | | | | | | |
| 2A1 2. Industria A. Mineral Products; | CO2 | 2'524.68 | 1'787.11 | 2.0 | 2.0 | 2.8 | 0.100 | -0.0138 | 0.0351 | -0.03 | 0.10 | 0.10 |
| 2A2 2. Industria A. Mineral Products; | CO2 | 53.35 | 54.26 | 1.4 | 1.4 | 2.0 | 0.002 | 0.0000 | 0.0011 | 0.00 | 0.00 | 0.00 |
| 2A3 2. Industria A. Mineral Products; | CO2 | 150.39 | 98.48 | 1.4 | 51.0 | 51.0 | 0.100 | -0.0010 | 0.0019 | -0.05 | 0.00 | 0.05 |
| 2A7 2. Industria A. Mineral Products; | CO2 | 15.30 | 7.68 | 1.4 | 1.4 | 2.0 | 0.000 | -0.0001 | 0.0002 | 0.00 | 0.00 | 0.00 |
| 2B 1. Industria B. Chemical Industry | CH4 | 1.54 | 2.39 | 7.1 | 29.2 | 30.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 2B 2. Industria B. Chemical Industry | CO2 | 111.22 | 110.47 | 7.1 | 7.1 | 10.0 | 0.022 | 0.0000 | 0.0022 | 0.00 | 0.02 | 0.02 |
| 2B 2. Industria B. Chemical Industry | N2O | 68.13 | 53.57 | 7.1 | 40.4 | 41.0 | 0.044 | -0.0003 | 0.0011 | -0.01 | 0.01 | 0.02 |
| 2C 2. Industria C. Metal Production; | SF6 | 0.00 | 0.00 | 10.6 | 10.6 | 15.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 2C 2. Industria C. Metal Production; | SF6 | 0.00 | 31.72 | 10.6 | 10.6 | 15.0 | 0.009 | 0.0006 | 0.0006 | 0.01 | 0.01 | 0.01 |
| 2C1 2. Industria C. Metal Production; | CO2 | 9.20 | 9.89 | 5.0 | 40.0 | 40.3 | 0.008 | 0.0000 | 0.0002 | 0.00 | 0.00 | 0.00 |
| 2C3 2. Industria C. Metal Production; | CO2 | 139.26 | 0.00 | 7.1 | 7.1 | 10.0 | 0.000 | -0.0027 | 0.0000 | -0.02 | 0.00 | 0.02 |
| 2C3 2. Industria C. Metal Production; | PF6 | 100.17 | 0.00 | 13.0 | 13.0 | 18.4 | 0.000 | -0.0019 | 0.0000 | -0.03 | 0.00 | 0.03 |
| 2C5 2. Industria C. Metal Production; | CO2 | 1.65 | 1.36 | 7.1 | 7.1 | 10.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 2F1 2. Industria F. Consumption of H | HFC | 0.02 | 1'137.81 | 8.5 | 8.5 | 12.0 | 0.271 | 0.0223 | 0.0223 | 0.19 | 0.27 | 0.33 |
| 2F1 2. Industria F. Consumption of H | PF6 | 0.04 | 6.14 | 8.5 | 8.5 | 12.0 | 0.001 | 0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 |
| 2F2 2. Industria F. Consumption of H | HFC | 0.00 | 13.27 | 34.6 | 34.6 | 49.0 | 0.013 | 0.0003 | 0.0003 | 0.01 | 0.01 | 0.02 |
| 2F4 2. Industria F. Consumption of H | HFC | 0.00 | 13.22 | 28.3 | 28.3 | 40.0 | 0.011 | 0.0003 | 0.0003 | 0.01 | 0.01 | 0.01 |
| 2F5 2. Industria F. Consumption of H | HFC | 0.00 | 4.59 | 1.4 | 1.4 | 2.0 | 0.000 | 0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 |
| 2F5 2. Industria F. Consumption of H | PF6 | 0.00 | 7.29 | 1.4 | 1.4 | 2.0 | 0.000 | 0.0001 | 0.0001 | 0.00 | 0.00 | 0.00 |
| 2F7 2. Industria F. Consumption of H | PF6 | 0.00 | 6.54 | 28.3 | 28.3 | 40.0 | 0.005 | 0.0001 | 0.0001 | 0.00 | 0.01 | 0.01 |
| 2F7 2. Industria F. Consumption of H | SF6 | 0.00 | 19.55 | 28.3 | 28.3 | 40.0 | 0.016 | 0.0004 | 0.0004 | 0.01 | 0.02 | 0.02 |
| 2F8 2. Industria F. Consumption of H | SF6 | 64.04 | 36.81 | 7.1 | 7.1 | 10.0 | 0.007 | -0.0005 | 0.0007 | 0.00 | 0.01 | 0.01 |
| 2F9 2. Industria F. Consumption of H | HFC | 0.00 | 76.14 | 56.6 | 56.6 | 80.0 | 0.121 | 0.0015 | 0.0015 | 0.08 | 0.12 | 0.15 |
| 2F9 2. Industria F. Consumption of H | PF6 | 0.00 | 13.10 | 56.6 | 56.6 | 80.0 | 0.021 | 0.0003 | 0.0003 | 0.01 | 0.02 | 0.03 |
| 2F9 2. Industria F. Consumption of H | SF6 | 79.58 | 135.91 | 56.6 | 56.6 | 80.0 | 0.216 | 0.0011 | 0.0027 | 0.06 | 0.21 | 0.22 |
| 2G 2. Industria G. Other | CO2 | 1.04 | 0.91 | 7.1 | 7.1 | 10.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 3 3. Solvent ; | CO2 | 360.04 | 155.28 | 35.4 | 35.4 | 50.0 | 0.154 | -0.0039 | 0.0030 | -0.14 | 0.15 | 0.21 |
| 3 3. Solvent ; | N2O | 110.14 | 44.62 | 35.4 | 71.8 | 80.0 | 0.071 | -0.0013 | 0.0009 | -0.09 | 0.04 | 0.10 |
| 4A 4. Agricultura A. Enteric Fermentati | CH4 | 2'635.45 | 2'496.98 | 6.4 | 17.2 | 18.3 | 0.910 | -0.0021 | 0.0490 | -0.04 | 0.45 | 0.45 |
| 4B 4. Agricultura B. Manure Managem | CH4 | 671.61 | 646.11 | 6.4 | 54.1 | 54.5 | 0.700 | -0.0003 | 0.0127 | -0.02 | 0.12 | 0.12 |
| 4B 4. Agricultura B. Manure Managem | N2O | 454.68 | 335.81 | 29.5 | 56.3 | 63.5 | 0.424 | -0.0022 | 0.0066 | -0.12 | 0.28 | 0.30 |
| 4D1 4. Agricultura D. Agricultural Soils; | N2O | 1'351.48 | 1'143.10 | 20.4 | 80.6 | 83.2 | 1.889 | -0.0038 | 0.0224 | -0.30 | 0.65 | 0.71 |
| 4D2 4. Agricultura D. Agricultural Soils; | N2O | 128.10 | 220.79 | 57.3 | 62.5 | 84.8 | 0.372 | 0.0019 | 0.0043 | 0.12 | 0.35 | 0.37 |
| 4D3 4. Agricultura D. Agricultural Soils; | N2O | 822.48 | 674.93 | 31.8 | 163.0 | 166.1 | 2.227 | -0.0027 | 0.0132 | -0.44 | 0.59 | 0.74 |
| 4D4 4. Agricultura D. Agricultural Soils; | N2O | 28.30 | 20.85 | 8.1 | 80.0 | 80.4 | 0.033 | -0.0001 | 0.0004 | -0.01 | 0.00 | 0.01 |
| 6A 6. Waste A. Solid Waste Dispc | CH4 | 688.16 | 158.26 | 30.0 | 50.0 | 58.3 | 0.183 | -0.0102 | 0.0031 | -0.51 | 0.13 | 0.53 |
| 6A 6. Waste A. Solid Waste Dispc | CO2 | 9.24 | 0.00 | 30.0 | 26.5 | 40.0 | 0.000 | -0.0002 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 6B 6. Waste B. Wastewater Hand | CH4 | 4.65 | 9.28 | 1.3 | 30.0 | 30.0 | 0.006 | 0.0001 | 0.0002 | 0.00 | 0.00 | 0.00 |
| 6B 6. Waste B. Wastewater Hand | N2O | 184.72 | 240.28 | 1.3 | 50.0 | 50.0 | 0.239 | 0.0011 | 0.0047 | 0.06 | 0.01 | 0.06 |
| 6C 6. Waste C. Waste Incinerator | CH4 | 11.58 | 6.14 | 28.3 | 52.9 | 60.0 | 0.007 | -0.0001 | 0.0001 | -0.01 | 0.00 | 0.01 |
| 6C 6. Waste C. Waste Incinerator | CO2 | 54.10 | 12.34 | 28.3 | 28.3 | 40.0 | 0.010 | -0.0008 | 0.0002 | -0.02 | 0.01 | 0.02 |
| 6C 6. Waste C. Waste Incinerator | N2O | 19.06 | 25.86 | 28.3 | 28.3 | 40.0 | 0.021 | 0.0001 | 0.0005 | 0.00 | 0.02 | 0.02 |
| 6D 6. Waste D. Other | CH4 | 29.94 | 113.76 | 10.0 | 100.0 | 100.5 | 0.227 | 0.0017 | 0.0022 | 0.17 | 0.03 | 0.17 |
| 6D 6. Waste D. Other | CO2 | 0.00 | 0.00 | 10.0 | 0.0 | 10.0 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 6D 6. Waste D. Other | N2O | 5.82 | 25.55 | 10.0 | 79.4 | 80.0 | 0.041 | 0.0004 | 0.0005 | 0.03 | 0.01 | 0.03 |
| 7 7. Other | CH4 | 0.55 | 0.57 | 28.3 | 74.8 | 80.0 | 0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 7 7. Other | CO2 | 10.96 | 12.92 | 28.3 | 28.3 | 40.0 | 0.010 | 0.0000 | 0.0003 | 0.00 | 0.01 | 0.01 |
| 7 7. Other | N2O | 0.62 | 0.62 | 28.3 | 147.3 | 150.0 | 0.002 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| LULUCF | | | | | | | | | | | | |
| 5E1 5. LULUCF. Settlements | CO2 | 3.60 | 33.69 | 5.0 | 50.0 | 50.2 | 0.034 | 0.0006 | 0.0007 | 0.03 | 0.00 | 0.03 |
| 5E2 5. LULUCF. Settlements | CO2 | 382.71 | 302.93 | 5.0 | 50.0 | 50.2 | 0.303 | -0.0015 | 0.0059 | -0.07 | 0.04 | 0.08 |
| 5A1 5. LULUCF. Forest Land | CH4 | 20.90 | 0.42 | 10.0 | 70.0 | 70.7 | 0.001 | -0.0004 | 0.0000 | -0.03 | 0.00 | 0.03 |
| 5A1 5. LULUCF. Forest Land | CO2 | -2'416.89 | -2'134.56 | 2.0 | 62.8 | 62.8 | -2.665 | 0.0049 | -0.0419 | 0.31 | -0.12 | 0.33 |
| 5A1 5. LULUCF. Forest Land | CO2 | 25.36 | 0.51 | 10.0 | 70.0 | 70.7 | 0.001 | -0.0005 | 0.0000 | -0.03 | 0.00 | 0.03 |
| 5A1 5. LULUCF. Forest Land | N2O | 4.77 | 0.09 | 10.0 | 70.0 | 70.7 | 0.000 | -0.0001 | 0.0000 | -0.01 | 0.00 | 0.01 |
| 5A2 5. LULUCF. Forest Land | CO2 | -621.57 | -518.61 | 2.0 | 62.8 | 62.8 | -0.648 | 0.0019 | -0.0102 | 0.12 | -0.03 | 0.12 |
| 5B1 5. LULUCF. Cropland | CO2 | 345.17 | 707.27 | 5.0 | 109.8 | 109.9 | 1.545 | 0.0072 | 0.0139 | 0.79 | 0.10 | 0.80 |
| 5B2 5. LULUCF. Cropland | CO2 | 43.33 | 22.26 | 6.0 | 143.4 | 143.5 | 0.063 | -0.0004 | 0.0004 | -0.06 | 0.00 | 0.06 |
| 5B2 5. LULUCF. Cropland | N2O | 5.58 | 3.58 | 6.0 | 90.0 | 90.2 | 0.006 | 0.0000 | 0.0001 | 0.00 | 0.00 | 0.00 |
| 5C1 5. LULUCF. Grassland | CH4 | 0.41 | 0.00 | 6.0 | 70.0 | 70.3 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 5C1 5. LULUCF. Grassland | CO2 | 107.09 | 134.74 | 6.0 | 2084.3 | 2084.3 | 5.581 | 0.0006 | 0.0026 | 1.19 | 0.02 | 1.19 |
| 5C1 5. LULUCF. Grassland | N2O | 0.19 | 0.00 | 6.0 | 70.0 | 70.3 | 0.000 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 5C2 5. LULUCF. Grassland | CO2 | 59.85 | 169.07 | 6.0 | 67.3 | 67.6 | 0.227 | 0.0022 | 0.0033 | 0.15 | 0.03 | 0.15 |
| 5D1 5. LULUCF. Wetlands | CO2 | -2.87 | -0.56 | 30.0 | 100.0 | 104.4 | -0.001 | 0.0000 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 5D2 5. LULUCF. Land converted to 5(ii) Non-CO2 emissions from draina | CH4 | 9.03 | 9.03 | 10.0 | 70.0 | 70.7 | 0.013 | 0.0000 | 0.0002 | 0.00 | 0.00 | 0.00 |
| 5D2 5. LULUCF. Wetlands | CO2 | 20.06 | 28.95 | 5.0 | 50.0 | 50.2 | 0.029 | 0.0002 | 0.0006 | 0.01 | 0.00 | 0.01 |
| 5F2 5. LULUCF. Other Land | CO2 | 91.98 | 112.28 | 4.0 | 50.0 | 50.2 | 0.112 | 0.0004 | 0.0022 | 0.02 | 0.01 | 0.02 |

Table 1-16 continued. Tier 1 uncertainty results for sources in Switzerland 2012 (IPCC 2000, Table 6.1, IPCC 2003).

Table 6.1 (CONTINUED)
Tier 1 Uncertainty Calculation and Reporting

| A (continued) | | | B | N | O | P | Q |
|----------------------|---|---|-----|--|--|------------------------------------|-----------------------------|
| IPCC Source category | | | Gas | Emission factor quality indicator IPCC Default, Measurement based, national Referenced data | Activity data quality indicator IPCC Default, Measurement based, national Referenced data | Expert judgement reference numbers | Reference to section in NIR |
| 1A1 | 1. Energy | A. Fuel Comb 1. Energy Indt Gaseous Fuels | CO2 | M | D | | Section 3.2.6 |
| 1A1 | 1. Energy | A. Fuel Comb 1. Energy Indt Liquid Fuels | CO2 | M | R | | Section 3.2.6 |
| 1A1 | 1. Energy | A. Fuel Comb 1. Energy Indt Other Fuels | CO2 | R | R | | Section 3.2.6 |
| 1A2 | 1. Energy | A. Fuel Comb 2. Manufactur Gaseous Fuels | CO2 | M | D | | Section 3.2.7 |
| 1A2 | 1. Energy | A. Fuel Comb 2. Manufactur Liquid Fuels | CO2 | M | R | | Section 3.2.7 |
| 1A2 | 1. Energy | A. Fuel Comb 2. Manufactur Solid Fuels | CO2 | D | D, R | | Section 3.2.7 |
| 1A2 | 1. Energy | A. Fuel Comb 2. Manufactur Other Fuels | CO2 | R | R | | Section 3.2.7 |
| 1A3a | 1. Energy | A. Fuel Comb 3. Transport; Civil Aviation | CO2 | M | R | | Section 3.2.8 |
| 1A3b | 1. Energy | A. Fuel Comb 3. Transport; I Diesel | CO2 | M | R | | Section 3.2.8 |
| 1A3b | 1. Energy | A. Fuel Comb 3. Transport; I Gasoline | CO2 | M | R | | Section 3.2.8 |
| 1A4a | 1. Energy | A. Fuel Comb 4. Other Sectr Gaseous Fuels | CO2 | M | D | | Section 3.2.9 |
| 1A4a | 1. Energy | A. Fuel Comb 4. Other Sectr Liquid Fuels | CO2 | M | R | | Section 3.2.9 |
| 1A4b | 1. Energy | A. Fuel Comb 4. Other Sectr Gaseous Fuels | CO2 | M | D | | Section 3.2.9 |
| 1A4b | 1. Energy | A. Fuel Comb 4. Other Sectr Liquid Fuels | CO2 | M | R | | Section 3.2.9 |
| 1A4b | 1. Energy | A. Fuel Comb 4. Other Sectr Biomass | CO2 | M | R | | Section 3.2.9 |
| 1A4c | 1. Energy | A. Fuel Comb 4. Other Sectr Liquid Fuels | CO2 | M | R | | Section 3.2.9 |
| 1A5 | 1. Energy | A. Fuel Comb 5. Other Liquid and Gaseous | CO2 | M | R | | Section 3.3.10 |
| 1A1 | 1. Energy | A. Fuel Comb 1. Energy Ind. Other Fuels | N2O | R | R | | Section 3.2.6 |
| 1A3b | 1. Energy | A. Fuel Comb 3b. Road Trar Gasoline | N2O | R | R | | Section 3.2.8 |
| 1A3b | 1. Energy | A. Fuel Comb 3b. Road Trar Gasoline | CH4 | R | R | | Section 3.2.8 |
| 1B2 | 1. Energy | B. Fugitive En 2. Oil and Natural Gas | CH4 | D | D | | Section 3.3.2 |
| 2A1 | 2. Industrial P A. Mineral Products; Cement Production-CO2 | | CO2 | D | D | | Section 4.2.3 |
| 2C | 2. Industrial P C. Metal Production without Aluminium Product | | CO2 | R | R | | Section 4.4.3 |
| 2F1 | 2. Industrial P F. Consumption of Halocarbons and SF6 | | PFC | R | R | | Section 4.7.3 |
| 2F9 | 2. Industrial P F. Consumption of Halocarbons and SF6; Ref | | HFC | R | R | | Section 4.7.3 |
| 2F9 | 2. Industrial P F. Consumption of Halocarbons and SF6; Ref | | SF6 | R | R | | Section 4.7.3 |
| 3 | 3. Solvent and Other Product Use | | CO2 | R | R | | Section 5.2.3 |
| 3 | 3. Solvent and Other Product Use | | N2O | R | R | | Section 5.2.3 |
| 4A | 4. Agriculture A. Enteric Fermentation | | CH4 | R | R | | Section 6.2.3 |
| 4B | 4. Agriculture B. Manure Management | | CH4 | R | R | | Section 6.3.3 |
| 4B | 4. Agriculture B. Manure Management | | N2O | D | R | | Section 6.3.3 |
| 4D1 | 4. Agriculture D. Agricultural Soils; Direct Soil Emissions | | N2O | D | R | | Section 6.5.3 |
| 4D2 | 4. Agriculture D. Agricultural Soils; Pasture, Range and Pad | | N2O | D | R | | Section 6.5.3 |
| 4D3 | 4. Agriculture D. Agricultural Soils; Indirect Emissions | | N2O | D | R | | Section 6.5.3 |
| 5A1 | 5. LULUCF A. Forest Land 1. Forest Land remaining Fores | | CO2 | R | R | | Section 7.3.5 |
| 5A2 | 5. LULUCF A. Forest Land 2. Land converted to Forest Land | | CO2 | R | R | | Section 7.3.5 |
| 5B1 | 5. LULUCF B. Cropland 1. Cropland remaining Cropland | | CO2 | M | R | | Section 7.4.5 |
| 5C1 | 5. LULUCF B. Grassland 1. Grassland remaining Grassland | | CO2 | R | R | | Section 7.5.5 |
| 5C2 | 5. LULUCF B. Grassland 2. Land converted to Grassland | | CO2 | R | R | | Section 7.5.5 |
| 5E2 | 5. LULUCF E. Settlement: 2. Land converted to Settlement | | CO2 | R | R | | Section 7.7.5 |
| 6A | 6. Waste A. Solid Waste Disposal on Land | | CH4 | R | R | | Section 8.2.3 |
| 6B | 6. Waste B. Wastewater Handling | | N2O | R | R | | Section 8.3.3 |
| 6D | 6. Waste D. Other | | CH4 | R | R | | Section 8.5.3 |
| Rest of sources | | | CO2 | R | R | | Exp. est. |

1.7.1.4 Results of Tier 2 Uncertainty Evaluation (Monte Carlo)

A Tier 2 uncertainty analysis for Switzerland's GHG Inventory was carried out for the inventory submitted in 2012 (FOEN 2012) and contained a level uncertainty for 2010 and a trend uncertainty for the period 1990-2010. For the inventory year 2012 (i.e. the current submission) the Monte Carlo simulation has been updated.

The principle of Monte Carlo analysis is to select random values for emission factor and activity data from within their individual probability distributions, and to calculate the corresponding emission values. This procedure is repeated until an adequately stable result has been found. The results of all iterations yield the overall emission probability distribution.

In the present analysis, Monte Carlo simulations were performed to estimate uncertainties both in emissions 2012 and in emission trends 1990–2012, at the source category level as well as for the inventory as a whole (excluding and including LULUCF). The simulations were run with the commercial software package Crystal Ball (® Decisioneering). This tool generates random numbers within user defined probability ranges and probability distributions. As a result, selected statistics are produced for the forecast variables.

The main Monte Carlo results for level and trend analyses are:

Uncertainties without LULUCF

The total uncertainty level of Switzerland's 2012 emissions is **3.86%** of the total GHG emissions without LULUCF. The 95% confidence interval is slightly asymmetric and lies between **96.37% and 104.09% of the total GHG emissions without LULUCF**.

The change in total emissions between 1990 and 2012 is -2.72%. With a probability of 95%, the change lies within the range of **-6.11% to +0.12%**, corresponding to a trend uncertainty of **3.11%**.

Uncertainties with LULUCF

The total uncertainty level of the 2012 Swiss emissions is **7.51%** of the total GHG emissions with LULUCF. The 95% confidence interval is almost symmetric and lies between **92.57% and 107.59%**.

The change in total emissions between 1990 and 2012 is -1.27%. With a probability of 95%, the change lies within the range of **-10.23% to 7.41%**, corresponding to a trend uncertainty of **8.82%**.




Figure 1-4: Probability distributions of the simulated total emissions for the base year 1990 (in orange) and year t=2012 (in green). On the horizontal axis the total emissions (without LULUCF) are given in Gg CO₂ eq. The number of Monte Carlo runs is 118'900. The vertical lines show simulated mean and percentile values. Note that the simulated values slightly deviate from the reported inventory values Table A - 37.

The uncertainties are also evaluated by gas with the following results of the Monte Carlo simulation. CO₂ emissions have the highest precision or the lowest uncertainties (2.17%) among the Kyoto gases, as expected.

Table 1-17 Level uncertainties by gas 2012 for the total national emissions without LULUCF.

| Gas | Emission 2012 excl. LULUCF Gg CO ₂ eq | Lower bound 2.5 percentile Gg CO ₂ eq | Upper bound 97.5 percentile Gg CO ₂ eq | Mean absolute uncertainty Gg CO ₂ eq | Mean relative uncertainty % |
|------------------|--|--|---|---|-----------------------------------|
| CO ₂ | 43'251 | 42'322 | 44'200 | 939 | 2.17% |
| CH ₄ | 3'689 | 3'088 | 4'299 | 605 | 16.41% |
| N ₂ O | 3'007 | 1'712 | 4'621 | 1'455 | 48.38% |
| HFC | 1'245 | 1'095 | 1'394 | 150 | 12.03% |
| PFC | 33 | 22 | 44 | 11 | 32.68% |
| SF ₆ | 224 | 115 | 333 | 109 | 48.70% |
| Total | 51'449 | 49'582 | 53'552 | 1'985 | 3.86% |

Table 1-18 shows the Tier 2 uncertainty results in the structure of table 6.2 of IPCC Good Practice Guidance and Uncertainty Management (IPCC 2000).

The Tornado (Figure A-10 in Annex 7) shows that the very high uncertainty of the emissions of category 5C1 Grasland remaining Grasland (CO₂) cause the most important contribution to the total uncertainty. Further important contributions stem from 5A1 Forestland remaining Forestland (CO₂), 4D1 Direct Soil Emission, Synthetic Fertilizers (N₂O), 4D3 Indirect Emissions, Leaching and Runoff (N₂O).

Table 1-18 Level and trend Tier 2 uncertainties by gas 2012 for the total national emissions including LULUCF.

| IPPC Source Category | | | | Gas | Base year (1990) emissions | Year t (2012) emissions | Uncertainty in year t emissions as % of emissions in the category | | Unc. introduced on national total in year t | % change in emissions betw. year t and base year | Range of likely % change between year t and base year | | |
|----------------------|--------------------|---|-------|----------------|----------------------------------|-------------------------------|--|-----------------------|---|--|---|-----------------------|-------|
| | | | | | Gg CO2 eq | Gg CO2 eq | % below 2.5 perc. | % above 97.5 perc. | (%) | (%) | % below 2.5 perc. | % above 97.5 perc. | |
| 1. Energy | A. Fuel Combustion | 1. Energy Industries | 1A1 | Gaseous Fuels | CH4 | 0.65 | 1.12 | 70.0 | 129.9 | 0.00 | 41.9 | 12 | 132 |
| | | | 1A1 | Liquid Fuels | CH4 | 0.49 | 0.67 | 70.0 | 129.9 | 0.00 | 25.7 | -16 | 85 |
| | | | 1A1 | Solid Fuels | CH4 | 0.10 | - | NO | NO | - | NO | -130 | -70 |
| | | | 1A1 | Biomass | CH4 | 0.33 | 0.43 | 71.8 | 131.6 | 0.00 | 22.2 | -20 | 78 |
| | | | 1A1 | Gaseous Fuels | CO2 | 289.73 | 498.74 | 95.0 | 105.0 | 0.05 | 41.9 | 62 | 82 |
| | | | 1A1 | Liquid Fuels | CO2 | 693.69 | 805.16 | 98.6 | 101.4 | 0.02 | 13.8 | 14 | 18 |
| | | | 1A1 | Solid Fuels | CO2 | 44.84 | - | NO | NO | - | NO | -108 | -92 |
| | | | 1A1 | Other Fuels | CO2 | 1'519.73 | 2'714.50 | 68.8 | 132.6 | 1.71 | 44.0 | 29 | 130 |
| | | | 1A1 | Gaseous Fuels | N2O | 0.16 | 0.28 | 19.9 | 180.2 | 0.00 | 41.9 | -87 | 231 |
| | | | 1A1 | Liquid Fuels | N2O | 2.15 | 2.86 | 19.8 | 180.0 | 0.00 | 24.6 | -100 | 165 |
| | | | 1A1 | Solid Fuels | N2O | 0.24 | - | NO | NO | - | NO | -181 | -20 |
| | | | 1A1 | Biomass | N2O | 27.72 | 19.75 | 22.2 | 183.7 | 0.03 | -40.4 | -129 | 69 |
| | | | 1A1 | Other Fuels | N2O | 20.85 | 20.96 | 20.1 | 181.2 | 0.03 | 0.5 | -113 | 114 |
| | | 2. Manufacturing Industries and Construction | 1A2 | Gaseous Fuels | CH4 | 2.66 | 4.74 | 70.1 | 130.1 | 0.00 | 43.8 | 17 | 139 |
| | | | 1A2 | Liquid Fuels | CH4 | 2.32 | 1.04 | 69.7 | 130.2 | 0.00 | -122.5 | -88 | -22 |
| | | | 1A2 | Solid Fuels | CH4 | 0.40 | 0.14 | 70.2 | 130.4 | 0.00 | -174.8 | -96 | -32 |
| | | | 1A2 | Biomass | CH4 | 2.46 | 1.56 | 71.4 | 131.9 | 0.00 | -57.5 | -73 | -2 |
| | | | 1A2 | Other Fuels | CH4 | 0.54 | 0.37 | 70.7 | 130.7 | 0.00 | -46.8 | -69 | 4 |
| | | | 1A2 | Gaseous Fuels | CO2 | 1'074.09 | 2'096.41 | 95.0 | 105.1 | 0.21 | 48.8 | 85 | 106 |
| | | | 1A2 | Liquid Fuels | CO2 | 3'692.22 | 2'640.18 | 98.6 | 101.4 | 0.07 | -39.8 | -30 | -27 |
| | | | 1A2 | Solid Fuels | CO2 | 1'204.47 | 454.87 | 92.4 | 107.9 | 0.07 | -164.8 | -71 | -54 |
| | | | 1A2 | Other Fuels | CO2 | 134.15 | 288.60 | 68.9 | 132.3 | 0.18 | 53.5 | 41 | 191 |
| | | | 1A2 | Gaseous Fuels | N2O | 0.59 | 1.15 | 20.2 | 179.9 | 0.00 | 48.7 | -81 | 270 |
| | | | 1A2 | Liquid Fuels | N2O | 13.92 | 11.76 | 20.3 | 180.0 | 0.02 | -18.3 | -120 | 89 |
| | | | 1A2 | Solid Fuels | N2O | 6.44 | 2.41 | 19.9 | 180.3 | 0.00 | -166.9 | -148 | 23 |
| | | | 1A2 | Biomass | N2O | 1.68 | 6.85 | 22.0 | 183.2 | 0.01 | 75.5 | -20 | 657 |
| | | | 1A2 | Other Fuels | N2O | 2.28 | 5.24 | 20.5 | 180.7 | 0.01 | 56.4 | -69 | 332 |
| | | 3. Transport; Civil Aviation | 1A3a | | CH4 | 0.24 | 0.25 | 39.9 | 159.9 | 0.00 | 2.1 | -84 | 88 |
| | | | 1A3a | | CO2 | 252.55 | 136.65 | 97.5 | 102.5 | 0.01 | -84.8 | -49 | -43 |
| | | | 1A3a | | N2O | 2.49 | 1.35 | -50.2 | 251.5 | 0.00 | -84.8 | -217 | 125 |
| | | 3. Transport; Road Transportation | 1A3b | Natural Gas | CH4 | - | 0.10 | 69.9 | 130.0 | 0.00 | 100.0 | NO | NO |
| | | | 1A3b | Diesel | CH4 | 1.38 | 0.55 | 79.9 | 120.2 | 0.00 | -149.0 | -82 | -38 |
| | | | 1A3b | Gasoline | CH4 | 97.47 | 18.82 | 62.9 | 137.0 | 0.01 | -418.0 | -119 | -43 |
| | | | 1A3b | Biomass | CH4 | - | -0.02 | 162.9 | 42.4 | 0.00 | 100.0 | NO | NO |
| | | | 1A3b | Diesel | CO2 | 2'587.68 | 6'767.05 | 97.8 | 102.2 | 0.30 | 61.8 | 156 | 167 |
| | | | 1A3b | Gasoline | CO2 | 11'335.3 | 9'016.6 | 97.5 | 102.6 | 0.46 | -25.7 | -23 | -18 |
| | | | 1A3b | Natural Gas | CO2 | - | 83.59 | 95.1 | 105.0 | 0.01 | 100.0 | NO | NO |
| | | | 1A3b | Diesel | N2O | 5.74 | 66.06 | 77.9 | 122.2 | 0.03 | 91.3 | 795 | 1'307 |
| | | | 1A3b | Gasoline | N2O | 142.38 | 27.52 | 50.2 | 150.3 | 0.03 | -417.3 | -132 | -30 |
| | | | 1A3b | Natural Gas | N2O | - | 3.47 | 20.1 | 180.3 | 0.01 | 100.0 | NO | NO |
| | | | 1A3b | Biomass | N2O | - | 0.70 | -47.9 | 254.9 | 0.00 | 100.0 | NO | NO |
| | | 3. Transport; Railways | 1A3c | Liquid Fuels | CH4 | 0.01 | 0.01 | 70.0 | 130.0 | 0.00 | 26.6 | -14 | 87 |
| | | | 1A3c | Liquid Fuels | CO2 | 28.69 | 39.69 | 98.6 | 101.4 | 0.00 | 27.7 | 36 | 41 |
| | | | 1A3c | Liquid Fuels | N2O | 0.38 | 0.52 | -49.2 | 249.8 | 0.00 | 26.9 | -218 | 290 |
| | | 3. Transport; Navigation | 1A3d | Gas/Diesel Oil | CH4 | 0.01 | 0.02 | 69.9 | 129.8 | 0.00 | 6.8 | -37 | 51 |
| | | | 1A3d | Gasoline | CH4 | 0.58 | 0.54 | 70.0 | 130.0 | 0.00 | -8.6 | -49 | 33 |
| | | | 1A3d | | CO2 | 111.93 | 121.14 | 97.8 | 102.3 | 0.01 | 7.6 | 5 | 12 |
| | | | 1A3d | Gas/Diesel Oil | N2O | 0.66 | 0.82 | -49.0 | 250.0 | 0.00 | 19.6 | -215 | 265 |
| | | | 1A3d | Gasoline | N2O | 0.60 | 0.55 | -50.4 | 249.5 | 0.00 | -7.9 | -213 | 197 |
| | | 3. Transport; Other non- specified | 1A3ei | | CH4 | 0.06 | 0.03 | 64.7 | 135.4 | 0.00 | -72.8 | -83 | -1 |
| | | | 1A3ei | | CO2 | 31.42 | 45.44 | 95.0 | 105.0 | 0.00 | 30.9 | 36 | 53 |
| | | | 1A3ei | | N2O | 0.02 | 0.03 | 19.6 | 179.8 | 0.00 | 30.9 | -97 | 185 |
| | | 4. Other Sectors; Commercial/Insti- tutional | 1A4a | Gaseous Fuels | CH4 | 2.41 | 3.83 | 70.3 | 130.0 | 0.00 | 36.9 | 3 | 115 |
| | | | 1A4a | Liquid Fuels | CH4 | 3.06 | 1.40 | 70.0 | 129.8 | 0.00 | -119.1 | -87 | -22 |
| | | | 1A4a | Biomass | CH4 | 9.74 | 4.13 | 71.8 | 131.6 | 0.00 | -135.6 | -91 | -26 |
| | | | 1A4a | Gaseous Fuels | CO2 | 987.24 | 1'482.76 | 95.0 | 105.0 | 0.15 | 33.4 | 41 | 59 |
| | | | 1A4a | Liquid Fuels | CO2 | 4'606.43 | 3'038.51 | 98.6 | 101.4 | 0.08 | -51.6 | -35 | -33 |
| | | | 1A4a | Gaseous Fuels | N2O | 0.55 | 0.82 | 20.1 | 180.7 | 0.00 | 33.4 | -95 | 195 |
| | | | 1A4a | Liquid Fuels | N2O | 11.70 | 7.80 | 20.2 | 180.1 | 0.01 | -50.0 | -130 | 63 |
| | | | 1A4a | Biomass | N2O | 1.45 | 3.23 | 22.2 | 184.3 | 0.01 | 55.0 | -70 | 324 |

cont'd on next page

| IPPC Source Category | | | | Gas | Base year (1990) emissions | Year t (2012) emissions | Uncertainty in year t emissions as % of emissions in the category | | Unc. introduced on national total in year t | % change in emissions between year t and base year | Range of likely % change between year t and base year | | |
|----------------------|---|---|------|--------------|----------------------------------|-------------------------------|--|-----------------------|---|--|---|-----------------------|-----|
| | | | | | Gg CO2 eq | Gg CO2 eq | % below 2.5 perc. | % above 97.5 perc. | (%) | (%) | % below 2.5 perc. | % above 97.5 perc. | |
| 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | 1A4b | Gaseous Fuel | CH4 | 3.24 | 6.10 | 70.1 | 130.2 | 0.00 | 47.0 | 25 | 153 |
| | | | 1A4b | Liquid Fuels | CH4 | 6.00 | 2.26 | 70.2 | 130.1 | 0.00 | -165.3 | -94 | -30 |
| | | | 1A4b | Solid Fuels | CH4 | 3.71 | 2.28 | 70.2 | 130.3 | 0.00 | -62.5 | -74 | -3 |
| | | | 1A4b | Biomass | CH4 | 97.87 | 33.78 | 38.8 | 167.0 | 0.04 | -189.7 | -135 | 0 |
| | | | 1A4b | Gaseous Fuel | CO2 | 1'424.38 | 2'649.60 | 95.0 | 105.0 | 0.26 | 46.2 | 76 | 96 |
| | | | 1A4b | Liquid Fuels | CO2 | 10'248.8 | 7'374.50 | 98.6 | 101.4 | 0.20 | -39.0 | -29 | -27 |
| | | | 1A4b | Solid Fuels | CO2 | 54.59 | 33.60 | 92.4 | 107.9 | 0.01 | -62.5 | -48 | -29 |
| | | | 1A4b | Gaseous Fuel | N2O | 0.79 | 1.46 | 20.1 | 179.5 | 0.00 | 46.2 | -83 | 255 |
| | | | 1A4b | Liquid Fuels | N2O | 25.94 | 18.69 | 19.2 | 180.3 | 0.03 | -38.7 | -127 | 72 |
| | | | 1A4b | Solid Fuels | N2O | 0.29 | 0.18 | 20.4 | 180.5 | 0.00 | -62.5 | -133 | 55 |
| | | | 1A4b | Biomass | N2O | 10.79 | 9.07 | 22.7 | 183.6 | 0.01 | -19.0 | -123 | 88 |
| | | 4. Other Sectors; Agriculture / Forestry | 1A4c | Gaseous Fuel | CH4 | 0.09 | 0.04 | 70.0 | 130.2 | 0.00 | -124.3 | -88 | -23 |
| | | | 1A4c | Liquid Fuels | CH4 | 1.62 | 1.43 | 70.1 | 129.9 | 0.00 | -13.1 | -52 | 29 |
| | | | 1A4c | Biomass | CH4 | 0.80 | 0.15 | 71.6 | 131.6 | 0.00 | -450.2 | -114 | -53 |
| | | | 1A4c | Gaseous Fuel | CO2 | 41.45 | 18.48 | 95.0 | 105.1 | 0.00 | -124.3 | -61 | -50 |
| | | | 1A4c | Liquid Fuels | CO2 | 547.34 | 540.01 | 98.6 | 101.4 | 0.01 | -1.4 | -3 | 1 |
| | | | 1A4c | Gaseous Fuel | N2O | 0.02 | 0.01 | 19.6 | 179.8 | 0.00 | -124.3 | -143 | 32 |
| | | | 1A4c | Liquid Fuels | N2O | 4.96 | 5.54 | 19.5 | 180.6 | 0.01 | 10.5 | -108 | 131 |
| | | | 1A4c | Biomass | N2O | 0.21 | 0.35 | 22.1 | 183.3 | 0.00 | 40.1 | -87 | 228 |
| | | 5. Other | 1A5 | Liquid Fuels | CH4 | 0.16 | 0.12 | 69.9 | 129.9 | 0.00 | -38.0 | -65 | 9 |
| | | | 1A5 | Liquid Fuels | CO2 | 203.58 | 114.80 | 98.6 | 101.4 | 0.00 | -77.3 | -45 | -42 |
| | | | 1A5 | Liquid Fuels | N2O | 2.01 | 1.13 | -49.6 | 249.6 | 0.00 | -77.1 | -215 | 129 |
| | B. Fugitive Emissions from Fuels | 2. Oil and Natural Gas | 1B2 | | CH4 | 263.72 | 169.45 | 69.8 | 130.2 | 0.10 | -55.6 | -71 | -0 |
| | | | 1B2 | | CO2 | 84.62 | 39.29 | 70.1 | 130.0 | 0.02 | -115.4 | -87 | -21 |
| | | | 1B2 | | N2O | 0.62 | 0.68 | 71.5 | 131.6 | 0.00 | 8.3 | -36 | 54 |
| 2. Industrial Proc. | A. Mineral Products; Cement Production-CO2 | | 2A1 | | CO2 | 2'524.68 | 1'787.11 | 97.2 | 102.8 | 0.10 | -41.3 | -33 | -26 |
| | A. Mineral Products; Lime Production-CO2 | | 2A2 | | CO2 | 53.35 | 54.26 | 98.0 | 102.0 | 0.00 | 1.7 | -1 | 5 |
| | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | 2A3 | | CO2 | 150.39 | 98.48 | 49.1 | 150.8 | 0.10 | -52.7 | -95 | 26 |
| | A. Mineral Products; Other non-specified-CO2 | | 2A7 | | CO2 | 15.30 | 7.68 | 98.0 | 102.0 | 0.00 | -99.1 | -52 | -48 |
| | B. Chemical Industry | 2B | | CH4 | 1.54 | 2.39 | 70.0 | 129.9 | 0.00 | 35.8 | 0 | 111 | |
| | | 2B | | CO2 | 111.22 | 110.47 | 90.0 | 110.0 | 0.02 | -0.7 | -15 | 13 | |
| | | 2B | | N2O | 68.13 | 53.57 | 58.9 | 140.9 | 0.04 | -27.2 | -73 | 31 | |
| | C. Metal Production; Magnesium Foundries | | 2C | | SF6 | - | 31.72 | 85.0 | 114.9 | 0.01 | 100.0 | NO | NO |
| | C. Metal Production; Steel Production | | 2C1 | | CO2 | 9.20 | 9.89 | 59.8 | 140.3 | 0.01 | 7.0 | -52 | 66 |
| | C. Metal Production; Aluminium Production-CO2 | | 2C3 | | CO2 | 139.26 | - | NO | NO | - | NO | -110 | -90 |
| | C. Metal Production; Aluminium Production-PFC | | 2C3 | | PFC | 100.17 | - | NO | NO | - | NO | -119 | -82 |
| | C. Metal Production; Non- ferrous metals-CO2 | | 2C5 | | CO2 | 1.65 | 1.36 | 90.1 | 110.2 | 0.00 | -21.2 | -30 | -5 |

cont'd on next page

| IPPC Source Category | | | | Gas | Base year (1990) emissions | Year t (2012) emissions | Uncertainty in year t emissions as % of emissions in the category | | Uncertainty introduced on national total in year t | % change in emissions between year t and base year | Range of likely % change between year t and base year | |
|----------------------------------|---|-----|------------------------|--------|----------------------------------|-------------------------------|--|-----------------------|--|--|---|-----------------------|
| | | | | | Gg CO2 eq | Gg CO2 eq | % below 2.5 perc. | % above 97.5 perc. | (%) | (%) | % below 2.5 perc. | % above 97.5 perc. |
| 2. Industrial Proc. | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | 2F1 | | HFC | 0.02 | 1'137.81 | 88.0 | 112.0 | 0.27 | 100.0 | 4.4E+06 | 5.7E+06 |
| | F. Consumption of Halocarbons and SF6; Refrigeration | 2F1 | | PFC | 0.04 | 6.14 | 88.0 | 111.9 | 0.00 | 99.3 | 1.3E+04 | 1.6E+04 |
| | F. Consumption of Halocarbons and SF6; Hard Foam | 2F2 | | HFC | - | 13.27 | 50.7 | 149.1 | 0.01 | 100.0 | NO | NO |
| | F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and Other | 2F4 | | HFC | - | 13.22 | 60.1 | 140.1 | 0.01 | 100.0 | NO | NO |
| | F. Consumption of Halocarbons and SF6; Solvents | 2F5 | | HFC | - | 4.59 | 98.0 | 102.0 | 0.00 | 100.0 | NO | NO |
| | | 2F5 | | PFC | - | 7.29 | 98.0 | 102.0 | 0.00 | 100.0 | NO | NO |
| | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | 2F7 | | PFC | - | 6.54 | 59.9 | 140.2 | 0.01 | 100.0 | NO | NO |
| | | 2F7 | | SF6 | - | 19.55 | 59.9 | 140.0 | 0.02 | 100.0 | NO | NO |
| | F. Consumption of Halocarbons and SF6; Electrical Eq. | 2F8 | | SF6 | 64.04 | 36.81 | 90.0 | 110.0 | 0.01 | -74.0 | -54 | -31 |
| | F. Consumption of Halocarbons and SF6; Other | 2F9 | | HFC | - | 76.14 | 19.7 | 180.1 | 0.12 | 100.0 | NO | NO |
| 2F9 | | | PFC | - | 13.10 | 20.0 | 179.8 | 0.02 | 100.0 | NO | NO | |
| 2F9 | | | SF6 | 79.58 | 135.91 | 20.1 | 179.9 | 0.21 | 41.4 | -88 | 229 | |
| 3. Solvent and Other Product Use | G. Other | 2G | | CO2 | 1.04 | 0.91 | 90.0 | 110.0 | 0.00 | -14.6 | -26 | 1 |
| | 3 | | CO2 | 360.04 | 155.28 | 54.2 | 155.0 | 0.15 | -131.9 | -115 | -6 | |
| | 3 | | N2O | 110.14 | 44.62 | 20.0 | 180.2 | 0.07 | -146.8 | -146 | 27 | |
| 4. Agriculture | A. Enteric Fermentation | 4A | | CH4 | 2'635.45 | 2'496.98 | 81.9 | 118.5 | 0.90 | -5.5 | -12 | 1 |
| | | 4B | | CH4 | 671.61 | 646.11 | 45.8 | 155.1 | 0.70 | -3.9 | -11 | 2 |
| | B. Manure Management | 4B | liquid | N2O | 43.47 | 36.79 | 15.4 | 169.3 | 0.04 | -18.2 | -42 | 16 |
| | | 4B | solid | N2O | 411.21 | 299.02 | 39.3 | 158.0 | 0.31 | -37.5 | -49 | -7 |
| | D. Agricultural Soils; Direct Soil Emissions | 4D1 | fertilizer | N2O | 1'279.41 | 1'073.89 | -5.6 | 242.0 | 2.62 | -19.1 | -118 | 69 |
| | | 4D1 | organic soils | N2O | 72.07 | 69.21 | 40.8 | 170.6 | 0.09 | -4.1 | -52 | 43 |
| | D. Agricultural Soils; Pasture, Range and Paddock Manure | 4D2 | - | N2O | 128.10 | 220.79 | 40.2 | 151.2 | 0.21 | 42.0 | 28 | 105 |
| | D. Agricultural Soils; Indirect Emissions | 4D3 | deposition | N2O | 286.78 | 236.85 | 37.1 | 181.5 | 0.36 | -21.1 | -70 | 23 |
| | | 4D3 | leaching and runoff | N2O | 535.69 | 438.08 | 39.1 | 223.6 | 1.72 | -22.3 | -153 | 42 |
| | D. Agricultural Soils; Sewage sludge and compost | 4D4 | - | N2O | 28.30 | 20.85 | 19.9 | 180.0 | 0.03 | -35.7 | -48 | -5 |
| 6. Waste | A. Solid Waste Disposal on Land | 6A | | CH4 | 688.16 | 158.26 | 41.5 | 158.1 | 0.18 | -334.8 | -142 | -21 |
| | | 6A | | CO2 | 9.24 | - | NO | NO | - | NO | -140 | -60 |
| | B. Wastewater Handling | 6B | | CH4 | 4.65 | 9.28 | 70.1 | 130.1 | 0.01 | 49.9 | 33 | 167 |
| | | 6B | | N2O | 184.72 | 240.28 | 50.0 | 149.8 | 0.24 | 23.1 | -52 | 112 |
| | C. Waste Incineration | 6C | | CH4 | 11.58 | 6.14 | 43.8 | 164.8 | 0.01 | -88.5 | -119 | 19 |
| | | 6C | | CO2 | 54.10 | 12.34 | 62.7 | 143.2 | 0.01 | -338.3 | -121 | -39 |
| | | 6C | | N2O | 19.06 | 25.86 | 62.7 | 143.2 | 0.02 | 26.3 | -31 | 105 |
| | D. Other | 6D | | CH4 | 29.94 | 113.76 | -0.6 | 200.6 | 0.23 | 73.7 | -114 | 673 |
| | | 6D | | N2O | 5.82 | 25.55 | 19.4 | 180.6 | 0.04 | 77.2 | -23 | 698 |
| 7. Other | 7 | | CH4 | 0.55 | 0.57 | 20.3 | 179.4 | 0.00 | 2.4 | -113 | 117 | |
| | 7 | | CO2 | 10.96 | 12.92 | 60.1 | 140.2 | 0.01 | 15.1 | -44 | 80 | |
| | 7 | | N2O | 0.62 | 0.62 | -50.8 | 250.9 | 0.00 | 0.0 | -213 | 212 | |

cont'd on next page

| IPPC Source Category | | | | | Gas | Base year (1990) emissions | Year t (2012) emissions | Uncertainty in year t emissions as % of emissions in the category | | Unc. introduced on national total in year t | % change in emissions between year t and base year | Range of likely % change between year t and base year | |
|----------------------|--------------------------------------|--|-----|----------------------------------|-------|----------------------------------|-------------------------------|--|-----------------------|---|--|---|-----------------------|
| | | | | | | Gg CO2 eq | Gg CO2 eq | % below 2.5 perc. | % above 97.5 perc. | (%) | (%) | % below 2.5 perc. | % above 97.5 perc. |
| LULUCF | | | | | | | | | | | | | |
| 5. LULUCF | E. Settlement s | 1. Settlements remaining Settlements | 5E1 | | CO2 | 3.60 | 33.69 | 49.6 | 150.1 | 0.03 | 89.3 | 361 | 1'310 |
| | | 2. Land converted to Settlements | 5E2 | | CO2 | 382.71 | 302.93 | 50.3 | 150.2 | 0.30 | -26.3 | -85 | 43 |
| | A. Forest Land | 1. Forest Land remaining Forest Land | 5A1 | | CH4 | 20.90 | 0.42 | 28.9 | 170.9 | 0.00 | -4901.0 | -169 | -28 |
| | | | 5A1 | | CO2 | -2'416.89 | -2'134.56 | 162.7 | 37.2 | 2.64 | -13.2 | 71 | -96 |
| | | | 5A1 | Biomass Burning, Wildfires | CO2 | 25.36 | 0.51 | 29.7 | 171.5 | 0.00 | -4911.5 | -169 | -28 |
| | | | 5A1 | | N2O | 4.77 | 0.09 | 29.2 | 170.8 | 0.00 | -5033.3 | -169 | -27 |
| | | 2. Land converted to Forest Land | 5A2 | | CO2 | -621.57 | -518.61 | 163.2 | 37.2 | 0.64 | -19.9 | 66 | -98 |
| | B. Cropland | 1. Cropland remaining Cropland | 5B1 | | CO2 | 345.17 | 707.27 | -9.7 | 209.8 | 1.53 | 51.2 | -145 | 357 |
| | | 2. Land converted to Cropland | 5B2 | | CO2 | 43.33 | 22.26 | -44.7 | 243.5 | 0.06 | -94.7 | -210 | 113 |
| | | | 5B2 | | N2O | 5.58 | 3.58 | 9.8 | 190.3 | 0.01 | -56.1 | -143 | 70 |
| | C. Grassland | 1. Grassland remaining Grassland | 5C1 | | CH4 | 0.41 | 0.00 | 30.2 | 170.8 | 0.00 | -14445.2 | -170 | -30 |
| | | | 5C1 | | CO2 | 107.09 | 134.74 | -1962.8 | 2064.7 | 5.52 | 20.5 | -3'300 | 3'380 |
| | | | 5C1 | | N2O | 0.19 | 0.00 | 30.0 | 170.2 | 0.00 | -14439.4 | -170 | -29 |
| | 2. Land converted to Grassland | 5C2 | | CO2 | 59.85 | 169.07 | 32.4 | 168.0 | 0.23 | 64.6 | -20 | 386 | |
| | D. Wetlands | 1. Wetlands remaining Wetlands | 5D1 | | CO2 | -2.87 | -0.56 | 204.5 | -4.5 | 0.00 | -408.8 | 27 | -187 |
| | D. Land converted to Wetlands | 5(II) Non-CO2 emissions from drainage of soils and wetlands, Flooded Lands | 5D2 | | CH4 | 9.03 | 9.03 | 29.9 | 170.9 | 0.01 | 0.0 | -100 | 100 |
| | D. Wetlands | 2. Land converted to Wetlands | 5D2 | | CO2 | 20.06 | 28.95 | 49.3 | 150.7 | 0.03 | 30.7 | -44 | 133 |
| | F. Other Land | 2. Land converted to Other Land | 5F2 | | CO2 | 91.98 | 112.28 | 49.9 | 150.2 | 0.11 | 18.1 | -58 | 101 |
| Total | without LULUCF | | | | - | 52'890 | 51'449 | 96.37 | 104.09 | 3.86 | -2.87 | -6.11 | 0.12 |
| | with LULUCF | | | | - | 50'969 | 50'320 | 92.57 | 107.59 | 7.51 | -1.40 | -10.23 | 7.41 |

Assumptions and further results to the Monte Carlo simulation are shown in Annex 7.

1.7.1.5 Comparison of Tier 1 and Tier 2 Results

In the GHG inventory, some of the uncertainties may become large, their statistical distribution may clearly deviate from normal distributions, and they can be correlated. Tier 1 uncertainty analysis is based on simple error propagation, which assumes only small, normally distributed and uncorrelated uncertainties. The application of the Tier 1 method is therefore not the optimal instrument for determining the uncertainties of a GHG inventory. The more appropriate choice is the Monte Carlo simulation, which is designed for uncertainties of any shape, for any size of uncertainties, any correlated figures and which is recommended by the IPCC Good Practice Guidance (IPCC 2000) as the Tier 2 method. The results of the Monte Carlo simulation are therefore considered to provide a more realistic picture of the uncertainties than the results of the Tier 1 method.

Tier 2 uncertainty analysis produces an overall level uncertainty, which is slightly larger than the result of Tier 1 uncertainty analysis (T2: 3.86%, T1: 3.65%). The correct treating of large uncertainties, asymmetric distributions for agricultural sources, and – mainly – the existence of relevant positive correlations do all together increase the level uncertainty. This statement holds for the analyses with and without LULUCF.

The trend uncertainty of Tier 2 analysis is also larger than in Tier 1 analysis (T2: 2.87%, T1: 1.87%). Again, positive correlations for activity data and emission factors between of the base year and 2012 tend to increase the trend uncertainty (as may be seen from equation A1.8 of IPCC Good Practice Guidance (IPCC 2000) with $r > 0$). This statement holds again for the analyses with and without LULUCF.

1.7.2 KP-LULUCF Inventory

Uncertainty estimates for KP-LULUCF activities are presented in chapter 11.3.1.5.

1.8 Completeness Assessment

1.8.1 GHG Inventory

For all known sources, complete estimates are accomplished for all gases. Based on current knowledge, the Swiss inventory under the UNFCCC is complete.

1.8.2 KP-LULUCF Inventory

For all known sources and sinks, complete estimates are accomplished for the current submission. The Swiss LULUCF inventory under the Kyoto Protocol is complete.

2 Trends in Greenhouse Gas Emissions and Removals

This chapter provides an overview of Switzerland's GHG emissions/removals and trends for the period 1990–2012. Numbers in the chapters 2.1-2.4 are relevant for reporting under the UNFCCC, whereas numbers in chapter 2.5 refer to accounting under the KP.

2.1 Aggregated Greenhouse Gas Emissions 2012 (UNFCCC)

In 2012, Switzerland emitted 51'449 Gg CO₂ equivalent (excluding LULUCF and international bunkers) to the atmosphere or 6.43 tonnes CO₂ equivalent per capita (inhabitants 2012: 7.997 million, SFSO 2013a). The largest contributing gas was CO₂ (excluding LULUCF and international bunkers) with 43'251 Gg (5.41 tonnes per capita), and the most important source was sector 1 Energy, 41'477 Gg CO₂ equivalent (Table 2-1). A breakdown of Switzerland's total emissions by gas (excluding LULUCF) is given in Figure 2-1. Figure 2-2 charts the relative contributions of the individual sectors (excluding LULUCF) to the emission of each GHG.

Table 2-1 Switzerland's GHG emissions in CO₂ equivalent (Gg) by gas and sector in 2012.

| Emissions 2012 | CO ₂ | CH ₄ | N ₂ O | HFCs | PFCs | SF ₆ | Total | Share |
|---------------------------------------|---------------------------------|-----------------|------------------|--------------|-----------|-----------------|---------------|---------------|
| | CO ₂ equivalent (Gg) | | | | | | | |
| 1 Energy | 41'000 | 255 | 221 | | | | 41'477 | 80.6% |
| 2 Industrial Processes | 2'070 | 2 | 54 | 1'245 | 33 | 224 | 3'628 | 7.1% |
| 3 Solvent and Other Product Use | 155 | NA | 45 | | | | 200 | 0.4% |
| 4 Agriculture | NA | 3'143 | 2'395 | | | | 5'539 | 10.8% |
| 6 Waste | 12 | 287 | 292 | | | | 591 | 1.1% |
| 7 Other | 13 | 1 | 1 | | | | 14 | 0.0% |
| Total (excluding LULUCF) | 43'251 | 3'689 | 3'007 | 1'245 | 33 | 224 | 51'449 | 100.0% |
| 5 LULUCF | -1'142 | 9 | 4 | | | | -1'129 | -2.2% |
| Total (including LULUCF) | 42'109 | 3'698 | 3'011 | 1'245 | 33 | 224 | 50'320 | 97.8% |
| <i>International Aviation Bunkers</i> | 4'658 | 1 | 45.9 | | | | 4'705 | |
| <i>International Marine Bunkers</i> | 28 | 0.0048 | 0.3 | | | | 28 | |




Figure 2-1 Contribution of individual gases to Switzerland's GHG emissions (excluding LULUCF) in 2012. 100% correspond to 51'449 CO₂ eq (Gg). (Numbers may not add to total due to rounding.)




Figure 2-2 Relative contributions of the individual sectors (excluding LULUCF) to GHG emissions in 2012.

Fuel combustion within the energy sector was by far the largest source of emissions of CO₂ in 2012. Emissions of CH₄ and N₂O originated mainly from agriculture, and the F-gas emissions stemmed by definition from industrial processes.

2.2 Emission Trends by Gas

Emission trends by gas for the period 1990–2012 are summarized in Table 2-2.

Table 2-2 Switzerland's GHG emissions in CO₂ equivalent (Gg) by gas; 1990–2012 (corresponds to CRF-table 10s5/, 10s5.2, 10s5.3, upper half). The column below on the far right (digits in italics) indicates the percentage change in emissions in 2012 as compared to the base year 1990. HFCs increased by 5'526'548% when compared to 1990 levels (0.02 Gg CO₂ equivalent).

| Greenhouse Gas Emissions | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 42'677 | 44'356 | 44'111 | 40'386 | 40'550 | 40'519 | 41'588 | 40'257 | 41'833 | 42'724 |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 44'639 | 46'369 | 46'258 | 43'718 | 42'981 | 43'683 | 44'340 | 43'514 | 44'812 | 44'894 |
| CH ₄ emissions including CH ₄ from LULUCF | 4'576 | 4'563 | 4'471 | 4'377 | 4'330 | 4'328 | 4'283 | 4'225 | 4'170 | 4'073 |
| CH ₄ emissions excluding CH ₄ from LULUCF | 4'546 | 4'551 | 4'461 | 4'367 | 4'315 | 4'311 | 4'270 | 4'187 | 4'156 | 4'063 |
| N ₂ O emissions including N ₂ O from LULUCF | 3'472 | 3'463 | 3'433 | 3'345 | 3'307 | 3'295 | 3'289 | 3'177 | 3'161 | 3'124 |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'461 | 3'457 | 3'427 | 3'339 | 3'300 | 3'287 | 3'283 | 3'166 | 3'155 | 3'119 |
| HFCs | 0 | 0 | 7 | 15 | 34 | 182 | 228 | 301 | 358 | 421 |
| PFCs | 100 | 85 | 69 | 30 | 18 | 15 | 17 | 20 | 23 | 36 |
| SF ₆ | 144 | 146 | 148 | 126 | 112 | 98 | 94 | 131 | 160 | 147 |
| Total (including LULUCF) | 50'969 | 52'613 | 52'239 | 48'279 | 48'350 | 48'436 | 49'501 | 48'112 | 49'704 | 50'524 |
| Total (excluding LULUCF) | 52'890 | 54'607 | 54'370 | 51'595 | 50'760 | 51'576 | 52'233 | 51'319 | 52'663 | 52'680 |

| Greenhouse Gas Emissions | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 43'949 | 46'171 | 44'805 | 43'584 | 42'967 | 44'337 | 44'238 | 42'145 | 44'658 | 43'453 |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 43'952 | 44'904 | 43'844 | 45'025 | 45'650 | 46'290 | 45'911 | 43'931 | 45'447 | 44'280 |
| CH ₄ emissions including CH ₄ from LULUCF | 3'995 | 3'983 | 3'939 | 3'849 | 3'806 | 3'801 | 3'804 | 3'801 | 3'850 | 3'787 |
| CH ₄ emissions excluding CH ₄ from LULUCF | 3'985 | 3'973 | 3'922 | 3'829 | 3'797 | 3'791 | 3'793 | 3'787 | 3'840 | 3'777 |
| N ₂ O emissions including N ₂ O from LULUCF | 3'114 | 3'133 | 3'106 | 3'047 | 2'997 | 2'981 | 2'977 | 3'006 | 3'043 | 3'008 |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'109 | 3'128 | 3'100 | 3'040 | 2'993 | 2'977 | 2'972 | 3'001 | 3'039 | 3'004 |
| HFCs | 501 | 597 | 635 | 710 | 820 | 905 | 936 | 976 | 1'042 | 1'083 |
| PFCs | 69 | 45 | 40 | 57 | 53 | 33 | 33 | 29 | 39 | 36 |
| SF ₆ | 158 | 157 | 168 | 174 | 190 | 213 | 201 | 186 | 245 | 187 |
| Total (including LULUCF) | 51'787 | 54'085 | 52'694 | 51'421 | 50'834 | 52'271 | 52'189 | 50'144 | 52'878 | 51'554 |
| Total (excluding LULUCF) | 51'775 | 52'805 | 51'710 | 52'835 | 53'503 | 54'209 | 53'846 | 51'910 | 53'653 | 52'366 |

| Greenhouse Gas Emissions | 2010 | 2011 | 2012 | Change baseyear to 2012 (%) |
|---|---------------------------------|---------------|---------------|-----------------------------|
| | CO ₂ equivalent (Gg) | | | |
| CO ₂ emissions including net CO ₂ from LULUCF | 44'976 | 39'934 | 42'109 | -1.3% |
| CO ₂ emissions excluding net CO ₂ from LULUCF | 45'923 | 41'848 | 43'251 | -3.1% |
| CH ₄ emissions including CH ₄ from LULUCF | 3'773 | 3'723 | 3'698 | -19.2% |
| CH ₄ emissions excluding CH ₄ from LULUCF | 3'764 | 3'711 | 3'689 | -18.8% |
| N ₂ O emissions including N ₂ O from LULUCF | 3'082 | 3'019 | 3'011 | -13.3% |
| N ₂ O emissions excluding N ₂ O from LULUCF | 3'078 | 3'015 | 3'007 | -13.1% |
| HFCs | 1'138 | 1'195 | 1'245 | see caption |
| PFCs | 37 | 40 | 33 | -67.0% |
| SF ₆ | 155 | 164 | 224 | 56.0% |
| Total (including LULUCF) | 53'161 | 48'076 | 50'320 | -1.3% |
| Total (excluding LULUCF) | 54'095 | 49'973 | 51'449 | -2.7% |

The emission trends for individual gases are as follows (see Table 2-2 above, Table 2-3 and Figure 2-3 below):

- Total emissions (excluding LULUCF) show a minimum of 94.5% in 2011 and a maximum of 103.2% in 1991 (100%: value of base year 1990). In 2012, the total emissions were 2.7% lower than the emissions recorded in the base year 1990. CO₂ contributed the largest share of emissions, accounting for 84.1% of the total in 2012.
- Total emissions (including LULUCF) in 2012 show a decrease of 1.3% compared to the emissions recorded in the base year 1990. The net CO₂ emissions/removals from LULUCF show considerable variability from year to year, because heavy storms in 1990 and 1999 ("Lothar") and other factors influence the wood harvesting and tree mortality rates in forests. In the period 1990-2012, wood harvesting generally increased but is still exceeded by the growth of living biomass. This led to reductions in net removals within the LULUCF sector between 1990 and 2012. Within the first commitment period 2008-2012, the total net CO₂ sink steadily increased.
- A comparison of CO₂ emissions with the number of heating degree days in the period 1990–2012 (see Figure 2-7 below) indicates a strong correlation between CO₂ emissions and winter climatic conditions. In the last few years, an increase in heating degree days did not proportionally translate into an equal increase in CO₂ emissions. For a definition of heating degree days see footnote 2 displayed with Figure 2-7.
- Between 1990 and 2012, CH₄ (excluding LULUCF) decreased by 18.8%, which was mainly attributable to a reduction of livestock that led to a reduction of emissions from enteric fermentation. Moreover, from 2000, a change in waste legislation, banning the disposal of municipal solid waste in landfills, contributed to this trend. The CH₄ share of total GHG emissions decreased from 8.6% in 1990 to 7.2% in 2012.
- In parallel to the reduction of CH₄ due to decreases in livestock populations, N₂O emissions from manure management and agricultural soils declined. Total N₂O emissions (excluding LULUCF) dropped by 13.1% between 1990 and 2012 and accounts now for 5.8% of total emissions.
- HFC emissions increased significantly due to their application as substitutes for CFCs, while PFC emissions declined by 67.0%. SF₆ emissions have shown relatively large fluctuations between 94.4 and 244.7 Gg CO₂ eq since 1990. In 2012, SF₆ emissions increased by 56.0% compared to 1990. The share of all F-gases (HFCs, PFCs and SF₆) in total emissions (excluding LULUCF) increased from 0.5% in 1990 to 2.9% in 2012.

Table 2-3 Switzerland's total GHG emissions (excluding LULUCF) in CO₂ equivalent (Gg), selected years.

| Greenhouse Gas Emissions (excluding LULUCF) | 1990 | | 1995 | | 2000 | | 2005 | |
|--|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| CO ₂ | 44'639 | 84.4% | 43'683 | 84.7% | 43'952 | 84.9% | 46'290 | 85.4% |
| CH ₄ | 4'546 | 8.6% | 4'311 | 8.4% | 3'985 | 7.7% | 3'791 | 7.0% |
| N ₂ O | 3'461 | 6.5% | 3'287 | 6.4% | 3'109 | 6.0% | 2'977 | 5.5% |
| HFCs | 0 | 0.0% | 182 | 0.4% | 501 | 1.0% | 905 | 1.7% |
| PFCs | 100 | 0.2% | 15 | 0.0% | 69 | 0.1% | 33 | 0.1% |
| SF ₆ | 144 | 0.3% | 98 | 0.2% | 158 | 0.3% | 213 | 0.4% |
| Total (excluding LULUCF) | 52'890 | 100% | 51'576 | 100% | 51'775 | 100% | 54'209 | 100% |

| Greenhouse Gas Emissions (excluding LULUCF) | 2009 | | 2010 | | 2011 | | 2012 | |
|--|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| CO ₂ | 44'280 | 84.6% | 45'923 | 84.9% | 41'848 | 83.7% | 43'251 | 84.1% |
| CH ₄ | 3'777 | 7.2% | 3'764 | 7.0% | 3'711 | 7.4% | 3'689 | 7.2% |
| N ₂ O | 3'004 | 5.7% | 3'078 | 5.7% | 3'015 | 6.0% | 3'007 | 5.8% |
| HFCs | 1'083 | 2.1% | 1'138 | 2.1% | 1'195 | 2.4% | 1'245 | 2.4% |
| PFCs | 36 | 0.1% | 37 | 0.1% | 40 | 0.1% | 33 | 0.1% |
| SF ₆ | 187 | 0.4% | 155 | 0.3% | 164 | 0.3% | 224 | 0.4% |
| Total (excluding LULUCF) | 52'366 | 100% | 54'095 | 100% | 49'973 | 100% | 51'449 | 100% |

Figure 2-3 shows Switzerland's relative GHG emission trends by gas. The base year 1990 is set to 100%.




Figure 2-3 Relative trend of Switzerland's GHG emissions excluding LULUCF by gas, 1990–2012 (base year 1990: 100%). The increase of the F-gases is not shown (616% in 2012, compared to 1990).

2.3 Emission Trends by Sources and Sinks

Table 2-4 shows the emission trends for all major sources and sink categories. As the largest share of emissions originated from the energy sector, the table also includes the contributions of the energy sub-sectors.

Table 2-4 Switzerland's GHG emissions in CO₂ equivalent (Gg) by sources and sinks, 1990–2012. The column below on the far right (digits in *italics*) indicates the percentage change in emissions in 2012 as compared to the base year 1990.

| Source and Sink Categories | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| 1. Energy | 41'989 | 44'121 | 44'191 | 41'915 | 41'029 | 41'916 | 42'787 | 42'147 | 43'474 | 43'533 |
| 1A1 Energy Industries | 2'601 | 2'859 | 2'939 | 2'584 | 2'613 | 2'643 | 2'855 | 2'813 | 3'132 | 3'165 |
| 1A2 Manufacturing Industries and Construction | 6'138 | 6'326 | 5'988 | 5'909 | 5'911 | 6'106 | 5'885 | 5'784 | 5'969 | 5'954 |
| 1A3 Transport | 14'600 | 15'094 | 15'418 | 14'352 | 14'539 | 14'225 | 14'287 | 14'844 | 15'056 | 15'663 |
| 1A4 Other Sectors | 18'095 | 19'273 | 19'276 | 18'500 | 17'407 | 18'393 | 19'207 | 18'138 | 18'745 | 18'227 |
| 1A5 Other (Military) | 206 | 188 | 180 | 171 | 166 | 148 | 137 | 147 | 146 | 132 |
| 1B Fugitive emissions from oil and natural gas | 349 | 381 | 390 | 399 | 393 | 400 | 416 | 420 | 426 | 392 |
| 2. Industrial Processes | 3'320 | 2'957 | 2'793 | 2'484 | 2'649 | 2'626 | 2'498 | 2'431 | 2'517 | 2'580 |
| 3. Solvent and Other Product Use | 470 | 444 | 420 | 392 | 374 | 354 | 331 | 308 | 286 | 273 |
| 4. Agriculture | 6'092 | 6'069 | 5'979 | 5'877 | 5'843 | 5'819 | 5'780 | 5'606 | 5'578 | 5'511 |
| 6. Waste | 1'007 | 1'004 | 976 | 914 | 852 | 847 | 823 | 814 | 795 | 768 |
| 7. Other | 12 | 12 | 13 | 13 | 13 | 13 | 13 | 13 | 14 | 14 |
| Total (excluding LULUCF) | 52'890 | 54'607 | 54'370 | 51'595 | 50'760 | 51'576 | 52'233 | 51'319 | 52'663 | 52'680 |
| 5. Land Use, Land-Use Change and Forestry | -1'921 | -1'995 | -2'131 | -3'316 | -2'410 | -3'140 | -2'732 | -3'207 | -2'959 | -2'155 |
| Total (including LULUCF) | 50'969 | 52'613 | 52'239 | 48'279 | 48'350 | 48'436 | 49'501 | 48'112 | 49'704 | 50'524 |

| Source and Sink Categories | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | CO ₂ equivalent (Gg) | | | | | | | | | |
| 1. Energy | 42'429 | 43'330 | 42'297 | 43'497 | 43'922 | 44'432 | 44'084 | 42'078 | 43'628 | 42'506 |
| 1A1 Energy Industries | 3'064 | 3'187 | 3'268 | 3'295 | 3'617 | 3'804 | 4'078 | 3'835 | 4'025 | 3'949 |
| 1A2 Manufacturing Industries and Construction | 5'839 | 6'125 | 5'865 | 6'001 | 6'120 | 6'173 | 6'320 | 6'159 | 6'173 | 5'798 |
| 1A3 Transport | 15'896 | 15'597 | 15'522 | 15'689 | 15'767 | 15'827 | 15'939 | 16'257 | 16'624 | 16'427 |
| 1A4 Other Sectors | 17'133 | 17'946 | 17'186 | 18'092 | 18'021 | 18'239 | 17'360 | 15'462 | 16'450 | 15'983 |
| 1A5 Other (Military) | 136 | 134 | 140 | 125 | 114 | 124 | 127 | 120 | 115 | 116 |
| 1B Fugitive emissions from oil and natural gas | 362 | 341 | 317 | 295 | 283 | 265 | 260 | 245 | 242 | 232 |
| 2. Industrial Processes | 2'836 | 2'938 | 2'922 | 2'970 | 3'235 | 3'425 | 3'397 | 3'421 | 3'535 | 3'446 |
| 3. Solvent and Other Product Use | 259 | 245 | 234 | 225 | 212 | 211 | 206 | 205 | 202 | 201 |
| 4. Agriculture | 5'496 | 5'561 | 5'536 | 5'461 | 5'447 | 5'474 | 5'493 | 5'556 | 5'645 | 5'587 |
| 6. Waste | 741 | 717 | 708 | 669 | 673 | 654 | 653 | 637 | 629 | 612 |
| 7. Other | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 | 14 |
| Total (excluding LULUCF) | 51'775 | 52'805 | 51'710 | 52'835 | 53'503 | 54'209 | 53'846 | 51'910 | 53'653 | 52'366 |
| 5. Land Use, Land-Use Change and Forestry | 12 | 1'281 | 984 | -1'414 | -2'669 | -1'939 | -1'657 | -1'766 | -775 | -813 |
| Total (including LULUCF) | 51'787 | 54'085 | 52'694 | 51'421 | 50'834 | 52'271 | 52'189 | 50'144 | 52'878 | 51'554 |


| Source and Sink Categories | 2010 | 2011 | 2012 | 2012/1990 |
|--|---------------------------------|--------|--------|-----------|
| | CO ₂ equivalent (Gg) | | | % |
| 1. Energy | 44'004 | 39'945 | 41'477 | -1.2% |
| 1A1 Energy Industries | 4'180 | 3'968 | 4'064 | 56.3% |
| 1A2 Manufacturing Industries and Construction | 5'954 | 5'449 | 5'515 | -10.1% |
| 1A3 Transport | 16'321 | 16'205 | 16'331 | 11.9% |
| 1A4 Other Sectors | 17'193 | 13'994 | 15'240 | -15.8% |
| 1A5 Other (Military) | 121 | 108 | 116 | -43.6% |
| 1B Fugitive emissions from oil and natural gas | 234 | 220 | 209 | -40.0% |
| 2. Industrial Processes | 3'634 | 3'642 | 3'628 | 9.3% |
| 3. Solvent and Other Product Use | 199 | 202 | 200 | -57.5% |
| 4. Agriculture | 5'637 | 5'572 | 5'539 | -9.1% |
| 6. Waste | 608 | 598 | 591 | -41.3% |
| 7. Other | 14 | 14 | 14 | 16.2% |
| Total (excluding LULUCF) | 54'095 | 49'973 | 51'449 | -2.7% |
| 5. Land Use, Land-Use Change and Forestry | -934 | -1'897 | -1'129 | -41.2% |
| Total (including LULUCF) | 53'161 | 48'076 | 50'320 | -1.3% |

The percentage shares of source categories are shown for selected years in Table 2-5. Figure 2-4 to Figure 2-6 are graphical representations of Table 2-4 data. For the time series of the sub-sectors of 1 Energy see Chapter 3.

Table 2-5 Switzerland's total gross GHG emissions (excluding LULUCF) in CO₂ equivalent (Gg) and the contribution of individual source categories for selected years.

| Source and Sink Categories | 1990 | | 1995 | | 2000 | | 2005 | | 2007 | |
|--|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| 1. Energy | 41'989 | 79.4% | 41'916 | 81.3% | 42'429 | 81.9% | 44'432 | 82.0% | 42'078 | 81.1% |
| 1A1 Energy Industries | 2'601 | 4.9% | 2'643 | 5.1% | 3'064 | 5.9% | 3'804 | 7.0% | 3'835 | 7.4% |
| 1A2 Manufacturing Industries and Construction | 6'138 | 11.6% | 6'106 | 11.8% | 5'839 | 11.3% | 6'173 | 11.4% | 6'159 | 11.9% |
| 1A3 Transport | 14'600 | 27.6% | 14'225 | 27.6% | 15'896 | 30.7% | 15'827 | 29.2% | 16'257 | 31.3% |
| 1A4 Other Sectors | 18'095 | 34.2% | 18'393 | 35.7% | 17'133 | 33.1% | 18'239 | 33.6% | 15'462 | 29.8% |
| 1A5 Other (Military) | 206 | 0.4% | 148 | 0.3% | 136 | 0.3% | 124 | 0.2% | 120 | 0.2% |
| 1B Fugitive emissions from oil and natural gas | 349 | 0.7% | 400 | 0.8% | 362 | 0.7% | 265 | 0.5% | 245 | 0.5% |
| 2. Industrial Processes | 3'320 | 6.3% | 2'626 | 5.1% | 2'836 | 5.5% | 3'425 | 6.3% | 3'421 | 6.6% |
| 3. Solvent and Other Product Use | 470 | 0.9% | 354 | 0.7% | 259 | 0.5% | 211 | 0.4% | 205 | 0.4% |
| 4. Agriculture | 6'092 | 11.5% | 5'819 | 11.3% | 5'496 | 10.6% | 5'474 | 10.1% | 5'556 | 10.7% |
| 6. Waste | 1'007 | 1.9% | 847 | 1.6% | 741 | 1.4% | 654 | 1.2% | 637 | 1.2% |
| 7. Other | 12 | 0.0% | 13 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% |
| Total (excluding LULUCF) | 52'890 | 100.0% | 51'576 | 100.0% | 51'775 | 100.0% | 54'209 | 100.0% | 51'910 | 100.0% |

| Source and Sink Categories | 2008 | | 2009 | | 2010 | | 2011 | | 2012 | |
|--|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|-----------------------|--------|
| | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % | Gg CO ₂ eq | % |
| 1. Energy | 43'628 | 81.3% | 42'506 | 81.2% | 44'004 | 81.3% | 39'945 | 79.9% | 41'477 | 80.6% |
| 1A1 Energy Industries | 4'025 | 7.5% | 3'949 | 7.5% | 4'180 | 7.7% | 3'968 | 7.9% | 4'064 | 7.9% |
| 1A2 Manufacturing Industries and Construction | 6'173 | 11.5% | 5'798 | 11.1% | 5'954 | 11.0% | 5'449 | 10.9% | 5'515 | 10.7% |
| 1A3 Transport | 16'624 | 31.0% | 16'427 | 31.4% | 16'321 | 30.2% | 16'205 | 32.4% | 16'331 | 31.7% |
| 1A4 Other Sectors | 16'450 | 30.7% | 15'983 | 30.5% | 17'193 | 31.8% | 13'994 | 28.0% | 15'240 | 29.6% |
| 1A5 Other (Military) | 115 | 0.2% | 116 | 0.2% | 121 | 0.2% | 108 | 0.2% | 116 | 0.2% |
| 1B Fugitive emissions from oil and natural gas | 242 | 0.5% | 232 | 0.4% | 234 | 0.4% | 220 | 0.4% | 209 | 0.4% |
| 2. Industrial Processes | 3'535 | 6.6% | 3'446 | 6.6% | 3'634 | 6.7% | 3'642 | 7.3% | 3'628 | 7.1% |
| 3. Solvent and Other Product Use | 202 | 0.4% | 201 | 0.4% | 199 | 0.4% | 202 | 0.4% | 200 | 0.4% |
| 4. Agriculture | 5'645 | 10.5% | 5'587 | 10.7% | 5'637 | 10.4% | 5'572 | 11.1% | 5'539 | 10.8% |
| 6. Waste | 629 | 1.2% | 612 | 1.2% | 608 | 1.1% | 598 | 1.2% | 591 | 1.1% |
| 7. Other | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% | 14 | 0.0% |
| Total (excluding LULUCF) | 53'653 | 100.0% | 52'366 | 100.0% | 54'095 | 100.0% | 49'973 | 100.0% | 51'449 | 100.0% |

Figure 2-4 Switzerland's GHG emissions in CO₂ equivalent (Gg) by sectors, 1990–2012 (excluding LULUCF).




Figure 2-5 Relative emission (CO₂ eq.) trends by main source categories (base year 1990 = 100%).

Emission trends for the various sectors are as follows:

- **1 Energy:** the variations can only be understood if the trends within individual source categories are considered separately. See Figure 2-6 and comments below.
- **2 Industrial Processes:** in line with economic development, overall emissions in the industry sector showed a decreasing trend in the early 90s and a gradual increase between 1998 and 2012, except for the economically difficult year 2009. Since 2005 the Ordinance on Chemical Risk Reduction (Swiss Confederation 2005) is in place and regulates the use of F-Gases, which led to an emission stabilization in this source category.
- **3 Solvent and Other Product Use:** there is a decreasing trend in overall emissions throughout all the years, which is however by far less pronounced since 2004. Whereas overall NMVOC emissions have decreased by 54.9% since 1990, direct CO₂ emissions from the post combustion of NMVOCs have increased. NMVOC emissions, the main source of indirect CO₂ emissions, have diminished between 1990 and 2004 due to their limitation brought by the Ordinance on Air Pollution Control (Swiss Confederation 1985) and due to the introduction of the VOC-tax in 2000 (Swiss Confederation 1997). Since 2004, emissions have remained relatively stable.
- **4 Agriculture:** declining populations of cattle and swine and reduced fertilizer use have led to a decrease in CO₂ equivalent emissions until 2000. Since then, CH₄ emissions slightly increased again.
- **6 Waste:** Total emissions from the source category Waste decreased steadily throughout the period 1990-2012. Since 2000, emissions have been further reduced by a change in legislation: disposal of combustible wastes in landfills has been banned, leading to an increasing amount of municipal solid waste being incinerated, with emissions reported under source 1A1 Energy Industries rather than sector 6 Waste. Altogether, “waste-related” emissions incl. emissions from waste management activities reported in sources 1A (Waste to energy) 4D (waste used as fertilizer) and 6 (waste treatment) have increased since 1990 by 34.25 % (see Figure 8-3 and 8-4 in Chapter 8).
- **7 Other:** The total emissions from sector 7 Other increased throughout the period 1990-2000. Since 2000 the emissions are stable. Please consider that emissions from sector

7 Other are not accounted for in the Kyoto Protocol and are only of minor importance (0.03% of total CO₂ equivalent emissions).

The main source categories within the Energy sector – representing the major sources of Switzerland's GHG emissions – are shown in Figure 2-6.




Figure 2-6 Emission trends (CO₂ eq.) for the source categories in sector 1 Energy/1A Fuel Combustion. The trend for the entire sector “1 Energy” is shown with a bold line. Not included in the figure is the trend for 1B Fugitive Emissions, which continuously decreased from 100% in 1990 to 60% in 2012.

It is noteworthy that, due to Switzerland's electricity production structure (about 94.5% generated by hydroelectric and nuclear power plants in 2012; see SFOE 2013: Table 24), sector 1A1 Energy Industries plays only a minor role – representing not classical thermal power stations but waste incineration plants in the Swiss GHG inventory. The following emission trends emerges within the Energy sector:

- Despite differing trends of individual source categories, the overall emissions from the energy sector remain at relatively constant level (bold line in Figure 2-6) until 2010 but noticeably decreased in 2011, mainly due to an exceptionally warm winter (see Figure 2-7). In 2012 a slight increase was detectable again.
- Overall emissions from source category 1A1 Energy Industry 2012 have increased by 56.3% since 1990. Fluctuations are caused by varying combustion activities in the petroleum refinery industry, waste incineration, new installations of district heating and weather related forcing of heating activities (see Figure 2-7). From 2010 to 2011, emissions from Gaseous Fuel consumption within source category 1A decreased by 11.3% due to the fact that 2011 was the warmest year measured since measurements started. Note that only approximately 10% of sector 1 Energy emissions stem from 1A1.
- The trend for sub-sector 1A3 Transport shows a slight increase over the period 1990–2008 by about 12%, but with fluctuations indicating a fairly strong correlation between this sector and overall economic development in Switzerland, with periods of stagnation (1991–1996, 2001–2003 and 2009) and growth (gross value-added) in 1997–2000 and 2004–2012 (except for 2009) (SFSO 2009a, SECO 2014). Since 2008 transport emissions show a slight decrease which points to a relative decoupling from overall economic development.

- The trend for sub-sector 1A4 Other Sectors reflects the impact of climatic variations on demand for heating. The strong correlation with the number of “heating degree days”² – used as an index of cold weather conditions – is apparent from Figure 2-7, which shows CO₂ emissions from sub-sector 1A4 Fuel Combustion – Other Sectors (only stationary sources) and the number of heating degree days. In 2012 heating degree days increased by 11.7% compared to 2011 and CO₂ emissions from fuel combustions in source category 1A4 Stationary Sources increased simultaneously by 9.3%. In the period 1990–2012, the number of buildings and apartments increased, as well as the average floor space per person and workplace. Both phenomena resulted in an increase in the total area heated by more than 30%. Over the same period, however, higher standards were specified for insulation and for combustion equipment efficiency for both new and renovated buildings, compensating for the emissions from the additional area heated.




Figure 2-7 Relative trend for CO₂ emissions from 1A4 Fuel Combustion - Other Sectors (stationary sources only) compared with the number of heating degree days.

Figure 2-8 shows the net emissions and removals from the LULUCF sector in Switzerland, which is dominated by biomass dynamics in forests. Except for 2001 and 2002 the removals in the LULUCF sector were higher than the emissions throughout the period 1990-2012. However, a strong year to year variation is evident over the whole period. The net removals decreased by 41.2% since 1990 but increased by 45.7% since 2008. The reason for the positive value in 2001 and 2002 is the winter storm “Lothar” end of 1999 which caused great damages in the forest stands and increased harvesting. The reduction of the removals from 2004 to 2008 is due to the reduction of dead wood as CO₂ sink (2005) towards a source in 2008.

² Heating degree days: Number of degrees per day calculated as the difference between 20°C (room temperature) and the daily average outdoor temperature for such days where the daily average temperature is below 12°C (e.g. daily outdoor average equals 7°C, then for that day 20 – 7 = 13). The number of degrees per day are summed up for a year t to yield the heating degree days of year t.




Figure 2-8 Switzerland's net CO₂ equivalent balance of sector Land Use, Land-Use Change and Forestry (LULUCF) 1990–2012 in Gg. Positive values refer to emissions, negative values refer to removals. Note that the annual contributions of CH₄ and N₂O emissions from LULUCF in this period are very small compared to the net CO₂ emissions and removals.

2.4 Emission Trends for Indirect Greenhouse Gases and SO₂

Emission trends for indirect greenhouse gases show a very pronounced decline (see Table 2-6 and Figure 2-9). A strict air pollution control policy and the implementation of a large number of emission reduction measures led to a decrease between 49% and 74% in emissions of respective air pollutants over the period 1990–2012. The main reduction measures were abatement of exhaust emissions from road vehicles and stationary combustion equipment, taxation of solvents and sulphured fuels, and voluntary agreements with industry sectors (FOEN 2010i, Swiss Confederation 1985, 1997).

Table 2-6 Switzerland's indirect GHG and SO₂ emissions (Gg), 1990–2012 (without NMVOC from LULUCF).

| Indirect Greenhouse Gases and SO ₂ | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|------|------|------|------|------|------|------|------|------|
| Gg | | | | | | | | | | |
| NO _x | 145 | 143 | 136 | 125 | 123 | 119 | 115 | 111 | 111 | 111 |
| CO | 800 | 765 | 712 | 627 | 576 | 536 | 518 | 488 | 467 | 454 |
| NMVOC | 306 | 289 | 268 | 240 | 219 | 203 | 190 | 177 | 164 | 155 |
| SO ₂ | 41 | 37 | 34 | 27 | 28 | 26 | 26 | 24 | 23 | 17 |
| Indirect Greenhouse Gases and SO ₂ | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| Gg | | | | | | | | | | |
| NO _x | 108 | 104 | 99 | 96 | 94 | 93 | 90 | 87 | 86 | 80 |
| CO | 427 | 405 | 378 | 366 | 347 | 330 | 307 | 289 | 279 | 263 |
| NMVOC | 146 | 137 | 126 | 118 | 108 | 104 | 100 | 97 | 95 | 93 |
| SO ₂ | 16 | 18 | 16 | 15 | 16 | 16 | 15 | 13 | 14 | 12 |
| Indirect Greenhouse Gases and SO ₂ | 2010 | 2011 | 2012 | | | | | | | |
| Gg | | | | | | | | | | |
| NO _x | 78 | 74 | 74 | | | | | | | |
| CO | 253 | 233 | 227 | | | | | | | |
| NMVOC | 91 | 88 | 86 | | | | | | | |
| SO ₂ | 12 | 10 | 11 | | | | | | | |




Figure 2-9 Relative trends for indirect GHG and SO₂ emissions (without NMVOC from LULUCF), 1990–2012 (base year 1990 = 100%).

The energy sector was by far the largest source of indirect greenhouse gas emissions (see Table 2-7), with the only exception being NMVOC, where sector 3 Solvent and Other Product Use accounted for 23.9% of the total. The total shown in Table 2-7 includes NMVOC emissions from LULUCF, which are estimated at constant 95.5 Gg per year (SAEFL 1996a). This corresponds to 52.7% of the total in 2012.

Table 2-7 Indirect GHG and SO₂ emissions (Gg) by source, 2012. The total NMVOC emissions including NMVOC from LULUCF.

| Sources | NO _x | CO | NMVOC | SO ₂ |
|---------------------------------|---------------------|---------------|---------------|-----------------|
| | Emissions 2012 (Gg) | | | |
| 1 Energy | 68.78 | 218.65 | 29.89 | 9.68 |
| 2 Industrial Processes | 0.40 | 4.85 | 6.73 | 0.89 |
| 3 Solvent and Other Product Use | 0.00 | 0.01 | 43.27 | 0.01 |
| 4 Agriculture | 4.08 | NO | 3.91 | NO |
| 5 LULUCF | IE, NE | IE, NE | 95.52 | NE |
| 6 Waste | 0.38 | 2.58 | 1.91 | 0.08 |
| 7 Other | 0.07 | 0.80 | 0.13 | 0.01 |
| Total | 73.71 | 226.90 | 181.36 | 10.67 |

Figure 2-10 shows the relative contributions (excluding LULUCF) of the various sectors for each individual gas (data from Table 2-7). The energy sector can clearly be identified as the main source of NO_x, CO and SO₂.




Figure 2-10 Relative contributions of individual sectors to indirect GHG and SO₂ emissions in 2012 (without NMVOC from LULUCF).

2.5 Emission Trends (Kyoto Protocol)

Relevant emission and removals under the Kyoto Protocol are shown in Table 2-8 and Table 2-9, sorted by sectors and gases respectively. Base year emissions for the first commitment period are fixed at the value reported in the Initial Report 2006 (FOEN 2006h, UNFCCC 2007a).

Table 2-8 Summary of Switzerland's GHG emissions in CO₂ equivalent (Gg), 1990–2012 excluding emissions from LULUCF, Other and International Bunkers.

| Annex A sources | Sector | Base year initial report | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| Annex A sources | 1 Energy | 42'134 | 41'989 | 44'121 | 44'191 | 41'915 | 41'029 | 41'916 | 42'787 | 42'147 | 43'474 |
| | 2 Industrial Processes | 3'258 | 3'320 | 2'957 | 2'793 | 2'484 | 2'649 | 2'626 | 2'498 | 2'431 | 2'517 |
| | 3 Solvent and Other Product Use | 466 | 470 | 444 | 420 | 392 | 374 | 354 | 331 | 308 | 286 |
| | 4 Agriculture | 5'903 | 6'092 | 6'069 | 5'979 | 5'877 | 5'843 | 5'819 | 5'780 | 5'606 | 5'578 |
| | 6 Waste | 1'030 | 1'007 | 1'004 | 976 | 914 | 852 | 847 | 823 | 814 | 795 |
| | Total (Annex A sources) | 52'791 | 52'878 | 54'595 | 54'358 | 51'582 | 50'747 | 51'563 | 52'220 | 51'305 | 52'650 |

| Annex A sources | Sector | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | | | | | | | | | | |
| Annex A sources | 1 Energy | 43'533 | 42'429 | 43'330 | 42'297 | 43'497 | 43'922 | 44'432 | 44'084 | 42'078 | 43'628 |
| | 2 Industrial Processes | 2'580 | 2'836 | 2'938 | 2'922 | 2'970 | 3'235 | 3'425 | 3'397 | 3'421 | 3'535 |
| | 3 Solvent and Other Product Use | 273 | 259 | 245 | 234 | 225 | 212 | 211 | 206 | 205 | 202 |
| | 4 Agriculture | 5'511 | 5'496 | 5'561 | 5'536 | 5'461 | 5'447 | 5'474 | 5'493 | 5'556 | 5'645 |
| | 6 Waste | 768 | 741 | 717 | 708 | 669 | 673 | 654 | 653 | 637 | 629 |
| | Total (Annex A sources) | 52'666 | 51'761 | 52'790 | 51'696 | 52'821 | 53'489 | 54'195 | 53'832 | 51'896 | 53'639 |

| KP-LULUCF | Art. 3.3 | Sector | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------|-------------------------------|-------------------------------|---------------------------------|---------------|---------------|---------------|------------------|
| | | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | | |
| KP-LULUCF | Art. 3.3 | Afforestation & reforestation | -24 | -23 | -20 | -17 | |
| | | Deforestation | 187 | 220 | 221 | 222 | |
| | Art. 3.4 | Forest management | -1'419 | -2'020 | -2'064 | -2'236 | |
| | | Cropland management | NA | NA | NA | NA | |
| | | Grazing land management | NA | NA | NA | NA | |
| | | Revegetation | NA | NA | NA | NA | |
| | Total (Art. 3.3 + 3.4) | | -1'257 | -1'823 | -1'862 | -2'032 | |

| Annex A sources | Sector | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------------|---------------------------------|---------------------------------|---------------|---------------|---------------|------------------|
| | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | |
| Annex A sources | 1 Energy | 42'506 | 44'004 | 39'945 | 41'477 | -1.6% |
| | 2 Industrial Processes | 3'446 | 3'634 | 3'642 | 3'628 | 11.4% |
| | 3 Solvent and Other Product Use | 201 | 199 | 202 | 200 | -57.1% |
| | 4 Agriculture | 5'587 | 5'637 | 5'572 | 5'539 | -6.2% |
| | 6 Waste | 612 | 608 | 598 | 591 | -42.5% |
| | Total (Annex A sources) | 52'352 | 54'081 | 49'959 | 51'435 | -2.6% |

| KP-LULUCF | Art. 3.3 | Sector | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------|-------------------------------|-------------------------------|---------------------------------|---------------|---------------|---------------|------------------|
| | | | CO ₂ equivalent (Gg) | | | | % |
| | | | | | | | |
| KP-LULUCF | Art. 3.3 | Afforestation & reforestation | -24 | -23 | -20 | -17 | |
| | | Deforestation | 187 | 220 | 221 | 222 | |
| | Art. 3.4 | Forest management | -1'419 | -2'020 | -2'064 | -2'236 | |
| | | Cropland management | NA | NA | NA | NA | |
| | | Grazing land management | NA | NA | NA | NA | |
| | | Revegetation | NA | NA | NA | NA | |
| | Total (Art. 3.3 + 3.4) | | -1'257 | -1'823 | -1'862 | -2'032 | |

Table 2-9 Switzerland's total GHG emissions (excluding LULUCF, Other and International Bunkers) and the contribution of individual gases in CO₂ equivalent (Gg), 1990-2012.

| Annex A sources | GHG | Base year initial report | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|-----------------|--------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | CO ₂ | 44'553 | 44'628 | 46'357 | 46'247 | 43'706 | 42'969 | 43'672 | 44'328 | 43'502 |
| | CH ₄ | 4'370 | 4'545 | 4'551 | 4'460 | 4'367 | 4'315 | 4'310 | 4'269 | 4'186 | 4'155 |
| | N ₂ O | 3'623 | 3'461 | 3'456 | 3'426 | 3'339 | 3'299 | 3'287 | 3'283 | 3'165 | 3'154 |
| | HFCs | 0.0 | 0.0 | 0.2 | 7 | 15 | 34 | 182 | 228 | 301 | 358 |
| | PFCs | 100 | 100 | 85 | 69 | 30 | 18 | 15 | 17 | 20 | 23 |
| | SF ₆ | 144 | 144 | 146 | 148 | 126 | 112 | 98 | 94 | 131 | 160 |
| | Total (Annex A sources) | 52'791 | 52'878 | 54'595 | 54'358 | 51'582 | 50'747 | 51'563 | 52'220 | 51'305 | 52'650 |

| Annex A sources | GHG | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|-----------------|--------------------------------|---------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | CO ₂ equivalent (Gg) | | | | | | | | | |
| | | CO ₂ | 44'881 | 43'939 | 44'891 | 43'831 | 45'012 | 45'637 | 46'277 | 45'898 | 43'918 |
| | CH ₄ | 4'063 | 3'985 | 3'972 | 3'922 | 3'828 | 3'797 | 3'791 | 3'793 | 3'786 | 3'840 |
| | N ₂ O | 3'118 | 3'109 | 3'127 | 3'099 | 3'039 | 2'992 | 2'976 | 2'971 | 3'001 | 3'039 |
| | HFCs | 421 | 501 | 597 | 635 | 710 | 820 | 905 | 936 | 976 | 1'042 |
| | PFCs | 36 | 69 | 45 | 40 | 57 | 53 | 33 | 33 | 29 | 39 |
| | SF ₆ | 147 | 158 | 157 | 168 | 174 | 190 | 213 | 201 | 186 | 245 |
| | Total (Annex A sources) | 52'666 | 51'761 | 52'790 | 51'696 | 52'821 | 53'489 | 54'195 | 53'832 | 51'896 | 53'639 |

| KP-LULUCF | Art.3.3 | CO ₂ | | | | | | | | | 82 |
|-----------|-------------------------------|------------------|--|--|--|--|--|--|--|--|---------------|
| | | CH ₄ | | | | | | | | | NO |
| | | N ₂ O | | | | | | | | | 0.1 |
| Art.3.4 | CO ₂ | | | | | | | | | | -1'204 |
| | CH ₄ | | | | | | | | | | 1.0 |
| | N ₂ O | | | | | | | | | | 0.2 |
| | Total (Art. 3.3 + 3.4) | | | | | | | | | | -1'121 |

| Annex A sources | GHG | 2009 | 2010 | 2011 | 2012 | 2012 – base year |
|-----------------|--------------------------------|---------------------------------|---------------|---------------|---------------|---------------------|
| | | CO ₂ equivalent (Gg) | | | | % |
| | | CO ₂ | 44'267 | 45'910 | 41'835 | 43'238 |
| | CH ₄ | 3'776 | 3'763 | 3'710 | 3'688 | -15.6% |
| | N ₂ O | 3'003 | 3'077 | 3'014 | 3'007 | -17.0% |
| | HFCs | 1'083 | 1'138 | 1'195 | 1'245 | NA |
| | PFCs | 36 | 37 | 40 | 33 | -67.0% |
| | SF ₆ | 187 | 155 | 164 | 224 | 56.0% |
| | Total (Annex A sources) | 52'352 | 54'081 | 49'959 | 51'435 | -2.6% |

| KP-LULUCF | Art.3.3 | CO ₂ | 162 | 197 | 202 | 205 |
|-----------|-------------------------------|------------------|---------------|---------------|---------------|-----|
| | | CH ₄ | NO | NO | NO | NO |
| | | N ₂ O | 0.0 | 0.0 | 0.0 | 0.0 |
| Art.3.4 | CO ₂ | -1'420 | -2'021 | -2'068 | -2'237 | |
| | CH ₄ | 0.8 | 0.5 | 3.2 | 0.4 | |
| | N ₂ O | 0.2 | 0.1 | 0.7 | 0.1 | |
| | Total (Art. 3.3 + 3.4) | -1'257 | -1'823 | -1'862 | -2'032 | |

The reported total emissions differ from those reported under the UNFCCC, as sector Other – in addition to LULUCF and international bunkers – is not accounted for under the Kyoto Protocol. On the other hand, activities under article 3.3 (Afforestation, reforestation and Deforestation) and 3.4 (forest-, cropland- and grazing management and revegetation) are taken into account over the commitment period 2008-2012. Under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol, Switzerland only accounts for Forest Management.

3 Energy

3.1 Overview

This chapter provides information on the estimation of the greenhouse gas emissions from the energy sector. The following source categories are reported:

- 1A Fuel Combustion
- 1B Fugitive Emissions from Fuels

In Switzerland, the energy sector is the most relevant greenhouse gas source. In 2012, it emitted 41'477 Gg CO₂ equivalent which corresponds to 80.6% of total emissions (51'449 Gg CO₂ equivalent, national total without LULUCF). The emissions of the period 1990–2012 are depicted in Figure 3-1.




Figure 3-1 Switzerland's GHG emissions of sector 1 Energy 1990–2012 in CO₂ equivalent (Gg).

For the total emissions of the energy sector, there are fluctuations up to 6% (100% = emissions 1990) in the period 1990–2012 but no trends. The value 2012 is 1.2% lower than the value of the base year 1990. Three sub-categories dominate the emissions:

- 1A3 Transport and 1A4 Other Sectors are the main sources of the sector energy that cover 39.4% and 36.7% of total energy emissions in 2012, respectively.
- 1A1 Energy Industries and 1A2 Manufacturing Industries and Construction are of lesser importance. They contribute 9.8% and 13.3% to the total emissions of the sector energy in 2012, respectively.
- 1A5 Other (Military) and 1B Fugitive Emissions only play a minor role. In 2012, they cover 0.3% and 0.5%, respectively, of the total emissions of the sector energy.

The trends of the individual gases are given in the next table and figure:

- By far the most important gas emitted from the sector energy is CO₂. It accounts for 98.9% of the the total greenhouse gas emissions of the sector. Its fluctuations reflect *inter alia* the climatic variability in Switzerland (see Figure 2-7 and related comments).

- In 2012, CH₄ emissions contributed 0.62% to the total emissions of the sector energy. The decreasing trend since 1990 is the result of improved gas transmission network resulting in substantially lower fugitive emissions (12.56 Gg in 1990, 8.07 Gg in 2012) and reduced emissions from gasoline passenger cars due to catalytic converters. Furthermore improved combustion technologies in 1A4 Other Sectors also contributed to the decreasing trend.
- N₂O contributed 0.53% to the total emissions of the sector energy. The changes in N₂O emissions may mainly be explained by changes in the emission of road transportation and revised EFs for diesel and gasoline combustion by vehicles. The first generation of catalytic converters generated N₂O as undesirable by-product in the exhaust gases, leading to an increase of N₂O emissions until 2000. With new converter materials being used, the emission factors are decreasing since 2001. For further details see chapter 3.2.8.2 - 1A3b.

Table 3-1 GHG emissions of source category 1 Energy by gas in CO₂ equivalent (Gg), 1990–2012.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 41'199 | 43'297 | 43'360 | 41'101 | 40'223 | 41'098 | 41'948 | 41'315 | 42'637 | 42'730 |
| CH ₄ | 502 | 521 | 516 | 512 | 500 | 507 | 518 | 509 | 512 | 479 |
| N ₂ O | 288 | 304 | 314 | 302 | 306 | 310 | 321 | 323 | 325 | 325 |
| Sum | 41'989 | 44'121 | 44'191 | 41'915 | 41'029 | 41'916 | 42'787 | 42'147 | 43'474 | 43'533 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 41'675 | 42'611 | 41'627 | 42'860 | 43'357 | 43'888 | 43'563 | 41'579 | 43'126 | 42'018 |
| CH ₄ | 439 | 414 | 382 | 362 | 342 | 325 | 307 | 291 | 285 | 274 |
| N ₂ O | 315 | 305 | 288 | 275 | 223 | 219 | 214 | 209 | 216 | 215 |
| Sum | 42'429 | 43'330 | 42'297 | 43'497 | 43'922 | 44'432 | 44'084 | 42'078 | 43'628 | 42'506 |

| Gas | 2010 | 2011 | 2012 |
|---------------------------------|--------|--------|--------|
| CO ₂ equivalent (Gg) | | | |
| CO ₂ | 43'503 | 39'479 | 41'000 |
| CH ₄ | 281 | 254 | 255 |
| N ₂ O | 220 | 212 | 221 |
| Sum | 44'004 | 39'945 | 41'477 |




Figure 3-2 Relative trends of the greenhouse gases of source category 1 "Energy" in the period 1990–2012. The base year 1990 represents 100%.

The following table summarises the emissions of the sector energy in 2012. The table includes emissions from international bunkers (aviation and marine) as well as CO₂ emissions from biomass burning which both are not accounted for in the Kyoto Protocol but are contained in the CRF-tables.

Table 3-2 Summary of sector 1 Energy, emissions³ in 2012 in Gg CO₂ equivalent (Total: rounded values).

| Emissions 2012 | CO ₂ | CH ₄ | N ₂ O | Total |
|---|---------------------------------|-----------------|------------------|--------|
| | CO ₂ equivalent (Gg) | | | |
| 1 Energy | 41'000.2 | 255.3 | 221.2 | 41'477 |
| 1A Fuel Combustion | 40'960.9 | 85.9 | 220.6 | 41'267 |
| 1A1 Energy Industries | 4'018.4 | 2.2 | 43.8 | 4'064 |
| 1A2 Manufacturing Industries and Construction | 5'480.1 | 7.9 | 27.4 | 5'515 |
| 1A3 Transport | 16'210.1 | 20.3 | 101.0 | 16'331 |
| 1A4 Other Sectors | 15'137.5 | 55.4 | 47.2 | 15'240 |
| 1A5 Other (Military) | 114.8 | 0.1 | 1.1 | 116 |
| 1B Fugitive Emissions from Fuels | 39.3 | 169.5 | 0.7 | 209 |
| International Bunkers | 4'685.2 | 1.4 | 46.1 | 4'733 |
| CO ₂ Emissions from Biomass | 6'804.5 | IE | IE | 6'804 |

In 2012, the Swiss greenhouse gas inventory identifies in Tier 1 analysis 31 key sources (without LULUCF), 19 of which belong to the energy sector. The key categories from the energy sector are depicted in Figure 3-3. Most dominant are the CO₂ emissions from 1A3b Transport (gasoline, CO₂) and 1A4b Other Sectors (liquid fuels, CO₂).

³ For full biomass CO₂ emissions see Table 3-14.




Figure 3-3 Key sources in the Swiss GHG inventory from the energy sector. Top: Tier 1, bottom Tier 2 analysis.

3.2 Source Category 1A – Fuel Combustion Activities

3.2.1 Comparison of the Sectoral Approach with the Reference Approach

Two methods are applied for modelling CO₂ emissions from the energy sector, the Sectoral (or National) Approach and the Reference Approach. For the inventory under the Framework Convention on Climate Change and the Kyoto Protocol the Sectoral (National) Approach is used. The Reference Approach is only used for verification purposes (quality control activity).

Figure 3-4 depicts the two approaches including the input data used and disaggregation of fuel types that ultimately allows for comparing the two approaches.




Figure 3-4 Calculation of the Reference and Sectoral Approach. The input data for both approaches stems from the Swiss overall energy statistics (SFOE 2013) but while the Reference Approach considers the net import/export balance, the Sectoral Approach considers the fuel use of the SFOE.

The Sectoral Approach is based on sectoral energy consumption and uses both the energy consumption as reported in the Swiss overall energy statistics (SFOE 2013) and additional specific methods for the various source categories. This includes fossil fuel consumption statistics (top-down approach, Tier 1) and bottom-up modelling of fuel consumption (bottom-up, Tier 2 and Tier 3). A detailed description of the sectoral approach is provided in chapter 3.2.5.

The Reference Approach on the other hand corresponds to a top-down approach (Tier 1) based on net quantities of fuel imported into Switzerland. Accordingly the fossil fuel supply statistics is used in this approach: all imports and exports of primary fuels (crude oil, natural

gas, coal⁴), secondary fuels (gasoline, diesel oil etc.) and stock changes stem from the Swiss overall energy statistics (SFOE 2013). Subsequently the apparent consumption, the net carbon emissions, and the effective CO₂ emissions are calculated for the Reference Approach as reported in the CRF-tables 1A(b)–1A(d). Thus the Reference Approach covers the CO₂ emissions of all net imported primary fuels, emissions from crude oil refinement (secondary fuel production) in the two Swiss refineries and emissions of imported secondary fuels. In 2012 29% of all fossil fuels sold in Switzerland were produced in Swiss refineries from primary fuels (EV 2013).

All necessary data for calculating the Reference Approach is implemented in the EMIS database and all the data on import, export, bunkers, stock changes, apparent consumption, carbon emission factors, carbon stored and actual emissions are calculated within EMIS under the following conditions:

- The oxidation factor is consequently set to 1.0 due to the following reason: combustion installations in Switzerland have very good combustion properties; combined emissions of CO and unburnt VOC lie in the range of only 0.1 to 0.3 percent of CO₂ emissions for oil and gas combustion. Since most of the coal used in Switzerland goes to the cement industry, also for coal an oxidation factor of 1.0 was chosen⁵. For detailed description see chapter 3.2.5.2.
- For the Reference Approach gas oil and diesel are reported together, with a weighted average NCV (NCV gas oil 42.6 TJ/Gg, NCV diesel 42.8 TJ/Gg). In contrast, marine bunkers consist of diesel only and are reported using the country-specific NCV of 42.8 TJ/Gg.
- For the Reference Approach, Liechtenstein's fossil fuel consumption is subtracted from the input figures of fossil fuel consumption as stated in SFOE (2013), which originally include Liechtenstein's consumption except for natural gas (see also chapter 3.2.5).
- In the Reference Approach, carbon which is stored in feedstocks or non-energy fuel use has to be subtracted from fuel import in order to report the effective CO₂ emissions correctly (see also chapter 3.2.3).

On this basis the differences between Reference and Sectoral Approach are calculated within the EMIS system. The results 1990-2012 are shown in Table 3-3 and in Figure 3-5. The CO₂ emissions from both approaches (excluding non-energy use and feedstocks) concur very well. For all years the differences lie between 0.70% and 1.29% and the difference in 2012 is 0.89%. For the corresponding energy consumption (excluding non-energy use and feedstocks) the differences lie between 0.57% and 0.98% while the difference in 2012 is 0.62%.

The comparably small difference between Reference and Sectoral Approach is influenced by various effects. Amongst others, the energy and carbon content of crude oil varies over time. Furthermore the efficiency and amount of production of Swiss refineries varies from year to year.

⁴ Coking coal is included under other bituminous coal in the reference approach.

⁵ EC 2004, Annex VII, Section 2.1.1: "In cement kilns the incomplete combustion of fossil fuels is negligible, due to the very high combustion temperatures, long residence time in kilns and minimal residual carbon found in clinker. Carbon in all kiln fuels shall therefore be accounted for as fully oxidized (oxidation factor = 1.0)."

Table 3-3 Differences in energy consumption and CO₂ emissions between the Reference and the Sectoral (National) Approach from CRF-table1.A(c). The difference is calculated according to $[(RA-SA)/SA]$ 100% with RA = Reference Approach, SA = Sectoral (National) Approach. Energy consumption: excluding non-energy use and feedstocks.

| | Difference between Reference and Sectoral Approach | | | | | | | | | |
|---------------------------|--|------|------|------|------|------|------|------|------|------|
| | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | % | | | | | | | | | |
| Energy Consumption | 0.69 | 0.87 | 0.86 | 0.98 | 0.95 | 0.84 | 0.73 | 0.75 | 0.64 | 0.57 |
| CO ₂ Emissions | 0.75 | 0.91 | 0.93 | 1.10 | 1.11 | 0.93 | 0.88 | 0.98 | 0.89 | 0.79 |

| | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------|------|------|------|------|------|------|------|------|------|------|
| | % | | | | | | | | | |
| Energy Consumption | 0.61 | 0.67 | 0.61 | 0.57 | 0.68 | 0.75 | 0.93 | 0.70 | 0.68 | 0.79 |
| CO ₂ Emissions | 0.79 | 0.82 | 0.85 | 0.70 | 0.93 | 1.02 | 1.29 | 1.02 | 1.01 | 1.16 |

| | 2010 | 2011 | 2012 |
|---------------------------|------|------|------|
| | % | | |
| Energy Consumption | 0.73 | 0.70 | 0.62 |
| CO ₂ Emissions | 1.09 | 1.10 | 0.89 |




Figure 3-5 Time series for the differences between Reference and Sectoral Approach. Numbers are taken from Table 3-3. See caption in Table 3-3 for further information about how data is calculated.

After a major revision (INFRAS 2010a), the calculation of the Reference Approach has been further improved in consistency and in agreement with the Swiss Federal Office of Energy. The differences between the Reference Approach and numbers provided by the IEA statistics (IEA 2012) are discussed in Annex 4.

3.2.2 International Bunker Fuels

3.2.2.1 Source Category Description

By definition, GHG emissions from the use of International Bunker Fuels are not a key category (IPCC 2000).

For Switzerland, the sources of international bunker emissions are primarily aviation, but there are also some marine bunkers occurring from the use of diesel oil for navigation activities on the river Rhine between Basel (Switzerland) and Rotterdam (Netherlands) and on the two Swiss lakes with neighbouring countries (Lake of Geneva, Lake of Constance).

Table 3-4 Specification of Swiss source category International Bunkers.

| International Bunker Fuels | Specificaitons | Data Source |
|----------------------------|--|--|
| Civil Aviation | Country specific model (Tier 3a) | FOCA 2006-2013 |
| Marine Bunkers | Navigation on the Rhine river north of Basel (Tier 1). Naviagation on foreign territory on the Lake of Geneva and Lake of Constance (Bodensee). | CARBURA 2010, Schweiz. Bodenseeschiffahrt (SBS), Schiffahrt Untersee und Rhein (Urh), Compagne Générale de Navigation sur le lac Léman (CGN): INFRAS 2011a |

3.2.2.2 Methodological Issues

Civil Aviation

The emissions from civil aviation (domestic and international) are calculated with a Tier 3a method. The Tier 3a method follows standard modelling procedures on the level of single aircraft movements based on detailed movement statistics. The emission factors are country specific with the exception of N₂O, for which a IPCC default value is applied. The activity data of the bunker is summarised in Table 3-6 (see also Table 3-31). Further information on the methodology used for international aviation is provided in chapter 3.2.8.2 a).

Due to the detailed information about activity data available, the resulting fuel consumption is considered complete. In spite of this, there remain small differences between the fuel consumption modelled bottom-up and the total fuel sold (SFOE 2013). In 1990, the modelled consumption adds up to 1.01 million tons, whereas 1.05 million tons were sold. The difference of 4% is considered acceptable, because discrepancies up to 10% can easily result from fuelling strategies of airlines (FOCA investigation showed that airlines are calculating whether it is economically beneficial to refuel at a place with lower fuel prize). In order to match the bottom up calculation with the fuel quantity sold, any occurring difference is attributed to international bunker emissions. The factor between calculated international fuel consumption and adjusted international fuel consumption is used to scale the bunker emissions linearly. For instance in 1990, the bunker fuel consumption and the emissions had to be expanded by the factor 1.045. For 2006, they had to be reduced by the factor 0.974 (FOCA 2007). For 2012, the correction factor was 0.969 (FOCA 2013). For the more recent years, the modelled and actual total fuel sales are listed in Table 3-5.

Table 3-5 Comparison between modelled and actual fuel sales in bunker fuel consumption for civil aviation.

| Modelled and actual fuel sales | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|-----------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | t | | | | | | | |
| Modelled domestic fuel sales | 38'754 | 38'550 | 43'968 | 37'627 | 39'626 | 39'252 | 42'047 | 43'414 |
| Modelled international fuel sales | 1'152'614 | 1'196'731 | 1'287'062 | 1'391'656 | 1'345'919 | 1'395'428 | 1'511'279 | 1'527'522 |
| Total modelled fuel sales (FOCA) | 1'191'368 | 1'235'281 | 1'331'030 | 1'429'283 | 1'385'545 | 1'434'680 | 1'553'326 | 1'570'936 |
| Actual fuel sales (GEST) | 1'148'131 | 1'203'868 | 1'289'152 | 1'382'835 | 1'324'224 | 1'390'824 | 1'531'805 | 1'523'116 |
| Difference between FOCA and GEST | 3.8% | 2.6% | 3.2% | 3.4% | 4.6% | 3.2% | 1.4% | 3.1% |
| Correction factor | 0.962 | 0.974 | 0.968 | 0.966 | 0.954 | 0.969 | 0.986 | 0.969 |

Marine Bunkers

Emissions from marine bunkers are calculated with a Tier 1 method. The emission factors are country specific and in accordance with Table 3-9. Since marine bunkers consist of

diesel only a Swiss standard NCV of 42.8 TJ/Gg is used. Activity data of these bunkers is summarised in Table 3-6.

Since there is an exemption from the existing fuel taxation, activity data on marine river bunkers on the Rhine are well documented by the customs administration (Schiffahrt Untersee und Rhein Urh) as well as by CARBURA, the Swiss organisation for the compulsory stockpiling of oil products (CARBURA 2010). From the latter, coherent data series are used.

Activity data for the marine lake bunkers are not very well documented for the whole time series. Data from 1995 on have been provided by the three concerned companies (Schweizerische Bodenseeschiffahrt SBS, INFRAS 2011a). For older data proxies, such as passenger data on a national basis had to be consulted. As marine lake bunkers provide only a minor share of the total marine bunkers this approach seems to be justifiable.

Table 3-6 International bunker fuels. Consumption of kerosene and diesel oil in TJ. (Note that Liechtenstein's kerosene consumption is subtracted, see Chapter 3.2.5.)

| International Bunker Fuels | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Fuel consumption in TJ | | | | | | | | | | |
| Total international aviation(1A3ai) | 41'884 | 40'872 | 43'499 | 45'342 | 46'840 | 49'918 | 51'975 | 53'983 | 56'599 | 60'805 |
| Total marine bunkers (1A3di) | 812 | 750 | 765 | 763 | 826 | 755 | 671 | 666 | 544 | 559 |
| Total | 42'696 | 41'623 | 44'265 | 46'105 | 47'666 | 50'673 | 52'646 | 54'649 | 57'142 | 61'365 |
| 1990 = 100% | 100% | 97% | 104% | 108% | 112% | 119% | 123% | 128% | 134% | 144% |

| International Bunker Fuels | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Fuel consumption in TJ | | | | | | | | | | |
| Total international aviation(1A3ai) | 63'687 | 60'097 | 55'468 | 49'763 | 46'896 | 47'671 | 50'109 | 53'543 | 57'844 | 55'238 |
| Total marine bunkers | 525 | 426 | 350 | 422 | 426 | 498 | 460 | 474 | 456 | 432 |
| Total | 64'211 | 60'523 | 55'818 | 50'185 | 47'322 | 48'169 | 50'569 | 54'017 | 58'300 | 55'670 |
| 1990 = 100% | 150% | 142% | 131% | 118% | 111% | 113% | 118% | 127% | 137% | 130% |

| International Bunker Fuels | 2010 | 2011 | 2012 |
|-------------------------------------|--------|--------|--------|
| Fuel consumption in TJ | | | |
| Total international aviation(1A3ai) | 58'118 | 64'060 | 63'627 |
| Total marine bunkers | 468 | 418 | 376 |
| Total | 58'586 | 64'477 | 64'003 |
| 1990 = 100% | 137% | 151% | 150% |

3.2.2.3 Uncertainties and Time-Series Consistency

Civil Aviation

See remarks in Chapter 3.2.8.3, sections Civil Aviation (1A3a).

Marine Bunkers

A comparison with the data by the customs administration over a period of 13 years reveals very high correlation. Therefore, data on marine bunkers is considered to be consistent.

3.2.2.4 Source-Specific QA/QC and Verification

No source specific QA/QC activities are implemented. The International Bunker Fuels are proofed by the general QA/QC proceeding for all source categories (see below):

The entire time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables,
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013,
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013.

3.2.2.5 Source-Specific Recalculations

- 1A3ai: kerosene consumption (AD) for international flights (bunker fuels) has been recalculated in the Swiss energy statistics (SFOE 2012) for the year 2011. Therefore, emissions from all gases have been recalculated for 2011 as well.
- 1A3dii: activity data of the international marine bunkers have been updated for 2011.
- 1A3dii: CH₄ and N₂O emission factors have been recalculated for the year 2011. They are now based on an interpolation of the emission factors between 2010 and 2015 .
- 1A3di: recalculations due to new activity data in 2004, 2008 and 2011 have been made which also affects the "tank tourism" of diesel fuel in 1A3b.

3.2.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. To accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.2.3 Feedstocks and Non-Energy Use of Fuels

The Swiss Overall Energy Statistics (SFOE 2013) reports feedstocks and so-called non-energy fuel use on an aggregated level only and does not provide a detailed breakdown into specific petroleum products. Some breakdown is provided by the petroleum balance of the annual report of the Swiss Petroleum Association (EV 2013).

Feedstocks and non-energy use of fuels is reported in CRF-table 1.A(d) and separated in the following fuel types:

- Naphtha and liquefied petroleum gas: They are used in a single Swiss plant as feedstocks in the thermal cracking process for the production of ammonia and ethylene (see source categories 2B1 and 2B5). Since data for naphtha and liquefied petroleum gas use are confidential they are included in fuel type Other in CRF-table 1.A(d). For reviewers there is an additional version of this subchapter available including all confidential data and information.
- Lubricants: Primary use of lubricants is considered non-emissive. However, used oil is collected and serves as alternative fuel in the cement industry (1A2f, Other fuels, waste oil) and in special waste incineration plants. NMVOC emissions of lubricants are reported in source category 3D5 Other.
- Bitumen: this is the most important petroleum product which is used as feedstock in Switzerland. It is mainly used for road paving with asphalt and to a lower extent in asphalt roofing (see source categories 2A5 and 2A6).
- Other: Additionally to the above mentioned feedstock use of naphtha and liquefied petroleum gas this fuel category comprises all the rest of unspecified petroleum products which are used as non-energy fuels including gasoline/petrol (andere Benzine), kerosene (andere Petrole), paraffines, waxes and white spirit.

A re-assessment of the disaggregation of feedstocks is envisaged in the course of the implementation of the new reporting guidelines for the 2015 submission.

3.2.4 CO₂ Capture from Flue Gases and Subsequent CO₂ Storage

(Not applicable for Switzerland.)

3.2.5 Country-specific issues

In the following chapter, the general country specific approach of determining activity data and emission factors is presented. Specific information about each source category is included in the respective chapters 3.2.6 to 3.2.10.

3.2.5.1 Activity Data

The energy related activity data in the inventory corresponds to the energy balance provided in the Swiss overall energy statistics (SFOE 2013). It is updated annually and contains all relevant information about primary and final energy consumption. This includes annual aggregated consumption data for various fuels and main consumers such as households, transport, energy industries, and industry and services.

The aggregated data on fuel consumption in the Swiss Overall Energy Statistics is derived from the following sources:

- "Carbura" and Swiss Petroleum Association for data on import, export, sales, stocks of oil products and for processing of crude oil in refineries (EV 2013)
- Annual import data for natural gas from Swiss gas industry association
- Annual import data for coal from the customs administration
- Data provided by industry associations

Table 3-7 shows the energy balance in Switzerland in 2012. Energy flow charts for 2012 and for the base year 1990 are given in Annex 3.1.1.

A time series of the final energy consumption is depicted in Figure 3-6. The total consumption has increased by 10.5% in the period 1990-2012. Simultaneously significant substitutions occurred: heating fuel consumption decreased by 31.1%, natural gas and transport fuel consumption increased by 79.6% and 18.2%, respectively, and electricity by 26.6%.

Table 3-7 Energy balance for Switzerland 2012 (SFOE 2013) in TJ⁶.

| | Holzenergie | Kohle | Müll und Industrieabfälle | Rohöl | Erdölprodukte | Gas | Wasserkraft | Kernbrennstoffe | Übrige erneuerbare Energien | Elektrizität | Fernwärme | Total |
|--|-----------------|---------|---------------------------|--------------|---------------------|---------|---------------------|-------------------------|-------------------------------|--------------|--------------------|-----------|
| | Energie du bois | Charbon | Ord. mén. et déchets ind. | Pétrole brut | Produits pétroliers | Gaz | Energie hydraulique | Combustibles nucléaires | Autres énergies renouvelables | Electricité | Chaleur à distance | Total |
| | (1) | (2) | (3) | (4) | (5) | (6) | (7) | (8) | (9) | (10) | (11) | (12) |
| Inlandproduktion | (a) | | | | | | | | | | | |
| + Import | 39 200 | – | 56 320 | – | – | – | 143 660 | – | 19 530 | – | – | 258 710 |
| + Export | 1 180 | 5 630 | – | 147 260 | 351 500 | 122 520 | – | 265 580 | 250 | 312 570 | – | 1 206 490 |
| + Lagerveränderung ¹ | – 300 | 0 | – | – | – 13 190 | – | – | – | – | – 320 490 | – | – 333 980 |
| | (d) | – 160 | – | – 90 | 18 060 | – | – | – | – | – | – | 17 810 |
| = Bruttoverbrauch | (e) | 5 470 | 56 320 | 147 170 | 356 370 | 122 520 | 143 660 | 265 580 | 19 780 | – 7 920 | 0 | 1 149 030 |
| + Energieumwandlung: | | | | | | | | | | | | |
| • Wasserkraftwerke | – | – | – | – | – | – | – 143 660 | – | – | 143 660 | – | 0 |
| • Kernkraftwerke | – | – | – | – | – | – | – | – 265 580 | – | 87 640 | 1 370 | – 176 570 |
| • konventionell-thermische Kraft-, Fernheiz- und Fernheizkraftwerke | – 1 720 | – | – 46 060 | – | – 800 | – 8 030 | – | – | – | 10 250 | 17 000 | – 29 360 |
| • Gaswerke | – | – | – | – | – | – | – | – | – | – | – | 0 |
| • Raffinerien | – | – | – | – 147 170 | 145 870 | – | – | – | – | – | – | – 1 300 |
| • Diverse Erneuerbare | – 1 320 | – | – | – | – | 320 | – | – | – 3 460 | 3 320 | 0 | – 1 140 |
| + Eigenverbrauch des Energiesektors, Netzverluste, Verbrauch der Speicherungen | | | | | | | | | | | | |
| (l) | – | – | – | – | – 11 330 | – 860 | – | – | – | – 24 650 | – 1 490 | – 38 330 |
| + Nichtenergetischer Verbrauch | | | | | | | | | | | | |
| (m) | – | – | – | – | – 20 050 | – | – | – | – | – | – | – 20 050 |
| = Endverbrauch | (n) | 5 470 | 10 260 | 0 | 470 060 | 113 950 | 0 | 0 | 16 320 | 212 300 | 16 880 | 882 280 |
| Haushalte | 19 340 | 400 | – | – | 100 040 | 47 230 | – | – | 11 300 | 66 000 | 6 480 | 250 790 |
| Industrie | 10 120 | 5 070 | 10 260 | – | 27 790 | 35 630 | – | – | 1 370 | 68 500 | 6 520 | 165 260 |
| Dienstleistungen | 6 830 | – | – | – | 40 690 | 23 060 | – | – | 2 940 | 63 110 | 3 880 | 140 510 |
| Verkehr | – | – | – | – | 299 420 | 1 490 | – | – | 520 | 11 140 | – | 312 570 |
| Statistische Differenz inkl. Landwirtschaft | 750 | 0 | 0 | – | 2 120 | 6 540 | – | – | 190 | 3 550 | 0 | 13 150 |
| (s) | | | | | | | | | | | | |

¹ + Lagerabnahme
– Lagerzunahme

¹ + diminution de stock
– augmentation de stock

⁶ Note that Liechtenstein's consumption of liquid fuel is included in the numbers (see chapter below on Final Swiss energy consumption). Numbers of gas consumption are from Switzerland only.




Figure 3-6 Final energy consumption in Switzerland between 1990 and 2012 by fuel type (SFOE 2013). Note that Liechtenstein's consumption of fuel is included in the numbers (see chapter below on Final Swiss energy consumption). It corresponds to 0.62% and 0.63% respectively of the Swiss fuel consumption.

Final Swiss energy consumption

The fundamental data on final energy consumption is provided by the Swiss overall energy statistics (SFOE 2013). However, since Switzerland and Liechtenstein form a customs and monetary union governed by a customs treaty, data regarding liquid fuels in the Swiss overall energy statistics also cover liquid fuel consumption in Liechtenstein. In order to calculate the correct Swiss fuel consumption, Liechtenstein's energy consumption (see Table 3-4 in Liechtenstein's NIR (OE 2014)) is subtracted from the figures provided by the Swiss overall energy statistics. In 2012, the sum of fossil fuels used in Liechtenstein was 2'780 TJ, corresponding to approximately 0.6% of the Swiss consumption of that year.

Disaggregation of the energy consumption

For the elaboration of the greenhouse gas inventory, information about energy consumption is needed at a much more detailed level than provided by the Swiss overall energy statistics. While the total amount of fuel consumption is given by the Swiss overall energy statistics, additional information sources are used to disaggregate fuel consumption into the source categories defined in the CRF-tables. For the different source categories the following sources are used:

- 1A1 Energy Industry: the fuel consumption for source category 1A1 is provided by data from the Swiss overall energy statistics (SFOE 2013).
- 1A2 Manufacturing Industries and Construction: For the industry sector, data are provided by the following sources (see also 3.2.7):
1A2a – 1A2fi Stationary:




Figure 3-7 Schematic presentation of the sources used for the industrial sector 1A2a – fi. The overall fuel consumption of 1A2 and 1A4a corresponds to the fuel consumption reported in the Swiss overall energy statistics. The industry model (Prognos) distributes the fuel consumption into the different source categories. The coloured boxes show the specific industry information that is used in the different source categories. The fuel consumption for each source category is determined by the industry model (Prognos).

The Swiss overall energy statistics does not provide a specific category for industry fuel consumption, but only an aggregated level (SFOE 2013). Additionally, in 1999, a new classification of the economic sectors led to a change of the allocation of fuel consumption to industry and commercial sector together within the energy statistics as shown in the figure above. In order to provide a consistent time series for industrial (and commercial) fuel consumption, a model is used to split the consumption into the various source categories.

In order to separate these two categories (industrial and commercial), and to achieve the required disaggregation within the industry sector 1A2, a model that determines the fuel use in the different source categories of 1A2 is used (Industry model, Prognos 2013).

The model is based on 164 individual industrial processes and further 64 processes related to infrastructure in industry. Fuel consumption of a specific process is calculated as the product of the process activity data and the process specific fuel consumption factor. For example within the chocolate industry, the activity data would be tonnes of chocolate produced and the specific fuel consumption factor would be kWh natural gas per tonne of chocolate produced.

The model is adjusted and scaled to fit available energy data and statistics, including the following data sources:

- Swiss overall energy statistics (SFOE 2013)
- Statistic of energy consumption from industry and commercial sector, Data from soundings of Helbling Ltd. (since 1999) (SFOE 2013d)
- Swiss Statistics on Renewable Energy (SFOE 2013a)

- Statistics on combined heat and power generation in Switzerland (SFOE 2013c)
- Consumption data of the Federation of the Swiss Pulp, paper and board industry (ZPK 2013)
- Data from Cemsuisse for 1990 and 2000 to 2010 (Cemsuisse 2013)
- Fuel supply data from CARBURA for 1985 to 2010 (Carbura 2010)
- Data on full-time-jobs and on industrial production from SFSO (2013a)
- Expert estimates and industry data based on surveys and annual reports from industry associations (Prognos 2013)

The model provides energy consumption for each source category 1A2a to 1A2f and each fuel type. Total energy consumption in industry according to the model is then subtracted from the energy consumption of industry and services according to the Swiss overall energy statistics. The resulting difference in energy consumption is allocated to source category 1A4a Commercial and Institutional, ensuring completeness of reporting and consistency with the final energy consumption according to the Swiss overall energy statistics.

The fuel consumption determined by the model is considered as fuel used in boilers of each source category of 1A2.

For specific industries, plant-level information is available and considered in source categories 1A2a, 1A2b, 1A2c, 1A2d and 1A2fi, see chapter 3.2.7 (coloured boxes in Figure 3-7).

Additional detailed statistics are available for wood consumption from the Swiss Wood Energy Statistics (SFOE 2013b, see description in chapter 3.2.5.1), sewage and biogas from the Swiss Statistics on Renewable Energy (SFOE 2013a) and the Statistics on combined heat and power generation in Switzerland (SFOE 2013c)). Emissions from these sources are summarized under 1A2fi due to insufficient information regarding sectoral disaggregation.

These specific informations from industries and additional data sources constitute a subset of the fuel consumption allocated by the Prognos model. The fuel consumption of each source category is thus not changed and the sum of source category 1A2 and 1A4a corresponds to the fuel consumption as per the Swiss overall energy statistics of Switzerland (SFOE 2013).

The use of two different information sources from industry and the Prognos model result in a statistical stock change for natural gas, gas oil, residual fuel oil and liquefied petroleum gas in the Energy model. A statistical stock change is necessary to account for the difference between the total fuel consumption as per the model of Prognos based on the Swiss overall energy statistics and the integrated results from the industry data within the Energy model. If for example the fuel consumption of natural gas reported by industry is higher than the natural gas consumption provided by the model of Prognos, this difference is compensated through stock changes to reach the fuel consumption as per the Swiss overall energy statistics.

1A2fii Mobile: Emissions of off-road mobile machinery are modelled using a (territorial) emission model developed by INFRAS (2008). The emissions of all off-road categories like construction machines, railways, navigation etc (1A2fii, 1A3c, 1A3d, 1A4a, 1A4b, 1A4c, 1A4d, 1A4e, 1A4f, 1A4g, 1A4h, 1A4i, 1A4j, 1A4k, 1A4l, 1A4m, 1A4n, 1A4o, 1A4p, 1A4q, 1A4r, 1A4s, 1A4t, 1A4u, 1A4v, 1A4w, 1A4x, 1A4y, 1A4z, 1A5) are modelled by the same approach using a Tier 2 method. Some details of the emission modelling that hold for all off-road families are described in Annex A3.1.6. During the complete revision of the emissions of the off-road sector that took place between 2005 and 2008, activity data and emission factors were updated and a new database structured in analogy to the on-road database (INFRAS 2010) was developed for the emission calculation. Emissions are calculated in five-year intervals for 1990 up to 2030. For the years in-between, the emissions are interpolated linearly. A slight modification of the activity data was carried out in 2013 based on the latest numbers on growth of population and economy (Prognos 2012a).

- 1A3 Transport: For the transport sector, INFRAS developed an emission model for territorial road transportation (1A3b, INFRAS 2010; for details refer to Annex A3.1.5) and off-road transport (1A3c and 1A3d, INFRAS 2008, see paragraph on 1A2fii above). The Swiss overall energy statistics provides information on the amounts of fuel sold. From the amounts sold, the consumptions modelled by the territorial road and off-road models (INFRAS 2008, 2010) are subtracted. The differences to the amount of fuels sold represent tank tourism, i.e. the amount sold in Switzerland but consumed abroad.
- 1A4: Activity data of the Other Sectors is provided by the following sources:
 1A4ai, 1A4bi Other Sectors, Stationary: The Swiss overall energy statistics does provide information on the consumption by industry and the commercial sector together. The information provided by the model of Prognos determines the fuel consumption in source category 1A2 Industry. The difference between the fuel consumption in the Swiss overall energy statistics and the fuel consumption by Prognos determines thus the fuel consumption of source category 1A4ai. Source category 1A4bi is provided by the Swiss overall energy statistics. The fuel used for co-generation in turbines and engines in the commercial sector and households is deducted from a model of stationary engines developed by Eicher + Pauli (Kaufmann 2013). This model builds one information source for the Statistics on combined heat and power generation in Switzerland (SFOE 2013c). This amount are part of the fuel consumptions of the Swiss overall energy statistics.
 For 1A4bi Other Sectors, grass drying (1A4ci), specific bottom-up industry information is available and is deducted from the fuel consumption of 1A2. The information on grass drying is documented in the respective EMIS comment.
 1A4aii/bii/cii 1A4bi Other Sectors, Mobile: In addition to energy consumption in stationary installations, also mobile sources are reported in source category 1A4aii/bii/cii, the emissions of which are calculated using the INFRAS off-road model (INFRAS 2008, see paragraph on 1A2fii above).
- 1A5 Military: In Switzerland military energy consumption concerns only the two source categories Mobile Military off-road and Military Aviation. For the Mobile Military off-road energy consumption the INFRAS off-road model is used (INFRAS 2008, see paragraph on 1A2fii above). The energy consumption of Military aviation is copied from the logbooks of the military aircrafts (VTG 2013).

The compilation of the different information and the resulting disaggregation is made in the so-called Energy model developed and annually updated by FOEN.

The following figures show the information and data sources used for the disaggregation of each fuel type in each source category in the Energy model.

Natural Gas




Figure 3-8 Schematic disaggregation of 1A Fuel Consumption for natural gas.

In addition to the information sources used as described above, the following specific informations are used for the disaggregation of the natural gas consumption:

- 1A1: The fuel consumption for Energy conversion from the Swiss overall energy statistics is corrected for losses in gas transportation as provided by the Swiss Gas and Water Industry Association. These fugitive emissions are reported under category 1B2. In the CRF-tables, the resulting fuel consumption under 1A1 for natural gas is reported under 1A1a Public Electricity and Heat Production.
- 1A3 Transport: Within the sector 1A3, natural gas consumption only occurs in source category 1A3b and 1A3e. As with other fuels, the consumption modelled by the territorial models is subtracted from the amount sold based on the energy statistics. The difference to the amount of fuel sold represents the tank tourism of natural gas, which is reported under 1A3bviii. Further information on the transport sector is provided in chapter 3.2.8.

Gas Oil




Figure 3-9 Schematic disaggregation of 1A Fuel Consumption for gas oil.

In addition to the information sources used as described above, the following specific informations are used for the disaggregation of the gas oil consumption:

- 1A1: Energy conversion and district heating is available within the Swiss overall energy statistics. As district heating is a sub category of energy conversion in the statistics, the respective fuel consumption is deducted from energy conversion to document boiler use in electricity production and in district heating separately within the activity data of source category 1A1a Public Electricity and Heat Production.
- 1A3: A small amount of gas oil is consumed by navigation 1A3d. It is subtracted from the total provided by the Swiss overall energy statistics before the rest is attributed to 1A4ai.
- 1A4ai: Within the commercial sector, the off-road gas oil consumption for 1A3d is deducted from the consumption of the commercial sector. As mentioned above, gas oil consumption of Liechtenstein is included in the total consumption of gas oil in the energy statistics, therefore, the amount used in Liechtenstein is subtracted from 1A4ai and 1A4bi.

Residual Fuel Oil




Figure 3-10 Schematic disaggregation of 1A Fuel Consumption for residual fuel oil.

In addition to the information sources used as described above, the following specific informations are used for the disaggregation of the residual fuel oil consumption:

- 1A1: Informations from the Swiss Petroleum Association (EV) is provided in addition to the information available from the Swiss overall energy statistics regarding district heating and energy conversion. The information of EV is used within source category 1A1a Public Electricity and Heat Production for the fuel use within boilers for electricity production (for the single fossil fuel power station that was operational from 1985 to 1994) and within 1A1b Petroleum refining for the use of fuel within the boilers of the refineries. As discussed under the section gas oil, residual fuel oil consumption from the Swiss overall energy statistics are reported for boiler use in electricity production and in district heating separately within the activity data of source category 1A1a Public Electricity and Heat Production.

Liquefied Petroleum Gas (LPG)




Figure 3-11 Schematic disaggregation of 1A Fuel Consumption for liquefied petroleum gas (LPG).

In addition to the information sources used as described above, the following specific informations are used for the disaggregation of the liquefied petroleum consumption:

- 1A1: Information on the refinery boilers is provided from the Swiss Petroleum Association (EV) and considered within source category 1A1b Petroleum refining. However, the characteristics of refinery liquefied petroleum gas is not identical to the characteristics of liquefied petroleum gas used in source category 1A2 and is therefore not deducted in source category 1A2.

Petroleum coke, Bituminous Coal and Lignite

For Petroleum coke, bituminous coal and lignite, the same approach as above described is used including the data from the Swiss overall energy statistics (SFOE 2013) and the Prognos model for the fuels used within the industry sector (Prognos 2013).

Wood




Figure 3-12 Schematic disaggregation of 1A Fuel Consumption for wood.

The Swiss wood energy statistics (SFOE 2013b) provides both the annual wood consumption for specified categories of combustion installations (Table K, categories 1-19) and the allocations of the combustion categories to the sectoral consumer categories (Table N, household, agriculture/forestry, industry, services, electricity and district heating). This allows to assign the annual wood consumption on the level of combustion installation categories, see Table 3-8, to the source categories 1A1a Public Electricity and Heat Production, 1A2fi Other, 1A4ai Commercial/Institutional, 1A4bi Residential and 1A4ci Agriculture/Forestry/Fisheries.

For some industries in source category 1A2fi, specific bottom-up information is available and included in the Energy model. Regarding wood consumption, the specific industry data is subtracted from the activity data of the respective combustion installation category in order to avoid double counting within source category 1A2fi. The information on the specific processes are documented in the respective EMIS comments (EMIS 2014/1A Holzfeuerungen).

Table 3-8 Categories of wood combustion installations based on SFOE 2013b.

| 1A Wood combustion, categories |
|---|
| Open fireplaces |
| Closed fireplaces, log wood stoves |
| Pellet stoves |
| Log wood hearths |
| Log wood boilers |
| Log wood dual chamber boilers |
| Automatic chip boilers < 50 kW |
| Automatic pellet boilers < 50 kW |
| Automatic chip boilers 50-500 kW w/o wood processing companies |
| Automatic pellet boilers 50-500 kW |
| Automatic chip boilers 50-500 kW within wood processing companies |
| Automatic chip boilers > 500 kW w/o wood processing companies |
| Automatic pellet boilers > 500 kW |
| Automatic chip boilers > 500 kW within wood processing companies |
| Combined chip heat and power plants |
| Plants for renewable waste from wood products |

Gasoline and Diesel Oil




Figure 3-13 Schematic disaggregation of 1A Fuel Consumption for gasoline and diesel oil.

Gasoline and diesel oil consumption is reported in several source categories. In addition to the information provided above, the following sources are used for the disaggregation of gasoline and diesel oil consumption:

- As with other fuels, the Swiss overall energy statistics provides information the amounts sold; from these the consumptions of gasoline and diesel oil modelled by the territorial models (INFRAS 2008 and 2010, see above) are subtracted. The differences to the amount of fuels sold represent the tank tourism of gasoline and diesel oil which are reported under 1A3bviii.
- The customs statistics allow to quantify the amount of marine bunker fuels of diesel oil consumed by navigation on the river Rhine between Basel (Switzerland) and Rotterdam (Netherlands) and on two lakes with international navigation. See Chapter 3.2.2 for further details.

Further information on the transport sector is provided in chapter 3.2.8.

3.2.5.2 Emission Factors

The different sources categories within source category 1A Fuel Combustion are characterised by rather similar industrial combustion processes and thus the same emission factors are applied throughout 1A for the main fuels. Emission factors for fuels that are only used in one particular source category are described in the context of that particular source category.

CO₂ Emission Factors

Table 3-9 CO₂ Emission Factors and NCV from 1990 to 2012.

| CO ₂ Emission Factors 1990 - 2012 | | | | |
|--|------------------------|------------|------|------------------------------|
| Fuel | t CO ₂ / TJ | NCV [GJ/t] | CS/D | Data Sources |
| Gasoline | 73.9 | 42.5 | CS | EMPA (1999) |
| Jet Kerosene | 73.2 | 43.0 | CS | EMPA (1999) |
| Diesel Oil | 73.6 | 42.8 | CS | EMPA (1999) |
| Gas Oil | 73.7 | 42.6 | CS | EMPA (1999) |
| Residual Fuel Oil | 77.0 | 41.2 | CS | EMPA (1999) |
| P-Coke | 91.4 | 31.8 | CS | FOEN (2011k) |
| Liquefied Petroleum Gas | 65.5 | 46.0 | CS | FOEN (2011k) |
| Natural Gas | 56.1 | 46.5 | D | IPCC Guidelines 2006 |
| Bituminous Coal | 92.7 | 25.5 | CS | FOEN (2011k) |
| Lignite | 96.1 | 23.6 | CS | FOEN (2011k) |
| Biofuel | t CO ₂ / TJ | NCV [GJ/t] | CS/D | Data Sources |
| Biodiesel | 73.6 | 42.8 | CS | assumed equal to diesel oil |
| Bioethanol | 73.9 | 42.5 | CS | assumed equal to gasoline |
| Biogas | 56.1 | 46.5 | D | assumed equal to natural gas |
| Wood | 92.0 | - | CS | SAEFL (2000) |

CO₂ emission factors and NCV values for gasoline, jet kerosene, diesel oil, gas oil and residual fuel oil are country specific and are based on measurement campaigns of NCV and carbon content of fuels (EMPA 1999, Intertek 2008, Intertek 2012). The values from the 1998 study of EMPA are used for the entire period since 1990. According to expert judgment from the Swiss Petroleum Association, the natural variability of the products is much larger than the measurement uncertainty provided in the studies. Accordingly adjusting the emission factors every couple of years seems inappropriate given the variability between the samples. Therefore, only spot checks are made periodically to verify that the values of the 1998 study are still applicable, without changing the values if considered compatible with the 1998 study. The latest measurements in 2011 revealed a few deviations from the 1998 values that seemed to exceed the expected range. However, based on the small sample number (10 samples) and the large variations observed, the changes were not statistically significant. The results triggered the launch of a new measurement campaign that was set up over the past year. This on-going measurement campaign will be based on a representative sample which covers summer and winter samples from the main import streams. The sampling started in July 2013 and will carry on for six months. Fortnightly samples are going to be taken from nine different sites (large-scale storage facilities and the two Swiss refineries). Preliminary information confirms that there is no change in the CO₂ emission factors. Final results are expected in summer 2014 and will be available for the 2015 submission. After completion of the entire campaign, the use of NCV and CO₂-EF will be re-assessed (both for the greenhouse gas inventory and the energy statistics of the SFOE).

For liquefied petroleum gas, the values are country specific and based on the CRC Handbook of Chemistry and Physics (see documentation in FOEN 2011k).

For natural gas, the default values of the 2006 IPCC guidelines are used (IPCC 2006).

CO₂ emission factors and NCV values of petroleum coke, bituminous coal and lignite are country specific and based on samples that were taken from Switzerland's cement plants. Cement plants are the largest consumer of solid fossil fuels in Switzerland. The samples from the individual plants were compiled over nine months and have been analysed for calorific value and carbon content by an independent analytical laboratory. The original data is compiled in an internal document from cemsuisse. The results from the individual plants were weighted according to the relative contribution of each plant. The CO₂ emission factors are lower than the IPCC default values (IPCC 2006), however, they all lie within the range provided by the IPCC (see documentation in FOEN 2011k).

Regarding the small amount of biofuels used in Switzerland, the CO₂ emission factors and NCV values are taken from the respective fossil fuels that are being substituted by the respective biofuel. Therefore the values for biodiesel, bioethanol and biogas are assumed to be equal to the ones of diesel oil, gasoline and natural gas, respectively.

CO₂ emission factor for wood combustion is country specific and provided from the handbook of emission factors for stationary sources (SAEFL 2000) as documented in the respective EMIS comment (EMIS 2014/1A Holzfeuerungen). Since the NCV of wood depends on the wood product used as fuel (log wood, wood chips, pellets) it is not displayed in the table above but is reported in the Swiss wood energy statistics (SFOE 2013b).

For off-road activities the emission factors for CO₂ are country specific and assumed to be constant in the period 1990-2012 with values 73.6 t/TJ for diesel oil, 73.9 t/TJ for gasoline and 56.1 t/TJ for CNG (equal to natural gas). See also Table 3-9.

Note that specific emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database INFRAS (2008)⁷.

Please note that the CO₂ emission factors and NCV values are constant over the whole period 1990-2012.

Uncertainty in CO₂ emission factors in fuel combustion (1A)

Liquid fuels:

Total uncertainty of net calorific values (NCVs) for liquid fuels is taken as a proxy for the uncertainty of the CO₂ emission factor of liquid fuels. Net calorific values are based on the determination of the gross calorific value and the calculation of the net calorific value by EMPA. To this aim, a set of fuel samples of different sources has been selected that is representative for the fuels traded in Switzerland in the year 1998. Assuming that this data on the uncertainty of the net calorific value is representative for the uncertainty of the emission factors in fuel combustion, a combined uncertainty of 0.51% (defined as two standard deviations, STD) results for the emission factor.

⁷ <http://www.bafu.admin.ch/luft/00596/06906/offroad-daten/index.html?lang=en> [24.01.2012]

Table 3-10 Results from the analysis of the net calorific values of liquid fuels in Switzerland (EMPA 1999).

| A | B | C | D | E | F | G |
|------------------|----------------------------------|---------------|------------|--------------------|------------------------|----------------|
| Fuel | Net calorific value liquid fuels | | | | | |
| | Mean [GJ/t] | STD [GJ/t] | STD [%] | Uncertainty [%] | =(C*G)^2 [GJ^2/t^2] | No. of samples |
| Heavy fuel oil | 41.2 | 0.85 | 2.06 | 4.13 | 0.000010 | 6 |
| Light fuel oil | 42.6 | 0.13 | 0.31 | 0.61 | 0.002891 | 10 |
| Diesel | 42.8 | 0.10 | 0.23 | 0.47 | 0.000707 | 10 |
| Gasoline | 42.5 | 0.29 | 0.68 | 1.36 | 0.007966 | 30 |
| Jet kerosene | 43.0 | 0.25 | 0.58 | 1.16 | 0.000004 | 10 |
| Sum | 42.6 | | | | 0.011579 | 66 |
| Combined STD/Unc | | 0.108 | 0.25 | 0.51 | | |

Gaseous fuels:

The uncertainty of the emission factor for CO₂ has been derived from data on measurements of the NCVs of natural gas in the grid. SGWA (2007) provides a range of -2.3% and +2.3%. Interpreting 2.3% as one standard deviation, an uncertainty of 4.6% results (i.e. two standard deviations).

Solid fuels:

For the uncertainty of the emission factor for CO₂, the IPCC Good Practice Guidance default value of 5% for countries with well-developed energy data systems is used (IPCC 2000: p. 2.15).

Other fuels (waste to energy):

The dominant factor influencing the uncertainty of CO₂ emissions from municipal solid waste incineration (1A1) is the fraction of fossil carbon in the waste. For the fraction of C in incinerated waste an uncertainty of 20% has been estimated, and for the fraction of fossil C in total C an uncertainty of 10% has been estimated, resulting in a preliminary uncertainty estimate of 30% for the waste incineration CO₂ emission factor (SAEFL 2005h).

Resulting uncertainty in CO₂ emissions in fuel combustion (1A)

The table below provides the results of the quantitative Tier 1 analysis (following Good Practice Guidance; IPCC (2000): p. 6.13ff) estimating uncertainties of CO₂ emissions from fuel combustion activities.

Table 3-11 Results from Tier 1 uncertainty calculation and reporting for CO₂ emissions in 1A Fuel Combustion.

| A | B | C | D | E | F | G | H | I | J | K | L | M |
|--|-----|--------------------------|---------------------|---------------------------|-----------------------------|----------------------|--|--------------------|--------------------|--|--|---|
| IPCC Source category | Gas | Base year emissions 1990 | Year 2012 emissions | Activity data uncertainty | Emission factor uncertainty | Combined uncertainty | Combined uncertainty as % of total national emission in year t | Type A sensitivity | Type B sensitivity | Uncertainty in trend in national emissions introduced by emission factor uncertainty | Uncertainty in trend in national emissions introduced by activity data uncertainty | Uncertainty introduced into the trend in total national emissions |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Gaseous Fuels | CO2 | 289.73 | 498.74 | 2.0 | 4.6 | 5.0 | 0.050 | 0.0042 | 0.0098 | 0.02 | 0.03 | 0.03 |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Liquid Fuels | CO2 | 693.69 | 805.16 | 1.3 | 0.5 | 1.4 | 0.022 | 0.0024 | 0.0158 | 0.00 | 0.03 | 0.03 |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Solid Fuels | CO2 | 44.84 | 0.00 | 5.9 | 5.0 | 7.7 | 0.000 | -0.0009 | 0.0000 | 0.00 | 0.00 | 0.00 |
| 1A1 1. Energy A. Fuel Combustion 1. Energy Industries Other Fuels | CO2 | 1519.73 | 2714.50 | 10.0 | 30.0 | 31.6 | 1.706 | 0.0238 | 0.0533 | 0.71 | 0.75 | 1.04 |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and Cons Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 2.0 | 4.6 | 5.0 | 0.209 | 0.0203 | 0.0411 | 0.09 | 0.12 | 0.15 |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and Cons Liquid Fuels | CO2 | 3692.22 | 2640.18 | 1.3 | 0.5 | 1.4 | 0.073 | -0.0197 | 0.0518 | -0.01 | 0.09 | 0.09 |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and Cons Solid Fuels | CO2 | 1204.47 | 454.87 | 5.9 | 5.0 | 7.7 | 0.070 | -0.0144 | 0.0089 | -0.07 | 0.07 | 0.10 |
| 1A2 1. Energy A. Fuel Combustion 2. Manufacturing Industries and Cons Other Fuels | CO2 | 134.15 | 288.60 | 10.0 | 30.0 | 31.6 | 0.181 | 0.0031 | 0.0057 | 0.09 | 0.08 | 0.12 |
| 1A3a 1. Energy A. Fuel Combustion 3. Transport; Civil Aviation | CO2 | 252.55 | 136.65 | 2.2 | 1.2 | 2.5 | 0.007 | -0.0022 | 0.0027 | 0.00 | 0.01 | 0.01 |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Diesel | CO2 | 2587.68 | 6767.05 | 2.2 | 0.5 | 2.2 | 0.301 | 0.0826 | 0.1328 | 0.04 | 0.41 | 0.41 |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Gasoline | CO2 | 11335.27 | 9016.58 | 2.2 | 1.4 | 2.6 | 0.462 | -0.0426 | 0.1769 | -0.06 | 0.55 | 0.55 |
| 1A3b 1. Energy A. Fuel Combustion 3. Transport; Road Transportation Natural Gas | CO2 | 0.00 | 83.59 | 3.5 | 3.5 | 5.0 | 0.008 | 0.0016 | 0.0016 | 0.01 | 0.01 | 0.01 |
| 1A3c 1. Energy A. Fuel Combustion 3. Transport; Railways | CO2 | 28.69 | 39.99 | 1.3 | 0.5 | 1.4 | 0.001 | 0.0002 | 0.0008 | 0.00 | 0.00 | 0.00 |
| 1A3d 1. Energy A. Fuel Combustion 3. Transport; Navigation | CO2 | 111.93 | 121.14 | 2.2 | 0.5 | 2.3 | 0.005 | 0.0002 | 0.0024 | 0.00 | 0.01 | 0.01 |
| 1A3e 1. Energy A. Fuel Combustion 3. Transport; Other non-specified | CO2 | 31.42 | 45.44 | 2.2 | 4.5 | 5.0 | 0.005 | 0.0003 | 0.0009 | 0.00 | 0.00 | 0.00 |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/Instit Gaseous Fuels | CO2 | 987.24 | 1482.76 | 2.0 | 4.6 | 5.0 | 0.148 | 0.0100 | 0.0291 | 0.05 | 0.08 | 0.09 |
| 1A4a 1. Energy A. Fuel Combustion 4. Other Sectors; Commercial/Instit Liquid Fuels | CO2 | 47606.43 | 3038.51 | 1.3 | 0.5 | 1.4 | 0.084 | -0.0296 | 0.0596 | -0.02 | 0.11 | 0.11 |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 2.0 | 4.6 | 5.0 | 0.264 | 0.0244 | 0.0520 | 0.11 | 0.15 | 0.18 |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Liquid Fuels | CO2 | 10248.79 | 7374.50 | 1.3 | 0.5 | 1.4 | 0.204 | -0.0537 | 0.1447 | -0.03 | 0.26 | 0.27 |
| 1A4b 1. Energy A. Fuel Combustion 4. Other Sectors; Residential Solid Fuels | CO2 | 54.59 | 33.60 | 5.9 | 5.0 | 7.7 | 0.005 | -0.0004 | 0.0007 | 0.00 | 0.01 | 0.01 |
| 1A4c 1. Energy A. Fuel Combustion 4. Other Sectors; Agriculture/Forestry Gaseous Fuels | CO2 | 41.45 | 18.48 | 2.0 | 4.6 | 5.0 | 0.002 | -0.0004 | 0.0004 | 0.00 | 0.00 | 0.00 |
| 1A4c 1. Energy A. Fuel Combustion 4. Other Sectors; Agriculture/Forestry Liquid Fuels | CO2 | 547.34 | 540.01 | 1.3 | 0.5 | 1.4 | 0.015 | 0.0000 | 0.0106 | 0.00 | 0.02 | 0.02 |
| 1A5 1. Energy A. Fuel Combustion 5. Other Liquid Fuels | CO2 | 203.58 | 114.80 | 1.3 | 0.5 | 1.4 | 0.003 | -0.0017 | 0.0023 | 0.00 | 0.00 | 0.00 |

CH₄ Emission Factors

Table 3-12 CH₄ Emission Factors from 1990 to 2012.

| CH ₄ Emission Factors 1990 - 2012 | | | |
|--|------------------------|------|-----------------------------------|
| Fuel | g CH ₄ / GJ | CS/D | Data Sources |
| Gas Oil | 1.0 | CS | SAEFL (2000) |
| Residual Fuel Oil | 4.0 | CS | SAEFL (2000) |
| P-Coke | 10.0 | D | * |
| Liquefied Petroleum Gas | 1.0 | D | IPCC Guidelines 2006 |
| Natural Gas | 6.0 | CS | SAEFL (2000) |
| Bituminous Coal | 10.0 | D | IPCC Guidelines 2006 |
| Lignite | 10.0 | D | IPCC Guidelines 2006 |
| Biofuel | g CH ₄ / GJ | CS/D | Data Sources |
| Wood | 2.8-230 | CS | Nussbaumer, T., Boogen, N. (2010) |

**This emission factor is still the same as for Coal because in previous submissions P-Coke was not reported separately. In Submission 2015 it has to be corrected to IPCC default value.*

CH₄ emission factors for gas oil, residual fuel oil, and natural gas are country specific and provided from the handbook of emission factors for stationary sources (SAEFL 2000).

For liquefied petroleum gas, P-coke, bituminous coal and lignite default values from the 2006 IPCC guidelines are used (IPCC 2006).

The CH₄ emission factors from wood combustion are country specific and have been modelled based on FID measurements from a series of wood combustion plants at various conditions (Nussbaumer, T., Boogen, N. 2010, unpublished). They vary between 2.8 and 230 g CH₄/GJ depending on the combustion installation, rated thermal input and technology used and thus decrease over time as result of improved technology.

For road transportation, the CH₄ emission factors are country specific; they are calculated as a fraction of hydrocarbon (HC) emissions, which in turn have been established through specialised measurement programmes (INFRAS 2010). The share of CH₄ in HC emissions is differentiated by emission concept (INFRAS 2004). For mobile off-road machinery, the CH₄ emission factor is country specific as well; it is based on VTT (2004) for diesel engines and EMPA (2004) for gasoline two- and four-stroke engines.

Please note that the CH₄ emission factors and NCV values are constant over the whole period 1990-2012. Only for wood combustion and road transportation, the CH₄ emission factors are not constant; they decrease over time as a result of improved technology (e.g. improved wood furnances).

N₂O Emission Factors

Table 3-13 N₂O Emission Factors from 1990 to 2012.

| N ₂ O Emission Factors 1990 - 2012 | | | |
|---|-------------------------|------|----------------------|
| Fuel | g N ₂ O / GJ | CS/D | Data Sources |
| Gas Oil | 0.6 | CS | SAEFL (2000) |
| Residual Fuel Oil | 0.8 | CS | SAEFL (2000) |
| P-Coke | 1.6 | CS | * |
| Liquefied Petroleum Gas | 0.1 | D | IPCC Guidelines 2006 |
| Natural Gas | 0.1 | D | IPCC Guidelines 2006 |
| Bituminous Coal | 1.6 | CS | SAEFL (2000) |
| Lignite | 1.6 | CS | SAEFL (2000) |
| Biofuel | g N ₂ O / GJ | CS/D | Data Sources |
| Wood | 1.6 | CS | SAEFL (2000) |

**This emission factor is still the same as for Coal because in previous submissions P-Coke was not reported separately. In Submission 2015 it has to be corrected to IPCC default value.*

The N₂O emission factors for gas oil, residual fuel oil, P-coke, bituminous coal, lignite and wood are country specific and provided from the handbook of emission factors for stationary sources (SAEFL 2000).

Default values from the 2006 IPCC guidelines are used for liquefied petroleum gas and natural gas (IPCC 2006).

For road transportation, N₂O emission factors are taken from the COPERT IV model (Gkatzoflias et al. 2012). For mobile off-road machinery, the (country-specific) N₂O emission factors are based on SAEFL (1996).

Please note that the N₂O emission factors and NCV values are constant over the whole period 1990-2012. Only for road transportation, the N₂O emission factors are not constant; they decrease over time as a result of technical improvements such as the further development of catalyst technologies.

Oxidation Factors

For the emission calculation, an oxidation factor of 100% is assumed for all fossil fuel combustion processes. A first reason for this is that technical standards for combustion installations in Switzerland are high. A second reason is that the small fraction of originally non-oxidised carbon retained in ash, particulates or soot is likely to be oxidized later naturally due to degradation processes.

As the fuel consumption of gaseous fuels strongly increased (1990 to 2012: +78.6% from 68'597 to 122'549 TJ), overestimation of oxidation factors for gaseous fuels would lead to overestimation of emission increase and would therefore be conservative. As the consumption of liquid fuels decreased (1990 to 2012: 10.6% from 465'212 to 416'051 TJ) overestimating of oxidation factors for liquid fuels would tend to overestimate emission reduction and would therefore not be conservative. Because of the reasons mentioned above for the assumption of an oxidation factor of 100%, the possible overestimation of emission decrease is considered to be of minor importance.

For coal, IPCC 1996 provides a global average oxidation factor of 98.0%. In Switzerland, the consumption of coal plays a minor role (0.8% of total energy consumption in 2012) and decreased significantly over the considered period (1990 to 2011 by 62.8% from 14'055 to 5'226 TJ). In case of a decrease, overestimating of oxidation factors may tend to overestimate emission decrease. The main remaining consumer of coal in Switzerland is the

cement industry that accounts for 80% of total Swiss coal consumption in 2012 (EMIS 2014/1A2fi Zementwerke Feuerung). According to EU guidelines, the oxidation factor in cement production is assumed to be 100% (EC 2004). Given the large share of coal used in cement production, and under the assumption of high efficiency boilers, the overestimation of emission decrease may become minor.

Therefore, for all fuel combustion activities, an oxidation factor of 100% is assumed in Switzerland. This is also confirmed by the EU and Swiss guidelines for the Emission Trading System, where a default oxidation factor of 100% is applied.

3.2.5.3 Emissions from Biomass

CO₂ emissions from biomass do not count for the national total emissions and are therefore a memo item only. The CO₂ emissions from biomass in the CRF-tables are incomplete and the following CO₂ emissions are not foreseen for reporting in the CRF-tables: 2D2 Food and Drink, 3D5 Consumption of tobacco, 6A Solid Waste Disposal on Land, 6B Wastewater Handling and 6D Composting and Fermentation of Waste.

The following table provides an overview of effective biomass CO₂ emissions in Switzerland in 2012 and their reporting in the CRF-tables (without land-use, land-use change and forestry). Data regarding waste incineration is provided by FOEN 2013j. Data for CO₂ emissions from wood combustion is provided from Swiss Wood Energy Statistics (SFOE 2013b) for activity data and from the handbook of emission factors for stationary sources (SAEFL 2000) and Nussbaumer, T., Boogen, N. 2010 for emission factors as documented in the chapter above. For 2D and 3D, the data is provided by industry and expert estimates as documented in the respective EMIS 2014. Also emissions from Solid waste disposal on land (6A) and Other waste (6D) are based on specific data as documented in the respective EMIS. Data on waste water handling is provided by SFOE 2013a. For further information on the biomass CO₂ emissions refer to the respective source category chapters below.

Table 3-14 Effective biomass CO₂ emissions in Switzerland in 2012 and their representation in the CRF-tables.

| Biomass CO ₂ emissions | Unit | 2012 | Note |
|--|------|-------|------------------------------|
| 1A1 Energy Industries (without MSW incineration) | Gg | 599 | Included in CRF |
| 1A1 Energy generation from MSW Incineration | Gg | 2'549 | Included in CRF |
| 1A2d Use of waste derived fuels in cellulose production | Gg | 0 | Included in CRF |
| 1A2f Manufacturing Industry and Construction | Gg | 1'209 | Included in CRF |
| thereof use of waste derived fuels in cement production | Gg | 49 | |
| 1A3 Transport | Gg | 44 | Included in CRF |
| 1A4 Other Sectors (Commercial/Institutional, Residential) | Gg | 2'403 | Included in CRF |
| 2D Food and Drink | Gg | 15 | Not included in CRF |
| 3D Consumption of tobacco | Gg | 13 | Not included in CRF |
| 6A Solid Waste Disposal on Land | Gg | 25 | Not included in CRF |
| 6B Wastewater Handling | Gg | 136 | Not included in CRF |
| 6C Waste Incineration (without MSW incineration) | Gg | 137 | Included in CRF |
| 6D Other Waste (compost and fermentation of waste) | Gg | 353 | Not included in CRF |
| Total biomass combustion CO ₂ emissions included in CRF | Gg | 6'941 | |
| Total energy related biomass combustion CO ₂ emissions included in CRF 1A | Gg | 6'804 | See table "Summary 2" in CRF |
| Total biomass CO ₂ emissions in Switzerland in 2011 | Gg | 7'483 | |

3.2.6 Source Category 1A1 - Energy Industries

3.2.6.1 Source Category Description

Tier 1 Key categories 1A1

CO₂ from the combustion of Liquid Fuels (level and trend)

CO₂ from the combustion of Gaseous Fuels (level and trend)

CO₂ from the combustion of Other Fuels (level and trend)

Tier 2 Key categories 1A1

CO₂ from the combustion of Other Fuels (level and trend)

According to IPCC guidelines, source category 1A1 Energy Industries comprises emissions from fuels combusted by fuel extraction and energy producing industries.

In Switzerland, fuel extraction is virtually not occurring (apart from a very small charcoal production activity (traditional and historic craft)). Source category 1A1 includes therefore primarily emissions from the production of heat and/or electricity for sale to the public. Energy Industries (source category 1A1) comprise:

- Public Electricity and Heat Production including heat and power production in municipal solid waste incineration plants and special waste incineration (1A1a)
- Petroleum Refining (1A1b)
- Charcoal production within Manufacture of Solid Fuels and Other Energy Industries (1A1c)

Emissions from the industry producing heat and/or electricity for their own use are included in category 1A2 Manufacturing Industries and Construction. Emissions from producers of heat and/or power for their own use in waste incineration plants, however, are included in 1A1a.

In Switzerland, electricity production is dominated by hydroelectric power plants (59%) and nuclear power stations (35%). Fossil fuelled combined heat and power generation (CHP) provide 3% of electricity production. Power generation from renewable energy sources as solar, wind and biomass account only for about 3% of the electricity generated in Switzerland (SFOE 2013; table 24; data for the year 2012).

Table 3-15 Specification of source category 1A1 Energy Industries.

EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1A1 | Source | Specification | Data Source |
|------|--|--|---|
| 1A1a | Public Electricity and Heat Production | Main source are waste incineration plants with heat and power generation (Other fuels) and public district heating systems, including a small fraction of combined heat and power. The only fossil fuelled public electricity generation unit "Vouvry" (300 MW _e ; no public heat production) ceased operation in 1999. | Waste incineration: AD: FOEN 2013j; EMIS 2014/1A1a EF: Mohn 2011; Mohn 2013; EMIS 2014/1A1a Other sources: AD: SFOE 2013; SFOE 2013a; SFOE 2013b; EV 2013; EMIS 2014/1A1a EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006; EMIS 2014/1A Holzfeuerungen; SAEFL 2000; Nussbaumer, T., Boogen, N. 2010; EMIS 2014/1A1a |
| 1A1b | Petroleum Refining | Combustion activities supporting the refining of petroleum products, excluding evaporative emissions. | AD: EV 2013, SFOE 2013 EF: Industry data |
| 1A1c | Manufacture of Solid Fuels and Other Energy Industries | Charcoal production | AD / EF: EMIS 2013/1A1c |

3.2.6.2 Methodological Issues

a) Public Electricity and Heat Production (1A1a)

The public electricity and heat production in Switzerland includes:

- Fossil fuel combustion of gas oil, residual fuel oil, natural gas and coal.
- Waste-to-energy through the incineration of municipal solid waste and special waste (Other fuels)
- Biomass combustion includes wood and renewable waste and biogas generation from co-generation of landfills and fermentation engines

Methodology

The method applied within Public Electricity and Heat Production (1A1a) is country specific.

As explained in chapter 3.2.5, a country specific Tier 2 top-down approach based on aggregated fuel consumption data from the Swiss overall energy statistics is used to calculate emissions. For waste incineration, direct data from the incineration plants is used.

Emissions of GHGs are calculated based on the following methodology:

- For fossil fuel combustion, GHGs are calculated by multiplying fuel consumption (in TJ) by emission factors.
- For heat and/or power generation in municipal solid waste and special waste incineration plants the GHG emissions are calculated by multiplying the waste quantity incinerated by emission factors.
- For fermentation engines and co-generation on landfills the GHG emissions are calculated by multiplying quantities of combusted CH₄ by emission factors.

- For wood combustion in district combined heat and power units and plants for renewable waste from wood products the GHG emissions are calculated by multiplying the used wood chips and wood waste quantities by emission factors.

Emission Factors

The following table presents the emission factors used in 1A1a:

Table 3-16 Emission Factors for 1A1a Public Electricity and Heat Production in Energy Industries in 2012. Grey shaded cells mark general emission factors as described in section 3.2.5.2.

| 1A1a Public Electricity/Heat | CO ₂ t/TJ | CO ₂ bio. t/TJ | CH ₄ kg/TJ | N ₂ O kg/TJ | NO _x kg/TJ | CO kg/TJ | NM VOC kg/TJ | SO ₂ kg/TJ |
|---|-------------------------|------------------------------|--------------------------|---------------------------|--------------------------|-------------|-----------------|--------------------------|
| Gas oil | 73.7 | | 1.0 | 0.6 | 35 | 7 | 2.0 | 22 |
| Residual fuel oil | 77.0 | 0 | 4.0 | 0.8 | 125 | 13 | 4.0 | 291.0 |
| Petroleum coke | NO | NO | NO | NO | NO | NO | NO | NO |
| Natural gas | 56.1 | | 6.0 | 0.1 | 19 | 11 | 2.0 | 0.5 |
| Other (waste-to-energy), fossil | 103.4 | | NA | 2.6 | 33 | 7 | 1.1 | 4 |
| Other (waste-to-energy), biogenic | | 110.6 | NA | 2.4 | | | | |
| Biomass (wood, renewable waste) | | 65.1 | 1.9 | 1.6 | 162 | 283 | 6.4 | 20 |
| Biogas (co-generation from landfills, fermentation engines) | | 99.7 | 2 | 0.100 | 49 | 67 | 2.9 | 15 |

Emission factors highlighted in grey are explained in section 3.2.5.2. The study mentioned in 3.2.5.2 also included emission factors for NO_x, CO, NMVOC and SO₂.

Specific emission factors for 1A1a Public Electricity and Heat Production in Energy Industries are:

(a) Municipal solid waste incineration ("Other fuels")

Emission factors of Other (waste-to-energy) corresponds to emission occurring in waste incineration with heat and/or power generation (reported under "Other fuels"). The emission factors for CO₂, N₂O, NO_x, CO, NMVOC and SO₂ emissions per ton of waste incinerated are country specific and based on measurements and expert estimates, as documented in EMIS 2014/1A1a Kehricht- und Sondermüllverbrennungsanlagen. Emission factors are taking into account flue gas cleaning standards in incineration plants. In addition, the burn-out efficiency in modern municipal solid and special waste incineration plants is very high.

CO₂ emission factors for other fuels fluctuate over the period 1990-2012. The emission factor of municipal waste incineration changes because of gradual changes in the biogenic fraction and small variations of the net calorific values of the waste. From 1990 till 1997, the emission factor gradually decreased and stabilised until 2005. From 2006 onwards, the emission factor is stable. Emission factor of special waste is constant over time. See respective documentation in EMIS 2014/1A1a Kehricht- und Sondermüllverbrennungsanlagen.

Regarding the percentage of fossil CO₂ within waste incineration, a study conducted by the Swiss Federal Laboratories for Materials Testing EMPA evaluated the waste incinerated in Switzerland (national and imported waste, Mohn 2011). Based on this information, the share of organic matter in the waste incinerated in MSW incineration plants is estimated to be 52.2% in 2012 (see documentation in EMIS 2014/1A1a Kehrichtverbrennungsanlagen).

Emissions of CH₄ are not occurring because of the high combustion temperatures in waste incineration plants.

N₂O emission factor is divided into fossil and biogenic emissions from waste incineration of municipal waste and special waste. The fossil emission factor is based on two processes: municipal waste incineration and special waste incineration. The biogenic emission factor includes only municipal waste incineration as special waste is considered not to be biogenic. In 2013, a study by EMPA has evaluated measurements that have been performed in the years 2010-2011 in five Swiss municipal waste incineration plants (MWIP, Mohn 2013).

Emission factors have been calculated according to the state of equipment of all the Swiss waste incineration plants (with two types of Denox-equipment (SCR, SNCR⁸) and without Denox-equipment). The emission factor is therefore not constant over time and decreases from 5.3 g/GJ in 1990 to 2.4 g/GJ in 2012 based on the installation of Denox-equipment in the Swiss municipal waste incineration plants. See respective documentation in EMIS 2014/1A1a Kehricht- und Sondermüllverbrennungsanlagen.

(b) Public Electricity and Heat Production from biogas

Emission factors for biogas is country specific and composed by four underlying processes: landfill engines, engines and boilers in agricultural fermentation plants and engines in industrial fermentation plants (see documentation in EMIS 2014/1 A 1 a und 6 D_Vergärung LW and EMIS 2014/1 A 1 a und 6 D_Vergärung IG). The emission factors displayed in Table 3-16 are a weighted mean value of these four processes.

CO₂ emission factor is provided by a study realised on behalf of SFOE based on measurements in several agricultural installations and a respective life-cycle assessment for electricity production from agricultural biogas (see documentation in EMIS 2014/1 A 1 a und 6 D_Vergärung LW and EMIS 2014/1 A 1 a und 6 D_Vergärung IG).

CH₄ emission factor for landfill engines corresponds to industrial gas engines reported in SAEFL 2000. The other CH₄ emission factor have been evaluated in 2009. Fermentation engines are based on a study realised on behalf of SFOE for the values until 2000 and expert judgement from 2000 onwards (see documentation in EMIS 2014/1 A 1 a und 6 D_Vergärung LW and EMIS 2014/1 A 1 a und 6 D_Vergärung IG). For agricultural fermentation boilers, the emission factor is provided by a study developed by Wolfgang Butz (see documentation in EMIS 2014/1 A 1 a und 6 D_Vergärung LW).

N₂O emission factors of biogas is considered to be the same as for natural gas engines in commercial and institutional buildings that corresponds to the IPCC default value for natural gas (IPCC 2006).

Alls emission factors described above are listed in Table 3-16.

Activity Data

Activity data for the different fuels is determined as described in chapter 3.2.5.1. This includes gas oil, residual fuel oil, natural gas and biomass. For the solid fuels bituminous coal and lignite, activity data is provided from the Swiss overall energy statistics (SFOE 2013). Other fuel is calculated from the annual amount of waste incinerated producing heat and/or electricity (see Table 3-18). Activity data for co-generation from landfills and fermentation engines is taken from the Swiss renewable energies statistics (SFOE 2013a).

⁸ SCR: Selective Catalytic Reduction SCR, SNCR: Selective Non Catalytic Reduction

Table 3-17 Activity data in 1A1a Public Electricity/Heat.

| 1A1a Public Electricity /Heat | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Total fuel consumption | TJ | 41'175 | 42'131 | 44'053 | 39'231 | 38'874 | 39'902 | 42'895 | 43'712 | 48'995 | 50'152 |
| Gas oil | TJ | 980 | 1'790 | 1'917 | 1'662 | 810 | 554 | 810 | 1'065 | 852 | 1'065 |
| Residual fuel oil | TJ | 3'195 | 5'006 | 6'336 | 1'748 | 1'541 | 1'791 | 2'420 | 1'063 | 4'093 | 1'227 |
| Petroleum coke | TJ | NO | NO | NO | NO | NO | NO | NO | NO | NO | NO |
| Natural gas | TJ | 5'165 | 5'249 | 5'239 | 5'229 | 5'472 | 6'137 | 7'468 | 7'828 | 7'701 | 9'998 |
| Bituminous coal | TJ | 484 | 102 | 102 | 51 | 76 | 51 | 0 | 0 | 0 | 0 |
| Lignite | TJ | NO | NO | NO | NO | NO | NO | NO | NO | NO | NO |
| Other (waste-to-energy), fossil | TJ | 13'874 | 13'363 | 13'718 | 13'379 | 13'842 | 14'029 | 14'492 | 15'396 | 16'899 | 17'514 |
| Other (waste-to-energy), biogenic | TJ | 16'895 | 16'006 | 15'967 | 16'216 | 16'038 | 16'235 | 16'419 | 17'265 | 18'386 | 19'270 |
| Biomass (wood, renewable waste) | TJ | 301 | 297 | 360 | 404 | 441 | 466 | 636 | 466 | 431 | 412 |
| Biogas (co-generation from landfills, fermentation engines) | TJ | 282 | 320 | 414 | 542 | 653 | 639 | 650 | 629 | 633 | 665 |

| 1A1a Public Electricity /Heat | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Total fuel consumption | TJ | 50'821 | 52'743 | 53'622 | 54'619 | 55'695 | 57'983 | 61'288 | 58'916 | 60'141 | 59'196 |
| Gas oil | TJ | 810 | 852 | 810 | 1'065 | 810 | 1'321 | 1'278 | 810 | 469 | 554 |
| Residual fuel oil | TJ | 314 | 371 | 377 | 455 | 350 | 289 | 297 | 242 | 158 | 124 |
| Petroleum coke | TJ | NO | NO | NO | NO | NO | NO | NO | NO | NO | NO |
| Natural gas | TJ | 9'109 | 9'493 | 9'691 | 10'470 | 10'471 | 10'560 | 9'379 | 8'571 | 9'160 | 8'739 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | NO | NO | NO | NO | NO | NO | NO | NO | NO | NO |
| Other (waste-to-energy), fossil | TJ | 18'564 | 18'995 | 19'321 | 19'559 | 20'492 | 21'551 | 24'023 | 23'506 | 24'144 | 23'675 |
| Other (waste-to-energy), biogenic | TJ | 20'807 | 21'814 | 22'164 | 21'740 | 22'294 | 22'956 | 24'857 | 23'699 | 23'199 | 22'427 |
| Biomass (wood, renewable waste) | TJ | 547 | 583 | 689 | 774 | 813 | 844 | 943 | 1'462 | 2'316 | 2'881 |
| Biogas (co-generation from landfills, fermentation engines) | TJ | 671 | 634 | 570 | 556 | 465 | 461 | 511 | 626 | 695 | 796 |

| 1A1a Public Electricity /Heat | Unit | 2010 | 2011 | 2012 |
|---|------|--------|--------|--------|
| Total fuel consumption | TJ | 63'489 | 61'525 | 65'417 |
| Gas oil | TJ | 512 | 426 | 810 |
| Residual fuel oil | TJ | 52 | 7 | 8 |
| Petroleum coke | TJ | NO | NO | NO |
| Natural gas | TJ | 10'715 | 8'170 | 8'890 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | NO | NO | NO |
| Other (waste-to-energy), fossil | TJ | 25'280 | 25'403 | 26'262 |
| Other (waste-to-energy), biogenic | TJ | 22'997 | 22'444 | 23'051 |
| Biomass (wood, renewable waste) | TJ | 2'964 | 3'988 | 5'025 |
| Biogas (co-generation from landfills, fermentation engines) | TJ | 969 | 1'086 | 1'370 |

The table above shows that in 2012 Other fuels are the major component with 75% of the total fuel consumptions. The fossil fuels contribute with a total of 15% while natural gas has the major contribution with 92% of the total fossil fuel contribution in this source category. Biomass and Biogas contribute with 10% to the total fuel consumption.

The table above documents the increase of Other Fuel consumption (fossil) by 89% from 1990 to 2012. Overall, Other Fuels increased by 60%. This increase is the reason for category 1A1 Other Fuels – CO₂ being a key category regarding trend. See further explanations on this source category below.

The consumption of natural gas increased by 72% and the consumption of liquid fuels decreased by 17% for gas oil and 100% for residual fuel oil. These developments are due to shift in fuel use of combined heat and power generation in Switzerland and show why the CO₂ emissions from liquid and gaseous are key categories regarding trend in this submission.

Municipal solid waste incineration ("Other fuels")

Figure 8-4 in Sector 6 Waste shows an overview of the type of treatment and amounts of waste fractions reported in the different sectors in Switzerland.

Municipal solid waste includes waste generated in households and waste from other sources of similar composition. Energy recovery from municipal solid waste incineration is mandatory in Switzerland and plants are equipped with energy recovery systems (Schwager 2005). The emissions from heat and/or power generation in municipal solid waste incineration plants are

therefore reported under category 1A1a. Included are also emissions from the incineration of special waste, because these plants are also equipped with energy recovery systems. Activity data for waste incineration is taken from FOEN 2013j and provided in the table below.

Special waste is composed by special waste with high calorific value, wastewater and sludge with organic load, inorganic solids and dusts, inorganic sludge containing heavy metals, acids and alkalis, PCB-containing wastes, non-metallic shredder residues, contaminated soil, filter materials and chemicals residues and others.

Table 3-18 Activity data for 1A1a Other fuels: municipal solid waste and special waste incinerated with heat and/or power generation 1990 to 2012.

| Source/fuel | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1A1a Other fuels | | | | | | | | | | | |
| Total Other fuels in 1A1a | Gg | 2'603 | 2'477 | 2'467 | 2'441 | 2'411 | 2'433 | 2'471 | 2'535 | 2'655 | 2'824 |
| Municipal solid waste | Gg | 2'470 | 2'340 | 2'310 | 2'310 | 2'250 | 2'270 | 2'290 | 2'337 | 2'419 | 2'586 |
| Special waste | Gg | 133 | 137 | 157 | 131 | 161 | 163 | 181 | 198 | 237 | 238 |

| Source/fuel | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1A1a Other fuels | | | | | | | | | | | |
| Total Other fuels in 1A1a | Gg | 3'040 | 3'163 | 3'258 | 3'226 | 3'366 | 3'527 | 3'896 | 3'816 | 3'865 | 3'827 |
| Municipal solid waste | Gg | 2'801 | 2'936 | 3'027 | 2'995 | 3'135 | 3'297 | 3'646 | 3'580 | 3'610 | 3'597 |
| Special waste | Gg | 239 | 227 | 232 | 231 | 231 | 230 | 250 | 236 | 255 | 230 |

| Source/fuel | Unit | 2010 | 2011 | 2012 |
|---------------------------|------|-------|-------|-------|
| 1A1a Other fuels | | | | |
| Total Other fuels in 1A1a | Gg | 3'968 | 3'924 | 4'104 |
| Municipal solid waste | Gg | 3'717 | 3'676 | 3'841 |
| Special waste | Gg | 252 | 247 | 263 |

The table above documents the increase by 56% of municipal solid waste and 97 % of special waste incinerated from 1990 to 2012. This is due to the fact that since 1st of January 2000, disposal on landfill sites of waste which can be incinerated is prohibited by law (TVA Art. 32). See also Chapter 8.4 on Waste Incineration. The increase is also partly due to municipal solid waste imported from neighbouring countries to optimize the load factor of MSW incineration plants.

This increase results in CO₂ emissions from Other fuels in category 1A1 being a key category regarding trend.

b) Petroleum Refining (1A1b)

Methodology

For fuel combustion in Petroleum Refining (1A1b), a country specific Tier 2 bottom-up method is used. The calculations are based on measurements and data from individual sources from the refining industry.

The emissions are calculated by multiplying the fuel consumption by the respective emission factor.

Emission Factors

The following table presents the emission factors used in 1A1b:

Table 3-19 Emission Factors for 1A1b Petroleum Refining in 2012.

| Source/fuel | CO ₂ t/TJ | CH ₄ kg/TJ | N ₂ O kg/TJ | NO _x kg/TJ | CO kg/TJ | NMVOC kg/TJ | SO ₂ kg/TJ |
|--------------------------|-------------------------|--------------------------|---------------------------|--------------------------|-------------|----------------|--------------------------|
| 1A1 b Petroleum Refining | | | | | | | |
| Residual fuel oil | 77 | 4 | 0.8 | 110 | 15 | 2.5 | 490 |
| Gas (refinery LPG) | 59.8 | 1 | 0.6 | 55 | 15 | 2.3 | 25 |
| P-Coke | 91.4 | 10 | 1.6 | 200 | 100 | 10.0 | 500 |

Emission factors of residual fuel oil and P-Coke (highlighted in grey) are explained in section 3.2.5.2. The study mentioned in this chapter also included emission factors for NO_x, CO, NMVOC and SO₂.

Regarding the emission factor of refinery gas, no regular measurements are available. In 2010, one refinery provided specific information on refinery gas composition over three successive years. Swiss value is higher than IPCC default, but lies in the given range.

Activity Data

Activity data on fuel combustion (TJ) for Petroleum Refining (1A1b) is extracted from the Annual Reports of the Swiss Petroleum Association (EV 2013).

Table 3-20 Activity data for 1A1b Petroleum Refining.

| Source/fuel | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|-------|-------|-------|-------|--------|--------|--------|--------|--------|--------|
| 1A1b Petroleum Refining Fuel Consumption | | | | | | | | | | | |
| Total | TJ | 5'906 | 8'670 | 8'137 | 9'290 | 10'679 | 10'317 | 11'092 | 10'693 | 11'022 | 11'353 |
| Residual fuel oil | TJ | 1'296 | 1'216 | 998 | 1'054 | 1'426 | 1'834 | 1'618 | 1'780 | 1'428 | 1'698 |
| Gas (refinery LPG) | TJ | 4'610 | 7'454 | 7'139 | 8'237 | 9'253 | 8'483 | 9'474 | 8'913 | 9'594 | 9'655 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Source/fuel | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1A1b Petroleum Refining Fuel Consumption | | | | | | | | | | | |
| Total | TJ | 10'091 | 10'909 | 11'447 | 10'525 | 14'257 | 14'395 | 15'814 | 13'482 | 14'841 | 14'200 |
| Residual fuel oil | TJ | 1'952 | 1'936 | 1'518 | 1'769 | 1'339 | 906 | 692 | 1'159 | 707 | 742 |
| Gas (refinery LPG) | TJ | 8'139 | 8'973 | 9'929 | 8'756 | 11'901 | 11'678 | 13'311 | 10'766 | 11'687 | 11'424 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 1'017 | 1'811 | 1'811 | 1'557 | 2'447 | 2'034 |

| Source/fuel | Unit | 2010 | 2011 | 2012 |
|--|------|--------|--------|--------|
| 1A1b Petroleum Refining Fuel Consumption | | | | |
| Total | TJ | 13'912 | 12'969 | 11'162 |
| Residual fuel oil | TJ | 895 | 776 | 1'228 |
| Gas (refinery LPG) | TJ | 11'015 | 10'508 | 8'154 |
| Petroleum coke | TJ | 2'002 | 1'684 | 1'780 |

The table above documents gas (refinery liquefied petroleum gas) as the major fuel used in source category 1A1b with a contribution of 73% in 2012. Because of the increase of consumption for petroleum refining by 77% from 1990 to 2012, gas (refinery liquefied petroleum gas) is key category regarding trend in the present submission. This is explained by the fact that in 1990 one of the two Swiss refineries operated at reduced capacity and in later years resumed full production, leading to higher fuel consumption in the following years.

Since 2004, one of the Swiss refineries is using petroleum coke as a fuel.

In 2012, one of the refineries was closed over six month based on the debt restructuring and the search for a new buyer, which explains the lower fuel consumption in 2012 (EV 2013).

c) Manufacture of Solid Fuels and Other Energy Industries (1A1c)

Methodology

In source category 1A1c Manufacture of Solid Fuels and Other Energy Industries the emissions from charcoal production are reported. A country specific Tier 2 bottom-up method is used. Emissions from charcoal production are calculated by multiplying the annual amount of produced charcoal by the corresponding emission factors.

Emission Factors

The following table presents the emission factors used in 1A1c:

Table 3-21 Emission Factor for 1A1c Manufacture of Solid Fuels and Other Energy Industries in 2012. CO₂ emission factor is biogenic.

| 1A1c Charcoal | Unit | CO ₂ biog. | CH ₄ | NO _x | CO | NMVOC |
|---------------------|-------|-----------------------|-----------------|-----------------|-------|-------|
| Charcoal production | kg/TJ | 16'900 | 1'000 | 10 | 7'000 | 1'700 |

CO₂ emission factor is based on literature (USEPA 1995) and CH₄, NO_x, CO and NMVOC emission factors are taken from Revised 1996 IPCC Guidelines as documented in EMIS 2014/1A1c.

Activity Data

The annual amount of charcoal produced in Switzerland base on data from the charcoal burner's association and single producers documented in EMIS 2014/1A1c. The value used differs from the data provided to the FAO. Nevertheless, the data used in the inventory is based on detailed queries with the few remaining sites where charcoal is produced. The main producer is the the Köhlerverein Romoos, small quantities are produced by individual traditional local trade shows (Karthause Ittingen, Freilichtmuseum Ballenberg). As the data is provided by detailed bottom-up information, this corresponds to the realistic value.

The production has increased by a factor of 2.6 between 1990 and 2012. This is due to two new charcoal production sites that started operation in 2004 as well as the reducing wood prices that increased production combined with a higher demand in Switzerland (Koehlerei, 2014).

Table 3-22 Activity data for 1A1c Manufacture of Solid Fuels and Other Energy Industries.

| 1A1c | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------|------|------|------|------|------|------|------|------|------|------|------|
| Charcoal production | TJ | 1.25 | 1.67 | 1.25 | 1.55 | 1.64 | 1.43 | 1.73 | 2.43 | 1.78 | 1.90 |

| 1A1c | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------|------|------|------|------|------|------|------|------|------|------|------|
| Charcoal production | TJ | 2.20 | 1.84 | 1.76 | 2.17 | 2.35 | 3.00 | 2.96 | 3.11 | 3.15 | 3.12 |

| 1A1c | Unit | 2010 | 2011 | 2012 |
|---------------------|------|------|------|------|
| Charcoal production | TJ | 2.82 | 2.93 | 3.25 |

3.2.6.3 Uncertainties and Time-Series Consistency

Overview of uncertainty in aggregated fuel consumption activity data (1A1 Fuel Combustion):

Details of uncertainty analysis of activity data (fuel consumption) in 1A1 are provided in the table below. For each fuel type, uncertainties of net import or net production data (column C) and uncertainties of stock changes (if applicable) have been estimated. From this, the combined uncertainty of final consumption of fuels has been calculated (column H).

Table 3-23 Details of uncertainty analysis of fuels in 1A1.

| A | B | C | D | E | F | G | H | I |
|-----------------------|--------------------------------------|--|--|---|---|-----------------------------------|--|---------|
| Fuel type (IPCC 2000) | Corresponding fuel type in SFOE 2013 | Net import/ net production Input data =Import-Export [TJ] | Import/ production data uncertainty Input data [%] | Correction for stock changes etc. =G-C [TJ] | Correction uncertainty Input data [%] | Consumption Input data [TJ] | Final consumption uncertainty = $\text{WURZEL}((C \cdot D)^2 + (E \cdot F)^2) / G$ [%] | Comment |
| Liquid fuels | Erdölprodukte | 400'503 | 1.0 | 18'060 | 20 | 418'563 | 1.3 | 1 |
| Gaseous fuels | Gas | 122'520 | 2 | 0 | 0 | 122'520 | 2.0 | 2 |
| Solid fuels | Kohle | 5'630 | 5 | -160 | 100 | 5'470 | 5.9 | 3 |
| Other fuels | Müll- und Industrieabfälle | 56'320 | 10 | 0 | 0 | 56'320 | 10.0 | 4 |

Comments:

- 1 Col. D: Expert estimate from carbura (email M. Ruffer 24.1.05; overall uncertainty has been doubled to account for 95% interval). - Col. F: Conservative interpretation of rough expert estimate from carbura ("one-digit uncertainty", i.e. 10% is one sigma, resulting in $\text{unc} = 2 \cdot \text{sigma} = 20\%$).
- 2 Col. D: 2% is GPG default value for developed countries siehe unten
- 3 Col. D: 5% is GPG default value for developed countries (IPCC 2000 p. 2.1). - Col. G: expert estimate
- 4 Col. D: An uncertainty of amount of waste of 10% is assumed (expert judgement), because waste input is reasonably well measured since the nineties.

Data on stock changes is taken from the Swiss overall energy statistics (SFOE 2013; Table 4). Accordingly, also net import/net production data were taken from the Swiss overall energy statistics for the present uncertainty analysis.

The uncertainty of CO₂ emission factors is described in section 3.2.5.2.

Uncertainty in emissions of non-CO₂ gases are estimated to be medium resulting in 30% for CH₄ and 80% for N₂O (see 1.7).

Consistency and Completeness in 1A1 Fuel Combustion

Consistency:

- Time series for 1A1 are all consistent.
- CO₂ emissions from biomass in 1 Energy (memo item) are only partly included in the CRF-tables, see Section 3.2.5.1.

Completeness:

- All estimates in the sector 1A1 are assumed to be complete.

3.2.6.4 Source-specific QA/QC and Verification

a) General

At the level of total energy-related CO₂ emissions, a quality control consists in the comparison of emissions modelled using the Sectoral Approach with emissions calculated from fuel consumption according to the Swiss overall energy statistics of SFOE. The differences in total CO₂ emissions for the years 1990–2012 are negligible - indicating the completeness of the inventory.

The cross-check of the Reference and Sectoral Approach is also used for an assessment of emissions related to the consumption of fuels in the energy sector. Again, a good agreement between the two approaches is found (see Chapter 3.2.1).

The quality control activities have been documented in checklists as described in Chapter 1.6

b) Specific Energy Industries

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of last submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of last submission 2013

In 2012, the emission factors of category 1A1 used in the Swiss Inventory have been compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (INFRAS 2012). The emission factor for CO₂ from Other Fuels from Other Fuels are higher than the emission factors in other countries. Please see section 3.2.6.2.

3.2.6.5 Source-Specific Recalculations

- **1A all fuels:** Activity data of all fuels of the overall time series have been recalculated based on the new data from SFOE 2013. This includes data on 2011 for fossil fuels and waste, data for biomass on 1997, 2009 – 2011 and data of other fuels 1990 – 2011.
- **1A wood consumption:** Activity data have been recalculated for the overall time series based on the new data from SFOE 2013b.
- **1A gas oil:** SO_x emission factor value has been updated for 2010 based on sulphur analyses of the gas oil for the year 2010 (Directorate General of Customs) resulting in a revised value for 2011 as well.
- **1A bituminous coal:** CO₂ emission factor has been corrected from 94t CO₂/TJ to 92.7 t CO₂/TJ for the entire time series.
- **1A1a biogas production:** Activity data of biogas production from solid waste disposal sites have been recalculated for 2011 based on new values in the Swiss statistics of renewable energies SFOE 2013a.
- **1A1a waste incineration:** Emission factor of N₂O has been updated for the entire time series based on a new study realized by EMPA (Mohn 2013).
- **1A1a biomass fermentation:** Emission factor of CH₄ has been corrected for the entire time series.
- **1A1a / 6D industrial and agricultural biogas:** Activity data for fermentation of biogenic waste has been updated for the years 1999 to 2011 based on new data from the Swiss statistics of renewable energies SFOE 2013a.
- **1A1a industrial and agricultural biogas:** N₂O default emission factor from the IPCC guidelines has been introduced for the whole time series for fermentation of biogenic waste.
- **1A1a bituminous coal:** Activity data has been corrected over the whole time series as it was not consistent with other years.
- **1A1a gas oil:** Activity data has been corrected over the whole time series based on data from the Swiss overall energy statistics (SFOE 2013).

- **1A1a other fuels:** Activity data has been corrected for 2007 and 2008 based on correction of the energy content of waste.
- **1A1b refinery boilers:** Activity data for 2011 has been corrected for residual fuel oil as there was a mistake in the database in Submission 2013.
- **1A1b refinery boilers:** CO₂ Emission factor of refinery liquefied petroleum gas for 1990 to 2011 has been updated based on measurements that led to an average emission factor of 59'800 g/GJ compared to the previous emission factor of 59'300 g/GJ that was based on expert judgement.
- **1A1c charcoal production:** Reporting of the biogenic CO₂ and precursor emissions has been shifted from source category 2D3 to 1A1c. (Please note that the reporting of the CH₄ emissions from the charcoal production has already been shifted from 2D3 to 1A1c within the resubmission of Switzerland's Greenhouse Gas Inventory 1990–2011(FOEN 2013g).)
- **1A1c charcoal production:** Activity data from 2004 to 2012 has been corrected based on corrected data from small producers of charcoal.
- **1A1c charcoal production:** Emission factors have been corrected over the whole timeseries. The emission factors are newly based on Revised 1996 IPCC Guidelines for CO, NMVOC, CH₄, NO_x and USEPA (1995) for CO₂.

3.2.6.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.2.7 Source Category 1A2 - Manufacturing Industries and Construction

3.2.7.1 Source Category Description

Tier 1 Key categories 1A2

CO₂ from the combustion of Liquid Fuels (level and trend)

CO₂ from the combustion of Solid Fuels (level and trend)

CO₂ from the combustion of Gaseous Fuels (level and trend)

CO₂ from the combustion of Other Fuels (level and trend)

Tier 2 Key categories 1A2

CO₂ from the combustion of Solid Fuels (level and trend)

CO₂ from the combustion of Gaseous Fuels (level and trend)

CO₂ from the combustion of Other Fuels (level and trend)

The source category 1A2 Manufacturing Industries and Construction comprises all emissions from the combustion of fuels in stationary boilers and cogeneration facilities within manufacturing industries and construction. This includes use of conventional fossil fuels as well as waste fuels and biomass. Use of fossil fuels as feedstocks or as so-called non-energy fuel as for example bitumen and lubricants are included in the reference approach and described in section 3.2.3.

In addition, this source category includes off-road construction and industrial vehicles and machinery in category 1A2fii, such as for example forklifts, diggers or industry tractors.

Table 3-24 Specification of source category 1A2 Manufacturing Industries and Construction.
 EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1A2 | Source | Specification | Data Source |
|--------|---|--|---|
| 1A2a | Iron and Steel | Iron and Steel industry | AD: SFOE 2013; Prognos, 2013 EMIS 2014/1A2a EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006 EMIS 2014/1A2a |
| 1A2b | Non-ferrous Metals | Non-ferrous Metals industry | AD: SFOE 2013; Prognos, 2013 EMIS 2014/1A2b EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006 EMIS 2014/1A2b |
| 1A2c | Chemicals | Chemical industry | AD: SFOE 2013; Prognos, 2013 EMIS 2014/1A2c EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006 EMIS 2014/1A2c |
| 1A2d | Pulp, Paper and Print | Pulp, Paper and Print industry | AD: SFOE 2013; Prognos, 2013 EMIS 2014/1A2d EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006 EMIS 2014/1A2d |
| 1A2e | Food Processing, Beverages and Tobacco | Food Processing, Beverages and Tobacco industry | AD: SFOE 2012; Prognos 2013 EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006 |
| 1A2fi | Other (Combustion Installations in Industries) | Category 1A2fi contains: Cement, Lime, Brick and tile, Fine ceramics, Asphalt concrete plants, Container glass, Glass, Glass wool, Mineral wool, Fibreboard Production, industrial biogas boilers and engines that do not provide heat or electricity to the public. | AD: SFOE 2013; Prognos 2013; SFOE 2013a; SFOE 2013b; SFOE 2013c EMIS 2014/1A2fi EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006; EMIS 2014/1A Holzfeuerungen; SAEFL 2000; Nussbaumer, T., Boogen, N. 2010; EMIS 2014/1A2fi |
| 1A2fii | | Category 1A2f ii contains: off-road construction and industrial vehicles and machinery. | AD, EF: INFRAS 2008 AD (partial update): Prognos 2012, Keller/INFRAS 2013 |

3.2.7.2 Methodological Issues

Methodology

For fuel combustion in source category 1A2 Manufacturing Industries and Construction, a country specific approach, as explained in chapter 3.2.5, is used combining Tier 2 and Tier 3 methods.

Emissions of GHGs are calculated by multiplying the level of activity (fuel consumption) by the respective emission factors.

Within 1A2f, also emissions from diesel and gasoline use in construction and industrial machinery (off-road) is included and accounted for within source category 1A2fii. They are calculated with a Tier 2 method. Some details of the emission modelling that hold for all off-road families are described in Annex A3.1.6 Off-road vehicles. Emission calculation was carried out in a database structured in analogy to the on-road database (INFRAS 2008).

Emission factors

The following table presents the emission factors used in source category 1A2 Manufacturing Industries and Construction:

Table 3-25 Emission Factors for 1A2 Manufacturing Industries and Construction in 2012.

| 1A2 Emission factors (mix of bottom-up and top-down approach (modelling)) for GHG | CO ₂ fossil | CO ₂ bio. | CH ₄ | N ₂ O |
|---|------------------------|----------------------|-----------------|------------------|
| | t/TJ | t/TJ | kg/TJ | kg/TJ |
| Gas oil | 73.7 | | 1.0 | 0.6 |
| Liquefied petroleum gas | 65.5 | | 1.0 | 0.1 |
| Residual fuel oil | 77.0 | | 3.2 | 0.8 |
| Petroleum coke | 91.4 | | 2.1 | 1.6 |
| Bituminous coal | 92.7 | | 1.6 | 1.6 |
| Lignite | 96.1 | | 0.8 | 1.6 |
| Natural gas | 56.1 | | 6.0 | 0.1 |
| Biomass | | 85.0 | 5.4 | 1.6 |
| Other fuels | 70.4 | | 4.3 | 9.5 |
| Diesel and gasoline for construction and industrial machinery | 73.6 | | 2.2 | 2.8 |

All emission factors highlighted in grey are explained in section 3.2.5.2.

Other fuels consist of various fossil wastes (see detailed description below in section Cement under Other – Stationary (1A2f i)). The CO₂ emission factor is an implied emission factor based on the fossil waste fuel mix in cement production plants. The CH₄ emission factor includes the overall CH₄ emissions of the cement industry based on direct exhaust measurements at the chimneys of the cement plants (see respective documentation in EMIS 2014/1A2fi Zementwerke_Feuerung, Cemsuisse 2010a). CH₄ emission factors for residual fuel oil, petroleum coke, bituminous coal and lignite are lower than the emission factors explained in section 3.2.5.2. This is because the overall CH₄ emissions of the cement industry are reported under Other fuels (see explanation above).

The emission factors of the precursors NO_x, CO, NMVOC and SO₂ for all fuels in source category 1A2 are provided in Annex A3.1.3. Emission factors for CO and NO_x for natural gas and gas oil of boilers are country specific and result from a specific modeling based on real measurements of 200'000 firing controls in eight Swiss cantons (Leupro 2012). NMVOC and SO₂ emission factors are country specific and based on measurements as documented in the respective EMIS documentation.

The emission factors for all other gases are country specific and shown in Table A - 15 to Table A - 18 in the Annex A3.1.6 (INFRAS 2008). NMVOC emissions are calculated as the difference of VOC and CH₄ emissions .

For off-road activities the emission factors for CO₂ are country specific and assumed to be constant in the period 1990-2012 with values 73.6 t/TJ for diesel oil, 73.9 t/TJ for gasoline and 56.1t/TJ for CNG (equal to natural gas). See also Table 3 9.

Note that specific emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database INFRAS (2008), see footnote 7 on page 111.

For off-road activities SO₂ emission factors are country specific and further described in Table A - 6 in Annex A2.

Activity data

Activity data for the different fuels is determined as described in chapter 3.2.5.1.

Table 3-26 Activity data fuel consumption in 1A2 Manufacturing Industries and Construction 1990 to 2012.

| Source | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1A2 Manufacturing Industries and Constr. (Total) | TJ | 83'907 | 88'028 | 84'591 | 84'318 | 84'414 | 87'966 | 86'534 | 85'874 | 88'569 | 88'788 |
| Gas oil | TJ | 19'307 | 23'533 | 23'578 | 23'368 | 21'996 | 23'421 | 24'499 | 25'525 | 26'591 | 28'492 |
| Liquefied petroleum gas | TJ | 4'495 | 5'236 | 4'723 | 4'542 | 4'683 | 4'684 | 4'866 | 5'789 | 6'157 | 6'664 |
| Residual fuel oil | TJ | 18'524 | 17'037 | 16'487 | 14'008 | 14'547 | 13'547 | 10'954 | 9'625 | 10'232 | 8'151 |
| Petroleum coke | TJ | 1'617 | 1'239 | 650 | 1'387 | 1'702 | 1'275 | 1'082 | 394 | 564 | 484 |
| Bituminous coal | TJ | 12'676 | 10'548 | 7'243 | 5'987 | 6'083 | 6'687 | 4'984 | 3'849 | 3'220 | 3'146 |
| Lignite | TJ | 306 | 353 | 306 | 259 | 259 | 188 | 236 | 165 | 139 | 136 |
| Natural gas | TJ | 19'146 | 21'865 | 23'386 | 26'065 | 26'679 | 28'658 | 29'263 | 29'970 | 30'694 | 30'531 |
| Biomass | TJ | 5'801 | 6'110 | 6'042 | 6'029 | 5'986 | 6'345 | 6'905 | 6'814 | 7'054 | 7'348 |
| Other Fuels | TJ | 2'035 | 2'106 | 2'176 | 2'673 | 2'480 | 3'162 | 3'746 | 3'744 | 3'918 | 3'837 |
| 1A2a Iron and Steel | TJ | 3'212 | 3'807 | 3'345 | 4'036 | 3'247 | 2'444 | 2'353 | 2'558 | 2'757 | 2'860 |
| Gas oil | TJ | 400 | 413 | 455 | 432 | 412 | 256 | 248 | 291 | 304 | 464 |
| Liquefied petroleum gas | TJ | 604 | 632 | 575 | 512 | 509 | 287 | 286 | 361 | 407 | 463 |
| Residual fuel oil | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Petroleum coke | TJ | 346 | 350 | 364 | 370 | 367 | 131 | 128 | 139 | 151 | 7 |
| Bituminous coal | TJ | 433 | 347 | 328 | 259 | 263 | 289 | 247 | 253 | 273 | 271 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 1'429 | 2'067 | 1'622 | 2'464 | 1'695 | 1'481 | 1'443 | 1'513 | 1'622 | 1'654 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2b Non-Ferrous Metals | TJ | 2'337 | 2'338 | 2'059 | 1'718 | 1'629 | 1'994 | 1'662 | 2'150 | 1'845 | 1'394 |
| Gas oil | TJ | 598 | 618 | 418 | 391 | 355 | 417 | 337 | 364 | 358 | 311 |
| Liquefied petroleum gas | TJ | 40 | 40 | 33 | 23 | 22 | 26 | 22 | 34 | 30 | 24 |
| Residual fuel oil | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 1'699 | 1'680 | 1'608 | 1'304 | 1'252 | 1'552 | 1'303 | 1'751 | 1'458 | 1'060 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2c Chemicals | TJ | 15'638 | 15'047 | 15'036 | 14'017 | 14'999 | 16'432 | 16'100 | 14'845 | 15'071 | 15'036 |
| Gas oil | TJ | 4'020 | 3'787 | 3'073 | 3'307 | 3'168 | 3'978 | 3'908 | 3'344 | 2'948 | 5'295 |
| Liquefied petroleum gas | TJ | 323 | 323 | 321 | 317 | 319 | 319 | 319 | 321 | 323 | 320 |
| Residual fuel oil | TJ | 1'434 | 1'193 | 851 | 796 | 654 | 693 | 561 | 383 | 256 | 315 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 9'861 | 9'744 | 10'791 | 9'597 | 10'859 | 11'442 | 11'312 | 10'797 | 11'544 | 9'107 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2d Pulp, Paper and Print | TJ | 10'831 | 11'183 | 12'488 | 12'640 | 13'520 | 12'121 | 11'253 | 11'467 | 11'309 | 10'649 |
| Gas oil | TJ | 711 | 949 | 1'027 | 1'095 | 1'112 | 1'235 | 1'340 | 1'466 | 1'580 | 1'857 |
| Liquefied petroleum gas | TJ | 127 | 161 | 223 | 219 | 269 | 210 | 185 | 215 | 216 | 222 |
| Residual fuel oil | TJ | 5'250 | 4'904 | 4'136 | 3'667 | 3'228 | 3'061 | 2'867 | 2'885 | 2'739 | 2'023 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 2'657 | 3'271 | 5'392 | 6'136 | 7'470 | 6'257 | 5'418 | 5'375 | 5'164 | 4'853 |
| Biomass | TJ | 2'085 | 1'898 | 1'711 | 1'524 | 1'441 | 1'358 | 1'442 | 1'526 | 1'610 | 1'694 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2e Food Processing, Beverages and Tobacco | TJ | 10'382 | 12'331 | 12'192 | 12'403 | 10'434 | 10'198 | 12'558 | 11'742 | 11'678 | 11'892 |
| Gas oil | TJ | 7'903 | 9'703 | 9'497 | 9'488 | 7'413 | 6'920 | 9'036 | 8'274 | 8'016 | 7'927 |
| Liquefied petroleum gas | TJ | 301 | 379 | 375 | 380 | 429 | 456 | 516 | 613 | 690 | 827 |
| Residual fuel oil | TJ | 1'160 | 1'009 | 810 | 705 | 553 | 466 | 405 | 284 | 221 | 184 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 1'018 | 1'240 | 1'511 | 1'830 | 2'040 | 2'356 | 2'601 | 2'571 | 2'751 | 2'954 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2f Other | TJ | 47'019 | 49'023 | 45'361 | 45'584 | 46'854 | 51'235 | 49'188 | 49'814 | 52'731 | 53'902 |
| Gas oil | TJ | 5'675 | 8'063 | 9'109 | 8'655 | 9'536 | 10'615 | 9'629 | 11'786 | 13'385 | 12'638 |
| Liquefied petroleum gas | TJ | 3'099 | 3'701 | 3'196 | 3'092 | 3'135 | 3'387 | 3'538 | 4'244 | 4'492 | 4'808 |
| Residual fuel oil | TJ | 10'680 | 9'931 | 10'691 | 8'841 | 10'112 | 9'327 | 7'121 | 6'073 | 7'015 | 5'629 |
| Petroleum coke | TJ | 1'271 | 890 | 286 | 1'017 | 1'335 | 1'144 | 953 | 254 | 413 | 477 |
| Bituminous coal | TJ | 12'242 | 10'202 | 6'915 | 5'728 | 5'819 | 6'398 | 4'736 | 3'596 | 2'947 | 2'874 |
| Lignite | TJ | 306 | 353 | 306 | 259 | 259 | 188 | 236 | 165 | 139 | 136 |
| Natural gas | TJ | 2'481 | 3'864 | 2'461 | 4'734 | 3'364 | 5'569 | 7'186 | 7'963 | 8'155 | 10'904 |
| Biomass | TJ | 3'716 | 4'212 | 4'331 | 4'505 | 4'545 | 4'987 | 5'463 | 5'288 | 5'444 | 5'653 |
| Other Fuels | TJ | 2'035 | 2'106 | 2'176 | 2'673 | 2'480 | 3'162 | 3'746 | 3'744 | 3'918 | 3'837 |
| 1A2fi Diesel and gasoline for construction and industrial machinery | TJ | 5'512 | 5'701 | 5'890 | 6'080 | 6'269 | 6'458 | 6'580 | 6'701 | 6'823 | 6'945 |

Table 3-26 continued: Activity data fuel consumption in 1A2 Manufacturing Industries and Construction.

| Source | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|--------|--------|--------|--------|---------|---------|---------|---------|---------|--------|
| 1A2 Manufacturing Industries and Constr. (Total) | TJ | 93'846 | 98'832 | 95'600 | 98'467 | 100'342 | 100'830 | 103'061 | 101'697 | 102'442 | 96'034 |
| Gas oil | TJ | 26'216 | 27'786 | 27'499 | 28'933 | 28'140 | 28'710 | 27'338 | 25'863 | 25'738 | 24'260 |
| Liquefied petroleum gas | TJ | 5'928 | 5'562 | 6'160 | 5'331 | 5'194 | 4'596 | 5'054 | 4'548 | 4'319 | 4'595 |
| Residual fuel oil | TJ | 5'665 | 7'623 | 4'564 | 4'858 | 5'700 | 4'435 | 5'128 | 3'519 | 3'517 | 2'549 |
| Petroleum coke | TJ | 529 | 422 | 645 | 194 | 954 | 1'228 | 1'772 | 1'470 | 1'281 | 1'378 |
| Bituminous coal | TJ | 5'265 | 5'432 | 5'000 | 5'178 | 4'704 | 4'704 | 3'889 | 4'831 | 4'373 | 4'144 |
| Lignite | TJ | 136 | 71 | 94 | 94 | 94 | 777 | 2'026 | 2'003 | 1'767 | 1'555 |
| Natural gas | TJ | 31'496 | 32'250 | 31'254 | 32'722 | 33'817 | 34'754 | 36'077 | 36'928 | 38'338 | 34'904 |
| Biomass | TJ | 7'191 | 7'378 | 7'516 | 7'960 | 8'065 | 8'242 | 8'313 | 8'708 | 8'616 | 7'827 |
| Other Fuels | TJ | 11'419 | 12'308 | 12'869 | 13'197 | 13'673 | 13'384 | 13'463 | 13'828 | 14'493 | 14'823 |
| 1A2a Iron and Steel | TJ | 2'912 | 3'009 | 3'186 | 3'412 | 3'417 | 3'153 | 4'075 | 4'266 | 3'963 | 2'918 |
| Gas oil | TJ | 367 | 472 | 451 | 523 | 463 | 398 | 338 | 352 | 341 | 302 |
| Liquefied petroleum gas | TJ | 427 | 392 | 506 | 447 | 405 | 324 | 465 | 440 | 368 | 319 |
| Residual fuel oil | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Petroleum coke | TJ | 21 | 40 | 9 | 3 | 191 | 179 | 310 | 230 | 232 | 171 |
| Bituminous coal | TJ | 266 | 234 | 179 | 163 | 148 | 154 | 150 | 160 | 177 | 70 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 1'832 | 1'870 | 2'042 | 2'276 | 2'209 | 2'098 | 2'811 | 3'085 | 2'844 | 2'056 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2b Non-Ferrous Metals | TJ | 1'504 | 1'351 | 843 | 925 | 1'103 | 954 | 1'146 | 1'031 | 1'035 | 987 |
| Gas oil | TJ | 292 | 115 | 330 | 166 | 114 | 161 | 100 | 129 | 156 | 223 |
| Liquefied petroleum gas | TJ | 23 | 21 | 10 | 12 | 15 | 11 | 15 | 11 | 10 | 10 |
| Residual fuel oil | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 1'189 | 1'215 | 502 | 747 | 974 | 783 | 1'032 | 890 | 869 | 754 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2c Chemicals | TJ | 14'333 | 14'985 | 13'981 | 15'767 | 16'100 | 16'522 | 16'636 | 15'557 | 15'691 | 13'809 |
| Gas oil | TJ | 4'139 | 4'073 | 3'806 | 4'409 | 4'482 | 4'394 | 4'498 | 3'450 | 3'192 | 3'484 |
| Liquefied petroleum gas | TJ | 318 | 318 | 319 | 318 | 317 | 316 | 316 | 315 | 364 | 313 |
| Residual fuel oil | TJ | 253 | 282 | 289 | 311 | 345 | 168 | 431 | 37 | 365 | 415 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 9'623 | 10'313 | 9'567 | 10'729 | 10'957 | 11'645 | 11'391 | 11'756 | 11'770 | 9'598 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2d Pulp, Paper and Print | TJ | 10'388 | 11'381 | 10'860 | 11'600 | 10'691 | 10'776 | 9'636 | 8'942 | 7'197 | 4'855 |
| Gas oil | TJ | 1'443 | 1'617 | 1'458 | 1'487 | 1'309 | 1'325 | 1'164 | 935 | 781 | 757 |
| Liquefied petroleum gas | TJ | 220 | 210 | 255 | 207 | 164 | 149 | 118 | 106 | 90 | 93 |
| Residual fuel oil | TJ | 1'421 | 2'256 | 1'235 | 1'941 | 2'074 | 1'811 | 2'239 | 1'644 | 1'175 | 643 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 5'610 | 5'851 | 6'170 | 6'081 | 5'115 | 5'438 | 4'038 | 4'158 | 3'827 | 3'362 |
| Biomass | TJ | 1'694 | 1'447 | 1'741 | 1'885 | 2'029 | 2'053 | 2'076 | 2'099 | 1'324 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2e Food Processing, Beverages and Tobacco | TJ | 11'703 | 12'268 | 11'676 | 11'583 | 11'675 | 11'787 | 12'622 | 12'014 | 11'877 | 13'531 |
| Gas oil | TJ | 7'336 | 7'651 | 6'881 | 6'544 | 6'282 | 5'831 | 5'521 | 4'894 | 4'892 | 5'363 |
| Liquefied petroleum gas | TJ | 799 | 776 | 929 | 812 | 818 | 796 | 1'008 | 889 | 801 | 1'098 |
| Residual fuel oil | TJ | 137 | 141 | 113 | 96 | 114 | 0 | 0 | 0 | 0 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Natural gas | TJ | 3'430 | 3'700 | 3'753 | 4'132 | 4'462 | 5'160 | 6'093 | 6'232 | 6'185 | 7'070 |
| Biomass | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1A2fi Other | TJ | 53'005 | 55'837 | 55'053 | 55'180 | 57'355 | 57'638 | 58'945 | 59'887 | 62'679 | 59'933 |
| Gas oil | TJ | 12'638 | 13'858 | 14'573 | 15'805 | 15'491 | 16'601 | 15'717 | 16'104 | 16'376 | 14'133 |
| Liquefied petroleum gas | TJ | 4'141 | 3'844 | 4'141 | 3'536 | 3'475 | 3'002 | 3'132 | 2'787 | 2'687 | 2'761 |
| Residual fuel oil | TJ | 3'854 | 4'945 | 2'926 | 2'510 | 3'167 | 2'456 | 2'458 | 1'839 | 1'977 | 1'490 |
| Petroleum coke | TJ | 508 | 381 | 636 | 191 | 763 | 1'049 | 1'462 | 1'239 | 1'049 | 1'208 |
| Bituminous coal | TJ | 4'999 | 5'198 | 4'821 | 5'015 | 4'556 | 4'550 | 3'739 | 4'672 | 4'196 | 4'074 |
| Lignite | TJ | 136 | 71 | 94 | 94 | 94 | 777 | 2'026 | 2'003 | 1'767 | 1'555 |
| Natural gas | TJ | 9'812 | 9'302 | 9'220 | 8'758 | 10'101 | 9'630 | 10'711 | 10'808 | 12'843 | 12'063 |
| Biomass | TJ | 5'497 | 5'931 | 5'774 | 6'075 | 6'036 | 6'189 | 6'238 | 6'609 | 7'292 | 7'827 |
| Other Fuels | TJ | 4'353 | 5'204 | 5'729 | 6'020 | 6'460 | 6'133 | 6'136 | 6'426 | 7'015 | 7'268 |
| 1A2fii Diesel and gasoline for construction and industrial machinery | TJ | 7'066 | 7'103 | 7'140 | 7'177 | 7'214 | 7'250 | 7'326 | 7'402 | 7'478 | 7'554 |

Table 3-26 continued: Activity data fuel consumption in 1A2 Manufacturing Industries and Construction.

| Source | Unit | 2010 | 2011 | 2012 |
|---|------|---------|--------|--------|
| 1A2 Manufacturing Industries and Constr. (Total) | TJ | 102'217 | 94'983 | 96'594 |
| Gas oil | TJ | 24'347 | 19'714 | 21'486 |
| Liquefied petroleum gas | TJ | 4'181 | 4'136 | 3'998 |
| Residual fuel oil | TJ | 1'955 | 1'518 | 1'568 |
| Petroleum coke | TJ | 4'322 | 3'813 | 3'686 |
| Bituminous coal | TJ | 4'322 | 3'813 | 3'686 |
| Lignite | TJ | 1'461 | 1'626 | 1'178 |
| Natural gas | TJ | 38'016 | 36'907 | 37'369 |
| Biomass | TJ | 7'916 | 7'562 | 8'289 |
| Other Fuels | TJ | 15'696 | 15'895 | 15'335 |
| 1A2a Iron and Steel | TJ | 3'436 | 3'386 | 4'352 |
| Gas oil | TJ | 327 | 322 | 241 |
| Liquefied petroleum gas | TJ | 327 | 338 | 435 |
| Residual fuel oil | TJ | 149 | 1 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 |
| Bituminous coal | TJ | 64 | 73 | 55 |
| Lignite | TJ | 0 | 0 | 0 |
| Natural gas | TJ | 2'569 | 2'652 | 3'620 |
| Biomass | TJ | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 |
| 1A2b Non-Ferrous Metals | TJ | 1'175 | 1'215 | 1'241 |
| Gas oil | TJ | 157 | 136 | 211 |
| Liquefied petroleum gas | TJ | 12 | 12 | 11 |
| Residual fuel oil | TJ | 0 | 0 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 |
| Natural gas | TJ | 1'007 | 1'067 | 1'019 |
| Biomass | TJ | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 |
| 1A2c Chemicals | TJ | 13'755 | 12'749 | 12'536 |
| Gas oil | TJ | 3'276 | 3'030 | 2'787 |
| Liquefied petroleum gas | TJ | 311 | 331 | 380 |
| Residual fuel oil | TJ | 217 | 3 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 |
| Natural gas | TJ | 9'952 | 9'385 | 9'369 |
| Biomass | TJ | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 |
| 1A2d Pulp, Paper and Print | TJ | 4'599 | 4'339 | 3'463 |
| Gas oil | TJ | 478 | 349 | 387 |
| Liquefied petroleum gas | TJ | 91 | 92 | 67 |
| Residual fuel oil | TJ | 83 | 1 | 2 |
| Petroleum coke | TJ | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 |
| Natural gas | TJ | 3'947 | 3'897 | 3'008 |
| Biomass | TJ | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 |
| 1A2e Food Processing, Beverages and Tobacco | TJ | 14'430 | 14'437 | 13'980 |
| Gas oil | TJ | 5'723 | 5'520 | 5'324 |
| Liquefied petroleum gas | TJ | 984 | 1'008 | 929 |
| Residual fuel oil | TJ | 0 | 0 | 0 |
| Petroleum coke | TJ | 0 | 0 | 0 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 |
| Natural gas | TJ | 7'723 | 7'909 | 7'727 |
| Biomass | TJ | 0 | 0 | 0 |
| Other Fuels | TJ | 0 | 0 | 0 |
| 1A2fi Other | TJ | 62'141 | 56'315 | 58'703 |
| Gas oil | TJ | 14'385 | 10'357 | 12'537 |
| Liquefied petroleum gas | TJ | 2'455 | 2'355 | 2'176 |
| Residual fuel oil | TJ | 1'655 | 1'513 | 1'566 |
| Petroleum coke | TJ | 1'494 | 1'271 | 1'367 |
| Bituminous coal | TJ | 4'259 | 3'740 | 3'630 |
| Lignite | TJ | 1'461 | 1'626 | 1'178 |
| Natural gas | TJ | 12'820 | 11'997 | 12'626 |
| Biomass | TJ | 7'916 | 7'562 | 8'289 |
| Other Fuels | TJ | 8'066 | 8'350 | 7'875 |
| 1A2fii Diesel and gasoline for construction and industrial machinery | TJ | 7'630 | 7'545 | 7'460 |

In Table 3-26 the specific fuel consumptions of source categories 1A2a–1A2f are given. Source category 1A2f Other is the most important category within source category 1A2 Manufacturing Industries and Construction and accounted for 61% of the overall fuel consumption in 2012. Categories 1A2e Food Processing, Beverages and Tobacco and 1A2c Chemicals are the second and third most important fuel consumers with 14% and 13% respectively.

Regarding the fuels used within Swiss industry, natural gas consumption represents 39% of fuel consumption in 2012 followed by gas oil and other fuels (fossil waste) with shares of 22% and 16%, respectively.

The table also documents the fuel switch within Swiss industry. From 1990 to 2012 the use of residual fuel oil and bituminous coal has decreased by 92% and 71%, respectively. In the same period, natural gas and other fuel consumption increased by 95% and a factor of 7.5 (654%), respectively. These developments explain why liquid, solid, gaseous and other fuels are key categories regarding trend.

The following sections describe the different source categories of 1A2 Manufacturing Industries and Construction. Further information is documented in the respective EMIS documentation(EMIS 2014/1A2x).

Iron and Steel (1A2a)

The source category 1A2a Iron and Steel includes specific information from the industry as well as information from the Prognos model.

There is no primary iron and steel production in Switzerland. Only secondary steel and iron production using recycled steel scrap occurs.

Iron is produced in 14 iron foundries. About 75% of the iron is processed in induction furnaces and 25% in cupola furnaces. The share of induction furnaces increased since 1990 from 47% with a sharp increase in 2009 based on the closure of at least one cupola furnace. Induction furnaces use electricity for the melting process and therefore only process emissions occur, which are reported in source category 2C1 Iron and steel production. Within the production process iron foundries add bituminous coal in the production process to increase the carbon content of the raw material steel scrap to produce iron with higher carbon content. The use of bituminous coal decreased significantly from 434 TJ in 1990 to 55 TJ in 2012. This is due to the significant decrease of iron funding by 73% from 1990 to 2012 and the switch from cupola furnances to induction furnances.

Today, in Switzerland steel is produced in two steel production plants. Both plants use electric arc furnaces (EAF) with carbon electrodes for melting the steel scrap. Therefore only emissions from the heating furnaces are included in source category 1A2a. These furnaces use mainly natural gas for reheating the ingot moulds prior to the rolling mills. Process emissions from steel production are included in source category 2C1 Iron and steel production. Steel production and the related natural gas consumption was significantly reduced in 1995 through the closure of two steel companies. Since 1995, steel production increased continuously until 2004 to reach the same production level as 1990. Since then, steel production is constant. Only in 2009, the production was significantly lower based on economic crisis. One steel producer switched its production to high quality steel and therefore the specific energy use per tonne of steel produced increased between 1995 and 2000. This led to a higher natural gas consumption.

Fuel consumption of source category 1A2a represents 5% of overall industry fuel consumption in 2012. As displayed in

Table 3-26, natural gas (83%), liquefied petroleum gas (10%), gas oil (6%) and bituminous coal (1%) are used in source category 1A2a.

Fuel consumption increased within this source category by 35%. Nevertheless, there has been a major change in the fuels used in the processes. The consumption of bituminous coal decreased by 87% based on the reduced iron production and the switch from cupola to induction furnaces in iron foundries. Gas oil and liquefied petroleum gas consumption also decreased by 40% and 28% respectively. Natural gas consumption increased by a total of 135% due to the switch to high quality steel production. These changes in fuel consumption result in an increase of GHG emissions by approximately 17%, significantly less than the fuel increase.

Non-ferrous Metals (1A2b)

The source category 1A2b Non-ferrous metals is based on specific information from the industry as well as information from the Prognos model and includes aluminium remelting plants as well as non-ferrous metal foundries, producing mainly copper alloys.

Until 1993, aluminium remelting plants have been in operation using gas oil. On the other hand emissions from primary aluminium production in Switzerland are reported in source category 2C3 as induction furnaces used. Its last production site closed down in April 2006.

Regarding non-ferrous metal industry in Switzerland, only casting and no production of non-ferrous metals occur. There is one large and several small foundries which are organized within the Swiss foundries association (Schweizerischer Giessereiverband GVS) which both provide annual production data.

Fuel consumption of source category 1A2b represents only 1% of the overall industry fuel consumption in 2012. As shown in Table 3-26, the fuels consumed in 2012 are mainly natural gas (82%), gas oil (17%) and liquefied petroleum gas (1%). Fuel consumption within this source category 1A2b decreased by 47% from 1990 to 2012 (see Table 3-25).

This is due both to the closure of the aluminium remelting plants and the strong reduction of 78% of the non-ferrous metal production since 2000. In the same time the consumption of gas oil and LPG went down by 65% and 72% respectively. The consumption of natural gas decreased by 40%. These developments result in a reduction of GHG emissions of 1A2b by approximately 52%.

Chemicals (1A2c)

The source category 1A2c Chemicals includes specific information from the industry as well as information from the Prognos model.

In Switzerland, there are more than thirty chemical companies mainly producing fine chemicals and pharmaceuticals. Fossil fuels are mainly used for steam production and process heat. The process emissions from the production of basic chemicals, i.e. ammonia, nitric acid, ethylene, acetic acid and sulphuric acid as well as silicon carbide are reported in source category 2B, see Section 4.3.

There is one large company producing ammonia and ethylene by thermal cracking of liquefied petroleum gas and light virgin naphtha. As by-products from the cracking process, so-called heating gas (liquefied petroleum gas) and gas oil are produced which are used thermally for steam production within the same plant and are accounted for within source category 1A2c (see Section 3.2.3).

Fuel consumption within 1A2c accounts for 13% of the overall industry fuel consumption in 2012. The fuels consumed in 2012 included natural gas (75%), gas oil (22%) and LPG (3%) (see Table 3-25). Fuel consumption in this source category has decreased by 20% between 1990 and 2012. Together with a fuel switch from residual fuel oil (-100%) and gas oil (-31%) to LPG (+18%) and natural gas (+5%) the GHG emissions have decreased by 23%.

Pulp, Paper and Print (1A2d)

The source category 1A2d Pulp, Paper and Print is based on specific information from the industry as well as information from the Prognos model and includes the fuel emissions from the Swiss pulp and paper industry and printing facilities.

Around half a dozen paper producers and several printing facilities exist in Switzerland. The only cellulose production plant was closed in 2008. Thermal energy is mainly used for provision of steam used in the drying process within paper production.

Source category 1A2d represents 4% of the overall fuel consumption in source category 1A2 in 2012. The fuels used in 2012 are mainly natural gas (87%) and gas oil (11%) (see Table 3-25). In this category only biomass from cellulose production (until 2008) is included, based on data from the only production site. Biomass used in paper production is reported in source category 1A2fi, because no comprehensive data exists to distribute biomass consumption to the specific industries within 1A2 as explained in Section 3.2.5.3.

The overall fuel consumption within the Swiss pulp and paper industry has decreased by approximately 68%, basically due to the closure of the cellulose production plant in 2008 and the closure of several paper producers in the last years. Since 1990 liquid fuels such as residual fuel oil, LPG and gas oil have decreased by 100%, 47% and 46%, respectively. In the same time, natural gas consumption increased by 13%. Biomass consumption decreased by 100% as the cellulose production was in 2008. GHG emissions have also decreased by about 67%.

Food Processing, Beverages and Tobacco (1A2e)

The source category 1A2e Food Processing, Beverages and Tobacco is based on information from the Prognos model.

In Switzerland, the source category 1A2e Food, beverages and tobacco includes around 200 companies. According to the national food industry association, the major part of revenues is provided by meat production (22%), milk products (18%) and convenience food (13%). Other productions are chocolate, sugar or baby food (Fial 2013). There are also some tobacco production sites, which are usually rather small farms of not more than 2 ha. Fossil fuels are used for steam production and drying processes.

Source category 1A2e accounts for 14% of the overall fuel consumption in source category 1A2 Manufacturing Industries and Construction in 2012. The fuels used in this category in the year 2012 were mainly natural gas (55%), gas oil (38%) and LPG (7%) (see Table 3-25).

Source category 1A2e shows an increase in fuel consumption of 35% between 1990 and 2012. This is based on the increased production of this sector in Switzerland. The consumption of residual fuel oil and gas oil have decreased by 100% and 33% respectively, whereas LPG consumption increased by a factor 3 and natural gas consumption by a factor 7.6. This has led to a total increase of GHG emissions by 18%.

Other – Stationary (1A2fi)

The source category 1A2fi Other – Stationary is based on specific information from the industry as well as information from the Prognos model.

Source category 1A2fi Other stationary includes several large fuel consumers mainly from mineral industry, i.e. cement, brick and tile, container glass and rock wool production as well as fibre board production. Additionally, the use of biomass such as wood, biogas and sewage sludge from all industry sectors is entirely reported in this source category. 1A2 Other – Stationary accounts for approximately 61% of the overall fuel consumption in 1A2 Manufacturing Industries and Construction.

Fuel consumption in 2012 comprises mainly natural gas (22%), gas oil (21%), biomass (14%), other fuels (fossil waste, 13%) and bituminous coal (6%) (see Table 3-25).

The most important industry within this category is cement production with approximately 25% of the total fuel consumption in 1A2fi. The fuels consumed in this category are very diverse and depend on the fuel use within the specific industry (see detailed documentation below). Between 1990 and 2012 there has been a switch in fuel consumption from liquid and solid fuels with shares of 49% and 30%, respectively, to liquid fuels, biomass and natural gas with shares of 33%, 26% and 24%, respectively (Table 3-25).

The consumption of residual fuel oil and liquefied petroleum gas decreased by 85% and 30%, respectively whereas gas oil consumption increased by 121%. Solid fossil fuels consumption also decreased by 55%. Natural gas, other fuels and biomass consumption increased by factors 5, 3.8 and 2.2, respectively. This fuel switch results in a reduction of emissions from this category by 3% although the overall fuel consumption increased by 25%. Specific industry developments and information on fuel use are described in the following sub chapters.

Cement

In Switzerland, there are six plants producing clinker and cement. The Swiss plants are rather small and do not exceed a production capacity of 3'000 tonnes of clinker per day. All of them use modern dry process technology. Cement industry emissions stem from incineration of a wide variety of fossil and waste derived fuels used to generate high temperatures needed for the calcination process.

Emission Factors

Specific emission factors for the cement industry are shown in the tables below:

Table 3-27 Emission factors for cement industry in 2012. Source: EMIS data base (EMIS 2014/1A2f). Emission factors for CO₂ are fuel specific (see Table 3-28).

| Cement industry (part of 1A2f) | CO ₂ | N ₂ O | CH ₄ | NO _x | CO | NM VOC | SO ₂ |
|--------------------------------|-----------------|------------------|-----------------|-----------------|-------|--------|-----------------|
| | t/TJ | | g/t cement | | | | |
| Cement | fuel specific | | 4 | 832 | 1'400 | 45 | 359 |

As described above in Table 3-25, the CH₄ emission factor includes the overall CH₄ emissions of the cement industry based on direct exhaust measurements at the chimneys of the cement plants. Therefore these CH₄ emissions are reported under the category "Other Fuels" in the CRF-tables.

Table 3-28 CO₂ Emission factors and other characteristics of waste derived fuels (Other fuels and Biomass) used in the cement industry.

| | NCV | EF CO ₂ Tot. | Fraction biomass-C |
|--|-------|-------------------------|--------------------|
| | MJ/kg | kg CO ₂ /GJ | % |
| Waste derived fuel | | | |
| Waste oil | 32.48 | 74.35 | 0 |
| Waste coke from coke filters | 23.70 | 97.00 | 0 |
| Mixed industrial waste | 18.34 | 74.00 | 0 |
| Other fossil waste fuels | 20.85 | 97.00 | 0 |
| Solvents and residues from distillation | 23.63 | 73.99 | 1 |
| Waste Tyres and rubber | 26.40 | 84.00 | 27 |
| Plastics | 25.24 | 84.66 | 28 |
| Mix of special waste with saw dust (CSS) | 9.22 | 102.40 | 78 |
| Sewage sludge (dried) | 9.39 | 94.52 | 100 |
| Wood | 16.26 | 99.90 | 100 |
| Animal meal | 16.81 | 86.66 | 100 |
| Sawdust | 16.26 | 99.90 | 100 |
| Agricultural waste / other biomass | 12.72 | 110.00 | 100 |

The NCVs and CO₂ emission factors for waste oil, solvents, plastics, CSS sewage sludge, animal meal and sawdust are based on a study of Cemsuisse (Cemsuisse 2010a). The values for waste tyres are taken from Hackl, A., Mauschwitz, G. (2003). The biogenic fraction of waste tyres is also based on an Austrian study and published by the German Ministry of Environment (UBA 2006).

Activity Data

Fossil fuels used in cement industry are coal (bituminous coal and lignite), petroleum coke and, to a lesser extent, residual fuel oil, natural gas and gas oil. In addition, also fossil and biogenic waste derived fuels are used. Fossil wastes comprise solvents and residues from distillation, waste tyres and rubbers, plastics and waste oil whereas biogenic wastes contain mainly waste wood, animal residues and sewage sludge.

The amount of fossil and waste derived fuels consumed in cement industry is shown in Table 3-29. Data is provided by cemsuisse and documented in the respective EMIS/1A2fi Zementwerke Feuerung.

Table 3-29 Activity data: Overview on fuel use in 1A2fi cement industry.

| Source | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|------------------------------|------|--------|--------|--------|--------|--------|--------|--------|--------|-------|-------|
| Cement industry | | | | | | | | | | | |
| Cement, total incl. waste | TJ | 16'140 | 14'195 | 12'878 | 11'450 | 12'997 | 12'291 | 10'647 | 10'080 | 9'860 | 9'853 |
| Cement fossil without waste | TJ | 14'265 | 12'295 | 10'951 | 9'071 | 10'856 | 9'511 | 7'326 | 6'822 | 6'489 | 6'624 |
| Residual fuel oil | TJ | 1'907 | 2'957 | 4'377 | 3'263 | 4'589 | 2'825 | 3'507 | 3'206 | 3'168 | 3'260 |
| Petroleum coke | TJ | 900 | 670 | 50 | 500 | 980 | 830 | 550 | 240 | 410 | 466 |
| Bituminous coal | TJ | 10'790 | 8'300 | 6'150 | 5'000 | 5'000 | 5'500 | 3'000 | 3'200 | 2'710 | 2'640 |
| Lignite | TJ | 306 | 353 | 306 | 259 | 259 | 188 | 236 | 165 | 139 | 136 |
| Gas | TJ | 362 | 14 | 67 | 48 | 27 | 168 | 34 | 10 | 62 | 121 |
| Cement, waste derived fuel | TJ | 1'874 | 1'901 | 1'927 | 2'379 | 2'142 | 2'780 | 3'321 | 3'258 | 3'371 | 3'229 |
| Waste oil | TJ | 1'169 | 1'137 | 1'104 | 1'527 | 1'208 | 1'485 | 1'514 | 1'257 | 1'509 | 1'403 |
| Sewage sludge (dried) | TJ | 9 | 9 | 9 | 19 | 65 | 128 | 175 | 240 | 216 | 279 |
| Wood | TJ | 0 | 0 | 0 | 0 | 106 | 321 | 395 | 319 | 0 | 0 |
| Solvents and residues from | TJ | 284 | 378 | 473 | 284 | 127 | 181 | 274 | 410 | 375 | 272 |
| Waste tyres and rubber | TJ | 330 | 304 | 277 | 441 | 402 | 415 | 420 | 366 | 363 | 321 |
| Plastics | TJ | 0 | 0 | 0 | 0 | 27 | 55 | 177 | 274 | 508 | 553 |
| Animal meal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 197 | 233 | 223 | 211 |
| Mix of special waste with s | TJ | 23 | 14 | 5 | 51 | 147 | 136 | 111 | 100 | 118 | 132 |
| Waste coke from coke filte | TJ | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 | 59 |
| Sawdust | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Mixed industrial waste | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other fossil waste fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Agricultural waste / other b | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Source | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|------------------------------|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cement industry | | | | | | | | | | | |
| Cement, total incl. waste | TJ | 10'582 | 11'054 | 10'756 | 10'490 | 11'226 | 11'551 | 11'663 | 12'022 | 11'954 | 11'816 |
| Cement fossil without waste | TJ | 6'897 | 6'553 | 5'754 | 5'231 | 5'459 | 6'136 | 6'344 | 6'914 | 6'389 | 6'127 |
| Residual fuel oil | TJ | 1'530 | 1'194 | 1'079 | 621 | 754 | 637 | 220 | 175 | 135 | 100 |
| Petroleum coke | TJ | 458 | 327 | 590 | 187 | 515 | 638 | 903 | 912 | 1'036 | 994 |
| Bituminous coal | TJ | 4'750 | 4'950 | 3'980 | 4'330 | 4'080 | 4'120 | 3'383 | 4'033 | 3'618 | 3'650 |
| Lignite | TJ | 136 | 71 | 94 | 94 | 94 | 737 | 1'834 | 1'790 | 1'596 | 1'379 |
| Gas | TJ | 22 | 11 | 11 | 0 | 16 | 4 | 4 | 4 | 4 | 4 |
| Cement, waste derived fuel | TJ | 3'686 | 4'501 | 5'002 | 5'258 | 5'767 | 5'415 | 5'319 | 5'108 | 5'565 | 5'689 |
| Waste oil | TJ | 1'519 | 1'341 | 1'583 | 1'489 | 1'536 | 1'411 | 1'279 | 844 | 866 | 1'278 |
| Sewage sludge (dried) | TJ | 332 | 348 | 360 | 386 | 407 | 494 | 560 | 549 | 511 | 475 |
| Wood | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 61 |
| Solvents and residues from | TJ | 427 | 517 | 726 | 740 | 1'002 | 976 | 981 | 1'295 | 1'476 | 1'032 |
| Waste tyres and rubber | TJ | 421 | 476 | 460 | 568 | 519 | 645 | 568 | 525 | 794 | 828 |
| Plastics | TJ | 572 | 600 | 527 | 525 | 770 | 841 | 926 | 1'013 | 995 | 1'119 |
| Animal meal | TJ | 198 | 1'030 | 1'172 | 1'379 | 1'326 | 856 | 799 | 664 | 658 | 621 |
| Mix of special waste with s | TJ | 158 | 130 | 116 | 114 | 163 | 133 | 146 | 164 | 157 | 131 |
| Waste coke from coke filte | TJ | 59 | 59 | 59 | 59 | 46 | 58 | 60 | 0 | 0 | 0 |
| Sawdust | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Mixed industrial waste | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 1 |
| Other fossil waste fuels | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 48 | 105 | 137 |
| Agricultural waste / other b | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 2 | 7 |

| Source | Unit | 2010 | 2011 | 2012 |
|------------------------------|------|--------|--------|--------|
| Cement industry | | | | |
| Cement, total incl. waste | TJ | 12'382 | 12'187 | 11'495 |
| Cement fossil without waste | TJ | 6'273 | 5'858 | 5'566 |
| Residual fuel oil | TJ | 112 | 101 | 297 |
| Petroleum coke | TJ | 1'130 | 1'081 | 1'081 |
| Bituminous coal | TJ | 3'662 | 3'167 | 3'097 |
| Lignite | TJ | 1'348 | 1'493 | 1'081 |
| Gas | TJ | 21 | 16 | 11 |
| Cement, waste derived fuel | TJ | 6'109 | 6'329 | 5'929 |
| Waste oil | TJ | 1'253 | 1'170 | 839 |
| Sewage sludge (dried) | TJ | 477 | 483 | 527 |
| Wood | TJ | 292 | 409 | 586 |
| Solvents and residues from | TJ | 1'189 | 1'264 | 1'294 |
| Waste tyres and rubber | TJ | 842 | 1'033 | 964 |
| Plastics | TJ | 1'252 | 1'163 | 964 |
| Animal meal | TJ | 624 | 614 | 572 |
| Mix of special waste with s | TJ | 123 | 96 | 100 |
| Waste coke from coke filte | TJ | 0 | 0 | 0 |
| Sawdust | TJ | 6 | 24 | 17 |
| Mixed industrial waste | TJ | 0 | 0 | 0 |
| Other fossil waste fuels | TJ | 45 | 55 | 36 |
| Agricultural waste / other b | TJ | 7 | 18 | 28 |

As shown in Table 3-29, in 2012 the Swiss cement industry used about 48% fossil fuels and 36% fossil and 16% biogenic waste derived fuels. The most important fossil fuels in 2012 were bituminous coal (27%), petroleum coke (9%) and lignite (9%).

Fuel consumption in cement plants has decreased by 39% between 1990 and 2012. This is partly due to a decrease in production since 1990 by about 15% and an increase in energy efficiency. In the same period the fuel mix has changed significantly from the use of mainly fossil fuels (88%) and some fossil waste derived fuels (11%) to the above mentioned mix of fossil fuels (48%), fossil and biogenic waste derived fuels (36% and 16%, respectively). The fossil fuels used in 1990 were bituminous coal, residual fuel oil and petroleum coke with shares of 67%, 12% and 6%, respectively, of the total fuel consumption.

Please note that all fossil waste derived fuels are reported as so-called Other Fuels in the CRF-tables, whereas the biogenic waste derived fuels belong to Biomass.

Lime

In Switzerland there is only one plant producing lime. Fossil fuels are used for the burning process (calcination) of limestone.

Since 1994 fuel consumption in lime production is mainly based on residual fuel oil with a share of 99% in 2012. Since 1995, no petroleum coke is used anymore as it was replaced by residual fuel oil.

The fuel consumption of two sugar plants that autoproduce lime is reported in category 1A2e.

Fine ceramics

In Switzerland, the main production of fine ceramics is sanitary ware produced by one big and some small companies. In earlier years, also other ceramics were produced as for example glazed ceramics tiles, electrical porcelain and earthenware. Since 2001, only sanitary ware is produced.

Current fuel consumption within fine ceramics production is mainly natural gas (99%). In 2001 the fuel-mix was natural gas (62%) and gas oil (38%). Since then, fuel consumption has shifted strongly to the use of natural gas. Compared to the production of other fine ceramics, the production of sanitary ware is more energy-intensive. Therefore, the specific energy use per tonne of produced fine ceramics has increased since 1990. This results in a lower reduction of fuel consumption (68%) compared to the reduction in production (80%) between 1990 and 2012.

Rock wool

In Switzerland there is one single producer of rock wool. Fossil fuels are used for the melting of rocks at a temperature of 1500 °C in cupola furnaces.

Currently bituminous coal (83%) and natural gas (17%) are used in the production process. Until 2004 also gas oil and liquefied petroleum gas were used which were substituted by natural gas in 2005. Rock wool production has increased by approximately 48% from 1990 to 2012 whereas the fuel consumption has increased by 15% only.

Brick and tile

In Switzerland there are about 20 plants producing bricks and tiles. Fossil fuels are used for drying and burning of the clay blanks.

Fuels used in the brick and tile production in 2012 are natural gas (66%), residual fuel oil (22%) as well as small amounts of gas oil (4%), liquefied petroleum gas (3%) and residual

fuel oil (3%). Apart from a production recovery in the years around 2004 the production has gradually decreased since 1990 by about 36% which is also represented in the overall fuel consumption decrease of about 37%. Regarding the fuels used, there has been a considerable shift from residual fuel oil to natural gas from 1990 onwards as well as a minor shift from gas oil and liquefied petroleum gas to natural gas from 2004 onwards. Paper production residues, wood and animal grease are used since 2000.

The emission factors for wood and the biogenic waste, i.e. paper production residues, and animal grease used in brick and tile production are 92kg/GJ, 86 kg/GJ and 81kg/GJ, respectively for animal grease (see documentation EMIS/1A2fi Ziegeleien).

Mixed goods

The production of mixed goods mainly includes the production of bitumen for road paving. A total of 110 production sites are producing mixed goods at stationary production sites.

The main fuel used is gas oil (71%) and natural gas (22%). The specific fuel consumption per ton of mixed goods was assumed to be constant between 1990 and 2012 and production of mixed goods oscillates around five million tons per year. There has been a fuel switch from gas oil (reduction of 23%) to natural gas (increase of 40%) in this time.

Glass

In Switzerland glass production includes three types of glass: container glass, tableware glass and glass wool. Today there exist only one production plant for container glass and one for tableware glass. Glass wool is produced in two plants.

In 2012 fuel consumption for container glass production includes mainly residual fuel oil (80%) and natural gas (20%). Since 1990, fuel consumption for container glass has drastically decreased due to reduction in production. Until 2003 only residual fuel oil was used in container glass production. Since 2004 the share of natural gas has increased to reach a stable contribution of 20% from 2006 onwards.

Fuel consumption for tableware glass currently includes only liquefied petroleum gas. Since 1990 fuel consumption for tableware glass strongly decreased because of the closure of one production plant in 2002 and another one in 2006. In addition, the consumption of residual fuel oil is eliminated since 2000 (17% in 1990 and 21% in 1995).

Fuel consumption for glass wool production includes currently only natural gas. Production of glass wool has increased since 1990 by approximately 60%, but the natural gas consumption decreased by approximately 10%. This can be explained by an increase in energy efficiency in the production process, i.e. a decrease of the specific energy consumption from 8.5 GJ/t to 4.8 GJ/t from 1990 to 2012.

Fibre board

Fibre board is produced in two companies in Switzerland. Thermal energy is used for heating and drying processes.

Current fuels used for fibre board production are waste wood (85%), natural gas (7%), residual fuel oil (5%) and animal grease (3%). Since 1990 the production of fibre board and thus the fuel consumption have increased significantly by a factor of 5. The fuel mix has strongly shifted between 1990 and 2012 from fossil fuels to biomass, i.e. a reduction in fossil fuel share from 65% to about 50% between 1996–2002 and from about 50% to 12% from 2006 to 2012. In 1990 the share of natural gas, wood waste and residual fuel oil amounted to 48%, 35% and 13%, respectively. Since 2001 also animal grease is used for fibre board production.

Other – Mobile (1A2fii)

Source category 1A2fii accounts for diesel and gasoline for mobile construction and industrial machinery (off-road). The most relevant mobile construction machines in terms of fuel consumption are excavators, loaders, dump trucks and mobile compressors. In the industry sector, forklifts and snow groomers consume the most fuel, but the share of electrical forklifts has been gradually increasing. Almost all fuel consumed in the sector (98%) is diesel oil, the rest is gasoline and natural gas. Activity data are taken from INFRAS (2008) and Keller/INFRAS (2013).

3.2.7.3 Uncertainties and Time-Series Consistency

The uncertainty of CO₂ emissions from fuel combustions is described in the uncertainty analysis of the Energy Industries (1A1) in Chapter 3.2.6.3. Uncertainty in emissions of other non-CO₂ gases is estimated to be medium: 30% for CH₄ and 80% for N₂O (see Table 1-14).

Consistency and Completeness in 1A2 Fuel Combustion

Consistency:

- Time series for 1A2 are all consistent.

Completeness:

- All estimates in the sector 1A2 are assumed to be complete.

3.2.7.4 Source-specific QA/QC and Verification

a) General

See Chapter 3.2.6.4.

b) Specific: Manufacturing Industries and Construction (1A2)

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables.
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of last year submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of last year submission 2013.

3.2.7.5 Source-Specific Recalculations

- **1A all fuels:** Activity data of all fuels of the overall time series have been recalculated based on the new data from SFOE 2013. This includes data on 2011 for fossil fuels and waste, data for biomass on 1997, 2009 – 2011 and data of other fuels 1990 – 2011.
- **1A gaseous fuels:** Activity data have been recalculated based on new data of from SFOE 2013.
- **1A wood consumption:** Activity data have been recalculated for the overall time series based on the new data from SFOE 2013b.
- **1A gas oil:** SO_x emission factor value has been updated for 2010 based on sulphur analyses of the gas oil for the year 2010 (Directorate General of Customs) resulting in a revised value for 2011 as well.

- **1A bituminous coal:** CO₂ emission factor has been corrected from 94t CO₂/TJ to 92.7 t CO₂/TJ for the entire time series.
- **1A2 liquefied petroleum gas:** N₂O emission factor of liquefied petroleum gas has been changed for the whole time series from 0.6 g/GJ to the IPCC 2006 emission factor of 0.1 g/GJ. In previous submissions, the emission factor for gas oil was used for liquefied petroleum gas because they have been reported jointly. This has been corrected in the present submission.
- **1A2 gas oil:** Since SO_x emission factor values for 2010 and 2011 have been updated based on sulphur analyses of gas oil for the year 2010 (Directorate General of Customs) also the SO_x emission factor values of liquefied petroleum gas for 2010 and 2011 have been revised as for all air pollutants the same EF are assumed as for gas oil.
- **1A2a iron foundries:** Activity data of bituminous coal has been revised due to corrected production shares of cupola and electric furnaces in iron foundries for 2010 and 2011. This change resulted in new activity data of bituminous coal in heat furnances for the same years.
- **1A2c cracker-by-product:** Activity data of steam production from the cracker-by-products has been updated for 1990 - 1999 based on new net calorific values. For calculation of the light virgin naphtha consumed as cracker feedstock the so far used net calorific value of gasoline has been replaced by the value for naphtha according to the 2006 IPCC guidelines resulting in revised activity data for 1990-1999.
- **1A2c cracker process:** Activity data of liquefied petroleum gas has been updated for 1990-1999 and 2011 based on new activity data from the cracker process and resulting steam production.
- **1A2fi glasswool production:** Activity data of one of the two glass wool production plants have been revised for 1991-2004 based on effective production data for 1996-2004 resulting in revised gas consumption.
- **1A2fi brick and tile and glasswool production:** Activity data of all fuels for 1991-2011 have been updated based on new production data.
- **1A2fi gas oil and natural gas:** Activity data from gas oil and natural gas have been revised for 1990-2011 by the subtraction of nonroad from "Heizkessel GLD, HEL" and "Heizkessel GLD, Gas" instead of "Industrie Heizkessel weitere, HEL" and "Industrie Heizkessel weitere, Gas" within the Energy model.
- **1A2fi cement production:** Newly also CH₄ emissions are reported with a CH₄ emission factor of 5 g/t cement from 1990 to 1995 and 4 g/t cement from 2002 onwards.
- **1A2fi brick and tile production:** Activity data has been updated for 2001-2006 using effective production data instead of interpolated data. In addition, also so far interpolated fuel consumptions for these three plants have been replaced by effective values for 2000 and 2007-2011 resulting in overall revised fossil and biogenic fuel consumptions for 1991-2011 and 2000-2011.
- **1A2fi biogas from wastewater treatment plants:** Activity data has been updated for 2008, 2009 and 2011 based on recalculations in SFOE 2013a.
- **1A2fii Off-road:** Diesel and gasoline consumption is based on INFRAS (2008) and on an update carried out in 2013 based on the latest figures on economy and population (Prognos 2012a, Keller/INFRAS 2013). The consumption has been recalculated accordingly. Numbers from 2005 onwards are affected.

3.2.7.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.2.8 Source Category 1A3 - Transport

3.2.8.1 Source Category Description:

Tier 1 Key Categories 1A3

CO₂ from the combustion of gasoline (level and trend)
 CO₂ from the combustion of diesel (level and trend)
 CO₂ from the combustion of natural gas (trend)
 CH₄ from the combustion of gasoline (trend)
 N₂O from the combustion of gasoline (trend)

Tier 2 Key Categories 1A3

CO₂ from the combustion of gasoline (level and trend)
 CO₂ from the combustion of diesel (level and trend)
 CH₄ from the combustion of gasoline (trend)
 N₂O from the combustion of gasoline (trend)

The source category includes civil aviation, road transportation, railways, navigation and other transportation. Further off-road transportation is included in category 1A2 Manufacturing Industries and Construction, in 1A4 Other Sectors and 1A5 Other (Military). For information on bunker fuel emissions from international aviation and navigation, see Chapter 3.2.2.

Table 3-30 Specification of Swiss source category 1A3 Transport. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1A3 | Source | Specification | Data Source |
|------|---------------------------|---|--|
| 1A3a | Civil Aviation (National) | Large (jet, turboprop) and small (piston) aircrafts, helicopters | AD: SFOE 2013, FOCA 2006, 2006a, 2007, 2008, 2009, 2010, 2011, 2012, 2013 |
| 1A3b | Road Transportation | Light and heavy motor vehicles, coaches, two-wheelers | AD: SFOE 2013, SFCA 2013, SFSO 2013b; Method, EF: INFRAS 2010, INFRAS 2011, FOEN 2010a, Hausberger et al. 2009, EMPA 2009 AD: ARE 2012 |
| 1A3c | Railways | Diesel locomotives, abrasion by merchandise and person traffic | Method, AD, EF: INFRAS 2008, AD: Prognos 2012, Keller/INFRAS 2013 |
| 1A3d | Navigation (National) | Passenger ships, motor and sailing boats on the Swiss lakes and the river Rhine | Method, AD, EF: INFRAS 2008, AD: Prognos 2012, Keller/INFRAS 2013 |
| 1A3e | Pipeline Compressors | Compressor station in Ruswil, Lucerne | AD: SFOE 2013 EF: Battelle 1994, SAEFL 2000, SGWA 2007, Xinmin 2004 |

3.2.8.2 Methodological Issues

In Switzerland, Transport (1A3) contains the sub-categories

- Aviation (1A3a, national/domestic civil aviation),
- Road Transportation (1A3b),
- Railways (1A3c),
- Navigation (1A3d, national/domestic navigation).
- Compressor station for gas distribution (1A3e)

a) Aviation (1A3a)

Tier 1 Key Categories 1A3a

CO₂ from the combustion of fuel (kerosene) in civil aviation (trend)

Tier 2 Key Categories 1A3a

There are no Tier 2 Key categories in 1A3a

The emissions of civil aviation are modelled by a Tier 3a method developed by FOCA (2006). FOCA is represented in the emissions technical working group (CAEP WG3) and in the modelling and database group (CAEP MDG) of the International Civil Aviation Organisation (ICAO). FOCA is directly involved in the development of ICAO guidance material for the calculation of aircraft emissions and in the update of the IPCC guidelines (via the secretariat of ICAO CAEP (Committee on Aviation Environmental Protection)). The Tier 3a method applied for the emission modelling is in line with the methods developed in the working groups mentioned. Note that the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006) have been prepared by the IPCC Task Force on National Greenhouse Gas Inventories and have been adopted in April 2006 by the IPCC. Under the UNFCCC, they have not yet been adopted for mandatory use in reporting on GHG inventories. Formally, the method therefore should be considered as a country specific method until improvement. The modelling scheme for civil aviation starting with aircraft basic data, activity data, emission factors and ending with emissions imported into EMIS database is shown in Figure 3-14.

The Tier 3a method follows standard modelling procedures on the level of single movements based on detailed movement statistics. The primary key for all calculations is the aircraft tail number, which allows to calculate on the most precise level, namely on the level of the individual aircraft and engine type. Every aircraft is linked to the FOCA engine data base containing emission factors for more than 600 individual engines with different power settings. Emissions in the landing and take-off cycle (LTO) are calculated with aircraft category dependant flight times and corresponding power settings. Cruise emissions are calculated based on the individual aircraft type and the trip distance for every flight. For piston-engine powered aircraft and helicopters, to the knowledge of FOCA, it has been the only provider of publicly available engine data and a full methodology, so far. All piston engine data and study results have been published in 2007 (FOCA 2007a). The guidance on the determination of helicopter emissions has been published in 2009 (FOCA 2009a).

The movement database from Swiss Airports contains departure and destination airport. With this information, all flights from and to Swiss airports are separated into domestic (national) and international flights prior to the emission calculation. The emissions of domestic flights are reported under 1A3a Civil Aviation, the emissions of international flights are reported under international bunker emissions (memo items).

The emission factors used are country specific or are taken from the ICAO engine emissions databank, from EMEP/CORINAIR databases (EEA 2002), Swedish Defence Research Agency (FOI) and Swiss FOCA measurements (precursors). Cruise emission factors are generally calculated from the values of the ICAO engine emissions databank, adjusted to cruise conditions by using the Boeing Fuel Flow Method 2. For N₂O, the IPCC default emission factor is used. Activity data are derived from a detailed movement statistics.




Figure 3-14 Modelling scheme (activity data, emission factors, emissions) for civil aviation.

A complete emission modelling (LTO and cruise emissions for domestic and international flights) has been carried out by Swiss FOCA for 1990, 1995, 2000, 2002, 2004–2012. The results of the emission modelling have been transmitted from FOCA to FOEN in an aggregated form. FOEN (the NIC) calculated the implied emission factors 1990, 1995, 2000, 2002, 2004 and carried out a linear interpolation for the years in-between. The interpolated implied emission factors were multiplied with the annual fuel sold from Swiss overall energy statistics (SFOE in respective years), providing the missing emissions of civil aviation for the years 1991–1994, 1996–1999, 2001 and 2003.

Details of emission factors and activity data follow below. Further tables containing more information are also given in Annex A3.1.4, more detailed descriptions of the emission modelling may be found in FOCA (2006).

Emission Factors

Kyoto gases:

- **CO₂:** The value of 73.2 t/TJ is country specific and is based on measurements and analyses of fuel samples (see Table 3-9). Small yearly variations have been neglected so far.
- **CH₄, NMVOC (country specific; CORINAIR):** VOC emissions (see Precursors below) are split into CH₄ and NMVOC by a constant share of 0.1 (CH₄) and 0.9 (NMVOC)⁹. For CH₄, the average emission factor for domestic flights is 2.0 kg/TJ in 2012 average LTO is 3.7 kg/TJ (international airports only), cruise 0.74 kg/TJ (international airports only) (FOCA 2013).
- **N₂O:** The IPCC default value 2.3 kg/TJ is used for the whole period 1990–2012 (IPCC 1997b).

SO₂ (IPCC):

- The emission factor is taken from the IPCC Guidelines 1996, 23.0 kg/TJ, and is assumed to be constant over the period 1990–2012 (IPCC 1997c, chapter 1.4.2.6)

Precursors (country specific; CORINAIR):

- **Assignment of emission factors for 1990 and 1995:** The fleet that operated in and from Switzerland during those years has been analysed. The corresponding most frequent engines within an aircraft category (ICAO Code) have been assigned to every aircraft type.
- **Assignment of emission factors for the year 2000, 2002, 2004 to 2012:** The actual engine of every single aircraft operating in and from Switzerland has been assigned. FOCA uses the aircraft tail number as the key variable which links activity data and individual aircraft engine information (see Annex A3.1.4 Table A-11 Aircraft Engine Combinations).

FOCA determines the emission factors of different precursors such as NO_x, VOC, CO and other pollutants as follows:

⁹ The share of 0.1 for methane is maintained until general acceptance of necessary corrections is reached. Studies indicate that during cruise, Methane exhaust concentrations are lower than Methane ambient concentrations, see Wiesen et al. (1994), Spicer et al. (1994) and Knighton et al. (2009). A first remark has been made in Table 1-52 of the IPCC Guidelines 1996.

LTO:

The Swiss FOCA engine emissions database consists of more than 600 individual engine data sets. Jet engine factors for engines above 26.7 kN thrust (emission certificated) are identical to the ICAO engine emissions databank. Emission factors for lower thrust engines, piston engines and helicopters were taken from manufacturers or from own measurements. Emission factors for turboprops could be obtained in collaboration with the Swedish Defence Research Agency (FOI).

Cruise:

The fuel flows of the whole Airbus fleet (which produces a great portion of the Swiss inventory) have been modelled on the basis of real operational aircraft data from flight data recorders (FDR) of Swiss International Airlines. Pollutant emission factors have been modelled on the basis of the ICAO engine databank and corrected to cruise conditions using FDR engine parameters and the Boeing Fuel Flow Method 2. Part of the cruise emission factors are taken from EMEP/CORINAIR (EEA 2002) and from former CROSSAIR (FOCA 1991). Other missing aircraft types have been modelled on the basis of FOCA aircraft performance modelling and the ICAO engine emissions databank, using the Boeing Fuel Flow Method 2, as well. For piston engine aircraft and helicopters, Swiss FOCA has produced its own data, which were taken under real flight conditions (2005 data, FOCA 2009a).

Activity data**Scheduled and charter aviation**

The statistical basis has been extended after 1996. Therefore, the modelling details are not exactly the same for the years 1990-1995 as for the subsequent years. The source for the 1990 and 1995 modelling is the movement statistics, which records information for every movement on airline, number of seats, Swiss airport, arrival/departure, origin/destination, number of passengers, distance. From 1996 onwards, every movement in the FOCA statistics also contains the individual aircraft tail number (aircraft registration). This is the key variable to connect airport data and aircraft data. The statistics may contain more than one million records with individual tail numbers. All annual aircraft movements recorded are split into domestic and international flights (there are 455'422 aircraft movements in the total of scheduled and charter traffic in 2012 as given by FOCA 2013).

Non-scheduled, non-charter and General Aviation (including Helicopters)

- Airports and most of the airfields report individual aircraft data (aircraft registration). FOCA may therefore compute the inventory for small aircraft with Tier 3a method, too. However, for 1990 and 1995, the emissions data for non-scheduled, non-charter and General Aviation (helicopters etc.) could not be calculated with a Tier 3a method. Its fuel consumption is estimated to be 10% of the domestic fuel consumption. Data were taken from two FOCA studies (FOCA 1991, FOCA 1991a). For 2000-2007, all movements from airfields are known, which allows a more detailed modelling of the emissions (FOCA 2007a).
- Helicopter flights which do not take off from an official airport or airfield such as transport flights, flights for lumbering, animal transports, supply of alpine huts, heli-skiing and flight trainings in alpine regions cannot be recorded with the movement data base from airports and airfields. Although these helicopter movements only account for 0.1% of the total civil aviation emissions, these emissions are taken into account using the Unternehmensstatistik der Schweizer Helikopterunternehmen. This statistics is officially collected by FOCA and updated annually (see FOCA 2004 as illustrative example for all subsequent years).

Since 2007, the data of the Unternehmensstatistik der Schweizer Helikopterunternehmen (statistics about Swiss helicopter companies) is included electronically in the data warehouse of the model and undergoes first some plausibility checks (E-plaus software). In order to distinguish between single engine helicopters and twin engine helicopter a fix split of 87 % for single engine helicopters and 13 % for twin engine helicopters is applied for the entire commitment period based on investigations in 2004 (FOCA 2004). Note that all emissions from helicopter flights without using an official airport or an official airfield are considered as domestic emissions. There is also a helicopter base in the Principality of Liechtenstein consuming a certain very small amount of fuel contained in the Swiss statistics. Thus, its consumption leads to domestic instead of international bunker emissions (about 0.4 Gg CO₂). FOCA and FOEN decided to report these emissions as Swiss-domestic since it is a very small amount and the effort for a separation would be considerable.

Fuel consumption: Table 3-31 summarises the activity data for domestic aviation (1A3a). It also includes international aviation, which belongs to the memo items, international bunkers/aviation (see also Chapter 3.2.2).

Note that the fuel consumption reported in the CRF is identical to the consumption due to the fuel sales reported in the Swiss overall energy statistics (see e.g. SFOE 2013) while the consumptions of military aircraft and of Liechtenstein's helicopter consumption is subtracted (see section 3.2.5). In fact, the emission model run by FOCA overestimates fuel consumption by ca. 1.5%. However, the domestic fuel consumption is reported according to the modelled value (conservative estimation), whereas the international fuel consumption (bunker) is scaled downwards so that the sum of domestic and international fuel consumption becomes identical with the fuel sold, as reported in the Swiss overall energy statistics.

Table 3-31 Fuel consumption of civil aviation in TJ. The "domestic" consumption and the corresponding emissions are reported under 1A3a, the "international" consumption is reported under Memo items, international bunkers (FOCA 2007, 2007a, 2008-2013).

| 1A3a Civil Aviation | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-----------------------------|------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | Fuel consumption in TJ | | | | | | | | | |
| Total domestic (1A3aii) | 3'450 | 3'194 | 3'217 | 3'165 | 3'077 | 3'075 | 2'972 | 2'850 | 2'742 | 2'684 |
| Total international (1A3ai) | 41'884 | 40'872 | 43'499 | 45'342 | 46'840 | 49'918 | 51'975 | 53'983 | 56'599 | 60'805 |
| Sum | 45'334 | 44'067 | 46'717 | 48'508 | 49'917 | 52'993 | 54'946 | 56'833 | 59'341 | 63'489 |
| 1990 = 100% | 100% | 97% | 103% | 107% | 110% | 117% | 121% | 125% | 131% | 140% |

| 1A3a Civil Aviation | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-----------------------------|------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | Fuel consumption in TJ | | | | | | | | | |
| Total domestic (1A3aii) | 2'539 | 2'296 | 2'028 | 1'951 | 1'963 | 1'699 | 1'658 | 1'891 | 1'618 | 1'704 |
| Total international (1A3ai) | 63'687 | 60'097 | 55'468 | 49'763 | 46'896 | 47'671 | 50'109 | 53'543 | 57'844 | 55'238 |
| Sum | 66'225 | 62'393 | 57'495 | 51'714 | 48'859 | 49'370 | 51'766 | 55'434 | 59'462 | 56'942 |
| 1990 = 100% | 146% | 138% | 127% | 114% | 108% | 109% | 114% | 122% | 131% | 126% |

| 1A3a Civil Aviation | 2010 | 2011 | 2012 |
|-----------------------------|------------------------|--------|--------|
| | Fuel consumption in TJ | | |
| Total domestic (1A3aii) | 1'688 | 1'808 | 1'867 |
| Total international (1A3ai) | 58'118 | 64'060 | 63'627 |
| Sum | 59'805 | 65'868 | 65'494 |
| 1990 = 100% | 132% | 145% | 144% |

b) Road Transportation (1A3b)

Tier 1 Key categories 1A3b

CO₂ from the combustion of gasoline (level and trend)
CO₂ from the combustion of diesel (level and trend)
CO₂ from the combustion of natural gas (trend)
N₂O from the combustion of gasoline (trend)

Tier 2 Key categories 1A3bCO₂ from the combustion of gasoline (level and trend)CO₂ from the combustion of diesel (level and trend)CH₄ from the combustion of gasoline (trend)N₂O from the combustion of gasoline (trend)**Methodology**

The CO₂ emissions are calculated with a Tier 2 method (top-down) as suggested by IPCC Good Practice Guidance (IPCC 2003) using country specific emission factors. The emission factors are derived from the carbon content of fuels (see Table 3-9). The activity data corresponds to the amounts of gasoline and diesel fuel sold in Switzerland (sales principle). The numbers are taken from the national fuel statistics which is part of the Swiss overall energy statistics (SFOE 2013).

The consumption of biofuels is reported for Road Transportation as well. Fuels involved, emission factors and activity data are summarised in a comment to the EMIS database (EMIS 2013 1A3bi-viii Strassenverkehr). Most important data sources stem from the Swiss overall energy statistics (SFOE 2013) the Swiss renewable energy statistics (SFOE 2013a) and the Swiss Federal Customs Administrations (SFCA 2013).

Other gases

The other gases are modelled with a well-documented country specific method (SAEFL 1995, 2004, 2004a, FOEN 2010i, INFRAS 2010, INFRAS 2011, Hausberger et al. 2009). The approach corresponds methodologically to Box 1 in the decision tree of Figure 2.5 (p. 2.45) of IPCC Good Practice Guidance.

The emission computation is based on two sets of data:

- Traffic activity data: transport performance in vehicle kilometres (hot emissions), number of starts/stops and vehicle stock (cold start, evaporation emissions and running losses)
- Emission factors: specific pollutant emissions in grams per unit (vehicle kilometres, start/stop or vehicle)

For the calculation of emissions these two data sets are multiplied for all other gases as follows (further details of emission modelling are given in Annex A3.1.5):

$$\text{Emission (gram)} = \text{activity data (veh-km/a, starts/stops/a, vehicles)} * \text{emission factor (gram/veh-km, gram/start/stop, gram/vehicle),}$$

Activity Data

The activity data is derived from different data sources:

- Vehicle stock: The Federal vehicle registration database (SFSO 2013b) supplies the number of vehicles (including age distributions) per vehicle category¹⁰. With the help of a fleet turnover model the vehicle categories are split up into so called «sub-segments», which are used to link with the specific emission factors (vehicle category/size class/fuel type/emission concept (see also INFRAS 2010).

¹⁰ The vehicle registration in Switzerland delivers all inputs to build up the fleet composition 1990-2011 which is characterised e.g. by vehicle category, engine capacity, fuel type, total weight, vehicle age and exhaust technology.

- **Transport performance:** The transport performance (mileage) is calculated from the specific mileage per vehicle (based on household surveys/Mikrozensus ARE/SFSO 2005) times the number of vehicles. This figure is calibrated to the official statistics of traffic performance (SFSO 2009c and SFSO 2010c). Lately, a recalibration of the mileage per vehicle category has been performed (ARE 2012).
- **Numbers of starts/stops:** Derived from vehicles stock, with data on trip length distributions and parking time distributions (ARE/SFSO 2005).

For the determination of the non-CO₂ greenhouse gases and the precursors, the transport performance must be attributed to so called “traffic situations” (characteristic patterns of driving behaviour) which serve as a key to select the appropriate emission factor. The relative shares of these traffic situations is derived from a national road traffic model (operated by the Federal Office of Spatial Development, see ARE 2010). The traffic model is based on an origin-destination matrix that is assigned to a network of about 20'000 road segments. The model is calibrated partly bottom-up and partly top-down: Bottom-up by a number of traffic counts from the national traffic-counter network (333 stations all over Switzerland, FEDRO 2010), and top-down by the total of the mileage per vehicle category. Furthermore, it supplies the attributes needed for assigning a “traffic situation” to each road segment.

Due to fuel price differences in the vicinity of the national borders, gasoline stations sell relevant amounts of gasoline to foreign car owners. This amount of fuel is mainly consumed abroad (“tank tourism”) but the whole amount must be reported as national under 1A3b Road Transportation. For the CO₂ emissions, the amount of tank-tourism is irrelevant since it is included in the sales principle. The non-CO₂ emissions related to the “tank tourism”, however, are not captured by the traffic model. For the purpose of assuring completeness within the GHG inventory, these emissions are quantified on the basis of the difference between fuel consumption according to the Swiss overall energy statistics (sales principle) and fuel consumption derived from the traffic model. The resulting amount of “tank tourism” fuel is multiplied with mean emission factors to determine the related emissions of CH₄, N₂O, NO_x, CO, NMVOC, and SO₂. For CO₂, which dominates the emissions by a factor of approx. 1'000-10'000, the use of Swiss mean factors is correct, since the carbon content constitutes the emission factor. For CH₄ and N₂O there are differences between the Swiss mean factors and the implied emission factors of the four neighbouring countries Austria, France, Germany, Italy, as a comparison with their implied emission factors for 1990 and 2004 has shown. The differences are small between Switzerland, Austria, and Germany because all three countries use the same emission factors (SAEFL 2004a), whereas there are some differences when compared to France and Italy that use other emission factors (COPERT¹¹). Nevertheless, the use of the mean Swiss emission factors seems to be the consistent approach.

The N₂O emissions from natural gas combustion for road transportation originate from two vehicle categories: Biofuel CNG/petrol passenger cars and urban buses running purely on CNG. The same data as for the estimation of other gases (e.g. CO₂) were used for these two vehicle categories. As for all other vehicle categories, a residual of the total activity is assigned to tank tourism.

Emission Factors

The emission factors for fossil CO₂ and other gases are country specific and based on measurements and analyses of fuel samples (Table 3-9). Emission factors for the further gases are country specific derived from “emission functions” which are determined from a compilation of measurements from various European countries with programs using similar

¹¹ see European Environment Agency <http://www.eea.europa.eu/publications/TEC05> [14.02.2013]

driving cycles (legislative as well as standardized real-world cycles, like “Common Artemis Driving Cycle” (CADC). The method has been developed in 1990-1995 and has been extended and updated in 2000, 2004 and 2010. These emission factors are compiled in a so called “Handbook of Emission Factors for Road Transport” (SAEFL 1995, 2004, 2004a, FOEN 2010i, INFRAS 2004, 2004a, 2010, 2011). The latest version (3.1) is presented and documented on the website <http://www.hbefa.net/>. Several reports may be downloaded from there:

- Documentation of the general emission factor methodology (INFRAS 2011; forthcoming in German),
- Emission Factors for Passenger Cars and Light Duty Vehicles Switzerland, Germany, Austria, Norway and Sweden (INFRAS 2010; in English),

The resulting emission factors are published on CD ROM (“Handbook of emission factors for Road Transport”, INFRAS 2010). The underlying database contains a dynamic fleet compositions model simulating the release of new exhaust technologies and the fading out of old technologies. Corrective factors are provided to account for future technologies. Further details are shown in Annex A3.1.5.

The following tables present a selection of mean emission factors. The CO₂ factors are constant over the whole period 1990–2012. The carbon content of the fuels has not changed. However, the increasing portion of biofuels to the fuels is encompassed by the data time series - see next chapter for the emission factors of biofuels. For the other gases, more or less pronounced decreases of the emission factors occur due to new emission regulations and subsequent new exhaust technologies (mandatory use of catalytic converters for gasoline cars and lower limits for sulphur content in diesel fuels). Early models of catalytic converters have been substantial sources of N₂O, leading to an emission increase until 1998. Recent converter technologies have overcome this problem resulting in a decrease of the (mean) emission factor.

As of Submission 2013 of the National Inventory Report, N₂O emission factors in g/km differentiated by vehicle category and technology from the Handbook of Emission Factors (INFRAS 2010) have been applied, in contrast to previous submissions that applied a constant value in g/TJ fuel consumption. This results in a more realistic change pattern over time of N₂O emissions from road transportation than in earlier submissions.

In contrast to the N₂O emission factors, the measurement sample for CH₄ emission factors remained the same. However, due to updates in the vehicles fleet composition, the implied emission factors changed eventually. Further detailed description of how the emission factors for CH₄ are estimated is provided in the Annex A3.1.5.

As of Submission 2013, N₂O emission factors for gaseous fuels have been applied. No country-specific EFs for N₂O are available. Therefore, emissions have been estimated using the EFs for alternative fuel vehicles provided in table 3.2.4 on page 3.23 of Volume 2 of the 2006 IPCC Guidelines (IPCC 1997b). The value of 101 mg/km from the 2006 IPCC Guidelines was used for urban buses running on CNG only. For the bi-fuel passenger cars, it is assumed that they use gasoline mainly during the start but otherwise run on CNG; therefore the respective CNG emission factor for light duty vehicles of 27 mg/km from the same source was applied. As for all other fuel categories, the emission factor used for tank tourism corresponds to the weighted average of the national transport mix.

Emission factors from the combustion of biofuels

In lieu of reviewed emission factors for biofuels the following assumption were made.

- Biodiesel: The implied emission factors 1A3b for fossil diesel are used. Values for 2012:
CO₂ 73.6 t/TJ; CH₄ 0.28 kg/TJ; N₂O 2.29 kg/TJ
- Bio ethanol: The implied emission factors 1A3b for gasoline are used. Values for 2012:
CO₂ 73.9 t/TJ; CH₄ 5.69 kg/TJ; N₂O 0.71 kg/TJ
- Biogas: The implied emission factors 1A3b for CNG are used. Values for 2012:
CO₂ 56.1 t/TJ; CH₄ 3.78 kg/TJ; N₂O 8.37 kg/TJ

Overview over mean emission factors

Table 3-32 Mean emission factors for road transport for passenger cars. For more details see Annex A3.1.5.

| Gas | Fuel | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-----------------------|----------|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Passenger Cars | | t/TJ | | | | | | | | | |
| CO₂ | Gasoline | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 |
| | Diesel | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 |
| | CNG | | | | | | | | | | |
| CH₄ | Gasoline | 0.0266 | 0.0237 | 0.0213 | 0.0196 | 0.0178 | 0.0166 | 0.0155 | 0.0143 | 0.0132 | 0.0123 |
| | Diesel | 0.0015 | 0.0015 | 0.0013 | 0.0012 | 0.0013 | 0.0012 | 0.0011 | 0.0010 | 0.0009 | 0.0009 |
| | CNG | | | | | | | | | | |
| N₂O | Gasoline | 0.0031 | 0.0033 | 0.0035 | 0.0037 | 0.0039 | 0.0041 | 0.0042 | 0.0042 | 0.0041 | 0.0039 |
| | Diesel | 0.0002 | 0.0003 | 0.0004 | 0.0005 | 0.0007 | 0.0008 | 0.0009 | 0.0010 | 0.0012 | 0.0014 |
| NO_x | Gasoline | 0.3449 | 0.3139 | 0.2832 | 0.2657 | 0.2568 | 0.2512 | 0.2469 | 0.2377 | 0.2274 | 0.2163 |
| | Diesel | 0.2527 | 0.2558 | 0.2463 | 0.2395 | 0.2457 | 0.2412 | 0.2404 | 0.2402 | 0.2422 | 0.2464 |
| | CNG | | | | | | | | | | |
| CO | Gasoline | 3.1952 | 2.7878 | 2.4249 | 2.1754 | 1.9608 | 1.7979 | 1.6596 | 1.5308 | 1.4167 | 1.3193 |
| | Diesel | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 |
| | CNG | | | | | | | | | | |
| NM VOC | Gasoline | 0.5008 | 0.4402 | 0.3865 | 0.3485 | 0.3131 | 0.2861 | 0.2623 | 0.2396 | 0.2192 | 0.2013 |
| | Diesel | 0.0608 | 0.0617 | 0.0547 | 0.0504 | 0.0515 | 0.0471 | 0.0445 | 0.0410 | 0.0380 | 0.0352 |
| | CNG | | | | | | | | | | |
| SO₂ | Gasoline | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 | 0.0094 |
| | Diesel | 0.3678 | 0.3651 | 0.3348 | 0.3076 | 0.2860 | 0.2656 | 0.2555 | 0.2365 | 0.2222 | 0.2079 |
| | CNG | | | | | | | | | | |

| Gas | Fuel | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-----------------------|----------|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Passenger Cars | | t/TJ | | | | | | | | | |
| CO₂ | Gasoline | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 | 73.9 |
| | Diesel | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 |
| | CNG | | | | | | | | 56.1 | 56.1 | 56.1 |
| CH₄ | Gasoline | 0.0114 | 0.0106 | 0.0098 | 0.0092 | 0.0086 | 0.0082 | 0.0076 | 0.0073 | 0.0069 | 0.0064 |
| | Diesel | 0.0008 | 0.0007 | 0.0006 | 0.0006 | 0.0006 | 0.0005 | 0.0004 | 0.0004 | 0.0004 | 0.0003 |
| | CNG | | | | | | | | 0.0045 | 0.0043 | 0.0042 |
| N₂O | Gasoline | 0.0037 | 0.0034 | 0.0032 | 0.0029 | 0.0017 | 0.0016 | 0.0013 | 0.0013 | 0.0011 | 0.0010 |
| | Diesel | 0.0015 | 0.0017 | 0.0018 | 0.0019 | 0.0019 | 0.0020 | 0.0020 | 0.0020 | 0.0021 | 0.0021 |
| NO_x | Gasoline | 0.2050 | 0.1925 | 0.1772 | 0.1646 | 0.1528 | 0.1433 | 0.1263 | 0.1192 | 0.1075 | 0.0964 |
| | Diesel | 0.2543 | 0.2653 | 0.2765 | 0.2873 | 0.2922 | 0.2905 | 0.2774 | 0.2677 | 0.2594 | 0.2544 |
| | CNG | | | | | | | | 0.0236 | 0.0232 | 0.0232 |
| CO | Gasoline | 1.2355 | 1.1774 | 1.1157 | 1.0698 | 1.0252 | 0.9934 | 0.9310 | 0.9065 | 0.8612 | 0.8144 |
| | Diesel | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 |
| | CNG | | | | | | | | 0.0842 | 0.0837 | 0.0841 |
| NM VOC | Gasoline | 0.1846 | 0.1722 | 0.1590 | 0.1486 | 0.1396 | 0.1334 | 0.1233 | 0.1200 | 0.1131 | 0.1063 |
| | Diesel | 0.0324 | 0.0293 | 0.0261 | 0.0243 | 0.0224 | 0.0205 | 0.0177 | 0.0164 | 0.0151 | 0.0140 |
| | CNG | | | | | | | | 0.0004 | 0.0004 | 0.0004 |
| SO₂ | Gasoline | 0.0067 | 0.0057 | 0.0048 | 0.0038 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 |
| | Diesel | 0.1832 | 0.1642 | 0.1448 | 0.1322 | 0.1131 | 0.1032 | 0.0903 | 0.0844 | 0.0785 | 0.0731 |
| | CNG | | | | | | | | 0.0000 | 0.0000 | 0.0000 |

| Gas | Fuel | 2010 | 2011 | 2012 |
|-----------------------|----------|-------------|--------|--------|
| Passenger Cars | | t/TJ | | |
| CO₂ | Gasoline | 73.9 | 73.9 | 73.9 |
| | Diesel | 73.6 | 73.6 | 73.6 |
| | CNG | 56.1 | 56.1 | 56.1 |
| CH₄ | Gasoline | 0.0060 | 0.0059 | 0.0057 |
| | Diesel | 0.0003 | 0.0003 | 0.0003 |
| | CNG | 0.0041 | 0.0054 | 0.0054 |
| N₂O | Gasoline | 0.0009 | 0.0008 | 0.0007 |
| | Diesel | 0.0021 | 0.0021 | 0.0022 |
| NO_x | Gasoline | 0.0861 | 0.0791 | 0.0726 |
| | Diesel | 0.2509 | 0.2487 | 0.2475 |
| | CNG | 0.0231 | 0.0221 | 0.0221 |
| CO | Gasoline | 0.7755 | 0.7530 | 0.7299 |
| | Diesel | 0.0000 | 0.0000 | 0.0000 |
| | CNG | 0.0839 | 0.1578 | 0.1570 |
| NM VOC | Gasoline | 0.1013 | 0.0991 | 0.0965 |
| | Diesel | 0.0133 | 0.0130 | 0.0127 |
| | CNG | 0.0004 | 0.0005 | 0.0005 |
| SO₂ | Gasoline | 0.0004 | 0.0004 | 0.0004 |
| | Diesel | 0.0693 | 0.0673 | 0.0652 |
| | CNG | 0.0000 | 0.0000 | 0.0000 |

Table 3-33 Mean emission factors for road transport for heavy duty vehicles. For more details see Annex A3.1.5.

| Gas | Fuel | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------------------|--------|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Heavy duty vehicles | | t/TJ | | | | | | | | | |
| CO ₂ | Diesel | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 |
| CH ₄ | Diesel | 0.0019 | 0.0019 | 0.0018 | 0.0018 | 0.0017 | 0.0016 | 0.0016 | 0.0015 | 0.0015 | 0.0014 |
| N ₂ O | Diesel | 0.0007 | 0.0007 | 0.0008 | 0.0008 | 0.0008 | 0.0008 | 0.0008 | 0.0008 | 0.0009 | 0.0009 |
| NO _x | Diesel | 1.029 | 1.028 | 1.025 | 1.016 | 0.986 | 0.956 | 0.935 | 0.920 | 0.911 | 0.902 |
| CO | Diesel | 0.219 | 0.218 | 0.218 | 0.214 | 0.206 | 0.201 | 0.196 | 0.191 | 0.184 | 0.178 |
| NM VOC | Diesel | 0.076 | 0.075 | 0.075 | 0.073 | 0.068 | 0.066 | 0.065 | 0.062 | 0.059 | 0.056 |
| SO ₂ | Diesel | 0.065 | 0.061 | 0.056 | 0.047 | 0.020 | 0.016 | 0.017 | 0.016 | 0.019 | 0.021 |

| Gas | Fuel | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------------------|--------|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Heavy duty vehicles | | t/TJ | | | | | | | | | |
| CO ₂ | Diesel | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 | 73.6 |
| CH ₄ | Diesel | 0.0013 | 0.0011 | 0.0010 | 0.0010 | 0.0009 | 0.0009 | 0.0008 | 0.0007 | 0.0005 | 0.0004 |
| N ₂ O | Diesel | 0.0009 | 0.0008 | 0.0008 | 0.0008 | 0.0007 | 0.0007 | 0.0009 | 0.0012 | 0.0017 | 0.0024 |
| NO _x | Diesel | 0.879 | 0.833 | 0.795 | 0.757 | 0.716 | 0.699 | 0.667 | 0.626 | 0.552 | 0.490 |
| CO | Diesel | 0.171 | 0.162 | 0.158 | 0.157 | 0.151 | 0.150 | 0.147 | 0.144 | 0.140 | 0.138 |
| NM VOC | Diesel | 0.053 | 0.046 | 0.043 | 0.040 | 0.036 | 0.035 | 0.031 | 0.028 | 0.022 | 0.017 |
| SO ₂ | Diesel | 0.0127 | 0.0117 | 0.0110 | 0.0093 | 0.0005 | 0.0005 | 0.0005 | 0.0005 | 0.0005 | 0.0005 |

| Gas | Fuel | 2010 | 2011 | 2012 |
|----------------------------|--------|-------------|--------|--------|
| Heavy duty vehicles | | t/TJ | | |
| CO ₂ | Diesel | 73.6 | 73.6 | 73.6 |
| CH ₄ | Diesel | 0.0004 | 0.0003 | 0.0002 |
| N ₂ O | Diesel | 0.0028 | 0.0032 | 0.0034 |
| NO _x | Diesel | 0.451 | 0.416 | 0.390 |
| CO | Diesel | 0.136 | 0.135 | 0.134 |
| NM VOC | Diesel | 0.015 | 0.012 | 0.010 |
| SO ₂ | Diesel | 0.0005 | 0.0005 | 0.0005 |

Activity data

The amount of gasoline and diesel fuel sold in Switzerland serves as the activity data for the calculation of the CO₂ emissions: The Swiss overall energy statistics gives the amount of gasoline and diesel oil sold (SFOE 2013). From these numbers, the off-road consumption and the fugitive emissions from transmission, storage and fuelling of gasoline (reported under 1B2av Distribution of oil products) are subtracted. The result gives the inventory-relevant consumption for estimating the CO₂ emissions. It contains the fuel consumption due to the traffic model plus the amount of “tank tourism” (see above). The following table shows the details.

Table 3-34 Upper and middle part of table: Split of fuel sales into territorial on-road (model), off-road (model) and tank tourism (residual value to sales amounts) for gasoline and diesel oil in PJ. (Numbers may not add to totals due to rounding.)

Lower part of table: Consumption of biofuels for road transportation. Consumption starts in 1997.

Note that the unit is TJ (not PJ like fossil fuels in the upper and middle part of the table) and that Vegetable/Waste oil is included in the numbers of Biodiesel as well as separately depicted. However no double counting occurs in the total sum and shares of total fuel consumption.

| Activity data | Source category | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| on-road and off-road categories | | PJ | | | | | | | | | |
| Gasoline | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 135.6 | 139.0 | 137.6 | 134.5 | 137.6 | 140.8 | 142.3 | 142.9 | 143.9 | 145.3 |
| "tank tourism" | 1A3b | 17.8 | 20.8 | 28.1 | 19.0 | 16.1 | 8.1 | 10.5 | 16.0 | 16.3 | 20.4 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 2.4 | 2.4 | 2.4 | 2.4 | 2.4 | 2.4 | 2.3 | 2.3 | 2.3 | 2.3 |
| Gasoline sold in Switzerland | | 155.8 | 162.2 | 168.1 | 155.9 | 156.1 | 151.3 | 155.2 | 161.2 | 162.5 | 168.0 |
| Diesel | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 36.5 | 37.4 | 38.3 | 38.1 | 39.0 | 39.8 | 39.7 | 40.0 | 41.1 | 42.7 |
| "tank tourism" | 1A3b | -1.3 | -1.8 | -4.5 | -6.2 | -4.6 | -4.8 | -7.7 | -6.5 | -5.8 | -4.7 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 11.6 | 11.8 | 12.1 | 12.3 | 12.6 | 12.8 | 13.0 | 13.2 | 13.4 | 13.6 |
| Diesel sold in Switzerland | | 46.7 | 47.4 | 45.9 | 44.2 | 46.9 | 47.8 | 44.9 | 46.7 | 48.7 | 51.6 |
| Total | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 172.0 | 176.4 | 175.9 | 172.6 | 176.6 | 180.6 | 182.0 | 182.9 | 185.0 | 188.0 |
| "tank tourism" | 1A3b | 16.5 | 19.0 | 23.7 | 12.9 | 11.5 | 3.3 | 2.8 | 9.5 | 10.5 | 15.7 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 14.0 | 14.2 | 14.5 | 14.7 | 14.9 | 15.2 | 15.3 | 15.5 | 15.7 | 15.9 |
| Gasoline and Diesel sold in Switzerland | | 202.5 | 209.6 | 214.0 | 200.1 | 203.0 | 199.1 | 200.1 | 207.9 | 211.1 | 219.6 |

| Activity data | Source category | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| on-road and off-road categories | | PJ | | | | | | | | | |
| Gasoline | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 146.9 | 145.5 | 144.5 | 141.9 | 139.1 | 136.1 | 132.1 | 129.1 | 126.4 | 122.8 |
| "tank tourism" | 1A3b | 18.9 | 15.6 | 13.4 | 15.3 | 15.4 | 13.6 | 12.9 | 14.5 | 14.0 | 13.7 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 2.3 | 2.3 | 2.2 | 2.2 | 2.2 | 2.2 | 2.2 | 2.2 | 2.2 | 2.2 |
| Gasoline sold in Switzerland | | 168.0 | 163.4 | 160.2 | 159.5 | 156.6 | 151.8 | 147.2 | 145.8 | 142.6 | 138.7 |
| Diesel | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 44.9 | 45.8 | 47.4 | 50.5 | 54.1 | 57.4 | 61.1 | 65.3 | 68.2 | 70.8 |
| "tank tourism" | 1A3b | -3.5 | -3.3 | -3.0 | -2.7 | -1.7 | 0.9 | 3.1 | 4.5 | 9.8 | 8.6 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 13.7 | 13.8 | 13.9 | 13.9 | 14.0 | 14.1 | 14.2 | 14.4 | 14.6 | 14.7 |
| Diesel sold in Switzerland | | 55.1 | 56.2 | 58.3 | 61.7 | 66.3 | 72.4 | 78.4 | 84.2 | 92.6 | 94.1 |
| Total | | | | | | | | | | | |
| on-road consumption (model) | 1A3b | 191.7 | 191.3 | 191.9 | 192.4 | 193.1 | 193.5 | 193.2 | 194.4 | 194.6 | 193.6 |
| "tank tourism" | 1A3b | 15.4 | 12.3 | 10.5 | 12.6 | 13.7 | 14.5 | 16.0 | 19.0 | 23.9 | 22.3 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 16.0 | 16.1 | 16.1 | 16.1 | 16.2 | 16.2 | 16.4 | 16.6 | 16.7 | 16.9 |
| Gasoline and Diesel sold in Switzerland | | 223.2 | 219.6 | 218.5 | 221.2 | 223.0 | 224.3 | 225.6 | 229.9 | 235.2 | 232.9 |

| Activity data | Source category | 2010 | 2011 | 2012 |
|---|--|-------|-------|-------|
| on-road and off-road categories | | PJ | | |
| Gasoline | | | | |
| on-road consumption (model) | 1A3b | 119.3 | 116.0 | 112.7 |
| "tank tourism" | 1A3b | 12.4 | 10.6 | 9.3 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 2.2 | 2.1 | 2.1 |
| Gasoline sold in Switzerland | | 133.8 | 128.7 | 124.1 |
| Diesel | | | | |
| on-road consumption (model) | 1A3b | 73.9 | 75.1 | 76.3 |
| "tank tourism" | 1A3b | 9.0 | 10.6 | 15.6 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 14.9 | 14.8 | 14.7 |
| Diesel sold in Switzerland | | 97.8 | 100.5 | 106.6 |
| Total | | | | |
| on-road consumption (model) | 1A3b | 193.2 | 191.1 | 189.0 |
| "tank tourism" | 1A3b | 21.4 | 21.1 | 25.0 |
| off-road consumption (models) | 1A2fii;1A3aii,c,d,e;1A4aii,bii,cii;1A5 | 17.1 | 16.9 | 16.8 |
| Gasoline and Diesel sold in Switzerland | | 231.6 | 229.1 | 230.7 |

| Biofuels | 1990-1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|-----------------------------------|-----------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | TJ | | | | | | | | | | | | | | | | |
| Biodiesel | 0 | 57 | 51 | 48 | 56 | 60.4 | 55.0 | 72.3 | 100.7 | 196.4 | 272.7 | 304.7 | 368.2 | 232.0 | 287.9 | 316.8 | 382.6 |
| Bioethanol | 0 | 0 | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.0 | 22.3 | 67.1 | 69.2 | 31.2 | 54.6 | 85.2 | 97.3 |
| Biogas | 0 | 0 | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 30.2 | 30.8 | 32.1 | 50.7 | 105.9 | 161.1 | |
| Sum | 0 | 57.2 | 51.4 | 48.3 | 56.4 | 60.4 | 55.0 | 72.3 | 100.7 | 215.4 | 295.0 | 402.0 | 468.2 | 295.3 | 393.2 | 508.0 | 640.9 |
| Share of total fuel consump. 1A3b | 0.0% | 0.3% | 0.3% | 0.3% | 0.3% | 0.3% | 0.3% | 0.4% | 0.5% | 1.1% | 1.5% | 2.1% | 2.4% | 1.5% | 2.0% | 2.7% | 3.4% |

Further activity data needed for modelling the non-CO₂ emissions are the mileages (vehicle kilometres) per vehicle category in Table 3-35. Note that the activity data have been recalculated due based on the latest figures on population and economy (Prognos 2012a, ARE 2012).

Table 3-35 Mileages in millions of vehicle kilometres. PC: passenger cars, LDV: light duty vehicles, HDV: heavy duty vehicles).

| Veh. category | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | million vehicle-km | | | | | | | | | |
| PC | 42'650 | 43'745 | 43'178 | 42'259 | 43'199 | 43'824 | 44'063 | 44'675 | 45'570 | 46'702 |
| LDV | 2'758 | 2'742 | 2'867 | 2'632 | 2'669 | 2'746 | 2'767 | 2'786 | 2'831 | 2'903 |
| HDV | 1'992 | 2'015 | 2'036 | 2'025 | 2'109 | 2'107 | 2'055 | 2'072 | 2'126 | 2'200 |
| Coaches | 108 | 108 | 109 | 109 | 110 | 110 | 109 | 108 | 101 | 98 |
| Urban Bus | 174 | 186 | 188 | 190 | 190 | 192 | 188 | 188 | 192 | 195 |
| 2-Wheelers | 2'025 | 1'947 | 1'866 | 1'792 | 1'717 | 1'744 | 1'756 | 1'823 | 1'872 | 1'941 |
| Sum | 49'707 | 50'743 | 50'244 | 49'007 | 49'993 | 50'724 | 50'939 | 51'653 | 52'692 | 54'039 |
| (1990=100%) | 100% | 102% | 101% | 99% | 101% | 102% | 102% | 104% | 106% | 109% |

| Veh. category | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | million vehicle-km | | | | | | | | | |
| PC | 48'063 | 48'509 | 49'062 | 49'527 | 50'019 | 50'465 | 50'812 | 51'208 | 51'949 | 52'852 |
| LDV | 2'978 | 3'059 | 3'119 | 3'149 | 3'215 | 3'300 | 3'374 | 3'473 | 3'529 | 3'584 |
| HDV | 2'273 | 2'165 | 2'109 | 2'115 | 2'144 | 2'127 | 2'189 | 2'203 | 2'223 | 2'172 |
| Coaches | 99 | 95 | 93 | 95 | 98 | 106 | 118 | 120 | 114 | 119 |
| Urban Bus | 200 | 205 | 211 | 215 | 220 | 229 | 233 | 240 | 245 | 249 |
| 2-Wheelers | 1'999 | 2'048 | 2'098 | 2'152 | 2'190 | 2'204 | 2'262 | 2'300 | 2'366 | 2'385 |
| Sum | 55'612 | 56'082 | 56'693 | 57'253 | 57'886 | 58'432 | 58'989 | 59'544 | 60'426 | 61'361 |
| (1990=100%) | 112% | 113% | 114% | 115% | 116% | 118% | 119% | 120% | 122% | 123% |

| Veh. category | 2010 | 2011 | 2012 |
|---------------|--------------------|--------|--------|
| | million vehicle-km | | |
| PC | 53'341 | 54'000 | 54'730 |
| LDV | 3'621 | 3'663 | 3'701 |
| HDV | 2'210 | 2'250 | 2'290 |
| Coaches | 119 | 119 | 118 |
| Urban Bus | 251 | 254 | 257 |
| 2-Wheelers | 2'407 | 2'436 | 2'465 |
| Sum | 61'950 | 62'722 | 63'562 |
| (1990=100%) | 125% | 126% | 128% |

In 2012, 86.1% of total vehicle kilometres are driven by passenger cars, 5.8% and 3.6% by light and heavy duty vehicles, respectively. The mileages increased for all vehicle categories (except coaches), totalling 28% in the period 1990–2012. In the same period, fuel consumption increased less strongly, by 13.9%, indicating improved fuel efficiency. This effect is also reflected in Table 3-36 that depicts the specific fuel consumption per vehicle-km. For most vehicle categories, the specific consumption has decreased in the period 1990–2012 (between 3% and 23%). Consumption of light duty vehicles remained indifferent while two-wheelers (15%) have increased their average specific consumption. Concerning the whole car fleet, a decrease of 19% in specific consumption has been reached between 1990 and 2012.

Table 3-36 Fuel consumption of road transport, not including "tank tourism" (PC: passenger cars, LDV: light duty vehicles, HDV: heavy duty vehicles).

| Veh. cat. | Fuel | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------|----------|---------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | | specific fuel consumption (MJ/veh-km) | | | | | | | | | |
| PC | Gasoline | 3.18 | 3.20 | 3.22 | 3.23 | 3.23 | 3.23 | 3.22 | 3.21 | 3.19 | 3.17 |
| | Diesel | 2.91 | 2.91 | 2.92 | 2.98 | 2.90 | 2.90 | 2.90 | 2.91 | 2.89 | 2.86 |
| | CNG | | | | | | | | | | |
| LDV | Gasoline | 3.17 | 3.18 | 3.17 | 3.18 | 3.18 | 3.18 | 3.18 | 3.17 | 3.17 | 3.18 |
| | Diesel | 3.86 | 3.87 | 3.87 | 3.88 | 3.87 | 3.86 | 3.83 | 3.81 | 3.79 | 3.77 |
| HDV | Diesel | 10.91 | 10.95 | 10.98 | 10.92 | 10.97 | 10.85 | 10.71 | 10.58 | 10.46 | 10.38 |
| Coach | Diesel | 11.84 | 11.85 | 11.87 | 11.81 | 11.75 | 11.69 | 11.62 | 11.55 | 11.48 | 11.42 |
| Urban Bus | Diesel | 16.22 | 16.29 | 16.33 | 16.34 | 16.32 | 16.29 | 16.20 | 16.10 | 16.02 | 15.90 |
| | CNG | | | | | | | | | | |
| 2-Wheeler | Gasoline | 1.11 | 1.14 | 1.17 | 1.19 | 1.21 | 1.22 | 1.22 | 1.24 | 1.24 | 1.24 |
| Average | | 3.45 | 3.47 | 3.49 | 3.51 | 3.51 | 3.49 | 3.45 | 3.42 | 3.39 | 3.36 |
| | | 100% | 100% | 101% | 102% | 102% | 101% | 100% | 99% | 98% | 97% |

| Veh. cat. | Fuel | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------|----------|---------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | | specific fuel consumption (MJ/veh-km) | | | | | | | | | |
| PC | Gasoline | 3.14 | 3.13 | 3.11 | 3.09 | 3.06 | 3.04 | 2.99 | 2.97 | 2.93 | 2.90 |
| | Diesel | 2.80 | 2.72 | 2.66 | 2.58 | 2.52 | 2.46 | 2.41 | 2.40 | 2.35 | 2.33 |
| | CNG | | | | | | | | 2.91 | 2.88 | 2.85 |
| LDV | Gasoline | 3.18 | 3.17 | 3.18 | 3.19 | 3.19 | 3.19 | 3.21 | 3.21 | 3.20 | 3.19 |
| | Diesel | 3.75 | 3.71 | 3.63 | 3.56 | 3.48 | 3.42 | 3.37 | 3.34 | 3.32 | 3.31 |
| HDV | Diesel | 10.33 | 10.56 | 10.62 | 10.63 | 10.61 | 10.77 | 10.71 | 10.73 | 10.65 | 10.59 |
| Coach | Diesel | 11.33 | 11.25 | 11.21 | 11.19 | 11.21 | 11.22 | 11.23 | 11.22 | 11.18 | 11.16 |
| Urban Bus | Diesel | 15.80 | 15.71 | 15.60 | 15.45 | 15.45 | 15.37 | 15.24 | 15.23 | 15.05 | 14.94 |
| | CNG | | | | | | | | 20.34 | 20.32 | 20.36 |
| 2-Wheeler | Gasoline | 1.25 | 1.25 | 1.24 | 1.25 | 1.27 | 1.28 | 1.29 | 1.31 | 1.33 | 1.35 |
| Average | | 3.32 | 3.29 | 3.24 | 3.20 | 3.15 | 3.11 | 3.06 | 3.02 | 2.96 | 2.93 |
| | | 96% | 95% | 94% | 93% | 91% | 90% | 89% | 87% | 86% | 85% |

| Veh. cat. | Fuel | 2010 | 2011 | 2012 |
|----------------|----------|-------------|-------------|-------------|
| | | MJ/veh-km | | |
| PC | Gasoline | 2.86 | 2.81 | 2.77 |
| | Diesel | 2.30 | 2.28 | 2.24 |
| | CNG | 2.83 | 2.53 | 2.51 |
| LDV | Gasoline | 3.18 | 3.17 | 3.15 |
| | Diesel | 3.31 | 3.30 | 3.29 |
| HDV | Diesel | 10.55 | 10.50 | 10.46 |
| Coach | Diesel | 11.16 | 11.15 | 11.14 |
| Urban Bus | Diesel | 14.81 | 14.76 | 14.72 |
| | CNG | 20.58 | 20.52 | 20.46 |
| 2-Wheeler | Gasoline | 1.34 | 1.34 | 1.34 |
| Average | | 2.89 | 2.86 | 2.82 |
| | | 84% | 83% | 82% |

For modelling of cold start and evaporative emissions of passenger cars and light duty vehicles, also vehicle stock and start numbers are used for activity data. The corresponding numbers are summarised in the next table. Vehicle stock figures correspond to registration data. The starts per vehicle are based on specific household surveys (ARE/SFSO 2005).

Table 3-37 Vehicle stock numbers and average number of starts per vehicle per day (PC: passenger cars, LDV: light duty vehicles).

| Veh. Category | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------|-----------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | stock in 1000 vehicles | | | | | | | | | |
| PC | 2'985 | 3'058 | 3'091 | 3'110 | 3'165 | 3'229 | 3'268 | 3'323 | 3'383 | 3'467 |
| LDV | 221 | 228 | 229 | 228 | 232 | 238 | 241 | 243 | 247 | 254 |
| 2-Wheelers | 764 | 747 | 729 | 720 | 708 | 704 | 699 | 709 | 718 | 728 |
| | starts per vehicle per day | | | | | | | | | |
| PC | 2.61 | 2.60 | 2.58 | 2.56 | 2.54 | 2.53 | 2.53 | 2.51 | 2.49 | 2.47 |
| LDV | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 | 1.97 |
| 2-Wheelers | 1.59 | 1.58 | 1.57 | 1.56 | 1.55 | 1.54 | 1.54 | 1.53 | 1.52 | 1.51 |

| Veh. Category | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------|-----------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | stock in 1000 vehicles | | | | | | | | | |
| PC | 3'545 | 3'630 | 3'701 | 3'754 | 3'811 | 3'862 | 3'894 | 3'956 | 3'990 | 4'010 |
| LDV | 260 | 268 | 274 | 278 | 284 | 291 | 298 | 307 | 312 | 317 |
| 2-Wheelers | 732 | 740 | 753 | 763 | 771 | 770 | 784 | 789 | 804 | 807 |
| | starts per vehicle per day | | | | | | | | | |
| PC | 2.46 | 2.45 | 2.44 | 2.43 | 2.41 | 2.40 | 2.39 | 2.38 | 2.37 | 2.35 |
| LDV | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 | 1.96 |
| 2-Wheelers | 1.50 | 1.51 | 1.52 | 1.52 | 1.53 | 1.54 | 1.54 | 1.55 | 1.56 | 1.56 |

| Veh. Category | 2010 | 2011 | 2012 |
|---------------|-----------------------------------|-------|-------|
| | stock in 1000 vehicles | | |
| PC | 4'076 | 4'195 | 4'302 |
| LDV | 326 | 328 | 331 |
| 2-Wheelers | 816 | 815 | 815 |
| | starts per vehicle per day | | |
| PC | 2.34 | 2.34 | 2.33 |
| LDV | 1.96 | 1.96 | 1.96 |
| 2-Wheelers | 1.57 | 1.57 | 1.57 |

c) Railways (1A3c)

Methodology

The entire Swiss railway system is electrified. Electric locomotives are used in passenger as well as freight railway traffic. Diesel locomotives are used for shunting purposes in marshalling yards and for construction activities only.

Emissions of diesel rail vehicles are modelled using the off-road model developed by INFRAS (2008). For details refer to Chapter 3.2.5.1 (paragraph on source category 1A2fii).

Emission Factors

Only diesel oil is being used as fuel, therefore all emission factors refer to diesel oil.

- The emission factor for CO₂ is country specific and assumed to be constant in the period 1990-2012 with value 73.6 t/TJ (diesel oil, see Table 3-9, SFOE 2001).
- For SO₂ the emission factors are country specific. They are depicted in Table A - 6 in Annex A2, row diesel oil: Continuous decrease from 65.4 kg/TJ in 1990 to 12.7 kg/TJ in 2000 and to 0.47 kg/TJ in 2010.

The emission factors for all other gases are country specific and are shown in Table A - 19 in Annex A3.1.6. Note that NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference of VOC and CH₄ emissions. Note that emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database (INFRAS 2008; see footnote 7 on page 111).

Activity data

The fuel consumption is calculated like emission modelling but with consumption factors using instead of emission factors (see Table A - 19). The operating hours depend on the number of vehicles per age and size class. In 2005 e.g., 1'260 vehicles were operating 0.77 million hours per year with an average number of 611 operating hours per year per vehicle (INFRAS 2008). As mentioned above, a slight update was carried out in 2013 based on the latest figures on population and economy (Prognos 2012a). The diesel consumption has been recalculated accordingly. Numbers from 2005 onwards are affected. The resulting fuel consumption is shown in Table 3-38.

Table 3-38 Activity data (diesel oil consumption) for railways.

| 1A3c Railways | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Diesel | TJ | 390 | 400 | 410 | 420 | 430 | 441 | 443 | 446 | 449 | 452 |
| 1990=100% | | 100.0% | 102.6% | 105.2% | 107.8% | 110.4% | 113.0% | 113.8% | 114.5% | 115.2% | 116.0% |

| 1A3c Railways | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Diesel | TJ | 455 | 460 | 466 | 472 | 477 | 483 | 494 | 504 | 515 | 526 |
| 1990=100% | | 116.7% | 118.1% | 119.6% | 121.0% | 122.4% | 123.8% | 126.6% | 129.4% | 132.2% | 135.0% |

| 1A3c Railways | Unit | 2010 | 2011 | 2012 |
|---------------|------|--------|--------|--------|
| Diesel | TJ | 537 | 538 | 539 |
| 1990=100% | | 137.8% | 138.1% | 138.3% |

d) Navigation (1A3d)

Methodology

There are passenger ships, dredgers, fishing boats, motor and sailing boats on the lakes of Switzerland and on the river Rhine. Every boat is registered at the cantonal authorities.

Emissions of ships and boats are calculated using the off-road model developed by INFRAS (2008). For details refer to Chapter 3.2.5.1 (paragraph on source category 1A2fii).

On the river Rhine as well as on the lakes Geneva and Constance, some of the boats cross the border and go abroad (Germany, France). Fuels bought in Switzerland will therefore become bunker fuel. The amount of bunker diesel oil is evaluated in Section 3.2.2.

Emission Factors

- The emission factor for CO₂ is country specific and is assumed to be constant in the period 1990-2012 with value 73.6 t/TJ for diesel oil, 73.9 t/TJ for gasoline and 73.7 t/TJ for gas oil (Table 3-9, SFOE 2001).
- For SO₂ the emission factors are country specific and are given in Table A - 6 in Annex A2 (diesel oil, gasoline, gas oil).
- The emission factors for all other gases are country specific and are shown in Table A - 20 to Table A - 23 in Annex A3.1.6. Note that NMVOC is not modelled bottom-up. The NMVOC emissions are calculated as the difference of VOC and CH₄ emissions.

Note that emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database INFRAS (2008), see footnote 7 on page 111.

Activity data

The numbers of vehicles and of operating hours are given in Annex A3.1.6 (INFRAS 2008). Table 3-39 shows the domestic fuel consumption. In 2012, the fuel-split was 52%, 38% and 10% for diesel oil, gasoline and gas oil. A slight modification of the activity data was carried out in 2013 based on the latest figures of population and economy (Prognos 2012a, Keller/INFRAS 2013). The consumption of liquid fuels has been recalculated accordingly.

Table 3-39 Fuel consumption of (domestic) navigation.

| 1A3d Navigation | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-----------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Diesel | TJ | 705 | 703 | 701 | 698 | 696 | 694 | 708 | 723 | 737 | 752 |
| Gasoline | TJ | 701 | 692 | 683 | 673 | 664 | 654 | 647 | 639 | 631 | 623 |
| Gas oil | TJ | 111 | 117 | 122 | 128 | 134 | 140 | 141 | 143 | 145 | 146 |
| Sum | TJ | 1'518 | 1'512 | 1'506 | 1'500 | 1'494 | 1'488 | 1'496 | 1'505 | 1'513 | 1'522 |
| 1990 = 100% | | 100% | 100% | 99% | 99% | 98% | 98% | 99% | 99% | 100% | 100% |

| 1A3d Navigation | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-----------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Diesel | TJ | 766 | 770 | 774 | 778 | 782 | 786 | 800 | 815 | 830 | 844 |
| Gasoline | TJ | 616 | 614 | 613 | 612 | 611 | 609 | 615 | 620 | 626 | 631 |
| Gas oil | TJ | 148 | 150 | 153 | 156 | 158 | 161 | 162 | 164 | 166 | 167 |
| Sum | TJ | 1'530 | 1'535 | 1'540 | 1'546 | 1'551 | 1'556 | 1'578 | 1'599 | 1'621 | 1'643 |
| 1990 = 100% | | 101% | 101% | 101% | 102% | 102% | 103% | 104% | 105% | 107% | 108% |

| 1A3d Navigation | Unit | 2010 | 2011 | 2012 |
|-----------------|------|-------|-------|-------|
| Diesel | TJ | 859 | 853 | 847 |
| Gasoline | TJ | 637 | 632 | 626 |
| Gas oil | TJ | 169 | 169 | 170 |
| Sum | TJ | 1'665 | 1'654 | 1'643 |
| 1990 = 100% | | 110% | 109% | 108% |

e) Pipeline Transportation (1A3e)

Source 1A3e includes emissions of CO₂, CH₄, N₂O, NO_x, CO, NMVOC and SO₂ from a compressor station located in Ruswil. The compressor station uses a centrifugal compressor according to Transigas AG (the company operating the compressor station and the pipeline network).

Emission Factors

The emission factors for CO₂, CH₄ and N₂O for 1A3e correspond to the ones used for gas turbines in Switzerland (SAEFL 2000) as suggested by expert judgement (see also Battelle 1994 and Xinmin 2004). With regard to CH₄, the EF was assumed to be 5 g/GJ up to 1995 and 2 g/GJ from 2000 onwards, with linear interpolation in between. This corresponds to the assumption that a catalyst was fitted to the system, which reduced the CH₄ emissions of the gas turbine.

Activity data

The data on fuel consumption for the operation of the compressor station in Ruswil is based on the Swiss overall energy statistics (SFOE 2013), see also Figure 3-8.

3.2.8.3 Uncertainties and Time-Series Consistency

a) General

For a general description of the uncertainty analysis and time series consistency of the Energy Sector see Chapter 3.2.6.3 a).

b) Specific: Uncertainties for CH₄ and N₂O in 1A3b Road Transportation

Due to a study for the road transportation in Germany (IFEU/INFRAS 2009), where the same handbook of emission factors is used as in Switzerland, the uncertainties for the CH₄ and N₂O emission factors have been adopted:

- CH₄: 37% (gasoline) and 20% (diesel),
- N₂O: 50% (gasoline) and 22% (diesel).

For the CH₄ emissions of CNG the qualitative uncertainty "medium" (30%) is taken and for biomass the uncertainty "high" (60%) according to Table 1-3.

For the N₂O emissions of CNG the qualitative uncertainty "medium" (80%) is taken and for biomass the uncertainty "high" (150%) according to Table 1-3.

Consistency and Completeness in 1A3 Fuel Combustion

- Time series for 1A3 are all consistent.
- All estimates in the sector 1A3 are assumed to be complete.

3.2.8.4 Source-specific QA/QC and Verification**a) General**

See chapter 3.2.6.4.

The emission factors of category 1A3b for CO₂, CH₄ and N₂O used in the Swiss Inventory were compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available for submission 2012 (INFRAS 2012). Switzerland's diesel and gasoline CO₂ emission factor lie in the midfield of the other countries. Furthermore CH₄ and N₂O emission factors for gasoline are significantly lower. For further explanations see Sat. Pap in Chapter 16.

b) Specific: Civil Aviation (1A3a)**Emissions**

Total calculated emissions for domestic and international flights have been compared between different years. The development of total emissions with time is consistent with a fleet renewal of former Swissair in the early nineties, the technological improvements and changes in fleet composition.

Emission factors

- From total fuel burnt, total distance, number of passenger (without freight) per aircraft type, the fuel consumption per 100 passenger km has been calculated (backward calculation). The result of 2 to 10 kg fuel/100 passenger km is in line with expectations for 1990 passenger fleets.
- The implied emission factors were calculated for 2012 and compared with previous years.

Activity data

- In an independent Tier 3b calculation, EUROCONTROL performed a fuel calculation for Switzerland's international flights, based on collected flight plan data and single movements. The results for the years 2004, 2005 and 2007 matched the FOCA

calculations by more than 97.4%. The FOCA results were generally 1% to 2% higher but included the total number of actual flight movements of all flights, including VFR (visual flight rules) and non-scheduled flights such as helicopter movements in alpine regions.

- Comparison between total movement numbers in the calculation and in the corresponding published statistics. Example: In 1990 calculation, FOCA considered all flights for which there was a form 'Traffic report to the airport authorities' filled in (total heavy aircraft). The total number of movements in 1990 is 263'951 (without Basel). The published number of movements for scheduled and charter flights in 1990 is: 263'952 (without Basel).
- The bottom-up calculation of total fuel matches the total fuel sold within a few percents. The remaining difference can be attributed to fuelling.
- Real-world fuel consumption was compared with modelled consumption for selected aircrafts of four Swiss airlines. The difference between the two methods was smaller than 1%.

c) Specific: Road Transportation (1A3b)

The international project for the update of the emission factors for road vehicles is overseen by a group of external and international experts that guarantees an independent quality control. For the update of the modelling of Switzerland's road transport emissions, which has been carried out between 2008 and 2010, several experts from the federal administration have conducted the project. The results have undergone large plausibility checks and comparisons with earlier estimates.

The emission factors CH₄ and N₂O used for the modelling of 1A3b Road Transportation are taken from the handbook of emission factors (INFRAS 2010), which is also applied in Germany, Austria, Netherlands, and Sweden. The Swiss emission factors for CH₄ and N₂O used in 1A3b were additionally compared with those depicted in the CRF from Germany and a good match was found. Possible small differences might result from a varying fleet composition. For gasoline, the activity data is easily verified due to the fact, that 98.3% (2012) of the gasoline sold in Switzerland is consumed by 1A3b Road Transportation itself. Therefore the amount of gasoline reported in the Swiss overall energy statistics is a strong control and verification parameter for the activity data of 1A3b. For diesel, the same control is carried out and the amount of diesel consumed by 1A3b Road Transportation is 86.3% (2012) compared to the amount sold.

3.2.8.5 Source-Specific Recalculations

- 1A3b: The entire time series has been recalculated following an update based on the latest figures on population growth and economy (Prognos 2012a, ARE 2012). Vehicle kilometres from 1993 are slightly lower in total; fleet compositions have changed, with slight impacts on implied emission factors; fuel consumption in tank tourism has been recalculated; the modelled share of biofuels has been reduced to be consistent with real-world developments. The overall impacts of these recalculations on emissions are low.
- 1A3c, 1A3d: The activity data have so far been taken from INFRAS (2008). For this submission, the latest numbers on growth of population and economy (Prognos 2012a, Keller/INFRAS 2013) have been integrated in the off-road model.

3.2.8.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. To accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.2.9 Source Category 1A4 - Other Sectors (Commercial/Institutional, Residential, Agriculture/Forestry/Fisheries)

3.2.9.1 Source Category Description

Tier 1 Key categories 1A4

CO₂ from the combustion of Liquid Fuels in the Commercial/Institutional Sector (level and trend)

CO₂ from the combustion of Gaseous Fuels in the Commercial/Institutional Sector (level and trend)

CO₂ from the combustion of Liquid Fuels in the Residential Sector (level and trend)

CO₂ from the combustion of Gaseous Fuels in the Residential Sector (level and trend)

CO₂ from the combustion of Liquid Fuels in the Agriculture/Forestry/Fisheries Sector (level)

Tier 2 Key categories 1A4

CO₂ from the combustion of Liquid Fuels in the Commercial/Institutional Sector (trend)

CO₂ from the combustion of Gaseous Fuels in the Commercial/Institutional Sector (level and trend)

CO₂ from the combustion of Liquid Fuels in the Residential Sector (level and trend)

CO₂ from the combustion of Gaseous Fuels in the Residential Sector (level and trend)

CH₄ from the combustion of Biomass in the Residential Sector (trend)

Table 3-40 Specification of source category 1A4 Other sectors.

EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1A4 | Source | Specification | Data Source |
|------|--------------------------------------|---|---|
| 1A4a | Commercial/ Institutional | Emission from stationary fuel combustion in commercial and institutional buildings (1A4ai) and from mobile off-road machinery (professional gardening) and motorised equipment (1A4aii) | AD: SFOE 2013, INFRAS 2008, Prognos 2012, Keller/INFRAS 2013, EMIS 2014/1A4div. EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006; EMIS 2014/1A Holzfeuerungen; SAEFL 2000; Nussbaumer, T., Boogen, N. 2010 EMIS 2014/1A4a INFRAS 2008, EMIS 2014/1A4div. |
| 1A4b | Residential | Emissions from stationary fuel combustion in households (1A4bi) and from mobile machinery (hobby gardening) and motorised equipment (1A4bii) | AD: SFOE 2013, INFRAS 2008, Prognos 2012, Keller/INFRAS 2013, EMIS 2014/1A4div. EF: EMPA 1999; Intertek 2008; Intertek 2012; FOEN 2011k; IPCC 2006; EMIS 2014/1A Holzfeuerungen; SAEFL 2000; Nussbaumer, T., Boogen, N. 2010 EMIS 2014/1A4b INFRAS 2008, EMIS 2014/1A4div. |
| 1A4c | Agriculture/ Forestry/ Fishing | Comprises stationary fuel combustion for heating in forestry and agriculture and grass drying (1A4ci) and mobile machinery (off-road) in agriculture and forestry (1A4cii) | Grass drying: EMIS 2013/1A4ci Off-road machinery: INFRAS 2008, Prognos 2012, Keller/INFRAS 2013 Wood heating: EMIS 2014/1A Holzfeuerungen |

3.2.9.2 Methodological Issues

As explained in chapter 3.2.5, a country specific Tier 2 approach based on aggregated fuel consumption data from the Swiss overall energy statistics is used to calculate emissions (SFOE 2013). Source category 1A4b also includes charcoal use and bonfires in Switzerland.

Emissions of GHGs are calculated by multiplying levels of activity by emission factors.

For mobile off-road sources (1A4aii, 1A4bii and 1A4cii) the emissions are calculated by the same approach as all other off-road categories using the off-road model developed by INFRAS (2008). For details refer to Chapter 3.2.5.1 (paragraph on source category 1A2fii).

Emission Factors

The following table presents the emission factors used in 1A4a/b:

Table 3-41 Emission Factors for 1A4a/b Other Sectors Commercial/Institutional and Residential in 2012.

| Source/fuel | CO ₂ | CO ₂ biog. | CH ₄ | N ₂ O | NO _x | CO | NM VOC | SO ₂ |
|--|-----------------|-----------------------|-----------------|------------------|-----------------|---------|--------|-----------------|
| | t/TJ | | kg/TJ | | | | | |
| 1A4 a Other Sectors: Commercial/Institutional | | | | | | | | |
| Gas oil (weighted average) | 73.7 | | 1.0 | 0.6 | 35.2 | 6.9 | 6.0 | 22.4 |
| Gas oil (heat only boilers) | 73.7 | | 1.0 | 0.6 | 35.2 | 6.8 | 6.0 | 22.4 |
| Gas oil (turbines) | NO | | NO | NO | NO | NO | NO | NO |
| Gas oil (engines) | 73.7 | | 0.6 | 0.6 | 40.0 | 30.0 | 8.0 | 21.0 |
| Natural gas (weighted average) | 56.1 | | 6.9 | 0.1 | 25.2 | 13.7 | 3.8 | 0.5 |
| NG (heat only boilers) | 56.1 | | 6.0 | 0.1 | 18.4 | 10.6 | 4.0 | 0.5 |
| NG (turbines) | 56.1 | | 2.0 | 0.1 | 60.0 | 15.0 | 0.1 | 0.5 |
| NG (engines) | 56.1 | | 20.0 | 0.1 | 124.7 | 58.3 | 1.0 | 0.5 |
| Bituminous coal | NO | | NO | NO | NO | NO | NO | NO |
| Lignite | NO | | NO | NO | NO | NO | NO | NO |
| Biomass (weighted average) | | 87.3 | 27.6 | 1.4 | 115.8 | 695.6 | 63.2 | 17.9 |
| Biomass (wood) | | 90.6 | 29.8 | 1.6 | 125.8 | 766.3 | 69.5 | 19.7 |
| Biomass (biogas) | | 56.1 | 6.0 | 0.1 | 18.4 | 10.6 | 2.0 | 0.5 |
| Gasoline (gardening professional) | 73.9 | | 91.7 | 2.1 | 157.7 | 23416.7 | 2201.1 | 0.4 |
| 1A4 b Other Sectors: Residential | | | | | | | | |
| Gas oil (weighted average) | 73.7 | | 1.0 | 0.6 | 37.0 | 12.6 | 6.0 | 22.4 |
| Gas oil (heat only boilers) | 73.7 | | 1.0 | 0.6 | 37.0 | 12.6 | 6.0 | 22.4 |
| Gas oil (turbines) | NO | | NO | NO | NO | NO | NO | NO |
| Gas oil (engines) | 73.7 | | 2.0 | 0.6 | 40.0 | 30.0 | 8.0 | 21.0 |
| Natural gas (weighted average) | 56.1 | | 6.2 | 0.1 | 17.6 | 14.1 | 4.0 | 0.5 |
| NG (heat only boilers) | 56.1 | | 6.0 | 0.1 | 17.4 | 13.6 | 4.0 | 0.5 |
| NG (turbines) | 56.1 | | 2.0 | 0.1 | 60.0 | 15.0 | 0.1 | 0.5 |
| NG (engines) | 56.1 | | 20.0 | 0.1 | 36.7 | 58.3 | 1.0 | 0.5 |
| Bituminous coal | 92.7 | | 300.0 | 1.6 | 65.0 | 2000.0 | 100.0 | 350.0 |
| Lignite | NO | | NO | NO | NO | NO | NO | NO |
| Biomass | | 90.3 | 83.4 | 1.6 | 91.9 | 1782.3 | 201.8 | 19.5 |
| Gasoline (gardening) | 73.9 | | 51.9 | 2.4 | 154.4 | 24215.4 | 1589.9 | 0.4 |

Emission factors highlighted in grey are explained in section 3.2.5.2.

Emission factors for CO₂, CH₄ and N₂O of charcoal use in the residential sector (1A4bi) are based on the 2006 IPCC guidelines, being the same as those of the revised 1996 IPCC guidelines. CO₂ emission factor for bonfires in the residential sector (1A4bi) is based on SAEFL 2000 and CH₄ and N₂O emission factors are the same as for charcoal use.

Table 3-42 Emission Factors for 1A4a/b Other Sectors Commercial/Institutional and Residential in 2012. CO₂ emission factor is biogenic.

| 1A4bi | Unit | CO ₂ biog. | CH ₄ | N ₂ O | NO _x | CO | NM VOC | SO ₂ |
|-----------------|-------|-----------------------|-----------------|------------------|-----------------|-------|--------|-----------------|
| use of charcoal | kg/TJ | 112'000 | 200 | 1 | 50 | 6'000 | 1'300 | 10 |
| bonfires | kg/TJ | 92'000 | 200 | 1 | 50 | 6'000 | 1'300 | 10 |

Emission factors for mobile off-road sources

- The emission factors for CO₂ are country specific and are assumed to be constant in the period 1990-2012 with values 73.6 t/TJ for diesel oil, 73.9 t/TJ for gasoline and 56.1 t/TJ for CNG (equal to natural gas). See Table 3-9.
- For SO₂ the emission factors are country specific and are given in Table A - 6 in Annex A2.
- The emission factors for all other gases are country specific and shown in Table A - 15 to Table A - 18 in the Annex A3.1.6 (INFRAS 2008). The NMVOC emissions are calculated as the difference of VOC and CH₄ emissions.

Note that emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database INFRAS (2008), see footnote 7 on page 111.

Activity Data

Activity data for the different fuels is determined as described in chapter 3.2.5.1. This includes gas oil, residual fuel oil, natural gas and biomass. For the solid fuels bituminous coal and lignite, activity data is provided from the Swiss overall energy statistics (SFOE 2013).

Table 3-43 Activity data in 1A4a Commercial/Institutional and 1A4b Residential.

| Source/Fuel | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--------------------------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1A4a Commercial/Institutional | TJ | 83'051 | 90'115 | 90'057 | 88'570 | 82'387 | 84'756 | 91'233 | 87'381 | 90'426 | 86'588 |
| Gas oil | TJ | 62'293 | 67'130 | 66'136 | 63'646 | 57'884 | 57'973 | 62'493 | 59'823 | 61'826 | 58'515 |
| Gas oil heat only boilers | TJ | 62'269 | 67'080 | 66'078 | 63'590 | 57'763 | 57'798 | 62'263 | 59'535 | 61'528 | 58'188 |
| Gas oil turbines | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Gas oil engines | TJ | 24 | 51 | 58 | 56 | 122 | 175 | 231 | 288 | 298 | 327 |
| Natural gas | TJ | 17'598 | 19'381 | 20'315 | 21'201 | 20'879 | 22'657 | 24'062 | 23'163 | 23'895 | 23'216 |
| NG heat only boilers | TJ | 17'321 | 18'946 | 19'755 | 20'575 | 20'057 | 21'486 | 22'651 | 21'698 | 22'293 | 21'505 |
| NG turbines | TJ | 85 | 114 | 109 | 106 | 107 | 78 | 21 | 5 | 12 | 4 |
| NG engines | TJ | 192 | 321 | 451 | 520 | 715 | 1'093 | 1'390 | 1'460 | 1'590 | 1'706 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Biomass | TJ | 2'952 | 3'382 | 3'373 | 3'479 | 3'368 | 3'858 | 4'399 | 4'105 | 4'405 | 4'547 |
| Biomass (wood) | TJ | 2'928 | 3'359 | 3'349 | 3'447 | 3'334 | 3'824 | 4'359 | 4'065 | 4'361 | 4'503 |
| Biomass (biogas) | TJ | 24 | 24 | 24 | 31 | 34 | 34 | 40 | 40 | 44 | 44 |
| Gasoline (gardening professional) | TJ | 209 | 221 | 233 | 244 | 256 | 267 | 278 | 289 | 300 | 311 |
| 1A4b Residential | TJ | 186'816 | 199'371 | 199'011 | 190'210 | 179'234 | 193'165 | 200'920 | 186'313 | 192'543 | 189'445 |
| Gas oil | TJ | 138'916 | 145'507 | 145'175 | 136'252 | 128'901 | 137'597 | 139'992 | 131'915 | 136'508 | 131'838 |
| Gas oil heat only boilers | TJ | 138'915 | 145'506 | 145'173 | 136'251 | 128'900 | 137'593 | 139'961 | 131'877 | 136'459 | 131'785 |
| Gas oil turbines | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Gas oil engines | TJ | 1 | 1 | 1 | 1 | 1 | 4 | 32 | 38 | 49 | 53 |
| Natural gas | TJ | 25'390 | 29'240 | 30'680 | 31'090 | 29'530 | 33'760 | 38'000 | 34'420 | 35'980 | 38'040 |
| NG heat only boilers | TJ | 25'330 | 29'138 | 30'536 | 30'922 | 29'326 | 33'502 | 37'693 | 34'107 | 35'630 | 37'635 |
| NG turbines | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| NG engines | TJ | 60 | 102 | 144 | 168 | 204 | 258 | 307 | 313 | 350 | 405 |
| Bituminous coal | TJ | 589 | 680 | 471 | 480 | 435 | 417 | 236 | 199 | 127 | 127 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Biomass | TJ | 21'922 | 23'945 | 22'685 | 22'388 | 20'368 | 21'391 | 22'692 | 19'779 | 19'928 | 19'441 |
| Gasoline (gardening) | TJ | 145 | 147 | 150 | 153 | 155 | 158 | 160 | 162 | 165 | 167 |

| Source/Fuel | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--------------------------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1A4a Commercial/Institutional | TJ | 84'398 | 88'139 | 82'656 | 86'877 | 86'101 | 87'220 | 82'030 | 73'800 | 79'215 | 76'286 |
| Gas oil | TJ | 55'485 | 57'502 | 53'946 | 56'415 | 54'631 | 54'629 | 51'065 | 44'249 | 47'255 | 44'626 |
| Gas oil heat only boilers | TJ | 55'135 | 57'135 | 53'594 | 56'082 | 54'306 | 54'311 | 50'772 | 44'067 | 47'085 | 44'472 |
| Gas oil turbines | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Gas oil engines | TJ | 351 | 367 | 352 | 333 | 326 | 318 | 293 | 181 | 169 | 154 |
| Natural gas | TJ | 24'257 | 25'609 | 23'812 | 25'127 | 26'026 | 26'986 | 24'719 | 23'468 | 25'283 | 24'670 |
| NG heat only boilers | TJ | 22'521 | 23'803 | 21'902 | 23'130 | 24'059 | 24'953 | 22'767 | 21'542 | 23'425 | 22'856 |
| NG turbines | TJ | 0 | 3 | 12 | 28 | 31 | 28 | 23 | 28 | 29 | 26 |
| NG engines | TJ | 1'737 | 1'803 | 1'899 | 1'968 | 1'937 | 2'004 | 1'929 | 1'898 | 1'829 | 1'787 |
| Bituminous coal | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Biomass | TJ | 4'334 | 4'713 | 4'590 | 5'034 | 5'148 | 5'318 | 5'959 | 5'796 | 6'391 | 6'705 |
| Biomass (wood) | TJ | 4'289 | 4'660 | 4'530 | 4'966 | 5'065 | 5'204 | 5'778 | 5'506 | 6'036 | 6'307 |
| Biomass (biogas) | TJ | 45 | 53 | 60 | 69 | 83 | 114 | 181 | 290 | 355 | 398 |
| Gasoline (gardening professional) | TJ | 321 | 315 | 308 | 301 | 295 | 288 | 287 | 286 | 286 | 285 |
| 1A4b Residential | TJ | 174'816 | 184'251 | 177'744 | 188'017 | 188'109 | 191'375 | 184'262 | 164'219 | 175'973 | 172'543 |
| Gas oil | TJ | 120'784 | 127'553 | 122'470 | 129'328 | 128'194 | 129'613 | 124'415 | 107'798 | 114'325 | 110'985 |
| Gas oil heat only boilers | TJ | 120'731 | 127'498 | 122'414 | 129'269 | 128'119 | 129'550 | 124'352 | 107'733 | 114'273 | 110'944 |
| Gas oil turbines | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Gas oil engines | TJ | 53 | 55 | 56 | 58 | 74 | 63 | 63 | 65 | 52 | 42 |
| Natural gas | TJ | 36'290 | 38'000 | 37'790 | 40'330 | 41'660 | 42'790 | 41'080 | 39'320 | 42'550 | 42'630 |
| NG heat only boilers | TJ | 35'851 | 37'539 | 37'325 | 39'813 | 41'153 | 42'260 | 40'538 | 38'775 | 42'009 | 42'092 |
| NG turbines | TJ | 0 | 0 | 5 | 3 | 2 | 0 | 0 | 3 | 3 | 0 |
| NG engines | TJ | 439 | 461 | 460 | 514 | 505 | 530 | 542 | 542 | 537 | 538 |
| Bituminous coal | TJ | 118 | 118 | 118 | 118 | 362 | 362 | 362 | 362 | 362 | 362 |
| Lignite | TJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Biomass | TJ | 17'624 | 18'581 | 17'366 | 18'241 | 17'893 | 18'610 | 18'405 | 16'739 | 18'736 | 18'565 |
| Gasoline (gardening) | TJ | 169 | 167 | 165 | 162 | 160 | 158 | 157 | 156 | 155 | 155 |

| Source/Fuel | Unit | 2010 | 2011 | 2012 |
|--------------------------------------|------|---------|---------|---------|
| 1A4a Commercial/Institutional | TJ | 82'310 | 68'483 | 74'788 |
| Gas oil | TJ | 47'763 | 39'177 | 40'950 |
| Gas oil heat only boilers | TJ | 47'640 | 39'066 | 40'847 |
| Gas oil turbines | TJ | 0 | 0 | 0 |
| Gas oil engines | TJ | 122 | 111 | 103 |
| Natural gas | TJ | 27'067 | 22'876 | 26'431 |
| NG heat only boilers | TJ | 25'313 | 21'202 | 24'737 |
| NG turbines | TJ | 23 | 17 | 5 |
| NG engines | TJ | 1'730 | 1'657 | 1'688 |
| Bituminous coal | TJ | 0 | 0 | 0 |
| Lignite | TJ | 0 | 0 | 0 |
| Biomass | TJ | 7'197 | 6'148 | 7'130 |
| Biomass (wood) | TJ | 6'709 | 5'602 | 6'463 |
| Biomass (biogas) | TJ | 488 | 546 | 667 |
| Gasoline (gardening professional) | TJ | 284 | 281 | 277 |
| 1A4b Residential | TJ | 187'067 | 150'376 | 165'972 |
| Gas oil | TJ | 118'021 | 92'168 | 99'913 |
| Gas oil heat only boilers | TJ | 117'984 | 92'133 | 99'878 |
| Gas oil turbines | TJ | 0 | 0 | 0 |
| Gas oil engines | TJ | 37 | 36 | 34 |
| Natural gas | TJ | 48'390 | 41'070 | 47'230 |
| NG heat only boilers | TJ | 47'870 | 40'571 | 46'722 |
| NG turbines | TJ | 0 | 0 | 0 |
| NG engines | TJ | 520 | 499 | 508 |
| Bituminous coal | TJ | 362 | 362 | 362 |
| Lignite | TJ | 0 | 0 | 0 |
| Biomass | TJ | 20'293 | 16'775 | 18'467 |
| Gasoline (gardening) | TJ | 154 | 151 | 148 |

The table above documents the use of gas oil (55%), natural gas (35%) and biomass (10%) as fuels consumed in source category 1A4a Commercial/Institutional. Since 1990, fuel consumption in this source category 1A4a reduced by 10%. Within the fuel consumption, a major fuel switch can be observed in this source category from gas oil (-34%) to natural gas (+50%) and biomass (+142%).

Regarding source category 1A4b Residential, the major fuels consumed are gas oil (60%), natural gas (28%) and biomass (11%). Since 1990, fuel consumption in this source category 1A4b reduced by 11%. Also in this source category, a fuel switch from gas oil (-28%) to natural gas (+86%) can be observed. Biomass consumption for heating purposes of residential buildings diminished from 1990 to 2012 by 16%.

This shift in fuel mix is the reason for CO₂ emissions from the use of natural gas and liquid fuels in category 1A4a and 1A4b being key categories regarding trend.

Underlying data for the activity data on mobile off-road sources (1A4aii and 1A4bii) like vehicle stock and operating hours are shown in Annex A3.1.6. A slight modification of the activity data was carried out in 2013 (Prognos 2012a, see also Chapter 3.2.9.5). The consumption of gasoline (gardening) has been recalculated accordingly. Numbers from 2005 onwards are affected.

Table 3-44 Activity data in 1A4bi charcoal consumption.

| 1A4bi | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-----------------|------|------|------|------|------|------|------|------|------|------|------|
| use of charcoal | TJ | 311 | 315 | 318 | 322 | 292 | 291 | 292 | 272 | 302 | 282 |
| bonfires | TJ | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 |

| 1A4bi | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-----------------|------|------|------|------|------|------|------|------|------|------|------|
| use of charcoal | TJ | 292 | 362 | 332 | 302 | 282 | 313 | 303 | 313 | 353 | 343 |
| bonfires | TJ | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 | 160 |

| 1A4bi | Unit | 2010 | 2011 | 2012 |
|-----------------|------|------|------|------|
| use of charcoal | TJ | 343 | 343 | 343 |
| bonfires | TJ | 160 | 160 | 160 |

Charcoal consumption has slightly increased from 1990 to 2009 and since 2009, a stable consumption of charcoal is assumed (SFOE 2013, see documentation in EMIS 2014/1A4bi Holzkohle-Verbrauch). Bonfires are assumed constant over the whole time series as there are no official numbers available. As charcoal consumption is more or less constant, it is assumed, that bonfires also are constant over time and correspond to an estimation of 2kg of wood for bonfires per habitant (see documentation in EMIS 2014/1A4bi Lagerfeuer).

For source category 1A4ci, the following activity data is reported:

Drying of grass: Activity data on grass drying (in tons of dried grass) is extracted from the EMIS database (EMIS 2014/1A4ci).

Off-road machinery: Activity data is shown in Annex A3.1.6 (INFRAS 2008, Prognos 2012a, Keller/INFRAS 2013).

Biomass: Activity data is based on Swiss wood energy statistics (SFOE 2013b) as explained in 3.2.5.1.

Table 3-45 Activity data in 1A4c Agriculture/Forestry.

| Source/Fuel | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1A4c Agriculture/Forestry | TJ | 8'597 | 8'623 | 8'568 | 8'546 | 8'489 | 8'494 | 8'518 | 8'423 | 8'415 | 8'391 |
| Drying of Grass | TJ | 1'895 | 1'828 | 1'748 | 1'683 | 1'620 | 1'544 | 1'482 | 1'409 | 1'349 | 1'291 |
| gas oil | TJ | 1'156 | 1'115 | 1'066 | 1'027 | 988 | 942 | 904 | 860 | 823 | 787 |
| natural gas | TJ | 739 | 713 | 682 | 657 | 632 | 602 | 578 | 550 | 526 | 503 |
| Machinery (diesel, gasoline) | TJ | 6'275 | 6'308 | 6'342 | 6'375 | 6'409 | 6'443 | 6'471 | 6'500 | 6'529 | 6'558 |
| Biomass | TJ | 427 | 487 | 478 | 488 | 461 | 508 | 564 | 514 | 536 | 542 |

| Source/Fuel | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1A4c Agriculture/Forestry | TJ | 8'316 | 8'211 | 8'174 | 8'211 | 8'212 | 8'240 | 8'099 | 8'233 | 8'237 | 8'344 |
| Drying of Grass | TJ | 1'223 | 1'077 | 1'061 | 1'055 | 1'039 | 994 | 845 | 948 | 822 | 856 |
| gas oil | TJ | 746 | 657 | 647 | 644 | 634 | 607 | 516 | 579 | 502 | 522 |
| natural gas | TJ | 477 | 420 | 414 | 412 | 405 | 388 | 330 | 370 | 321 | 334 |
| Machinery (diesel, gasoline) | TJ | 6'587 | 6'588 | 6'589 | 6'590 | 6'591 | 6'592 | 6'658 | 6'724 | 6'791 | 6'857 |
| Biomass | TJ | 506 | 546 | 524 | 566 | 582 | 654 | 596 | 560 | 624 | 631 |

| Source/Fuel | Unit | 2010 | 2011 | 2012 |
|----------------------------------|------|-------|-------|-------|
| 1A4c Agriculture/Forestry | TJ | 8'361 | 8'332 | 8'376 |
| Drying of Grass | TJ | 739 | 891 | 845 |
| gas oil | TJ | 451 | 543 | 515 |
| natural gas | TJ | 288 | 347 | 329 |
| Machinery (diesel, gasoline) | TJ | 6'923 | 6'871 | 6'818 |
| Biomass | TJ | 699 | 571 | 714 |

The table above documents the fuel use in source category 1A4c Agriculture/Forestry. Machinery is the major source with 81% fuel consumption of the source category 1A4c compared to grass drying with 10% and biomass with 9%. Fuel consumption in machinery increased by 8.7% since 1990. For grass drying, fuel consumption is divided in gas oil (61%) and natural gas (39%). Since 1990, the fuel consumption significantly decreased by 55% for gas oil as well as natural gas. Biomass consumption increased by 67%.

In the last submission in 2013 a mistake occurred in the table above as the year 2011 had the same values as in the year 2001 for the line "gas oil", "natural gas" and "machinery" and there was one process not calculated under "machinery". This error is now corrected in current Table 3-44.

3.2.9.3 Uncertainties and Time-Series Consistency

The uncertainty of CO₂ emissions from fuel combustions is described in the uncertainty analysis of the Energy Industries (1A1) in Chapter 3.2.6.3. Uncertainty in emissions of other non-CO₂ gases are estimated to be medium: 30% for CH₄ and 80% for N₂O (see Table 1-14).

A general description of the time series consistency of the Energy Sector is provided in Chapter 3.2.6.3.

3.2.9.4 Source-specific QA/QC and Verification

a) General

See Chapter 3.2.6.4.

b) Specific: Other sectors (1A4)

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables.

- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.

For A4ci grass drying: The fuel consumption was verified in 2003 by a statistical analysis of 20 typical grass drying plants (VSTB 2003).

3.2.9.5 Source-Specific Recalculations

- **1A all fuels:** Activity data of all fuels of the overall time series have been recalculated based on the new data from SFOE 2013. This includes data on 2011 for fossil fuels and waste, data for biomass on 1997, 2009 – 2011 and data of other fuels 1990 – 2011.
- **1A gaseous fuels:** Activity data have been recalculated based on new data of from SFOE 2013.
- **1A wood consumption:** Activity data have been recalculated for the overall time series based on the new data from SFOE 2013b.
- **1A gas oil:** SO_x emission factor value has been updated for 2010 based on sulphur analyses of the gas oil for the year 2010 (Directorate General of Customs) resulting in a revised value for 2011 as well.
- **1A bituminous coal:** CO₂ emission factor has been corrected from 94t CO₂/TJ to 92.7 t CO₂/TJ for the entire time series.
- **1A4a/1A4b engines and gasturbines:** Activity data in 1A4 households and services have been updated for 2011 based on updated statistical data.
- **1A4a engines, natural gas:** Activity data for natural gas in the commercial and institutional sector have been updated for 1990, 1995, 1997, 1998 and 2011 based on recalculations in SFOE 2013.
- **1A4a engines, natural gas:** Activity data for natural gas have been updated based on a recalculation of SFOE 2013 for the year 2011.
- **1A4a engines, gas oil and natural gas:** Activity data for gas oil and natural gas consumption have been updated for the overall time series based on changes in the energy model resulting from changes in the non-road transport model for boats and natural gas consumption in industry.
- **1A4b engines, natural gas:** Activity data have been updated for 1990, 1995, 1997, 1998 and 2011 based on recalculations in SFOE 2013.
- **1A4b engines, gas oil:** Activity data for gas oil have been updated for the whole time series based on recalculations in the energy consumption.
- **1A4b gas oil and natural gas:** CO and SO₂ emission factors for gas oil and natural gas of 2011 have been corrected in the energy model.
- **1A4b charcoal consumption:** Reporting of charcoal consumption has been shifted to sector 1A4b. Before, it was reported under 2D3. (Please note that the reporting of the CH₄ emissions from the charcoal production has already been shifted from 2D3 to 1A1c within the resubmission of Switzerland's Greenhouse Gas Inventory 1990–2011(FOEN2013g).)
- **1A4b bonfires:** A new process has been introduced in this submission.
- **1A4aii, 1A4bii, 1A4cii mobile off-road machinery:** The activity data have so far been taken from INFRAS (2008). For this submission, the latest numbers on growth of population and economy (Prognos 2012a, Keller/INFRAS 2013) have been integrated

in the off-road model. This leads to an increase of the fuel consumption from 2005 onwards.

3.2.9.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.2.10 Source Category 1A5b - Military

3.2.10.1 Source Category Description

Tier 1 Key categories 1A5

CO₂ from the combustion of Liquid Fuels (trend)

In Switzerland, the source categories are defined according to the next table. The IPCC category structure distinguishes stationary (1A5a) and mobile (1A5b) sources. All of the Swiss sub-categories refer to mobile sources.

Table 3-46 Specification of Swiss source category 1A5 Other (Military).

| 1A5 | Source | Specification | Data Source |
|------|-------------------------------------|--|---|
| 1A5a | Stationary | Not occurring in Switzerland (NO) | |
| 1A5b | Mobile Military off-road sources | Tanks and similar off-road vehicles (emissions from military road vehicles are included in 1A3b Road Transportation) | Method, AD, EF: INFRAS 2008, AD: Prognos 2012, Keller/INFRAS 2013 |
| 1A5b | Military Aviation | | VTG 2013 |

3.2.10.2 Methodological Issues

a) Military off-road vehicles

The emissions of military off-road machinery (excluding aviation) are modelled by the same approach as all other mobile off-road sources using the off-road model developed by INFRAS (2008). For details refer to Chapter 3.2.5.1 (paragraph on source category 1A2fii).

b) Military aviation

To calculate the emissions from military aviation, a Tier 1 method is used. The fuel consumption 1990–2012 is known on an annual basis (VTG 2013). A very small fraction of fuel is consumed for training abroad and might be allocated under “International Bunkers” (less than 3% of total military aviation consumption). Since the exact number is not known, it is not subtracted from the total consumption but included under national military aviation, as recommended by the IPCC Good Practice Guidance (IPCC 2000, chapter 2.5.1.3). Emissions of NO_x, CO and VOC have been modelled in detail by the Federal Office for Military Aviation (Bundesamt für Betriebe der Luftwaffe) for 1990 and 1995. From these inputs, FOEN determined average emission factors 1990 and 1995. For 1991–1994 the emission factors are linearly interpolated between 1990 and 1995. For 1996–2011, the factors for 1995 are used. The emissions are then calculated yearly by multiplying the average emission factors with the activity data.

The extension of the emission modelling to CO₂, CH₄, N₂O, NMVOC and SO₂ is also accomplished by FOEN.

Emission factors for military off-road vehicles

- The emission factors for CO₂ are country specific and are assumed to be constant in the period 1990-2012 with values 73.6 t/TJ for diesel oil, 73.9 t/TJ for gasoline and 55.0 t/TJ for CNG (equal to natural gas), see Table 3-9.
- For SO₂ the emission factors are country specific and are given in Table A - 6 in Annex A2.
- The emission factors for all other gases are country specific and shown in Table A - 15 to Table A - 18 in the Annex A3.1.6 (INFRAS 2008) The NMVOC emissions are calculated as the difference of VOC and CH₄ emissions.

Note that emission factors in the unit of kg/h may be downloaded by query from the public part of the off-road database INFRAS (2008), see footnote 7 on page 111.

Emission factors for military aviation

- CO₂: The emission factor of 73.2 t/TJ is country specific and is based on measurements and analyses of fuel samples (see Table 3-9, SFOE 2001, FOEN 2011k, Intertek 2008).
- NO_x, VOC, CO: Engine producer information is used (CORINAIR, for details see SAEFL 1996: p. 202) for calculation of the emission factors in 1990 and 1995. For 1991-1994 the values are linearly interpolated between 1990 and 1995. For 1996-2012, the values 1995 are used.
- CH₄, NMVOC: For VOC, aircraft-specific information used for calculation of the emission factors in 1990 and 1995. For 1991-1994 the values are linearly interpolated between 1990 and 1995. For 1996-2012, the values 1995 are used. The division of VOC into CH₄ and NMVOC is carried out by a constant split of 53%: 47% (country specific).
- N₂O: The implemented emission factor for N₂O is 2.356 kg/TJ. By mistake, in previous submissions the emission factor was defined as 23 kg/TJ.
- SO₂: The emission factor is taken from the IPCC Guidelines 1996, 23.3 kg/TJ, and is assumed to be constant over the period 1990–2012 (IPCC 1997c, Table 1-50)

Activity data for military off-road vehicles and military aviation

Fuel consumption data is shown in Table 3-47. The underlying data for military off-road such as vehicle stock and operating hours are shown in Table A - 24 and Table A - 25 in Annex A3.1.6.

Fuel consumption of military aviation is copied from the logbooks of the military aircrafts and summed up yearly (VTG 2013).

Table 3-47 Activity data (fuel consumption) for military off-road vehicles and military aviation

| 1A5 | Fuel | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------|--------------|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | fuel consumption in TJ | | | | | | | | | |
| Military off-road | Diesel | 48 | 48 | 48 | 48 | 49 | 49 | 49 | 49 | 50 | 50 |
| Military off-road | Gasoline | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| Military aviation | Jet kerosene | 2'733 | 2'495 | 2'382 | 2'268 | 2'192 | 1'955 | 1'806 | 1'941 | 1'927 | 1'734 |

| 1A5 | Fuel | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------|--------------|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | fuel consumption in TJ | | | | | | | | | |
| Military off-road | Diesel | 50 | 50 | 49 | 49 | 48 | 48 | 48 | 48 | 48 | 48 |
| Military off-road | Gasoline | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| Military aviation | Jet kerosene | 1'793 | 1'755 | 1'837 | 1'641 | 1'488 | 1'621 | 1'672 | 1'572 | 1'500 | 1'524 |

| 1A5 | Fuel | 2010 | 2011 | 2012 |
|-------------------|--------------|------------------------|-------|-------|
| | | fuel consumption in TJ | | |
| Military off-road | Diesel | 48 | 48 | 47 |
| Military off-road | Gasoline | 0.6 | 0.6 | 0.6 |
| Military aviation | Jet kerosene | 1'586 | 1'414 | 1'521 |

3.2.10.3 Uncertainties and Time-Series Consistency

a) General

For a general description of the uncertainty analysis and time series consistency of the Energy Sector see Chapter 3.2.6.3 a).

b) Specific

See Chapter 3.2.6.3.

3.2.10.4 Source-specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF.
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013.

The activity data of military aviation (kerosene consumption) are provided by the Federal Department of Defence, Civil Protection and Sport. For a compatibility check with the emission data base of civil aviation, they are sent to the FOCA (office of the Federal Department of the Environment, Transport, Energy and Communications). A further compatibility check is carried out by the NIR authors of the energy chapter. No peculiarities have been detected by the specialists in the time series of the kerosene consumption of military aviation.

3.2.10.5 Source-Specific Recalculations

- 1A5b: The activity data have so far been taken from INFRAS (2008). For this submission, the latest numbers on growth of population and economy (Prognos 2012a, Keller/INFRAS 2013) have been integrated in the off-road model. This leads to an increase of the fuel consumption from 2005 onwards.

3.2.10.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. To accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.3 Source Category 1B – Fugitive Emissions from Fuels

Fugitive emissions arise from the production, processing, transmission, storage and use of fuels. According to IPCC guidelines, emissions from venting and flaring at oil and gas production facilities are included while emissions from vehicles are not included in 1B.

Source Category 1B Fugitive Emissions from Fuels comprises the following sub-categories:

- Solid fuels (1B1)
- Oil (1B2a)
- Natural Gas (1B2b)

Tier 1 Key categories 1B2

CH₄ from fugitive emissions of Oil and Natural Gas (level and trend)

Tier 2 Key categories 1B2

CH₄ from fugitive emissions of Oil and Natural Gas (trend)

3.3.1 Source Category 1B1 - Solid Fuels

Coal mining is not occurring in Switzerland.

3.3.2 Source Category 1B2a - Oil

3.3.2.1 Source Category Description

In Switzerland, oil production is not occurring. Fugitive emissions in the oil industry result from two refining companies and several fuel handling stations. Production from the refining companies cover around 40% of the oil consumption in Switzerland. The other 60% are imports of final products. Oil pipelines are very short in Switzerland (approximately 40km and 70km) and are mainly underground.

The following source categories occur in Switzerland:

- Transport (1B2a iii)
- Refining / Storage (1B2a iv)
- Distribution of Oil Products from storage tanks and gasoline stations (1B2a v)

Table 3-48 Specification of source category 1B2a Fugitive Emissions from Oil.

EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1B2 | Source | Specification | Data Source |
|-------|--------|---|---|
| 1B2 a | Oil | Emissions from refining/storage of oil and the distribution of oil products | AD: EV 2013, SFOE 2013 EF: IPCC 2006, EMIS 2014/1B2a |

3.3.2.2 Methodological Issues

For source 1B2a Oil, emissions of CO₂, CH₄ and NMVOC are reported. CO₂ emissions occur through the oxidation of volatile organic compounds in the atmosphere. CH₄ emissions only occur in 1B2a iii.

Emissions are calculated based on a Tier 1 approach. Fugitive emissions from fuels are calculated by multiplying level of activity by emission factor.

Emission factors

The following table presents the emission factors used in 1B2a Oil:

Table 3-49 Emission Factors for 1B2a Oil in 2012.

| Source/fuel | CO ₂ | CH ₄ | N ₂ O | NO _x | CO | NM VOC | SO ₂ |
|--------------------------|-----------------|-----------------|------------------|-----------------|-------|--------|-----------------|
| | g/t | g/t | kg/TJ | kg/TJ | kg/TJ | g/t | kg/TJ |
| 1B2a Oil Products | | | | | | | |
| Oil Transport | 0.60 | 6.59 | NA | NA | NA | 65.9 | NA |
| Oil Refining and Storage | 1356 | 45 | NA | NA | NA | 430 | NA |
| Gasoline storage tank | 893 | NA | NA | NA | NA | 283 | NA |
| Gasoline station | 1240 | NA | NA | NA | NA | 393 | NA |

For oil transport (1B2a iii), the default value from IPCC 2006 for pipeline transport is used to calculate emissions.

For oil refining and storage (1B2a iv), country specific emission factors for CH₄ and NMVOC are used. The conversion factor used to calculate the CO₂ emission factor bases on the NIR of the Netherlands.

For oil distribution from storage tanks and gasoline stations (1B2a v), CO₂ emission factors base on the conversion factor from the NIR of the Netherlands. NMVOC emission factor for oil distribution from tanks is taken from country specific calculations based on statistical information from the Energy Model based on data from SFOE 2013 (see documentation in EMIS 2014/1B2a v Benzinumschlag Tanklager). NMVOC emission factor from gasoline stations is calculated based on country specific information (see documentation in EMIS 2014/1B2a v Benzinumschlag Tankstellen).

Activity data

Table 3-50 Activity data (fuel consumption) for 1B2a Oil

| 1B2a Oil Products | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Crude oil | t | 3127315 | 4671296 | 4317130 | 4763889 | 4879630 | 4657407 | 5289352 | 4830324 | 5069907 | 5093056 |
| Gasoline transport | t | 3682727 | 3834439 | 3972485 | 3682707 | 3682344 | 3568755 | 3660597 | 3800552 | 3829912 | 3956876 |

| 1B2a Oil Products | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Crude oil | t | 4649074 | 4846065 | 4848380 | 4567130 | 5146296 | 4810417 | 5496528 | 4661806 | 5066898 | 4778472 |
| Gasoline transport | t | 3958527 | 3849300 | 3773347 | 3755292 | 3687960 | 3575627 | 3466291 | 3432200 | 3357092 | 3266530 |

| 1B2a Oil Products | Unit | 2010 | 2011 | 2012 |
|--------------------|------|---------|---------|---------|
| Crude oil | t | 4490741 | 4402315 | 3408796 |
| Gasoline transport | t | 3151135 | 3029864 | 2922408 |

For oil transport (1B2a iii), crude oil used is based on annual statistics of the Swiss Petroleum Association (EV 2013).

For oil refining and storage (1B2a iv), crude oil used is based on annual statistics of the Swiss Petroleum Association (EV 2013).

For oil distribution from tanks and gasoline stations (1B2a v), data is provided by the Energy Model based on SFOE 2013 (see Section 3.2.5.1). As the statistics include also oil consumption of Liechtenstein, this is subtracted from the data provided to represent oil consumption of Switzerland only.

Since 1990, Crude oil production increased slightly by 9% and Gasoline transport decreased by 21%.

3.3.2.3 Uncertainties and Time-Series Consistency

A preliminary uncertainty assessment of all other sources in source category 1B2 based on expert judgement results in medium confidence in the emissions estimate (see Table 1-14).

The time series is consistent.

3.3.2.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.

3.3.2.5 Source-Specific Recalculations

- **1B2a v distribution of oil products:** Emission factor for CO₂ and NMVOC were corrected to exclude emissions from Liechtenstein that overestimated emissions by 0.5%.

3.3.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.3.3 Source Category 1B2b - Natural Gas

3.3.3.1 Source Category Description

Emissions from natural gas production are only occurring for the years of operation of the single production plant in Switzerland from 1985 - 1994. Other emissions in this sector occur from natural gas transmission and distribution. Emission from transmission also include leakages from gas pipelines. Major accidents and isolated events are reported under other leakage.

The following source categories occur in Switzerland:

- Production (1B2b ii) (only 1990-1994)
- Transmission (1B2b iii)
- Distribution (1B2b iv)
- Other leakage (1B2b v) (isolated events in single years)

Table 3-51 Specification of source category 1B2b Fugitive Emissions from Natural Gas.

EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1B2 | Source | Specification | Data Source |
|-------|-------------|------------------------------|--|
| 1B2 b | Natural Gas | Emissions from gas pipelines | AD: SFOE 2013, Quantis 2014 EF: IPCC 2006, Quantis 2014 |

3.3.3.2 Methodological Issues

For source 1B2b Natural Gas, emissions of CO₂, CH₄ and NMVOC are reported for natural gas production as well as transmission, distribution and related accidents. Emissions are calculated based on annual production data which is consistent with the IPCC Tier 1 approach. Emission from leakages from gas pipelines are calculated with a country specific method. The method considers the length, type and pressure of the gas pipelines as well as the annual gas consumption. The distribution network components (regulators, shut off fittings and gas meters), the losses from maintenance and extension as well as the end user losses are separately taken into account.

Fugitive emissions from fuels are calculated by multiplying level of activity by emission factor.

Emission factors

CO₂, CH₄ and NMVOC Emission factors for natural gas production is provided by the default values of IPCC 2006 Guidelines as documented in EMIS 2014/1B2b Gasproduktion.

Emission factors for gas transport and distribution losses (source 1B2b iii and 1b2 iv) are based on a new study realized by Quantis (Quantis 2014). Emission factors are provided by literature and base mostly on the study of Batelle 1994 that provides specific emission factors for different sources of fugitive emissions based on measurements of 1998 in Germany. Specific data for Switzerland is provided by a study of Xinmin (2004), but also these emission factors are mostly based on Batelle (1994).

Table 3-52 Emission factors for 1B2b Natural Gas and 1B2d Other (only occurring in 2010).

| Source/fuel | CO ₂ g/GJ | CH ₄ g/GJ | N ₂ O g/GJ | NO _x g/GJ | CO g/GJ | NMVOC g/GJ | SO ₂ g/GJ |
|------------------|-------------------------|-------------------------|--------------------------|-------------------------|------------|---------------|-------------------------|
| 1B2b Natural Gas | | | | | | | |
| Production | NO | NO | NA | NA | NA | NO | NA |
| Transmission | 312 | 17'938 | NA | NA | NA | 17'938 | NA |
| Distribution | 312 | 17'938 | NA | NA | NA | 0.002 | NA |
| 1B2d Other | 312 | 17'938 | NA | NA | NA | 0.002 | NA |

Activity data

Table 3-53 Activity data (fuel consumption) for 1B2b Natural Gas and 1B2d Other (only occurring in 2010).

| 1B2b Natural Gas / 1B2d Other | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1B2b Natural Gas | | | | | | | | | | | |
| Production | GJ | 130'000 | 110'000 | 100'000 | 80'000 | 30'000 | 0 | 0 | 0 | 0 | 0 |
| Transmission | GJ | 13'237 | 13'269 | 13'313 | 13'347 | 13'156 | 13'254 | 13'261 | 13'261 | 13'257 | 13'160 |
| Distribution | GJ | 676'096 | 714'632 | 749'911 | 783'021 | 804'621 | 835'693 | 863'669 | 890'494 | 910'730 | 838'810 |
| 1B2d Other | GJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 1B2b Natural Gas / 1B2d Other | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1B2b Natural Gas | | | | | | | | | | | |
| Production | GJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Transmission | GJ | 13'138 | 13'113 | 13'087 | 13'061 | 13'036 | 13'014 | 12'993 | 13'291 | 13'310 | 13'301 |
| Distribution | GJ | 772'213 | 711'334 | 656'231 | 606'938 | 563'463 | 525'790 | 493'880 | 478'854 | 459'030 | 443'559 |
| 1B2d Other | GJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 1B2b Natural Gas / 1B2d Other | Unit | 2010 | 2011 | 2012 |
|-------------------------------|------|---------|---------|---------|
| 1B2b Natural Gas | | | | |
| Production | GJ | 0 | 0 | 0 |
| Transmission | GJ | 13'218 | 13'214 | 13'229 |
| Distribution | GJ | 431'079 | 426'425 | 426'145 |
| 1B2d Other | GJ | 28'899 | 0 | 0 |

Activity data for natural gas production are extracted from the Swiss overall energy statistics (SFOE 2013).

Activity data for gas transport and distribution losses (source 1B2b iii and 1B2b iv) are based on information from the Swiss Gas and Water Industry Association (SGWA) as documented in the study realized by Quantis (Quantis 2014). Since 1990, the natural gas net increased by 73% from 142'000 km in 1990 to 246'000 km in 2012. In the same period, the natural gas consumption increased by 62% from 21'000 to 34'000 GWh.

Within the different sectors, continuous leakage emissions from pipelines are the mayor emission source followed by fugitive emissions from the end user losses that are dominated by the emissions from industry and power stations using natural gas. Fugitive emissions from damages and ruptures of the pipelines, maintenance of the pipelines and the components are very small (Quantis 2014).

It can be observed in the table above, that the fugitive emissions from gas transport and distribution losses strongly decreased between 1990 and 2012. Total CH₄ emissions decreased by 36% based on the strong decrease of fugitive emissions from the continuous leakage by more than 50% which can be explained by the change of material used for the pipelines from cast-iron pipes to polyethylene pipelines. Fugitive emission from the end user losses increased slightly. This switch is also visible in the respective contribution of these two sources between 1990 and 2012. While fugitive emissions from continuous leakage contributed with more than 80% to the total CH₄ emissions in 1990, it only contributes with 60% in 2012.

3.3.3.3 Uncertainties and Time-Series Consistency

a) Uncertainty in fugitive CH₄ emissions from natural gas pipelines in 1B2

Following Good Practice Guidance (IPCC 2000: p. 2.92) overall uncertainty of bottom-up inventories of fugitive methane losses from gas activities are expected to result in errors of 25-50%. From this a conservative uncertainty of 50% is estimated for Switzerland.

b) Qualitative estimate of uncertainties of non-key category emissions in 1B2 Fugitive Emissions from Fuels

A preliminary uncertainty assessment of all other sources in source category 1B2 based on expert judgement results in medium confidence in the emissions estimate (see Table 1-14).

The time series is consistent.

3.3.3.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.

3.3.3.5 Source-Specific Recalculations

- **1B2b natural gas production:** As recommended in ARR 2012 and 2013 Switzerland has included in his inventory the emissions from the only small plant for natural gas production (Finsterwald) from 1985-1994. The default emission factors from IPCC Guidelines 2006 is used.
- **1B2b natural gas production:** Activity data from 1990-1997 has been updated based on data from SFOE 2013.
- **1B2b natural gas production:** A new model for emissions in swiss gas transport system based on a new study realized by Quantis has been developed (Quantis 2014).

3.3.3.6 Source-Specific Planned Improvements

Regarding the emission factor of CH₄ emissions of natural gas pipelines, an error has been detected during the internal review that leads to an underestimation of the emissions based on a too low emission factor used for the calculation. The error will be corrected for the next submission.

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

3.3.4 Source Category 1B2c - Venting

3.3.4.1 Source Category Description

In Switzerland, oil or natural gas production is not occurring. The fugitive emissions from venting result only from the torches in the two refining companies (1B2c i Flaring).

Table 3-54 Specification of source category 1B2c Fugitive Emissions from Venting.
EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 1B2 | Source | Specification | Data Source |
|-------|-------------------|--|-----------------------------------|
| 1B2 c | Venting / Flaring | The release/combustion of excess gas at the oil refinery | AD: EV 2013 EF: EMIS 2014/1B2c |

3.3.4.2 Methodological Issues

Fugitive emissions from fuels are calculated by multiplying level of activity by emission factor.

For source category 1B2c Venting/Flaring (Oil), CO₂ as well as CH₄, N₂O, NO_x, CO and NMVOC are considered. The emissions venting/flaring are calculated based on country specific annual production/consumption data.

Emission factors

Emission factors are based on data from the refining industry and expert estimates as documented in EMIS 2014/1B2c Raffinerie Abfackelung.

Table 3-55 Emission factors for 1B2c Venting/Flaring.

| Source/fuel | CO ₂ g/t | CH ₄ g/t | N ₂ O g/t | NO _x g/t | CO g/t | NMVOC g/t | SO ₂ g/t |
|------------------------|------------------------|------------------------|-------------------------|------------------------|-----------|--------------|------------------------|
| 1B2c Venting / Flaring | 8300 | 3.5 | 0.64 | 43 | 10 | 3.5 | 62 |

Activity data

Table 3-56 Activity data for 1B2c Venting/Flaring.

| 1B2c Venting/Flaring | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------------|------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| Crude Oil used | t | 3'127'315 | 4'671'296 | 4'317'130 | 4'763'889 | 4'879'630 | 4'657'407 | 5'289'352 | 4'830'324 | 5'069'907 | 5'093'056 |

| 1B2c Venting/Flaring | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------------|------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| Crude Oil used | t | 4'649'074 | 4'846'065 | 4'848'380 | 4'567'130 | 5'146'296 | 4'810'417 | 5'496'528 | 4'661'806 | 5'066'898 | 4'778'472 |

| 1B2c Venting/Flaring | Unit | 2010 | 2011 | 2012 |
|----------------------|------|-----------|-----------|-----------|
| Crude Oil used | t | 4'490'741 | 4'402'315 | 3'408'796 |

For source category 1B2c, crude oil used is based on annual statistics of the Swiss Petroleum Association (EV 2012). This is the same activity data as for 1B2a Oil.

3.3.4.3 Uncertainties and Time-Series Consistency

A preliminary uncertainty assessment of all other sources in source category 1B2 based on expert judgement results in medium confidence in the emissions estimate (see Table 1-14).

The time series is consistent.

3.3.4.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the last submission 2013.

3.3.4.5 Source-Specific Recalculations

No recalculations occurred in 1B2c.

3.3.4.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

4 Industrial Processes

4.1 Overview

This chapter provides information on the estimation of the greenhouse gas emissions from sector 2 Industrial Processes. The following source categories are reported:

- 2A Mineral Products
- 2B Chemical Industry
- 2C Metal Production
- 2D Other Production
- *2E Production of Halocarbons and SF₆ is not occurring*
- 2F Consumption of Halocarbons and SF₆
- 2G Other

Emissions within this sector comprise greenhouse gas emissions as by-products from industrial processes and also emissions of F-gases during production, use and disposal. Emissions from fuel combustion in industry are reported in source category 1A2 under sector 1 Energy. Figure 4-1 shows the development of greenhouse gas emissions in source category 2 between 1990 and 2012.

Please note that for several industrial processes within source categories 2A Mineral Products and 2B Chemical Products data and information of emission factors and activity data is classified as confidential (C). For reviewers there is an additional version of chapter 4 Industrial Processes available, including all confidential data and information.




Figure 4-1 Switzerland's greenhouse gas emissions of sector 2 Industrial Processes 1990-2012. Source category 2D Other Production emits no direct greenhouse gas. The emissions of source category 2G Other are very small (2012: 0.9 Gg CO₂ eq) and are therefore not visible in this figure.

2A Mineral Products remain the dominant source of sector 2 with a share of 53.7% of the greenhouse gas emissions in 2012 although they have decreased by 29% since 1990. 2B

Chemical Industry accounts for 4.6% and has decreased by 8% since 1990. 2C Metal Production has decreased by 82.8% and accounts for 1.2% in 2012. 2F Consumption of Halocarbons and SF₆ is of increasing importance: The emissions have increased by a factor of 10.2 since 1990 and are currently responsible for 40.5% of total greenhouse gas emissions in sector 2. This is primarily due to the replacement of HFC for CFC in many technical applications.

In Table 4-1 the development of greenhouse gas emissions in sector 2 Industrial Processes are given by gases. Dominant gases are CO₂ and synthetic gases with shares of 57.1% and 41.4%, respectively, of the emissions in 2012 whereas N₂O and CH₄ contribute with 1.5% and less than one tenth of a per cent, respectively. The relative trend of these gases referring to the base year 1990 is shown in Figure 4-2 and 4-3.

Table 4-1 Greenhouse gas emissions of sector 2 Industrial Processes by gases in Gg CO₂ equivalent for the period 1990-2012.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 3'006 | 2'662 | 2'513 | 2'260 | 2'424 | 2'271 | 2'100 | 1'926 | 1'920 | 1'920 |
| CH ₄ | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.8 | 1.5 | 1.3 |
| N ₂ O | 68 | 62 | 54 | 52 | 61 | 60 | 58 | 51 | 54 | 55 |
| Synth. gases | 244 | 231 | 224 | 171 | 164 | 294 | 340 | 452 | 541 | 604 |
| Sum | 3'320 | 2'957 | 2'793 | 2'484 | 2'649 | 2'626 | 2'498 | 2'431 | 2'517 | 2'580 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 2'046 | 2'073 | 2'011 | 1'968 | 2'108 | 2'219 | 2'165 | 2'169 | 2'139 | 2'081 |
| CH ₄ | 1.5 | 1.6 | 1.5 | 1.5 | 2.1 | 2.2 | 1.9 | 2.2 | 2.5 | 1.6 |
| N ₂ O | 60 | 63 | 66 | 59 | 62 | 52 | 60 | 58 | 67 | 58 |
| Synth. gases | 728 | 799 | 844 | 941 | 1'063 | 1'152 | 1'170 | 1'191 | 1'326 | 1'306 |
| Sum | 2'836 | 2'938 | 2'922 | 2'970 | 3'235 | 3'425 | 3'397 | 3'421 | 3'535 | 3'446 |

| Gas | 2010 | 2011 | 2012 |
|-------------------------|-------|-------|-------|
| CO ₂ eq (Gg) | | | |
| CO ₂ | 2'241 | 2'186 | 2'070 |
| CH ₄ | 2.3 | 2.3 | 2.4 |
| N ₂ O | 60 | 54 | 54 |
| Synth. gases | 1'330 | 1'400 | 1'502 |
| Sum | 3'634 | 3'642 | 3'628 |




Figure 4-2 Relative trends of the greenhouse gases of sector 2 Industrial Processes in the period 1990-2012. The base year 1990 represents 100%.

Figure 4-2 shows that in the period 1990-2012 the emissions of CO₂ and N₂O from sector 2 Industrial Processes have decreased by 31.1% and 21.4% respectively compared to the base year 1990. Emissions of CH₄ have increased by 55.7% in the same time span.

Figure 4-3 shows that the emissions of F-gases have increased sixfold compared to the year 1990.




Figure 4-3 Relative trends of the F-gases of sector 2 Industrial Processes in the period 1990-2012. The base year 1990 represents 100%.

4.2 Source Category 2A – Mineral Products

4.2.1 Source Category Description

Tier 1 Key category 2A1

CO₂ emissions from Cement Production (level and trend).

Tier 2 Key category 2A1

CO₂ emissions from Cement Production (level).

Tier 2 Key category 2A3

CO₂ emissions from Limestone and Dolomite Use (trend).

Source category 2A Mineral Products comprises process emissions from production of cement and lime, limestone and dolomite use, asphalt roofing, road paving with asphalt and from production of plaster and glass.

Table 4-2 Specification of source category 2A Mineral Products. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 2A | Source | Specification | Data Source |
|-----|-----------------------------|--|--|
| 2A1 | Cement Production | Geogenic CO ₂ emissions from calcination process in cement production Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting operations | AD, EF: EMIS 2014/2A1 Zementwerke Rohmaterial AD, EF: EMIS 2014/2A1 Zementwerke übriger Betrieb |
| 2A2 | Lime Production | Geogenic CO ₂ emissions from calcination process in lime production Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting operations | AD, EF: EMIS 2014/2A2 Kalkproduktion, Rohmaterial AD, EF: EMIS 2014/2A2 Kalkproduktion, übriger Betrieb |
| 2A3 | Limestone and Dolomite Use | Geogenic CO ₂ emissions from fine ceramics, rock wool, and brick and tile production | EF: IPCC 2006, EMIS 2014/2A3 Ziegeleien AD: EMIS 2014/2A3 Feinkeramik Produktion, EMIS 2014/2A3 Steinwolle Produktion, EMIS 2014/2A3 Ziegeleien |
| 2A4 | Soda Ash Production and Use | Production is not occurring in Switzerland. Geogenic CO ₂ emissions from the use of soda ash in fine ceramics and glass production is reported in 2A3 Limestone and Dolomite Use and 2A7 Other respectively | |
| 2A5 | Asphalt Roofing | Emissions of CO and NMVOC from asphalt roofing | AD, EF: EMIS 2014/2A5 Dachpappenproduktion und Verlegung |
| 2A6 | Road Paving with Asphalt | Emissions of NMVOC from road paving | AD, EF: EMIS 2014/2A6 Strassenbelagsarbeiten |
| 2A7 | Other | Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from the production of plaster Geogenic CO ₂ emissions from production of container and tableware glass, and glass wool | AD, EF: EMIS 2014/2A7 Gips-Produktion übriger Betrieb EF: IPCC 2006 AD, EF: EMIS 2014/2A7 Hohlglas Produktion, EMIS 2014/2A7 Glas übrige Produktion, EMIS 2014/2A7 Glaswolle Produktion Rohprodukt |

4.2.2 Methodological Issues

4.2.2.1 Cement production (2A1)

Emissions of geogenic CO₂ occur during the production of clinker which is an intermediate component in the cement manufacturing process. During the production of clinker, limestone, which is mainly calcium carbonate (CaCO₃) is heated (calcined) to produce lime (CaO) and CO₂ as by-product. The CaO reacts subsequently with minerals in the raw materials and yields clinker. During this reaction step no further CO₂ is emitted. Clinker is then mixed with other components such as gypsum to make cement.

In Switzerland there are six plants producing clinker and cement. The Swiss plants are rather small and do not exceed a capacity of 3'000 tonnes of clinker per day. All of them use modern dry process technology.

Blasting operations in the limestone quarries are another source of emissions for both CO₂ and indirect greenhouse gases such as NO_x, CO, NMVOC and SO₂.

Methodology

Calcination process:

The geogenic CO₂ emissions from the calcination process in cement production are determined by a Tier 2 approach according to 2000 IPCC good practice guidance (IPCC 2000, chapter 3.1.1 *Cement production*). For cement production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = M_{\text{Clinker}} \cdot EF_{\text{Clinker}} \cdot CKD_{\text{Correction Faktor}}$$

In Switzerland no long wet or long dry kilns but only modern preheater or precalciner kilns are used and also no so-called low-alkali cement is produced. Therefore there is no landfilling of calcined cement dust (CKD) in Switzerland. In the cement plants all the filter dust is collected in high performance electrostatic precipitator or bag filters (having an efficiency of more than 99.999%) and being recycled to the kiln feed. In some cases small portions of the CKD are added directly to the cement as filler. Due to the kiln technology used in Switzerland the decarbonating degree of the CKD is almost equal to that of the kiln feed, meaning, that this CKD has not been decarbonated yet. Therefore the CKD correction factor is 1.00.

Blasting operations:

The emissions resulting from blasting operations during the digging of limestone are included following a country specific method. Emissions of GHGs related to blasting operations are calculated by multiplying the annual *cement* output by emission factors. Please note that the CO₂ emissions from "blasting" are related to the usage of the explosive itself and are not related to fuel consumption of e.g. bulldozers etc. The amount of used explosive is reported to be 0.13 kg/t cement¹² (EMIS 2014/2A1 Zementwerke übriger Betrieb).

Total emissions reported for the production of cement are the sum of emissions from calcination process and blasting operations. The share of CO₂ emissions from blasting operations in limestone quarries is well below one tenth of a per cent of the geogenic CO₂ emissions from the calcination process.

¹² The CO₂ emission factor for the use of blasting agents amounts to 600 kg CO₂/t of blasting agent. For the average amount on blasting agent used per kg cement measurement data for the year 2002 were taken. Measurement data were available for four Swiss cement plants, covering more than 60% of the Swiss cement production. Therefore this information is regarded as representative for the Swiss situation. The average blasting agent input per ton of cement amounts to 0.13 kg. The emission factor for CO₂ per ton of cement therefore amounts to 78 g/t cement.

Emission Factors

Calcination process:

The emission factor for CO₂ for calcination is a country specific value depending on the composition of the raw material. The emission factors for the entire time series are listed in Table 4-3. In 2012 it amounts to 530.56 kg CO₂ per ton of *clinker* produced. The IPCC approach neglects CO₂ emissions from decomposition of MgCO₃, which are taken into account in this country-specific value.

Table 4-3 CO₂ emission factor for calcination in 2A1 Cement Production 1990 to 2012 (EMIS 2014/2A1 Zementwerke Rohmaterial).

| 2A1 Cement production | Unit | 1990 - 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 - 2012 |
|------------------------------|--------------|-------------|-------|-------|--------|--------|--------|--------|-------------|
| Calcination, CO ₂ | kg/t clinker | 525 | 530.6 | 527.9 | 528.58 | 529.26 | 531.00 | 532.15 | 530.56 |

Blasting operations:

The emission factors are country specific based on measurements and data from industry and expert estimates as documented in EMIS 2014/2A1 Zementwerke übriger Betrieb. They are given per ton of *cement*. All emission factors have been recalculated for the entire time series due to a calculation error resulting in about 20% lower values. For CO₂ the emission factor has been adjusted from 96 g/t cement to 78 g/t.

Table 4-4 Emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting operations in g/t cement from source category 2A1 Cement Production in 2012 (EMIS 2014/2A1 Zementwerke übriger Betrieb).

| 2A1 Cement production | Unit | CO ₂ | NO _x | CO | NMVOC | SO ₂ |
|-----------------------|------------|-----------------|-----------------|----|-------|-----------------|
| Blasting operations | g/t cement | 78 | 3 | 18 | 7.8 | 0.13 |

Activity Data

Activity data on annual clinker and cement production is provided by industry and documented in EMIS 2014/2A1 Zementwerke Rohmaterial and EMIS 2014/2A1 Zementwerke übriger Betrieb.

Table 4-5 Activity data of clinker and cement production in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2A1 Zementwerke Rohmaterial and EMIS 2014/2A1 Zementwerke übriger Betrieb).

| 2A1 Cement production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-----------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Cement production | Gg | 5'117 | 4'683 | 4'268 | 4'043 | 4'432 | 3'994 | 3'648 | 3'485 | 3'371 | 3'540 |
| Clinker production | Gg | 4'808 | 4'189 | 3'927 | 3'564 | 3'930 | 3'706 | 3'337 | 2'994 | 2'995 | 2'992 |

| 2A1 Cement production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-----------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Cement production | Gg | 3'754 | 3'891 | 3'771 | 3'592 | 3'957 | 4'136 | 4'143 | 4'243 | 4'284 | 4'303 |
| Clinker production | Gg | 3'214 | 3'275 | 3'150 | 3'081 | 3'265 | 3'442 | 3'452 | 3'512 | 3'461 | 3'443 |

| 2A1 Cement production | Unit | 2010 | 2011 | 2012 |
|-----------------------|------|-------|-------|-------|
| Cement production | Gg | 4'553 | 4'577 | 4'359 |
| Clinker production | Gg | 3'642 | 3'587 | 3'368 |

4.2.2.2 Lime production (2A2)

During the production of lime calcium carbonate (CaCO₃) is heated (calcined) yielding burnt lime (CaO) and CO₂ as by-product. In Switzerland there is only one plant producing lime. There is no industry in Switzerland producing lime for its own requirements. A request to the

sugar producing plants Aarberg and Frauenfeld confirmed that indeed they produce lime from limestone in own shaft kilns but that the CO₂ is re-captured in the sugar production process. Thus no CO₂ emissions occur.

Blasting operations in quarry is another source of emissions for both CO₂ and indirect emissions such as NO_x, CO, NMVOC and SO₂.

Methodology

Calcination process:

The geogenic CO₂ emissions from the calcination process in lime production are determined by a country specific approach according to 2000 IPCC good practice guidance (IPCC 2000, chapter 3.1.2 *Lime production*). For lime production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = M_{\text{Lime}} \cdot EF_{\text{Lime}}$$

Blasting operations:

The emissions resulting from blasting operations during the digging of limestone are included following a country specific method. They are calculated by multiplying the annual lime production by emission factors. Please note that the CO₂ emissions from "blasting" are related to the usage of the explosive itself and are not related to fuel consumption of e.g. bulldozers etc

Total emissions reported for the production of lime are the sum of emissions from calcination process and blasting operations. The share of CO₂ emissions from blasting operations in the quarry is well below one tenth of a per cent of the geogenic CO₂ emissions from the calcinations process.

Emission Factors

Calcination process:

The emission factor for CO₂ from calcination of limestone depends both on the purity of the limestone and the grade of calcination (i.e. amount of rest CO₂ remaining in the final lime). The plant specific value has been calculated based on industry declaration and is assumed to be constant over time (EMIS 2014/2A2 Kalkproduktion, Rohmaterial). The value is considered confidential, however, available to reviewers.

Blasting operations:

The emission factors are country specific as documented in EMIS 2014/2A1 Kalkproduktion, übriger Betrieb. The value is considered confidential, however, available to reviewers.

Table 4-6 CO₂ emission factor for calcination process and blasting operations in lime production in kg/t lime and emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting operations in g/t lime in 2012 (EMIS 2014/2A2 Kalkproduktion, Rohmaterial and EMIS 2014/2A1 Kalkproduktion übriger Betrieb).

| 2A2 Lime production | Unit | CO ₂ | NO _x | CO | NMVOC | SO ₂ |
|---------------------|------|-----------------|-----------------|----|-------|-----------------|
| Calcination | kg/t | C | NA | NA | NA | NA |
| Blasting operations | g/t | C | C | C | C | C |

Activity Data

Activity data on annual lime production is based on data from the one existing plant in Switzerland, documented in the EMIS database (EMIS 2014/2A2 Kalkproduktion, Rohmaterial and EMIS 2014/2A1 Kalkproduktion übriger Betrieb). Detailed activity data is not reported as it is considered confidential, however, available to reviewers.

4.2.2.3 Limestone and dolomite use (2A3)

In Switzerland limestone and dolomite are used as raw material in the production of

- fine ceramics,
- rock wool and
- bricks and tiles.

When using limestone and dolomite in such production processes geogenic CO₂ is released to the atmosphere. The three different production processes are discussed consecutively in the following.

The use of limestone and dolomite as raw materials in glass production is reported in source category 2A7 Glass Production.

Fine ceramics (2A3)

In Switzerland the main production of fine ceramics is sanitary ware. The carbonate containing raw materials limestone and dolomite are used in product glazes only. The glazes contain small amounts of soda ash (Na₂CO₃) as well. All information on the fine ceramics production is documented in EMIS 2014/2A3 Feinkeramik Produktion.

Methodology

The geogenic CO₂ emissions from fine ceramics production are determined by a Tier 2 approach according to 2006 IPCC guidelines (IPCC 2006, chapter 2.5 *Other process uses of carbonates*). For fine ceramics production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = (M_{\text{Limestone}} \cdot EF_{\text{Limestone}}) + (M_{\text{Dolomite}} \cdot EF_{\text{Dolomite}}) + (M_{\text{Soda Ash}} \cdot EF_{\text{Soda Ash}})$$

Emission Factors

For fine ceramics production in Switzerland the CO₂ emission factors of limestone, dolomite and soda ash are taken from IPCC 2006 (chapter 2.5 *Other process uses of carbonates*, Table 2.1). As these emission factors are material properties they remain constant over time.

Table 4-7 Geogenic CO₂ emission factors for 2012 used for fine ceramics, rock wool production and the production of brick and tile in g/t carbonate containing raw material and g/t product, respectively (IPCC 2006, EMIS 2014/2A3 Ziegeleien).

| 2A3 Limestone and dolomite use | Unit | CO ₂ geogenic |
|--|---------------|--------------------------|
| fine ceramics and rock wool production | | |
| limestone use | g/t limestone | 439'710 |
| dolomite use | g/t dolomite | 477'320 |
| soda use | g/t soda | 414'920 |
| brick and tile production | g/t | 117'000 |

Activity Data

Activity data for carbonate containing raw materials, i.e. limestone, dolomite and soda ash used in the glazes of the fine ceramics production are extrapolated values based on industry data from the largest fine ceramics production plant in Switzerland. Detailed activity data on the carbonate containing raw materials is considered confidential; however, it is available to the reviewers.

Rock wool production (2A3)

In Switzerland there is one single producer of rock wool. The plant uses dolomite as raw material. No other carbonate containing raw material is used in the production process. All information of the rock wool production is documented in EMIS 2014/2A3 Steinwolle Produktion.

Methodology

The geogenic CO₂ emissions from rock wool production are determined by a Tier 2 approach according to IPCC 2006 (chapter 2.5 *Other process uses of carbonates*). For rock wool production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = M_{\text{Dolomite}} \cdot \text{EF}_{\text{Dolomite}}$$

Emission Factors

For rock wool production in Switzerland the CO₂ emission factor of dolomite is taken from IPCC 2006 (chapter 2.5 *Other process uses of carbonates*, Table 2.1). As the emission factor is a material property it remains constant over time (see Table 4-7).

Activity Data

Activity data is based on industry data from the single rock wool production plant in Switzerland.

Table 4-8 Activity data for the use of limestone and dolomite in fine ceramics and rock wool production and of the brick and tile production in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2A3 Feinkeramik Produktion, EMIS 2014/2A3 Steinwolle Produktion, EMIS 2014/2A3 Ziegeleien).

| 2A3 Limestone and dolomite use | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|
| fine ceramics production | | | | | | | | | | | |
| limestone use | Gg | C | C | C | C | C | C | C | C | C | C |
| dolomite use | Gg | C | C | C | C | C | C | C | C | C | C |
| soda use | Gg | C | C | C | C | C | C | C | C | C | C |
| rock wool production | | | | | | | | | | | |
| dolomite use | Gg | 2.8 | 2.9 | 2.8 | 2.6 | 2.7 | 2.9 | 2.8 | 2.6 | 3.0 | 3.3 |
| brick and tile production | Gg | 1'271 | 1'240 | 1'208 | 1'177 | 1'146 | 1'115 | 1'084 | 1'052 | 1'021 | 990 |

| 2A3 Limestone and dolomite use | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--------------------------------|------|------|------|------|------|-------|-------|-------|------|------|------|
| fine ceramics production | | | | | | | | | | | |
| limestone use | Gg | C | C | C | C | C | C | C | C | C | C |
| dolomite use | Gg | C | C | C | C | C | C | C | C | C | C |
| soda use | Gg | C | C | C | C | C | C | C | C | C | C |
| rock wool production | | | | | | | | | | | |
| dolomite use | Gg | 3.7 | 2.7 | 4.9 | 5.6 | 4.9 | 1.7 | 0.3 | 3.7 | 4.7 | 4.7 |
| brick and tile production | Gg | 959 | 945 | 875 | 859 | 1'023 | 1'086 | 1'065 | 975 | 865 | 701 |

| 2A3 Limestone and dolomite use | Unit | 2010 | 2011 | 2012 |
|--------------------------------|------|------|------|------|
| fine ceramics production | | | | |
| limestone use | Gg | C | C | C |
| dolomite use | Gg | C | C | C |
| soda use | Gg | C | C | C |
| rock wool production | | | | |
| dolomite use | Gg | 5.9 | 5.9 | 6.4 |
| brick and tile production | Gg | 879 | 800 | 818 |

Brick and tile production (2A3)

In Switzerland there are about 20 plants producing bricks and tiles. The manufacturing process uses limestone containing clay as main raw material.

Methodology

Concerning the release of geogenic CO₂ emissions from brick and tile production there has been no specific information on the employed raw materials available from Swiss industry in the past years. In 2013, again, a request to the Swiss association of brick and tile industry (Verband Schweizerische Ziegelindustrie VSZ) was made. Unfortunately without success.

Therefore, for submission 2014 data from a comparison of geogenic CO₂ emissions based on analyses of the carbonate content of the clay used for brick and tile production in a number of plants in Switzerland and the European Union are applied. This study was carried out by the VSZ in 2012 (see EMIS 2014/2A3 Ziegeleien).

In order to estimate the geogenic CO₂ emission from brick and tile production in Switzerland the following formula was used:

$$\text{CO}_2 \text{ Emissions} = M_{\text{brick and tile}} \cdot EF_{\text{brick and tile}}$$

Emission Factors

According to this study, bricks emit a weighted average of 13.2% of geogenic CO₂ (variation range 5.4 - 24%) and roof tiles have a weighted average of 8.6% (variation range 5.6 - 13%). Based on the production shares of the largest Swiss brick producer a production ratio for bricks:tiles of 2:1 was assumed for the whole period from 1990 on. This results in an average geogenic CO₂ emission content of 11.7%.

For estimating the geogenic CO₂ emissions from Swiss brick and tile production, a constant emission factor of 117 kg CO₂/t brick and tile was assumed. This represents the mean value provided by the industry association as discussed above and is about 50% higher than the value of 80 kg CO₂/t brick and tile used before.

Activity Data

Activity Data is based on production data from the Swiss association of brick and tile industry (see Table 4-8).

4.2.2.4 Soda ash production and use (2A4)

There is no soda ash production in Switzerland.

The main use of soda ash is in the glass production which is reported separately in source category 2A7 Glass production. A very small amount of soda ash is also applied in glazes of fine ceramics and is thus included in source category 2A3.

4.2.2.5 Asphalt roofing (2A5)

This source category comprises emissions from production and use of asphalt roofing materials (saturated felt, roofing and siding shingles, roll roofing and sidings). These products are used in roofing and other building applications. From 2A5 Asphalt roofing only indirect greenhouse gas emissions of CO and NMVOC arise. CO is emitted during the production process of asphalt roofing materials whereas NMVOC emissions are released during the entire production and laying processes (primers included).

Methodology

Emissions of CO and NMVOC from asphalt roofing are calculated by multiplying the annual amounts of asphalt roofing products and primers produced and employed by the corresponding emission factors.

Emission Factors

The emission factors for CO and NMVOC emissions from asphalt roofing processes are country specific. They are based on measurements, industry data and expert estimates as documented in EMIS 2014/2A5 Dachpappenproduktion und Verlegung.

Table 4-9 Emission factors for 2012 for CO and NMVOC in kg/t asphalt sealing sheeting and asphalt concrete from 2A5 Asphalt roofing and 2A6 Road paving with asphalt, respectively (EMIS 2014/2A5 Dachpappenproduktion und Verlegung and EMIS 2014/2A6 Strassenbelagsarbeiten).

| | Unit | CO | NMVOC |
|----------------------------|-------------------------------|-----|-------|
| 2A5 Asphalt roofing | kg/t asphalt sealing sheeting | 121 | 21 |
| 2A6 Road paving | kg/t asphalt concrete | NA | 0.49 |

Activity Data

Activity data on asphalt roofing products and primers produced is based on industry and expert estimates as documented in EMIS 2014/2A5 Dachpappen Produktion und Verlegung.

Table 4-10 Activity data for asphalt roofing and road paving with asphalt for the period 1990-2012 in Gg (EMIS 2014/2A5 Dachpappenproduktion und Verlegung and EMIS 2014/2A6 Strassenbelagsarbeiten).

| | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 2A5 Asphalt roofing | | | | | | | | | | | |
| asphalt sealing sheeting | Gg | 50 | 49 | 48 | 47 | 46 | 45 | 44 | 42 | 41 | 41 |
| 2A6 Road paving with asphalt | | | | | | | | | | | |
| asphalt concrete | Gg | 5'500 | 5'360 | 5'220 | 5'080 | 4'940 | 4'800 | 4'763 | 4'727 | 4'690 | 5'070 |

| | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 2A5 Asphalt roofing | | | | | | | | | | | |
| asphalt sealing sheeting | Gg | 41 | 41 | 38 | 35 | 32 | 30 | 28 | 26 | 25 | 25 |
| 2A6 Road paving with asphalt | | | | | | | | | | | |
| asphalt concrete | Gg | 5'170 | 4'860 | 4'770 | 4'860 | 4'840 | 4'780 | 5'400 | 5'100 | 5'160 | 5'200 |

| | Unit | 2010 | 2011 | 2012 |
|-------------------------------------|------|-------|-------|-------|
| 2A5 Asphalt roofing | | | | |
| asphalt sealing sheeting | Gg | 25 | 24 | 24 |
| 2A6 Road paving with asphalt | | | | |
| asphalt concrete | Gg | 5'250 | 5'300 | 4'770 |

4.2.2.6 Road paving with asphalt (2A6)

Asphalt road surfaces are composed of compacted aggregate and asphalt binder. From road surfacing operations NMVOC emissions occur only.

Methodology

The NMVOC emissions are determined by a country specific method as documented in EMIS 2014/2A5 Strassenbelagsarbeiten and calculated by multiplying the annual amount of asphalt products used for road paving by the corresponding emission factor.

Emission Factors

The emission factor for NMVOC emissions from road paving with asphalt is country specific. It consists of a EF for the NMVOC emissions from the bitumen content of asphalt products which is decreasing since 1990 and a variable EF from prime coatings. The values are based on industry data from 1990, 1998, 2007 and 2010. All other years are interpolated and complemented with expert estimates as documented in EMIS 2014/2A6 Strassenbelagsarbeiten (see Table 4-9).

Activity Data

Activity data on annual production of asphalt concrete is provided by the industry association on a yearly basis from 1998 on and for 1990 and 1995 (with expert estimates for the years in between) as documented in EMIS 2014/2A6 Strassenbelagsarbeiten (see Table 4-10).

4.2.2.7 Other (2A7)

Source category 2A7 Other comprises emissions from plaster production and from the production of container and table ware glass as well as glass wool.

Plaster Production (2A7)

Methodology

The emissions of CO₂, NO_x, CO, NMVOC and SO₂ from 2A7 Plaster production refer to emissions from blasting operations during the mining of gypsum, i.e. the raw material for plaster production. The emissions are calculated by multiplying the annual amount of processed rock by the emission factors. There are two plaster production sites in Switzerland.

Emission Factors

As there are no specific emission factors for gypsum mining, the emission factors for cement raw material mining are taken instead (with a rough estimate that 1.5 t of rocks are needed for 1 t of cement). This approach is documented in EMIS 2014/2A7 Gips-Produktion übriger Betrieb.

Table 4-11 Emission factors for plaster production in g/t mined rocks for 2012 (EMIS 2014/2A7 Gips-Produktion übriger Betrieb).

| 2A7 Plaster production | Unit | CO ₂ | NO _x | CO | NMVOC | SO ₂ |
|------------------------|-----------|-----------------|-----------------|----|-------|-----------------|
| | g/t rocks | 144 | 5.6 | 33 | 14.4 | 0.24 |

Activity Data

The activity data of the annual amount of rocks processed in the plaster production is based on industry data and expert estimates as documented in EMIS 2014/2A7 Gips-Produktion übriger Betrieb.

Table 4-12 Activity data for the mining of gypsum in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2A7 Gips-Produktion übriger Betrieb).

| 2A7 Plaster production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|------------------------|------|------|------|------|------|------|------|------|------|------|------|
| rocks | Gg | 319 | 316 | 313 | 310 | 307 | 304 | 300 | 297 | 294 | 291 |

| 2A7 Plaster production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|------------------------|------|------|------|------|------|------|------|------|------|------|------|
| rocks | Gg | 288 | 285 | 290 | 296 | 301 | 327 | 323 | 314 | 295 | 293 |

| 2A7 Plaster production | Unit | 2010 | 2011 | 2012 |
|------------------------|------|------|------|------|
| rocks | Gg | 335 | 293 | 271 |

Glass production (2A7)

The carbonate containing raw materials in the glass production are soda ash, limestone and dolomite. In Switzerland the following three glass types are produced: container glass, tableware glass and glass wool. Today there is only one production plant for container glass and one for tableware glass in Switzerland after the other one closed in 2002 and 2006, respectively. Glass wool is produced in two plants.

Methodology

For determination of geogenic CO₂ emission from glass production a Tier 2 approach according to IPCC 2006 (chapter 2.4 *Glass production*) is used. For glass production in Switzerland this results in the following formula:

$$\text{CO}_2 \text{ Emissions} = M_{\text{Glass type}} \cdot \text{EF}_{\text{Glass type}} \cdot (1 - \text{cullet ratio})$$

The cullet ratio describes the share of recycled glass material which is used in the production. The melting of cullet causes no geogenic CO₂ emissions.

Emission Factors

The emission factor for glass production in Switzerland is taken from IPCC 2006 (chapter 2.4 *Glass production*, Table 2.6). For the production of container glass, tableware glass and glass wool the values for glass type *container*, *tableware* and *fibreglass* are taken, respectively. As the emission factors are material properties they remain constant over the time.

Table 4-13 Geogenic CO₂ emission factor for glass production in g/t glass (IPCC 2006).

| 2A7 Glass production | Unit | CO ₂ geogenic |
|-------------------------------------|------|--------------------------|
| container glass | g/t | 210'000 |
| glass wool (fibre glass insulation) | g/t | 250'000 |
| glass (speciality tableware) | g/t | 100'000 |

Activity Data and Cullet Ratios

Activity data is based on industry data from Swiss glass producers. Detailed information on activity data for container glass production and tableware production is considered confidential as there is only one producing plant respectively. However, the detailed data is available to the reviewers. Activity data for glass wool production is based on industry data from the two glass wool production plants in Switzerland.

Table 4-14 Glass production in Switzerland for the period 1990-2012 in Gg and cullet ratio in % (EMIS 2014/2A7 Hohlglas Produktion, EMIS 2014/2A7 Glas übrige Produktion and EMIS 2014/2A7 Glaswolle Produktion Rohprodukt).

| 2A7 Glass production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| container glass | | | | | | | | | | | |
| production | Gg | C | C | C | C | C | C | C | C | C | C |
| cullet ratio | % | C | C | C | C | C | C | C | C | C | C |
| glass (speciality tableware) | | | | | | | | | | | |
| production | Gg | C | C | C | C | C | C | C | C | C | C |
| cullet ratio | % | C | C | C | C | C | C | C | C | C | C |
| glass wool | | | | | | | | | | | |
| production | Gg | 24.3 | 22.8 | 23.4 | 21.7 | 24.9 | 24.2 | 19.9 | 25.6 | 27.5 | 32.1 |
| cullet ratio | % | 21 | 26 | 49 | 53 | 65 | 45 | 61 | 66 | 65 | 67 |

| 2A7 Glass production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| container glass | | | | | | | | | | | |
| production | Gg | C | C | C | C | C | C | C | C | C | C |
| cullet ratio | % | C | C | C | C | C | C | C | C | C | C |
| glass (speciality tableware) | | | | | | | | | | | |
| production | Gg | C | C | C | C | C | C | C | C | C | C |
| cullet ratio | % | C | C | C | C | C | C | C | C | C | C |
| glass wool | | | | | | | | | | | |
| production | Gg | 31.1 | 25.2 | 19.9 | 25.9 | 32.7 | 37.5 | 38.1 | 44.5 | 44.4 | 33.5 |
| cullet ratio | % | 69 | 65 | 59 | 62 | 65 | 65 | 73 | 71 | 69 | 69 |

| 2A7 Glass production | Unit | 2010 | 2011 | 2012 |
|------------------------------|-------------|-------------|-------------|-------------|
| container glass | | | | |
| production | Gg | C | C | C |
| cullet ratio | % | C | C | C |
| glass (speciality tableware) | | | | |
| production | Gg | C | C | C |
| cullet ratio | % | C | C | C |
| glass wool | | | | |
| production | Gg | 35.7 | 41.4 | 38.7 |
| cullet ratio | % | 71 | 72 | 61 |

4.2.3 Uncertainties and Time-Series Consistency

The uncertainty for CO₂ emissions in cement production (2A1) which is key category regarding level and trend amounts to 2.8%. The uncertainty of CO₂ emissions was calculated following the steps in Table 3.2 in 2000 IPCC good practice guidance (IPCC 2000, p. 3.15). As CO₂ emissions are calculated based on plant level CaO contents of the clinker (Tier 2) an uncertainty of 2% is assumed both for activity data and emission factor (step 3 in table 3.2).

For non-key categories, the NIR provides qualitative estimates of uncertainties only. The terms high, medium and low data quality is used and a quantitative relative uncertainty is assigned to every uncertainty category (see Table 1-14 Semi-quantitative uncertainties for non-key categories). The uncertainties for CO₂ emissions from source categories 2A2 and 2A7 are estimated to be low and thus amount to 2%. For CO₂ emission in source category 2A3 an overall uncertainty of 51% is taken. The uncertainty of 51% corresponds to the uncertainty of brick and tile and is the highest among the three processes included in source category 2A3.

The time series is consistent.

4.2.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

In submission 2012 the emission factor of category 2A1 used in the Swiss Inventory was compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value (INFRAS 2012). Switzerland's factor lies in the midfield of the other countries; see chpt. 4.2.2.1.

In submission 2012 the emission factor of category 2A2 used in the Swiss Inventory was compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value (INFRAS 2012). Switzerland's factor lies in the midfield of the other countries, see chpt. 4.2.2.2.

In submission 2012 the emission factor of category 2A3 used in the Swiss Inventory was compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (INFRAS 2012). Switzerland's factor lies beyond the other countries. This is due to the fact, that the Swiss factor includes emissions from three different sources: fine ceramics, rock wool and brick and tile. The most dominant process in category 2A3 is brick and tile production. The emission factor amounts to 0.08 t CO₂/t brick and tile. Comparing this value to the German value which is 0.029 t CO₂/t bricks and tiles, shows that the Swiss value is rather high. The Swiss brick and tile industry has determined very recently the carbonate content of the clay raw material at several pits. A first comparison with the carbonate content of clay from pits in other European countries confirms that the Swiss values are indeed rather high.

4.2.5 Source-Specific Recalculations

2A1 Cement production: For blasting operations a calculation error has been corrected resulting in about 20% lower EF values for CO₂, NO_x, CO, NMVOC and SO₂ for 1990-2011.

2A3 Brick and tile production: So far interpolated activity data of the three brick and tile production plants which are not member of the industry association have been replaced by effective production data for 2001-2006 resulting thus in revised AD for these years. The EF for CO₂ has been revised based on some specifications of the carbonate content of the raw material and an assumed brick and tile production share yielding an about 50% higher EF value for the entire time series.

2A5 Asphalt roofing: Activity data for 2008-2011 has been revised due to updated projection values for 2015.

2A7 Glasswool production: Activity data of one of the two production plants have been revised for 1991-2004 based on effective production data for 1996-2004 resulting in revised EF values of CO₂ for 1991-2004 as well due to different cullet ratios of the two plants.

4.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

4.3 Source Category 2B – Chemical Industry

4.3.1 Source Category Description

Source category 2B Chemical Industry comprises process emissions from the production of ammonia, nitric acid, silicon carbide, ethylene, acetic acid and sulphuric acid.

Table 4-15 Specification of source category 2B Chemical Industry. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 2B | Source | Specification | Data Source |
|-----|-------------------------|--|--|
| 2B1 | Ammonia Production | Emissions of CO ₂ and NMVOC are reported in 2B5 Ethylene production | AD, EF: EMIS 2014/2B1 Ammoniak-Produktion |
| 2B2 | Nitric Acid Production | Emissions of N ₂ O and NO _x from the production of nitric acid | AD, EF: EMIS 2014/2B2 Salpetersäure Produktion |
| 2B3 | Adipic Acid Production | Not occurring in Switzerland | |
| 2B4 | Carbide Production | Emissions of CO ₂ , CH ₄ and SO ₂ from the production of silicon carbide | EF: IPCC 2006, EMIS 2014/2B4 Graphit und Siliziumkarbid Produktion AD: EMIS 2014/2B4 Graphit und Siliziumkarbid Produktion |
| 2B5 | Other Chemical Industry | Emissions of CO ₂ and NMVOC from ethylene production Emissions of CO ₂ , CH ₄ , CO and NMVOC from acetic acid production SO ₂ emissions from sulphuric acid production | AD, EF: EMIS 2014/2B5 ethylene production AD, EF: EMIS 2014/2B5 Essigsäure-Produktion AD, EF: EMIS 2014/2B5 Schwefelsäure-Produktion |

4.3.2 Methodological Issues

4.3.2.1 Ammonia production (2B1)

Ammonia (NH₃) is produced in one single plant in Switzerland by catalytic reaction of nitrogen and synthetic hydrogen (see Figure 4-4). Ammonia is not produced in an isolated reaction plant but is part of an integrated production chain (see Figure 4-5).

The starting production process is the thermal cracking of liquefied petroleum gas (LPG) and light virgin naphtha yielding ethylene (ethene, C₂H₄), and a series of by-products such as e.g. synthetic hydrogen and methane, which are used as educts for further production steps.

According to the Swiss ammonia producer it is not possible to split and allocate the emissions of the cracking process (CO₂ and NMVOC) to every single product such as, e.g., ethylene, acetylene (ethyne, C₂H₂), cyanic acid or ammonia. **Therefore, all CO₂ and NMVOC emissions of the cracking process are allocated to the ethylene production** and are reported under the category 2B5 Ethylene production. Thus, for source category 2B1 Ammonia production, CO₂ and NMVOC emissions are reported as included elsewhere (IE). All information on the ammonia production and the cracking process is documented in EMIS 2014/2B1 Ammoniak-Produktion and EMIS 2014/2B5 Ethen-Produktion, respectively.




Figure 4-4 Process flow chart for the production of ammonia (NH₃) from nitrogen (N₂) and hydrogen (H₂, synth.). Hydrogen is derived from the thermal cracking process in the same plant (see Figure 4-5).




Figure 4-5 Process flow chart for the production of ethylene (C₂H₄) and acetylene (C₂H₂) by thermal cracking of liquefied petroleum gas (LPG) and light virgin naphta. The intermediate product H₂, synth. is used as educt in the ammonia production in the same plant (see Figure 4-4).

4.3.2.2 Nitric acid production (2B2)

In Switzerland there is one single plant producing nitric acid (HNO₃). Nitric acid is produced by catalytic oxidation of ammonia (NH₃) with air. At temperatures of 800 °C nitric monoxide (NO) is formed. During cooling, nitrogen monoxide reacts with excess oxygen to form nitrogen dioxide (NO₂). The nitrogen dioxide reacts with water to form 60% nitric acid (HNO₃). Today, two types of processes are used for nitric acid production: single pressure or dual pressure plants. In Switzerland a dual pressure plant is installed.

During the process nitrous oxide (N₂O) can be formed as an unintended by-product. In addition, also some nitrogen oxide (NO_x) is produced. In the Swiss production plant abatement of NO_x is done by selective catalytic reduction (SCR) which reduces NO_x to N₂ and O₂ (the SCR in this plant is also used for treatment of other flue gases and was not installed for the HNO₃ production specially). No additional abatement technique is installed to destroy N₂O. A decomposition of N₂O occurs, to some extent, simultaneously in the NO_x

reduction process. The production and abatement technology has essentially remained the same since 1990.

Methodology

The N₂O and NO_x emissions from nitric acid production are determined by a Tier 2 approach. The emissions are calculated by multiplying the annual nitric acid production output by the corresponding emission factors for N₂O and NO_x emissions respectively.

Emission Factors

The N₂O and NO_x emission factors for nitric acid production in Switzerland are based on measurements from the single nitric acid production plant. The measurement of N₂O was carried out according to the guideline VDI-Richtlinie 2469/Blatt 1 (Messen gasförmiger Emissionen - Messen von Distickstoffmonoxid - Manuelles gaschromatographisches Verfahren). The test gas is sucked in via a heated titanium sensor and then treated with a solution of potassium permanganate and hydrogen peroxide in order to remove nitrogen oxides and further disturbing components. The N₂O concentration is then measured using a gas chromatograph with an electron capture detector. The measurement uncertainty is $\pm 20\%$ (minimum $\pm 0.5 \text{ mg/m}^3$). On repeated enquires the plant confirmed that since a denitrification system and an automatic control system for the ammonia addition was installed in 1988 and 1990, respectively, no modification has been made in the production line. The values are documented in EMIS 2014/2B2 Salpetersäure Produktion. They are considered confidential, however, available to reviewers on request.

Table 4-16 Emission factors for N₂O and NO_x for nitric acid production in Switzerland in kg/t nitric acid for 2012. Data refers to 100% nitric acid (EMIS 2014/2B2 Salpetersäure Produktion).

| 2B2 Nitric acid production | Unit | N ₂ O | NO _x |
|----------------------------|------|------------------|-----------------|
| | kg/t | C | C |

Activity Data

Activity data on annual production of nitric acid (100%) is provided on a yearly basis by the Swiss production plant for the entire time period 1990-2012. The data is confidential but available for reviewers (see EMIS 2014/2B2 Salpetersäure Produktion).

4.3.2.3 Carbide production (2B4)

In Switzerland there is one single plant producing carbide. The plant produces silicon carbide which is used in abrasives, refractories, metallurgy and anti-skid flooring. The Swiss silicon carbide is produced in an electric furnace at temperatures above 2000 °C using the Acheson process. The starting materials are quartz sand (SiO₂), petroleum coke and anthracite (C) which yield silicon carbide (SiC) and carbon monoxide (CO). The CO is converted to CO₂ in excess oxygen and released to the atmosphere. Petroleum coke and anthracite – although to a lower portion – may contain volatile organic compounds which can form methane (CH₄) as an unintended by-product. There is no abatement techniques installed which could capture the CO₂ or CH₄ emissions.

Methodology

The CO₂, CH₄ and SO₂ emissions from silicon carbide production are determined by a Tier 2 approach. The emissions are calculated by multiplying the annual silicon carbide production output by the corresponding emission factors for CO₂, CH₄ and SO₂ emissions respectively.

Emission Factors

The CO₂, CH₄ and SO₂ emission factors are considered confidential, however, available to reviewers on request. The values base partly on measurements and data from the single silicon carbide production plant and are documented in EMIS 2014/2B2 Graphit und Siliziumkarbid Produktion.

Table 4-17 Emission factors for CO₂, CH₄ and SO₂ for carbide production in Switzerland for 2012 in kg/t silicon carbide respectively (EMIS 2014/2B4 Graphit und Siliziumkarbid Produktion).

| 2B4 Silicon carbide production | Unit | CO ₂ | CH ₄ | SO ₂ |
|--------------------------------|------|-----------------|-----------------|-----------------|
| | kg/t | C | C | C |

Activity Data

Activity data on annual production of silicon carbide is provided on a yearly basis from 1997 onwards by the Swiss production plant. For the time period 1990-1996 activity data bases on industry data for 1990 and 1995 and interpolated values in between. The data is confidential but available for reviewers (see EMIS 2014/2B4 Graphit und Siliziumkarbid Produktion).

4.3.2.4 Other (2B5)

Source category Other (2B5) comprises emissions from production of ethylene, acetic acid and sulphuric acid.

4.3.2.5 Ethylene production (2B5)

Ethylene (ethene, C₂H₄) is produced by a single plant in Switzerland by thermal cracking of liquefied petroleum gas (LPG) and virgin naphta. Ethylene is not produced in an isolated process but is co-processed together with several other products such as H₂, CH₄, and C₂H₂ (see flow chart in Figure 4-5 in section 4.3.2.1). From the thermal cracking process emissions of CO₂ and NMVOC are released. They are both allocated entirely to the production of ethylene which is the first product within the integrated production chain. CH₄ emissions to atmosphere do not occur since CH₄ is completely used as an educt in the downstream production of cyanic acid (HCN) in the same facility (again, see Figure 4-5 and for further information see EMIS 2014/2B5 Ethen-Produktion). Therefore CH₄ emissions are reported as NA for ethylene production and only CO₂ and NMVOC emissions are reported.

Methodology

The CO₂ and NMVOC emissions from ethylene production are determined by a country-specific approach. The emissions are calculated by multiplying the annual ethylene production output by the corresponding emission factors for CO₂ and NMVOC emissions respectively.

Emission Factors

The CO₂ and NMVOC emission factors for ethylene production are based on industry data from the single ethylene production plant in Switzerland. Annual emission data was only available from the year 2000 onwards. For the period 1990-1999 a constant value, i.e. the mean value of the years 2000-2009 was assumed. The emission factors for ethylene production are considered confidential; however, they are available to reviewers on request.

Table 4-18 Emission factors for CO₂ and NMVOC in ethylene production, CH₄, CO and NMVOC in acetic acid production and SO₂ in sulphuric acid production for 2012 in kg/t product (EMIS 2014/2B5 Ethen-Produktion, EMIS 2014/2B5 Essigsäure-Produktion and EMIS 2014/2B5 Schwefelsäure-Produktion).

| 2B5 Chemical industry, other | Unit | CO ₂ | CH ₄ | CO | NMVOC | SO ₂ |
|------------------------------|------|-----------------|-----------------|-------|-------|-----------------|
| ethylene production | kg/t | C | NA | NA | C | NA |
| acetic acid production | kg/t | 12 | 0.023 | 0.014 | 0.25 | NA |
| sulphuric acid production | kg/t | NA | NA | NA | NA | C |

Activity Data

Activity data on the annual production of ethylene is provided on a yearly basis by the single ethylene production plant in Switzerland for the entire time period 1990-2012. The data is considered confidential but available for reviewers on request.

Table 4-19 Activity data for the production of ethylene, acetic acid and sulphuric acid in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2B5 Ethen-Produktion, EMIS 2014/2B5 Essigsäure-Produktion and EMIS 2014/2B5 Schwefelsäure-Produktion).

| 2B5 Chemical industry, other | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|------------------------------|------|------|------|------|------|------|------|------|------|------|------|
| ethylene production | Gg | C | C | C | C | C | C | C | C | C | C |
| acetic acid production | Gg | 30 | 29 | 29 | 28 | 28 | 27 | 27 | 26 | 26 | 25 |
| sulphuric acid production | Gg | C | C | C | C | C | C | C | C | C | C |

| 2B5 Chemical industry, other | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|------------------------------|------|------|------|------|------|------|------|------|------|------|------|
| ethylene production | Gg | C | C | C | C | C | C | C | C | C | C |
| acetic acid production | Gg | 24 | 14 | 11 | 10 | 9 | 8 | 8 | 9 | 18 | 28 |
| sulphuric acid production | Gg | C | C | C | C | C | C | C | C | C | C |

| 2B5 Chemical industry, other | Unit | 2010 | 2011 | 2012 |
|------------------------------|------|------|------|------|
| ethylene production | Gg | C | C | C |
| acetic acid production | Gg | 20 | 18 | 12 |
| sulphuric acid production | Gg | C | C | C |

4.3.2.6 Acetic and sulphuric acid production (2B5)

In Switzerland there are two plants producing acetic acid (CH₃COOH). From acetic acid production emissions of CO₂, CH₄, CO and NMVOC occur. For this year's submission all three so far known plants producing acetic acid were contacted directly for information on production data, process emissions and manufacturing processes. Thereby it turned out that one plant does not produce acetic acid but acetic acid occurs as a by-product only of esterifications using acetic anhydride.

Sulphuric acid (H₂SO₄) is produced by one plant only in Switzerland. From this production process SO₂ is emitted.

Methodology

In order to determine emissions of CO₂, CH₄, CO and NMVOC as well as of SO₂ from acetic acid and sulphuric acid production, respectively, a country specific approach is used. The emissions are calculated by multiplying the annual production of acetic acid and sulphuric acid, respectively, by the corresponding emission factor.

Emission Factors

The emission factors for CO₂, CH₄, CO and NMVOC from acetic acid production and for SO₂ from sulphuric acid production in Switzerland are country specific and base on measurements and data from industry and expert estimates documented in EMIS 2014/2B5 Essigsäure-Produktion and EMIS 2014/2B5 Schwefelsäure-Produktion (see Table 4-18).

For this year's submission emission factors of the acetic acid production have been revised for the entire time series in co-operation with industry. From one production plant NMVOC emissions are emitted only, whereas the other one reports emissions of CO₂, CH₄, CO and NMVOC. Usually the process emissions are treated in a flue gas incineration. Thus, the reported emissions of CH₄, CO and NMVOC only occur in case of malfunction resulting in strongly fluctuating plant-specific emission factors. In addition the resulting implied emission factors based on the emissions of both plants are modulated by considerable production fluctuations of one of the plants from 2000 onwards.

The data for sulphuric acid production is confidential but available for reviewers on request.

Activity Data

The annual amount of produced acetic acid and sulphuric acid base on data from industry and expert estimates documented in EMIS 2014/2B5 Essigsäure-Produktion and EMIS 2014/2B5 Schwefelsäure-Produktion (see Table 4-19). The data for sulphuric acid production is confidential but available for reviewers.

4.3.3 Uncertainties and Time-Series Consistency

For non-key categories, the NIR provides qualitative estimates of uncertainties only. The terms high, medium and low data quality are used and a quantitative relative uncertainty is assigned to every uncertainty category (see Table 1-14 Semi-quantitative uncertainties for non-key categories). The uncertainties for CO₂ and CH₄ in source category 2B are both estimated to be medium, resulting in a relative uncertainty of 10% for CO₂ and of 30% for CH₄. For N₂O emissions from 2B2 Nitric Acid Production which has been a key category in previous submissions the uncertainty was calculated to be 41%.

The time series is consistent.

4.3.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

In submission 2012 the N₂O emission factor of source category 2B2 Nitric Acid Production used in the Swiss Inventory was compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value (INFRAS 2012). Switzerland's factor lies in the midfield of the other countries; see chpt. 4.3.2.2.

4.3.5 Source-Specific Recalculations

2B2 Nitric acid production: The EF values for NO_x have been revised for the entire time series based on measurement data of NO_x emissions for the years 2007, 2009 and 2012 from the production plant.

2B5 Acetic acid production: Based on information from the production plants the activity data has been revised from 1991-2011 (effective data from 2000 onwards and interpolated values 1991-1999). In addition also the EF values for CH_4 , CO and NMVOC are revised and the one of CO_2 is newly introduced for the entire time series (except for NMVOC only from 2000-2011) based on measurement data from 2000 onwards.

2B5 Sulphuric acid production: The activity data has been updated for 2009-2011 based on corrected and new production data from the plant for 2009-2010 and 2011, respectively. The EF values for SO_2 have been revised for the entire time series based on new information from the producer for the years 2009-2012.

4.3.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

4.4 Source Category 2C – Metal Production

4.4.1 Source Category Description

Source category 2C Metal Production comprises process emissions from the production of iron and steel and aluminium, from the use of SF₆ in aluminium and magnesium foundries, as well as from battery recycling and non-ferrous metal foundries.

Table 4-20 Specification of source category 2C Metal Production. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 2C | Source | Specification | Data Source |
|-----|---|--|--|
| 2C1 | Iron and Steel Production | Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from the production of iron and steel | AD, EF: EMIS 2014/2C1 Eisengiessereien Elektroschmelzöfen/übriger Betrieb, EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen/übriger Betrieb |
| 2C2 | Ferroalloys Production | Production is not occurring in Switzerland | |
| 2C3 | Aluminium Production | Emissions of PFC, CO ₂ , NO _x , CO, NMVOC and SO ₂ from the production of aluminium (ceased in 2006) | AD: EMIS 2014/2C3 Aluminium Produktion EF for PFC: Industry Data EF other gases: EMIS 2014/2C3 Aluminium Produktion |
| 2C4 | Use of SF ₆ in Aluminium and Magnesium Foundries | Emissions from use of SF ₆ in aluminium and magnesium foundries | AD: Industry Data EF: IPCC 2006 |
| 2C5 | Other | Emissions of CO ₂ , NO _x , CO and SO ₂ from battery recycling Emissions of CO and NMVOC from non-ferrous metal foundries | AD, EF: EMIS 2014/2C5e Batterie-Recycling, AD, EF: EMIS 2014/2C5e Buntmetallgiessereien Elektroöfen |

4.4.2 Methodological Issues

4.4.2.1 Iron and Steel production (2C1)

There is no primary iron and steel production in Switzerland. Only secondary steel production occurs, which is steel production from recycled steel scrap. After closing down of two steel plants in 1994 there remain two plants in Switzerland. Both plants use electric arc furnaces (EAF) with a carbon electrode for melting the steel scrap. During the melting process CO₂ emissions occur mainly from scrap, electrodes and carburization coal. Indirect emissions such as NO_x, CO, NMVOC and SO₂ occur as well.

In Switzerland no production of pig iron occurs but iron is processed in foundries only. 14 iron foundries exist in Switzerland today. About 75% of the iron is processed in induction furnaces and 25% in cupola furnaces. From induction furnaces only indirect emissions occur. From cupola furnaces also CO₂ emissions occur. Those CO₂ emissions are accounted for in source category 1A2a.

Methodology

For determination of CO₂ emission from iron and steel production a mixture of a Tier 2 and a Tier 3 approach according to IPCC 2006 (chapter 4.2 *Iron & steel and metallurgical coke production*) is used since this year's submission. For the years 2005-2011 plant specific information is available (Tier 3). From this information data for the other years are interpolated for calculating an implied emission factor.

For steel production in Switzerland this results in the following formula:

$$E_{\text{CO}_2, \text{ non-energy}} = \text{EAF} \cdot \text{EF}_{\text{EAF}}$$

whereas EAF is the quantity of EAF crude steel produced in tonnes and EF_{EAF} the emission factor in tonnes CO_2 /tonne steel produced. The same formula is also applied to calculate emissions of indirect greenhouse gases from iron and steel production. No CH_4 emissions occur in the Swiss EAF process.

Emission Factors

The emission factors for iron and steel production in Switzerland are country specific and base on measurement data from industry and expert estimates documented in EMIS 2014/2C1 Eisengiessereien Elektroschmelzofen/übriger Betrieb, EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen and EMIS 2014/2C1 Stahlwerke Walzwerke.

For this year's submission the CO_2 emission factor was completely revised as new data from the two Swiss steel plants were available. For the calculations all carbon sources (graphite electrodes, steel scrap, alloy coal, etc.) and carbon sinks (steel, filter dust and slag) for the years 2005-2011 were taken into account. From these mass flows emission factors were calculated for both steel plants and weighted according to the production amount to yield overall CO_2 emission factors for Swiss steel industry for the entire time series, see Table 4-27. On average the revised values are about a factor of 16 (range: 12-21) lower than the ones used so far. The latter were based on measurements at the flue gas chimneys including emissions from gas burners which already are reported in source category 1A2a as well.

Table 4-21 CO_2 emission factor of electric arc furnaces in 2C1 Steel Production for the period 1990-2012 in kg/t (EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen).

| 2C1 Steel production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| CO_2 | kg/t | 8.3 | 8.2 | 8.1 | 8.1 | 8.1 | 8.0 | 8.0 | 7.8 | 7.5 | 7.5 |

| 2C1 Steel production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| CO_2 | kg/t | 7.7 | 7.7 | 7.8 | 7.9 | 7.8 | 8.8 | 9.1 | 8.5 | 6.8 | 6.8 |

| 2C1 Steel production | Unit | 2010 | 2011 | 2012 |
|----------------------|------|------|------|------|
| CO_2 | kg/t | 7.6 | 7.1 | 7.9 |

Table 4-22 Emission factors for NO_x , CO and NMVOC in iron production, for CO_2 , NO_x , CO, NMVOC and SO_2 in steel production, for CO_2 , NO_x , CO and SO_2 in battery recycling and for CO and NMVOC in non-ferrous metal production for 2012 (EMIS 2014/2C1 Eisengiessereien Elektroschmelzofen/übriger Betrieb, EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen and EMIS 2014/2C1 Stahlwerke Walzwerke, EMIS 2014/2C5e Batterie-Recycling and 2014/2C5e Buntmetallgiessereien Elektroöfen).

| 2C Metal production | Unit | CO_2 | NO_x | CO | NMVOC | SO_2 |
|------------------------|------|---------------|---------------|------|-------|---------------|
| 2C1 Iron | kg/t | NA | 0.01 | 4.1 | 4.0 | NA |
| 2C1 Steel | kg/t | 7.9 | 0.19 | 0.8 | 0.1 | 0.017 |
| 2C5 Battery recycling | kg/t | 560 | 0.88 | 1.2 | NA | 0.01 |
| 2C5 Non-ferrous metals | kg/t | NA | NA | 0.24 | 0.05 | NA |

Activity Data

Activity data on annual production of iron and steel is provided on a yearly basis by the Swiss production plants and the foundry association. Data is given in the following table:

Table 4-23 Production of iron, steel, aluminium and non-ferrous metals as well as amount of batteries recycled in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2C1 Eisengiessereien Elektroschmelzöfen/ übriger Betrieb, EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen, EMIS 2014/2C3 Aluminium Produktion, EMIS 2014/2C5e Batterie-Recycling and 2014/2C5e Buntmetallgiessereien Elektroöfen).

| 2C Metal production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|------------------------|------|-------|-------|-------|-------|-------|------|------|------|------|------|
| 2C1 Iron | Gg | 170 | 140 | 136 | 110 | 115 | 130 | 111 | 114 | 123 | 122 |
| 2C1 Steel | Gg | 1'108 | 1'155 | 1'245 | 1'276 | 1'230 | 716 | 738 | 789 | 880 | 918 |
| 2C3 Aluminium | Gg | 87 | 82 | 75 | 36 | 24 | 21 | 27 | 27 | 32 | 34 |
| 2C5 Battery recycling | Gg | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| 2C5 Non-ferrous metals | Gg | 55 | 56 | 57 | 58 | 59 | 60 | 65 | 66 | 68 | 69 |

| 2C Metal production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|
| 2C1 Iron | Gg | 120 | 105 | 80 | 73 | 67 | 67 | 67 | 72 | 78 | 49 |
| 2C1 Steel | Gg | 1'022 | 1'048 | 1'125 | 1'143 | 1'226 | 1'159 | 1'254 | 1'267 | 1'315 | 935 |
| 2C3 Aluminium | Gg | 36 | 36 | 40 | 44 | 45 | 45 | 12 | 0 | 0 | 0 |
| 2C5 Battery recycling | Gg | 3.0 | 3.0 | 3.0 | 2.9 | 3.3 | 2.8 | 2.4 | 2.4 | 2.5 | 3.4 |
| 2C5 Non-ferrous metals | Gg | 70 | 60 | 49 | 43 | 38 | 33 | 30 | 28 | 21 | 15 |

| 2C Metal production | Unit | 2010 | 2011 | 2012 |
|------------------------|------|-------|-------|-------|
| 2C1 Iron | Gg | 53 | 61 | 46 |
| 2C1 Steel | Gg | 1'218 | 1'322 | 1'252 |
| 2C3 Aluminium | Gg | 0 | 0 | 0 |
| 2C5 Battery recycling | Gg | 3.3 | 2.4 | 2.4 |
| 2C5 Non-ferrous metals | Gg | 20 | 12 | 18 |

4.4.2.2 Aluminium Production (2C3)

Methodology

The last production site for aluminium in Switzerland closed down in April 2006. Both CO₂ and PFC emissions were based on a country specific approach. More specific for PFC emissions a Tier 3b approach according to 2000 IPCC good practice guidance (IPCC 2000) was used. Operating smelter emissions have been monitored periodically by the industry for selected years. The emissions were calculated by multiplying annual production by emission factors.

Emission Factors

The emission factor for CO₂ per ton of metal product is country specific. It is based on measurements and data from industry and expert estimates, documented in EMIS 2014/2C3 Aluminium Produktion. For CO₂ emissions from aluminium production, an emission factor of 1.6 ton CO₂ per ton of aluminium is used (EMIS 2014/2C3 Aluminium Produktion). This CO₂ stems from the oxidation of the anode in the electrolysis process. The value is based on an estimate of the amount of anode material used. In Switzerland only pre-backed anodes are used. The emissions for CO₂ are calculated with 0.43 tons of anode per ton of aluminium; it is assumed that the anode consists completely of carbon and that it is fully oxidized during the process (value from Swiss foundries, value for 1990, assumed to be constant over the time series).

For PFC emissions from aluminium production, operating smelter EF have been monitored periodically by the industry for selected years. The only Swiss factory provided own measurements for 1990, 1999 and 2000 yielding smaller EFs than the European average (by factors of 3.9, 4.7 and 5.1, respectively) (Alcan 2003). The comparison with these data and data from IAI (2005) on global PFC emissions from aluminium production showed that the emissions from the smelter in Switzerland are lower by a factor of about 4. This seems to be plausible because they used point feed prebake (PFPB) technology and it is known that this technology has the lowest emissions per tonne of aluminium. Therefore a "general reduction factor" of 4.0 for both PFC gases (CF₄ and C₂F₆) is adopted based on the average European

values as reported from the European Aluminium Association (Alcan 2002) for the years with no measured emission data available. The resulting emission factors for Switzerland are still within the uncertainty range according to IPCC GPG 2000. In order to calculate the emissions factors for the years 2001 to 2006 — without any measurements in Switzerland — the data has been interpolated from the European data. E.g. for the year 2006 a value of 0.035 kgPFC/tAL, results with a European average emission factor of 0.14 kgPFC/tAL and a correction factor of 0.25. For the ratio of CF₄ to C₂F₆ a value of 90% to 10% is applied. As it was not possible to perform industry independent measurements, and because of the fact that aluminium production was closed in 2006 it is not possible to redo any measurements or to collect any information about the process details retroactively. The emission factors have decreased by a factor of about 4.9 between 1990 and 2006 due to technical efforts to reduce emissions (Alcan 2003).

The factors according to Table 4-24 are used. The large difference between the emission factors of the year 1999 and 2000 is based on measured data given by the company.

Table 4-24 PFC emissions factors for aluminium production in Switzerland. Aluminium production in Switzerland ceased in the year 2006. Data beyond 2009 is not presented (emissions are not occurring).

| Gas | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------------------|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CF ₄ | kg/t | 0.1530 | 0.1373 | 0.1215 | 0.1058 | 0.0900 | 0.0833 | 0.0765 | 0.0698 | 0.0630 | 0.0540 |
| C ₂ F ₆ | kg/t | 0.0170 | 0.0153 | 0.0135 | 0.0118 | 0.0100 | 0.0093 | 0.0085 | 0.0078 | 0.0070 | 0.0060 |

| Gas | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------------------|------|--------|--------|--------|--------|--------|--------|--------|------|------|------|
| CF ₄ | kg/t | 0.0360 | 0.0360 | 0.0360 | 0.0360 | 0.0338 | 0.0315 | 0.0315 | NO | NO | NO |
| C ₂ F ₆ | kg/t | 0.0040 | 0.0040 | 0.0040 | 0.0040 | 0.0038 | 0.0035 | 0.0035 | NO | NO | NO |

| Gas | Unit | 2010 | 2011 | 2012 |
|-------------------------------|------|------|------|------|
| CF ₄ | kg/t | NO | NO | NO |
| C ₂ F ₆ | kg/t | NO | NO | NO |

Activity Data

In 2006 the last production site of aluminium in Switzerland was closed. Activity data on aluminium production from 1997 to 2006 is based on annual data published by the Swiss Aluminium Association. For earlier years, the data was provided directly by the aluminium industry. Activity data for aluminium production in Switzerland is given in Table 4-23.

4.4.2.3 Use of SF₆ in Aluminium and Magnesium Foundries (2C4)

Methodology

SF₆ is used in aluminium and magnesium foundries in the cleaning process as inert gas to fill casting forms. The Swiss Foundry Association (GVS) has not provided information on emission factors and hence a Tier 1 based approach is used. The inventory data on SF₆ used in aluminium and magnesium foundries (2C4) is based on the total imported amount of SF₆ according to the import statistic. It is assumed that the total imported amount is emitted within one year. For the inventory of any particular year the mean value of the imports in the present and the previous year is used to account for possible time lag between import and consumption (e.g. for 2012 inventory the mean value of 2012 and 2013 import data is used).

Emission Factors

For SF₆ used in aluminium and magnesium foundries (2C4) it is assumed that the total imported amount is emitted (IPCC 2006, default emission factor of 1000 kg per ton of imported substance).

Activity Data

Activity data on SF₆ used in aluminium and magnesium foundries (2C4) is based on import data. For the activity data of any particular year the mean value of the imports in the present and the previous year is used to account for possible time lag between import and consumption (e.g. for 2012 the mean value of 2011 and 2012 import data is used). SF₆ is used in Swiss aluminium and magnesium foundries since 1997. There have been two magnesium foundries known to be using SF₆. In 2007 one of them closed down production which led to a reduction in activity data for magnesium foundries by 25% from 2007 to 2008. The remaining magnesium foundry reported activity data for 2008 to 2012 to the SWISSMEM statistics. The fact that only one magnesium foundry uses SF₆ was confirmed by a survey which has been carried out in 2011 within members of the Swiss Foundry Association (GVS). Use of SF₆ in Aluminium foundries is not occurring in 2012. The import amount for aluminium cleaning is extrapolated from an estimate value given in the year 2003 by an import company. Details on the imported amount are not available for later years. A steady decrease since 2003 is assumed for import of SF₆ used for aluminium cleaning. This assumption is based on the above mentioned survey and on information which was obtained on other applications within the category 'others' from FOEN import statistics which indicates that decreasing amounts of SF₆ are used for aluminum cleaning.

4.4.2.4 Other (2C5)

Battery recycling and non-ferrous metal foundries (2C5)

There is one plant recycling batteries in Switzerland. The recycling is done applying the Sumitomo process. The batteries are first pyrolysed at temperatures of 700 °C in reducing atmosphere in a shaft kiln. The gas with the carbonised components then goes to a post-combustion step where it is completely oxidised at temperatures of 1000 °C. The flue gas is then led to flue gas cleaning. The metal fraction from the pyrolysis goes to a melting furnace where it is reduced by addition of coal and magnesium oxide. As reducing agent coke and Carburit is used.

In Switzerland there are one large and several small plants operating non-ferrous metal foundries. During the melting process emissions of CO and NMVOC occur.

Methodology

To determine emissions of CO₂, NO_x, CO and SO₂ from battery recycling and of CO and NMVOC from non-ferrous metal foundries, a country specific approach is used. The emissions are calculated by multiplying the annual amount of recycled batteries and produced non-ferrous metals by the corresponding emission factors.

Emission Factors

The emission factors of CO₂, NO_x, CO and SO₂ from battery recycling and of CO and NMVOC from non-ferrous metal foundries in Switzerland are country specific and base on measurements from industry and expert estimates documented in EMIS 2014/2C5e Batterie-Recycling and 2014/2C5e Buntmetallgiessereien Elektroöfen (see Table 4-22).

Activity Data

The annual amount of recycled batteries and produced non-ferrous metals in Switzerland is reported from industry and the foundry association as documented in EMIS 2014/2C5e Batterie-Recycling and 2014/2C5e Buntmetallgiessereien Elektroöfen (see Table 4-21).

4.4.3 Uncertainties and Time-Series Consistency

4.4.3.1 Uncertainty for key category 2C1 Iron and Steel Production

The uncertainty for CO₂ emissions in steel production amounts to 40.3 %. Production data of the steel industry has a high confidence and its uncertainty is estimated to 5% (EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen). The uncertainty for the CO₂ emission factor is estimated to be 40% (EMIS 2014/2C1 Stahl-Produktion Elektroschmelzöfen).

4.4.3.2 Uncertainty for source category 2C4 Use of SF₆ in Aluminium and Magnesium Foundries

For the use of SF₆ in Aluminium and Magnesium Foundries, an uncertainty of 15% (with normal distribution) is assumed, which is a result of a Monte Carlo simulation of the emissions of F-gases (Carbotech 2014).

4.4.3.3 Qualitative estimate of uncertainties for non-key category 2C5 Other

For non-key categories, the NIR provides qualitative estimates of uncertainties. The terms high, medium and low data quality are used and a quantitative relative uncertainty is assigned to every uncertainty category (see Table 1-14, semi-quantitative uncertainties for non-key categories). The uncertainty for CO₂ emissions from source category 2C5 is estimated to be medium and thus amounts to 10%.

The time series is consistent.

4.4.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

In submission 2012 the CO₂ emission factor of source category 2C1 Steel Production used in the Swiss Inventory was compared with the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and the IPCC default value (INFRAS 2012). Switzerland's factor lies in the lower end of the countries. This is due to the fact that in Switzerland only secondary steel making occurs; see chpt. 4.4.2.1.

For source category 2C4 Use of SF₆ in Aluminium and Magnesium Foundries the data received from SWISSMEM and import firms has been checked for double counting.

4.4.5 Source-Specific Recalculations

2C1 Iron foundries: The activity data has been revised due to corrected production shares of cupola and electric furnaces in iron foundries for 2010 and 2011. The EF value for NMVOC for 2012 has been updated resulting in revised interpolated values 1991-2011 as well.

2C1 Steel production: The EF values for CO₂ have been revised for the entire time series based on effective data of the carbon content of the consumed electrodes, the steel scrap and the alloying elements of the years 2005-2011 (emission reduction of about one order of magnitude).

2C5e Battery recycling: Activity data for 2011 has been revised due to corrected data from the plant operator.

4.4.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines..

4.5 Source Category 2D – Other Production

4.5.1 Source Category Description

Source category 2D Other Production comprises process emissions of indirect greenhouse gases from the production of pulp and paper including chipboard, fibreboard and cellulose, of food and drink as well as of charcoal. Biogenic CO₂ emissions from the production of beer, brandy, bread and wine within source category 2D2 Food and Drink are not reported. From this year's submission on charcoal production is no longer reported in source category 2D3 Wood Processing but in source category 1A1c (see section 3.2.6.2)¹³.

Table 4-25 Specification of source category 2D Other Production. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 2D | Source | Specification | Data Source |
|-----|----------------|--|-----------------------|
| 2D1 | Pulp and Paper | Emissions from NMVOC from pulp and paper including chipboard, fibreboard and cellulose production (ceased in 2008) | AD, EF: EMIS 2014/2D1 |
| 2D2 | Food and Drink | Emissions of CO and NMVOC from production of food and drink | AD, EF: EMIS 2014/2D2 |

4.5.2 Methodological Issues

4.5.2.1 Pulp and paper production (2D1)

Methodology

To determine NMVOC emissions from pulp and paper production a country specific approach is used. The emissions are calculated by multiplying the annual amount of processed pulp and paper by the corresponding emission factors. Please note that the cellulose production in Switzerland closed down in 2008.

¹³ The CH₄ emissions from charcoal production have already been shifted to sector 1A1c in the NIR submission from September 2013 (FOEN 2013g).

Emission Factors

The emission factors for NMVOC emissions from pulp and paper production in Switzerland are country specific and base on measurements and data from industry and expert estimates documented in EMIS 2014/2D1.

Table 4-26 Emission factors for CO and NMVOC in pulp and paper production, food and drink production and charcoal production for 2012 (EMIS 2014/2D1, EMIS 2014/2D2 and EMIS 2014/2D3).

| 2D Other production | Unit | CO | NMVOC |
|---|------|-----|-------|
| 2D1 Pulp and paper | g/t | NA | 580 |
| 2D2 Food and drink (exc. beer, wine, spirits) | g/t | 250 | 1'200 |
| 2D2 Food and drink (beer, wine, spirits) | g/m3 | NA | 360 |

Activity Data

The annual amount of pulp and paper produced in Switzerland bases on data from industry and expert estimates documented in EMIS 2014/2D1.

Table 4-27 Production of pulp and paper, food and drink and charcoal in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2D1, EMIS 2014/2D2 and EMIS 2014/2D3).

| 2D Other production | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 2D1 Pulp and paper | Gg | 604 | 608 | 663 | 668 | 632 | 593 | 567 | 586 | 615 | 629 |
| 2D2 Food and drink (exc. beer, wine, spirits) | Gg | 2'254 | 2'253 | 2110 | 2186 | 2092 | 2116 | 2240 | 2167 | 2177 | 2061 |
| 2D2 Food and drink (beer, wine, spirits) | m3 | 560'972 | 581'643 | 579'714 | 546'882 | 531'068 | 516'519 | 497'401 | 505'873 | 461'979 | 476'067 |

| 2D Other production | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 2D1 Pulp and paper | Gg | 641 | 640 | 645 | 634 | 652 | 660 | 696 | 752 | 728 | 514 |
| 2D2 Food and drink (exc. beer, wine, spirits) | Gg | 2'301 | 2'083 | 2'276 | 2'246 | 2'153 | 2'138 | 2'167 | 2'344 | 2'370 | 2'467 |
| 2D2 Food and drink (beer, wine, spirits) | m3 | 492'208 | 481'114 | 466'112 | 461'071 | 475'754 | 452'877 | 451'924 | 462'141 | 479'293 | 465'753 |

| 2D Other production | Unit | 2010 | 2011 | 2012 |
|---|------|---------|---------|---------|
| 2D1 Pulp and paper | Gg | 570 | 534 | 538 |
| 2D2 Food and drink (exc. beer, wine, spirits) | Gg | 2'433 | 2'484 | 2'415 |
| 2D2 Food and drink (beer, wine, spirits) | m3 | 467'699 | 462'446 | 454'903 |

4.5.2.2 Food and drink production (2D2)

Methodology

To determine CO and NMVOC emissions from food and drink production a country specific approach is used. The emissions are calculated by multiplying the annual amount of produced food and drink by the corresponding emission factors.

Emission Factors

The emission factors for CO and NMVOC emissions from food and drink production in Switzerland are country specific and base on measurements and data from industry and expert estimates documented in EMIS 2014/2D2 (see Table 4-26).

Activity Data

The annual amount of food and drink produced in Switzerland base on data from industry, farmers' association and expert estimates documented in EMIS 2014/2D2 (see Table 4-27).

4.5.3 Uncertainties and Time-Series Consistency

The time series is consistent.

4.5.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

4.5.5 Source-Specific Recalculations

2D2 Food and Drink: Activity data of meat smokehouses has been updated for the years 2007-2011.

4.5.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

4.6 Source Category 2E – Production of Halocarbons and SF₆

No emissions occurring in this sector within Switzerland. There is no production of HFC, PFC or SF₆ in Switzerland.

4.7 Source Category 2F – Consumption of Halocarbons and SF₆

4.7.1 Source Category Description

Tier 1 Key Category 2F1

HFC from the consumption of halocarbons and SF₆; Refrigeration and Air Conditioning Equipment (level and trend).

Tier 2 Key Category 2F1

HFC from the consumption of halocarbons and SF₆; Refrigeration and Air Conditioning Equipment (level and trend).

Tier 2 Key Categories 2F9

SF₆ from the consumption of halocarbons and SF₆; Other (level and trend).

HFC from the consumption of halocarbons and SF₆; Other (level and trend).

Source category 2F comprises HFC, PFC and SF₆ emissions from consumption of the applications listed below.

Table 4-28 Specification of source category 2F Consumption of Halocarbons and SF₆. Data source: Carbotech (2014).

| 2F | Source | Specification | Data Source |
|-----|--|---|---|
| 2F1 | Refrigeration and Air Conditioning Equipment | Emissions from Refrigeration and Air Conditioning Equipment | AD: Various national statistics [1] and industry data EF: Industry data and expert estimates |
| 2F2 | Foam Blowing | Emissions from Foam Blowing, incl. Polyurethane Spray | AD: Industry data and import statistics EF: Expert estimates |
| 2F3 | Fire Extinguishers | Not occurring in Switzerland | - |
| 2F4 | Aerosol / Metered Dose Inhalers | Emissions from use as aerosols, incl. metered dose inhalers | AD: Import statistics EF: IPCC default values |
| 2F5 | Solvents | Emissions from use as solvents | AD: Import statistics EF: IPCC default values |
| 2F6 | Other applications using ODS substitutes | Not occurring in Switzerland | - |
| 2F7 | Semiconductor Manufacturing | Emissions from use in semiconductor manufacturing | AD: Import statistics and industry data[2] EF: IPCC default values and industry data |
| 2F8 | Electrical | Emissions from use in | AD: Industry data EF: Industry data |
| 2F9 | Other | Emissions of SF ₆ which are not yet | AD: Import statistics and Industry data EF: Industry data and estimates |

[1] e.g. statistics on registration of cars and trucks, import statistics on F-gases (Carbotech 2014).

[2] e.g. import amount of some substance for specific company with known application type.

The following graph shows emissions in source category 2F by sub-sector and by different groups of gases. Refrigeration and air conditioning equipment account by far for the highest emissions in this source category with a share of 80% of the total emissions in the source category 2F.




Figure 4-6 Distribution of emissions under source category 2F Consumption of Halocarbons and SF₆ (2012 data).

4.7.2 Methodological Issues

The data models used for source category 2F are complex and therefore a comprehensive documentation of all relevant model parameters is not possible within the framework of the NIR. Annex A3.2 shows an illustrative example of the model structure and parameters used for calculating emissions from mobile air-conditioning in cars. Where possible, the most important assumptions for the data model are documented (e.g. Table 4-29). Detailed documentation of the individual data models is available from Carbotech (2014) as well as related background documents. This information is FOEN internal due to confidentiality of data, but is open for consultation by reviewers.

4.7.2.1 Refrigeration and Air Conditioning Equipment (2F1)

Methodology

The inventory under this sub-source category includes the following types of equipment: domestic refrigeration, commercial and industrial refrigeration, transport refrigeration, stationary air conditioning, mobile air conditioning, and heat pumps. For each of these types of equipment individual emission models are used for calculating actual emissions as per IPCC GPG Tier 2. In order to obtain the most reliable data for the calculations, two different approaches are applied to get the stock data needed for the model calculations: 'top down' using available statistics or estimations on the Swiss market from experts and associations

and 'bottom up' through questionnaires sent to companies active in importation, production and service of appliances.

The import data as reported to FOEN was adjusted for imported substances to be used in Liechtenstein. This is to eliminate double counting with the inventory data of Liechtenstein. Under source category 2F1 import data from the year 2008 onwards which is related to commercial and industrial refrigeration equipment are split between Switzerland and Liechtenstein. The split factor is based on the proportion of employees in the industrial and service sector (share of import for Liechtenstein < 1%). For other equipment types no scope for double counting with the inventory of Liechtenstein was identified and therefore no adjustment is required.

For the present submission also a number of minor improvements and corrections have been made to the model assumptions on emissions factors and activity data for source category 2F1. Further details can be seen from the section on recalculations.




Figure 4-7: Required data for the model calculation of refrigeration and Air Conditioning in Switzerland

Emission Factors

Emission factors for manufacturing, product life and disposal as well as average product life times are established on the basis of expert judgement and literature. Direct monitoring of the product life emission factors is only done at company level (i.e. retailers such as Coop and Migros). The product life factors are used to make the allocation of imported F-gases to new products and maintenance activities.

In 2008 a revised ordinance on chemical risk reduction (Swiss Confederation 2005) was introduced. As part of this revision an obligation for operators handling equipment containing more than 3 kg HFCs was introduced to provide information to FOEN on the date of operation start, type of equipment, type and amount of refrigerant and date of disposal. Today the statistics on equipment containing more than 3kg are comprehensive. However,

these figures only cover about 50% to 70% of the Refrigeration and Air Conditioning Equipment reported under source category 2F1, since there are many types of equipment containing less than 3kg HFCs. Furthermore, there is no information available from the statistics regarding the emissions due to operation losses from the registered equipment. This data source provides valuable information to improve the estimates used for modelling emissions under source category 2F. However, it will not allow to directly draw the stock data or emission factors for the national inventory from this database in the near future.

Table 4-29 displays the detailed model parameters used for the present inventory. For product life emission factors of some equipment types a dynamic model is applied which implies that emission losses improve linearly between 1995 and 2012 (respectively 2020 for some equipment types) due to better production technologies and the continuous sensitisation of service technicians. The start/end values are based on expert statements, UBA (2005, 2007) and Schwarz (2001, 2005). The charge at end of life for different applications has been analysed considering the technical minimal charge of equipment and the expected frequency of maintenance (UBA/Ökorecherche 2012).

Table 4-29 Typical values on life time, charge and emission factors used in model calculations for Refrigeration and Air Conditioning Equipment. Data between 1995 and 2012 respectively 2020 is linearly interpolated.

| Equipment type | Product life time | Initial charge of new product | Manufacturing emission factor | Product life emission factor | Charge at end of life [% of initial charge of new product] *) | Disposal loss emission factor |
|---|-------------------|---|-------------------------------|---|---|-------------------------------|
| | [a] | [kg] | [% of initial charge] | [% per annum] | | [% of remaining charge] |
| Domestic Refrigeration | 16 | 0.1 | NO | 0.5 | 92 | 19 **) |
| Commercial and Industrial Refrigeration | 10 | NR | 0.5 | 12 (2012: 7.2) (2020: 5) | 87 | 20 |
| Transport Refrigeration / Trucks | 10 | 1.8-7.8 (various types) | 1.5 | 15 | 86 | 20 |
| Transport Refrigeration / Railway | 16 | NR | NO | 10 | 100 | 20 |
| Stationary Air Conditioning (direct / indirect cooling system) | 15 | NR | direct: 3 indirect: 1 | direct: 10 (2010: 4) indirect: 6 (2010: 4) | 85-90 (direct 89%, indirect 86%) | direct: 28 indirect: 19 |
| Heat Pumps | 15 | (1999: 4.7-7.5) decreasing to (2010: 2.8-4.5) | 3 | 2 | 86 | 20 |
| Mobile Air Conditioning / Cars | 15 | 0.7 (0.84) ***) | NO | 8.5 | 58 | 50 ****) |
| Mobile Air Conditioning / Trucks | 12 | 8.5 | NO | 10 until year 2000 decreasing to 8.35 in 2011 | 69 | 50 ****) |
| Mobile Air Conditioning / Buses | 12 | 7.5 | NO | 10 until year 2010 8.5 for 2011 onwards | 100 | 50 ****) |
| Mobile Air Conditioning / Railway | 16 | NR | NO | 5.5 | 100 | 10 |

*) takes into account refill of losses during product life where applicable.

***) takes into account R134a content in foams, based on information from the national recycling organisation SENS.

****) Assumed constant since 2002. 0.84 kg in 1990. Linear interpolation between 1990 and 2002.

*****) HFC disposal losses occur from 2003 onwards for Trucks/Buses resp. from 2006 for Cars (introduction of HFCs in MAC from 1991 only and 12 resp. 15 years lifetime). Value of 50% is based on UBA 2005 and expert assumptions on share of total refrigerant loss, e.g. due to road accident.

NR = not relevant as only aggregate data is used

NO = Not occurring (only import of charged units)

Activity Data

Activity data is taken from industry information and national statistics such as for admission of new cars and trucks. Stock data is modelled dynamically. Due to the large number of sub-models used for modelling the total emissions for sub-source category 2F1, no table on time series of activity data is provided here, despite 2F1 being a key category. For illustration, the detailed calculation model for car air-conditioning including the time series for the activity data for this particular sub-model can be seen from Annex A3.2. Mobile air-conditioning accounts for approx. 32% of the total emissions (CO₂ eq) of sub-source category 2F1 Refrigeration and Air Conditioning Equipment.

For the inventory report 2012 (FOEN 2012) a cross check has been performed for results from model calculation and FOEN statistics on disposal and recycling of HFCs. This has indicated a significant gap. Some of the gap is explained by the onsite reuse and recycling of refrigerants, which is not reflected by the FOEN statistics and with other factors as for example the not accounted export of refrigeration equipment (only export of vehicles with air-conditioning considered).

To avoid double counting with the inventory data of Liechtenstein, the activity data for the equipment type commercial and industrial refrigeration from the year 2008 onward is reduced by 0.9%, based on the share of imports of substances to be used in Liechtenstein. The reduction factor is based on the proportion of employees in the industrial and service sector in these two countries. For other equipment types no scope for double counting with the inventory of Liechtenstein was identified and therefore no correction factor is applied.

4.7.2.2 Foam Blowing (2F2)

Methodology

In Switzerland no production of open cell foam based on HFCs is reported by the industry. Therefore only closed cell PU and XPS foams, PU spray applications and sandwich elements are relevant under this source category.

The emission model (Tier 2) for foam blowing has been developed 'top down' based on import statistics for products, industry information and expert assumptions for market volumes and emission factors. Emissions for sandwich elements have been calculated as residual balance between SAEFL import statistics and consumption in PU spray, PU and XPS foams.

Emission Factors

For emission factors and lifetime of XPS and PU foam, expert estimates and general default values according to IPCC are being used (IPCC 2000: p. 3.95). For PU spray, expert estimates and specific default values according to IPCC are being used (IPCC 2000: p. 3.96).

Table 4-30 Typical values on life time, charge and emission factors used in model calculations for foam blowing.

| | Product life time | Charge of new product | Manufacturing emission factor | Product life emission factor | Charge at end of life |
|-------------------|-------------------|-----------------------|-------------------------------|------------------------------|-------------------------|
| | years | % of product weight | % of initial charge | % per annum | % charge of new product |
| PU foam | 50 | 4.5 | NR | NR | NR |
| XPS foam HFC 134a | 50 | 6.5 | NR | 10 / 0.66** | 100 |
| HFC 152a | | | | 100 / 0** | 100 |
| PU spray | 50 | 13.6 / 0 * | 95 | 95 / 2.5 ** | 100 |
| Sandwich Elements | | | | | |
| HFC 134a, | 50 | NR | 10 | 0.7 | 100 |
| HFC 227ea, | | | | | |
| HFC 365 mfc | | | | | |
| HFC 152a | | | 100 | 0 | 100 |

* Data for 1990 / since 2009

** Data for 1st year / following years

NR Not relevant (PU foam: no substances according to this protocol have been used; XPS foam: emissions occur outside Switzerland; Sandwich elements: calculations are based on the remaining propellant import amount)..

Activity Data

HFCs have been used till 2008 in the Swiss production of PU spray. The export rate of PU spray from Swiss production is about 96.5% of total production volume. About one third of PU spray sold in Switzerland originates from local production, the rest is import. For PU rigid foams no HFCs are used as foam blowing agent (only Pentane and CO₂). From 2000 onwards until 2010 there is no production of XPS in Switzerland with HFC. XPS foams were 100% imported until 2010. In 2011 a new production facility was started which does not use HFCs. The HFC import not related to the main applications above has been allocated to the production of sandwich elements.

Detailed activity data for this sub-source category is available at FOEN but not reported due to confidentiality.

4.7.2.3 Fire Extinguishers (2F3)

No emissions occurring in this sector within Switzerland. The application of HFC, PFC and SF₆ in fire extinguishers is prohibited by law.

4.7.2.4 Aerosol / Metered Dose Inhalers (2F4)

Methodology

The Tier 2 emission model for Aerosol / MDI is based on a 'top down' approach using import statistics for HFCs.

Emission Factors

A manufacturing emission factor of 1% is applied. For product life emission factor the model assumes that 50% of the remaining substance is emitted in the first year and 50% in the second year respectively, which is in line with IPCC GPG. To account for variations in imports and stocks, the average figure from imports for the actual year (t) and for the past year (t-1) is reported. This emission model can lead to implied product life emission factors of > 100% in case of decreasing imports.

Activity Data

In most aerosol applications, HFC has been replaced already in the past years. According to the information of companies filling aerosol bottles for use in households, e.g. cosmetics, cloth care and paint, no HFC is being used. For special technical applications - especially metered dose inhalers (MDI) - HFC is still in use. Compared to the total amount of aerosol applied, the HFC use for MDI is considered to be irrelevant.

Activity data is based on import statistics. The export and import of filled products is unknown and assumed to be in a similar range. Detailed activity data for this sub-source category is available at FOEN but not reported due to confidentiality.

4.7.2.5 Solvents (2F5)

Methodology

HFC and PFC are used as solvents. Emissions are calculated according to Tier 1 method according to IPCC GPG on basis of a 'top down' approach using import statistics and industry information on allocation of the imported HFC and PFC amounts to different applications.

The import data as reported to FOEN was adjusted for imported substances to be used in Liechtenstein. This is to eliminate double counting with the inventory data of Liechtenstein. Under source category 2F5 import data from the year 2008 onwards are split between Switzerland and Liechtenstein. The split factor is based on the proportion of inhabitants.

Emission Factors

An emission factor of 50% in the first and in the second year, respectively, is applied in line with IPCC GPG.

Activity Data

Activity data is based on import statistics. Detailed activity data for this source category is available at FOEN but not reported due to confidentiality. For the inventory report of the year 2011 (FOEN 2011) interviews were made with industry to get in-depth information on allocation of imported HFC and PFC volumes to different applications. This resulted that most PFC import declared as Solvents (2F5) or Other (2F9) until 2010 are related to the semiconductor manufacturing and thus the model for allocation of imported PFC volumes was adjusted accordingly. Since 2011 imports for semiconductors manufacturing and further etching process are registered separately.

To account for double counting with the inventory data of Liechtenstein, the import data reported to FOEN which is assigned to source category 2F5 in the inventory of Switzerland is adjusted by 0.5%. The adjustment factor is based on the proportion of inhabitants in these two countries.

4.7.2.6 Other applications using ODS substitutes (2F6)

No emissions occurring in this sector within Switzerland.

4.7.2.7 Semiconductor Manufacturing (2F7)

Methodology

A Tier 2 approach with process gas-specific parameters was used for emission calculations. General default values for gas-specific transformation rate and general values for exhaust treatment were applied.

Up to the inventory report 2010 (FOEN 2010), HFC, PFC and SF₆ emissions under 2F7 Semiconductor Manufacturing were calculated according to Tier 1 method according to IPCC GPG on basis of a 'top down' approach using import statistics. For the inventory report 2011 (FOEN 2011) interviews had been made with industry to get in-depth information on allocation of imported PFC volumes to different applications and to obtain process specific information from consumers. This resulted that until 2010 most PFC import declared as Solvents (2F5) or Other (2F9) are related to the semiconductor manufacturing and thus the model for allocation of imported PFC volumes was adjusted accordingly which leads to increased emissions under source category 2F7 Semiconductor Manufacturing. Since 2011 PFCs import declarations have been improved and information is provided for the source category 2F7 separately.

Emission Factors

Default emission factors as per IPCC GPG are used. Since the inventory report 2011 (FOEN 2012) the rate of exhaust treatment is assumed to be higher due to legislation under the Chemical Risk Reduction Ordinance (Swiss Confederation 2005) which limits emissions for industrial applications such as semiconductor manufacturing to 5%. For some large users the presence of exhaust treatment was confirmed in a survey.

Activity Data

Activity data is based on import statistics and industry information.

4.7.2.8 Electrical Equipment (2F8)

Methodology

Under an agreement with FOEN, the industry association SWISSMEM is reporting actual emissions of SF₆ on basis of a mass balance approach (Tier 3a). The balance includes mainly data for the production, installation, operation and disposal of electrical equipment, but included in past years also small amounts of SF₆ for other applications (i.e. research, magnesium foundry). SWISSMEM is collecting data from its members and is cross-checking the reported SF₆ consumption data with data from importers of the SF₆. Installations in operation with electrical equipment containing SF₆ are periodically inspected for leakage and losses are refilled (topping up). The refilled quantities and any SF₆ charge required for during repair are reported as emissions at the time of filling. A product lifetime of 35 years is applied.

Emission Factors

Emission factors for this sub-source category are based on industry information and are calculated values based on the mass balance data. For 2012 the calculated product life

emission factor is 0.12%. The calculated product life emission factor is varying between 0.5%/a (2005) and 0.12%/a (2012). The discontinuity in emission factor from 2005 to 2006 data is partly due to the inspection intervals, optimised data collection system and technical optimisation of equipment. The continued trend for reduced emission factors can be linked to the existing agreement of SWISSMEM and FOEN on reduction of SF₆ emissions.

Activity Data

Activity data is based on industry information. The wide annual fluctuation of SF₆ emissions from electrical equipment is related to the annual fluctuation of market volumes for such equipment as well as variations in inspection intervals and equipment break-down requiring topping up of SF₆ charge in the equipment. Also for inventory report 2012 (FOEN 2012) the split factors for allocation of imported amounts to different applications were checked through industry interviews and in-depth analysis in order to eliminate double counting between SWISSMEM data and other import declarations.

4.7.2.9 Other (2F9)

Methodology

The emissions reported under 2F9 relate to a small amount of unallocated SF₆ from the FOEN import statistics and since 2003 to further applications of halocarbons such as laboratory and research use. In the past years an increasing amount of CF₄ and HFC 134a was registered to be used as trace gas, particularly in nuclear research. The unallocated difference for SF₆ between the FOEN import statistics and the SWISSMEM mass balance (see 2F8) have been assigned to cables and electrical control systems using a Tier 2 approach. Some imports of HFC 134a were declared for medical use, and small import amount of HFC 23 was declared for electronics and refrigeration technology.

Emission Factors

For the unallocated amount of SF₆ assigned to cables and electrical control systems the manufacturing emission factor is assumed at 4% and the product life emission factor at 1%. 1% of the remaining charge is emitted at time of disposal after 40 years lifetime. Because of the long life time the disposal emissions are not relevant for the given results.

According to the IPCC guidelines (IPCC 2000) the emission factors for HFC 134a (medical and research use) and for HFC 23 (electronics and refrigeration technology) were chosen as 50% in the first year and 50% in the second year.

For the CF₄ and SF₆ related to analytics, laboratory and research use a 50% lower emission factor was assumed considering a transformation and an exhaust treatment in some of the applications.

Activity Data

Activity data is based on import statistics and industry information. For the unallocated amount of SF₆ assigned to cables and electrical control systems an export rate of 80% was assumed comparable to electrical equipment 2F8. Also for inventory report 2012 (FOEN 2012) the split factors for allocation of imported amounts to different applications was checked through industry interviews and in-depth analysis in order to eliminate double counting between SWISSMEM data and other import declarations. The quality check of import declarations and information obtained from import companies and SWISSMEM lead to a shift of SF₆ within different applications.

4.7.3 Uncertainties and Time-Series Consistency

For refrigeration equipment, air-conditioning equipment as well as for the foam blowing source category, a Monte Carlo analysis according to IPCC Good Practice Guidance for the evaluation of uncertainties of model calculations according to Tier 2 has been carried out. The Monte Carlo Analysis was performed on the inventory data of the current GHG inventory (submission April 2014). For the purpose of the Monte Carlo Analysis, uncertainty of all relevant parameters (e.g. initial appliance charge, product life emission factor, import and export volumes, etc.) used in the emission models for the applications as per Table 4-31 below has been characterised by a statistical distribution. Frequently a triangular distribution was chosen, defined by the three parameters: minimum, maximum and most likely value. Some uniform distributions were chosen where the spectrum was assumed to have the same probability. In the other cases normal or lognormal distribution has been chosen. The analysis was carried out with 10'000 cycles. Some details on the distributions of parameters used (i.e. type of distribution, minimum, maximum, likeliest value) are documented in the report Carbotech (2014).

For the submission of 12 April 2006 the uncertainty for the import statistic data had been estimated for the first time. Discussions with the persons responsible for data collection in the years 1997–2012 led to the estimations given in Table 4-31.

Table 4-31 Estimated uncertainty for the data of the imported substances

| Year | Minimal | Maximal | Remarks |
|-------------|---------|---------|---|
| Up to 1999 | -10% | 30% | Assumed that the data is not complete |
| 2000 – 2003 | -10% | 15% | Data can be incomplete or possible double declaration |
| 2004 – 2012 | -10% | 10% | Data can be incomplete or possible double declaration |

The following table summarises the results for the application-specific emission models. The “value 2012” represents the actual emissions in Gg CO₂ equivalent for the specific application as used for calculating the 2012 CRF-tables. The average, median, uncertainty, minimum and maximum values are output values of the Monte Carlo Analysis. The uncertainty of the resulting total emissions from the consumption of halocarbons and SF₆ is about 10.4%. Higher values result for the contributions of single applications.

Uncertainties of more than 20% have been calculated for the following applications:

- Stationary Airconditioning
- Transport Refrigeration
- Domestic Refrigeration
- Foam Blowing
- Aerosols
- Solvents
- Semiconductors
- Others

Uncertainties of 15% to 20% have been calculated for the following applications:

- Mobile Airconditioning

- Commercial and Industrial refrigeration

Low uncertainties of less than 15% have been calculated for the following applications:

- Electric Equipment

For the model calculations of stocks, uncertainties result with a maximum of 35.2% for R134a in PU/XPS Foam Blowing. For the model calculations of stocks in domestic refrigeration no uncertainties value is given due to very asymmetric distribution. Calculation of stocks is not reported in detail here because the uncertainties for stock and new filled refrigerant related to the split of refrigerant on different applications is of less relevance for the overall emissions. This is because different applications show similar characteristics for the building of stocks and related emissions. Detailed data is available with FOEN.

Relevant parameters for the building of stock in PU-foam are the PU-foam import and export rate and the PU-Spray first year emission factor. The data base for PU-Sprays has been significantly improved with effect from the 2007 submission (FOEN 2007). This is attributed to improved models which are elaborated by the main producer and its blowing agent import firm. However, the following three factors lead to a small amount remaining in the stock with a relative high uncertainty: high import and export rate of PU-Spray, incompleteness of information on import volumes of PU-Spray and about propellant used in import products and finally high emission factor of the first year.

Table 4-32 Summary of results for model parameter “emissions” from Monte Carlo Analysis for 2012 data on selected emission sources.

| Application | Model parameter | Value 2012 Gg CO2 eq. | Average Gg CO2 eq. | Median Gg CO2 eq. | min. Gg CO2 eq. | max. Gg CO2 eq. | Uncertainty % |
|--|-------------------------|--------------------------|-----------------------|----------------------|--------------------|--------------------|------------------|
| 2F1 Refrigeration and Airconditioning | Emissions in Gg CO2 eq. | 1142 | 1165 | 1163 | 943 | 1770 | 11.2 |
| - Commercial / Industrial Refrigeration | | 602 | 621 | 618 | 422 | 852 | 19.4 |
| - Mobile Air-Conditioning | | 366 | 340 | 356 | 264 | 474 | 20 |
| - Stationary Air-Conditioning | | 134 | 149 | 149 | 100 | 679 | 25 |
| - Transport Conditioning | | 25.1 | 28.7 | 28.5 | 18.7 | 44.8 | 25 |
| - Domestic Refrigeration | | 15.5 | 8.7 | 6.7 | 0.2 | 35.9 *) | |
| 2F2 Foam Blowing | | 14.5 **) | 16.7 | 16.4 | 7.1 | 37.8 | 43.2 |
| 2F4 Aerosol | | 13.2 | 13.3 | 13.3 | 5 | 23 | 41 |
| 2F5 Solvents | | 11.9 | 12 | 12 | 8.8 | 15.9 | 23.2 |
| 2F7 Semiconductors | | 26 | 27.5 | 27.4 | 9.9 | 45.5 | 39.2 |
| 2F8 Electrical equipment | | 36.8 | 36.8 | 36.8 | 27 | 46 | 14.4 |
| 2F9 Other | | 212 | 190 | 206 | 113 | 282 | 41.6 |
| Total HFC, PFC and SF ₆ from 2F | | 1470 | 1473 | 1473 | 1174 | 2037 | 10.4 |

*) very asymmetric distribution, therefore no indication of a standard deviation.

**) incl. HFC 365mfc

The time series is consistent for all source categories, with exception of the sub-source category Electrical Equipment (2F8) where from 2000 onwards the data is based on a Tier 3a approach instead of model calculations according to Tier 2 as applied for data before 2000. Due to lack of basic information it is not possible to provide a consistent time series for category Electrical Equipment (2F8) retroactively.

4.7.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables.
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.

- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013.

Recalculations were identified and explained. Detailed controls of all modelling results produced by Carbotech (2014) have been carried out firstly by FOEN specialists and secondly by the author for the NIR chapters containing F-gases.

The assumption of decreasing emissions factors for the different equipment types under sub-source category 2F1 Refrigeration and Air Conditioning Equipment have been cross-checked with the inventories of Austria and Germany and have found to be in line with the assumptions made for these inventories.

The emission factor of category 2F used in the Swiss Inventory has been compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (INFRAS 2012). Concerning the consumption of halocarbons and SF₆ the following sources of emissions are deemed relevant: HFC-125, HFC-134a and HFC-143a from stationary and commercial refrigeration as well as mobile air-conditioning. The product life factor is relevant, since there is no production of neither halocarbons or SF₆ in Switzerland. For all these sources Switzerland's emission factors lie in the midfield of the other countries except for the life factor in mobile air-conditioning. However when compared to neighbouring countries such as Germany, very similar values are used. The Swiss product life factors are often lower than the average for the following reasons. Since 2005 the ordinance on Chemical Risk Reduction (Swiss Confederation 2005) is in place that ensures the proper handling and disposal of halocarbons and SF₆. Furthermore the decommissioning sector is well organized by the SENS foundation and recycling is taxed in advance. Finally servicing staff is well trained to proper handling and disposal of respective appliances.

The FOEN supports a monitoring campaign at the high altitude research station Jungfraujoch, where various greenhouse gases are measured continuously. The location of the research station normally provides for analysis of tropospheric background concentrations. However, under special meteorological conditions, an estimate of Swiss emissions can be derived from the measurements. For HFC-134a, HFC-125, HFC-152a HFC-143a and HFC-32, a comparison of the inventory data with the inferred emissions is presented in Annex A6.1.

Special effort was undertaken to verify the underlying reasons for the discontinuity in emission factor of SF₆ in source category 2F8 Electrical equipment from 2005 to 2006 data. It however was not possible to find new supporting facts. With the change of personell at the data supplier SWISSMEM the basis for historical in-depth investigations has ceased.

4.7.5 Source-Specific Recalculations

In the data files used for calculating the emissions from Mobile Air-Conditioning / Buses in source category 2F1 an error has been discovered. Accordingly all emissions in Submission 2013 related to the equipment type Mobile Air-Conditioning / Buses were not taken into account in the total figure for emissions under source category 2F1. The emissions from mobile air-conditioning have been recalculated now, including emissions from buses. The recalculation results in additional emissions of HFC 134a of the order of 25-28 Gg CO₂eq per year for the years 2008-2011.

Further Source-specific recalculations for the time series 1990 to 2011 are summarized in Table 4-33. The different improvements carried out in the present inventory are related to the sub-source categories with the highest emissions.

The recalculation of the emissions 2011 delivers about 3.8% higher total emissions under source category 2F than reported in the previous submission.

Table 4-33 Summary of recalculations in source category 2F.

| NFR code | Sector/ Process | AD/EF | Year | Gas | Specification |
|----------------------------|---|-------|-----------|---|--|
| 2 IIA F 1 (all sources) | Refrigeration and Air-Conditioning | AD | 1991-2011 | HFC 32 HFC 125 HFC 134a HFC 143a HFC 152a PFC 218 HFC23 | Improvement of model calculations of stock. Recharge of equipment considering minimal technical charge and related frequency of service (resulting average charge between initial charge and minimal technical charge). The improvement has an impact on the calculation of stocks, emissions from stock and on the required in bulk refrigerant used in different applications (calculation of remaining in bulk refrigerant for industrial/commercial refrigeration) |
| 2 IIA F 1 6 | Refrigeration: Mobile Air-conditioning | AD/EF | 1991-2011 | HFC 134a | Bus air-conditioning added to the calculation of mobile air-conditioning (related evaluations delivered for review process) |
| 2 IIA F 1 3 | Transport refrigeration | AD | 2000-2011 | HFC 125 HFC 134a HFC 143a PFKW 218 (=C3F8) | Export of retiring equipment included in model calculations of trucks, lifetime of railway elevated (so far first HFC containing equipment still in use) |
| 2 F 4 | Aerosols | AD/EF | 1998-2012 | HFC 134a HFC 152a | Use of spray products not considering the earlier aerosol loss of production (1% double counting of aerosol emissions). |

4.7.6 Source-Specific Planned Improvements

Gradual improvement of the data quality in co-operation with industry is on-going. As in the past years, methodologies and emission models will be updated during the yearly process of F-gas inquiry. The focus will be on improvements of HFC-emission calculations from refrigeration and air-conditioning equipment.

4.8 Source Category 2G – Other

4.8.1 Source Category Description

Source category 2G Other comprises process emissions from blasting and shooting and Claus units in refineries.

Table 4-34 Specification of source category 2G Other. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 2G | Source | Specification | Data Source |
|----|--------|--|--|
| 2G | Other | Emissions of CO ₂ , NO _x , CO, NMVOC and SO ₂ from blasting and shooting Emissions of SO ₂ from Claus units in refineries | AD, EF: EMIS 2014/2G Sprengen und Schiessen AD, EF: SFOE 2013, expert estimates |

4.8.2 Methodological Issues

Blasting and shooting and Claus units in refineries (2G)

Methodology

For determination of emissions of CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting a country specific method is used as documented in EMIS 2014/2G Sprengen und Schiessen. The emissions are calculated by multiplying the annual amount of used explosive by the corresponding emission factors. The SO₂ emissions from Claus units are calculated by multiplying the annual amount of processed crude oil by the emission factor.

Emission Factors

The emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting activities in Switzerland and for SO₂ emissions from Claus units in refineries are country specific and base on measurements and data from industry and expert estimates documented in EMIS 2014/2G Sprengen und Schiessen.

Table 4-35 Emission factors for CO₂, NO_x, CO, NMVOC and SO₂ from blasting and shooting and SO₂ from Claus units in refineries for 2012 (EMIS 2014/2G Sprengen und Schiessen).

| 2G Other | Unit | CO ₂ | NO _x | CO | NMVOC | SO ₂ |
|---------------------------|------|-----------------|-----------------|-----|-------|-----------------|
| Blasting and shooting | kg/t | 400 | 35 | 310 | 60 | 0.5 |
| Claus units in refineries | g/t | NA | NA | NA | NA | 38 |

Activity Data

The annual amount of used explosives and of processed crude oil in Clause units base on the Federal statistics on explosives as documented in EMIS 2014/2G Sprengen und Schiessen and the Swiss overall energy statistics (SFOE 2013), respectively.

Table 4-36 Amount of used explosives and processed crude oil in Switzerland for the period 1990-2012 in Gg (EMIS 2014/2G Sprengen und Schiessen and SFOE 2013).

| 2G Other | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Blasting and shooting | | | | | | | | | | | |
| blasting agent and powder | Gg | 2.6 | 2.3 | 2.1 | 1.8 | 1.6 | 1.3 | 0.5 | 0.8 | 1.1 | 1.6 |
| Claus units in refineries | | | | | | | | | | | |
| crude oil | Gg | 3'127 | 4'671 | 4'317 | 4'764 | 4'880 | 4'657 | 5'289 | 4'830 | 5'070 | 5'093 |

| 2G Other | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Blasting and shooting | | | | | | | | | | | |
| blasting agent and powder | Gg | 1.9 | 2.0 | 3.3 | 4.1 | 3.6 | 0.8 | 1.5 | 1.1 | 1.4 | 2.1 |
| Claus units in refineries | | | | | | | | | | | |
| crude oil | Gg | 4'649 | 4'846 | 4'848 | 4'567 | 5'146 | 4'810 | 5'497 | 4'662 | 5'067 | 4'778 |

| 2G Other | Unit | 2010 | 2011 | 2012 |
|---------------------------|------|-------|-------|-------|
| Blasting and shooting | | | | |
| blasting agent and powder | Gg | 2.4 | 2.9 | 2.3 |
| Claus units in refineries | | | | |
| crude oil | Gg | 4'491 | 4'402 | 3'409 |

4.8.3 Uncertainties and Time-Series Consistency

For non-key categories, the NIR provides qualitative estimates of uncertainties. The terms high, medium and low data quality are used and a quantitative relative uncertainty is assigned to every uncertainty category (see Table 1-14, semi-quantitative uncertainties for non-key categories). The uncertainty for CO₂ emissions from 2G is rated medium and thus amounts to 10%.

The time series is consistent.

4.8.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables

- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

4.8.5 Source-Specific Recalculations

2G Claus units in refineries: Activity data of the years 1990–1997 has been revised due to recalculations in the Swiss overall energy statistics.

4.8.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

5 Solvent and Other Product Use

5.1 Overview

This chapter provides information on the calculation of the greenhouse gas emissions from solvent and other product use. The emissions contain NMVOC emissions from the use of solvents in different applications. Also, it includes direct CO₂ emissions resulting from post-combustion of NMVOC to reduce NMVOC in exhaust gases and indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere. Further included are emissions of CO₂, NO_x, CO and SO₂ arising from the use of firework and N₂O emissions from medical and private use.

Emissions of biogenic CO₂ from the use of tobacco products are not reported. The disposal of solvents is reported in the waste sector (Chapter 8). Emissions from the use of halocarbons and sulphur hexafluoride are reported in the Industrial Processes Chapter under 2F.

Tier 1 Key category 3

CO₂ emissions from Solvent and Other Product Use (trend).

Tier 2 Key category 3

CO₂ emissions from Solvent and Other Product Use (trend).

N₂O emissions from Solvent and Other Product Use (trend).

5.1.1 Emissions of CO₂ and N₂O




Figure 5-1 Switzerland's GHG emissions of sector 3 Solvent and Other Product Use 1990-2012 in Gg CO₂ eq.

In 2012 200 Gg of CO₂ eq emissions were released from sector 3 Solvent and Other Product Use as shown in Figure 5-1 and Table 5-1. This is a decline of 57.5% between 1990 and 2012. Source category 3D Other remains the dominant source within sector 3 Solvent and Other Product Use although its emissions have decreased by 45.8% since 1990. Source category 3A Paint Application has decreased by 73.0% since 1990, source category 3B Degreasing and Dry Cleaning has decreased by 80.3% and source category 3C Chemical Products has decreased by 63.2%.

Table 5-1 Emissions of sector 3 Solvent and Other Product Use 1990-2012 in Gg CO₂ eq.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 360 | 338 | 318 | 295 | 283 | 267 | 250 | 233 | 216 | 209 |
| N ₂ O | 110 | 106 | 101 | 96 | 92 | 86 | 81 | 75 | 70 | 65 |
| Sum | 470 | 444 | 420 | 392 | 374 | 354 | 331 | 308 | 286 | 273 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 200 | 191 | 181 | 172 | 160 | 159 | 157 | 158 | 156 | 157 |
| N ₂ O | 59 | 54 | 52 | 53 | 51 | 52 | 49 | 47 | 46 | 45 |
| Sum | 259 | 245 | 234 | 225 | 212 | 211 | 206 | 205 | 202 | 201 |

| Gas | 2010 | 2011 | 2012 |
|--------------------------|------|------|------|
| CO ₂ eq. (Gg) | | | |
| CO ₂ | 153 | 158 | 155 |
| N ₂ O | 46 | 44 | 45 |
| Sum | 199 | 202 | 200 |

The relative trends of the emissions of CO₂ and N₂O are shown in Figure 5-2. The base year 1990 represents 100%.




Figure 5-2 Relative trends of the greenhouse gases of sector 3 Solvent and Other Product Use in the period 1990-2012.

5.1.2 Emissions of NMVOC

Due to the importance of NMVOC emissions in sector 3 Solvent and Other Product Use they are given separately in Table 5-2.

Table 5-2 Emissions of NMVOC in sector 3 Solvent and Other Product Use 1990-2012 in Gg.

| NMVOC | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|------|------|------|------|------|------|------|------|------|
| | Gg | | | | | | | | | |
| 3A Paint application | 54 | 51 | 47 | 44 | 40 | 37 | 34 | 30 | 27 | 26 |
| 3B Degreasing and dry cleaning | 12 | 11 | 10 | 9 | 8 | 7 | 7 | 6 | 5 | 5 |
| 3C Chemical products, manufacture and processing | 28 | 23 | 18 | 13 | 12 | 11 | 9 | 8 | 7 | 6 |
| 3D Other | 60 | 57 | 55 | 52 | 49 | 46 | 43 | 41 | 38 | 35 |
| Sum | 155 | 142 | 130 | 118 | 110 | 101 | 92 | 84 | 76 | 72 |

| NMVOC | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|------|------|------|------|------|------|------|------|------|
| | Gg | | | | | | | | | |
| 3A Paint application | 25 | 23 | 20 | 16 | 12 | 12 | 12 | 12 | 12 | 13 |
| 3B Degreasing and dry cleaning | 4 | 4 | 4 | 4 | 3 | 3 | 3 | 3 | 3 | 2 |
| 3C Chemical products, manufacture and processing | 6 | 5 | 5 | 5 | 4 | 4 | 4 | 4 | 4 | 4 |
| 3D Other | 32 | 29 | 28 | 27 | 26 | 25 | 24 | 24 | 24 | 24 |
| Sum | 67 | 62 | 57 | 51 | 45 | 44 | 43 | 43 | 43 | 43 |

| NMVOC | 2010 | 2011 | 2012 |
|--|------|------|------|
| | Gg | | |
| 3A Paint application | 13 | 13 | 12 |
| 3B Degreasing and dry cleaning | 2 | 2 | 2 |
| 3C Chemical products, manufacture and processing | 4 | 4 | 4 |
| 3D Other | 24 | 24 | 25 |
| Sum | 44 | 44 | 43 |

NMVOC emissions have decreased by 72.0 % between 1990 and 2012. This is mainly due to two reduction efforts: The introduction of NMVOC emission limit values by the ordinance on Air Pollution Control (Swiss Confederation 1985) and the introduction of the VOC-tax in 2000 (Swiss Confederation 1997).

5.2 Source Category 3A – Paint Application

5.2.1 Source Category Description

Source category 3A Paint Application comprises NMVOC emissions from paints, lacquers, thinners and related materials used in coatings in industrial, commercial and household applications. Also, it includes indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere and direct CO₂ emissions resulting from post-combustion of NMVOC in exhaust gases.

Table 5-3 Specification of source category 3A Paint Application. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| | Source | Specification | Data Source |
|----|-------------------|--|------------------------------------|
| 3A | Paint Application | Emissions of CO ₂ and NMVOC from paint application in households, industry and construction | AD, EF: EMIS 2014/3A1, 3A2 and 3A3 |

5.2.2 Methodological Issues

Methodology

For determination of NMVOC emissions from paint application a country specific method based on the consumption of paint and its solvent content is used. Switzerland's Informative Inventory Report 2014 contains a description of the country-specific methods used for estimating the NMVOC emissions from the most important sources within source category 3A (FOEN 2014e, section 5.2.2).

The indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere are calculated using a carbon content fraction of 0.6 according to the 2006 IPCC Guidelines (IPCC 2006).

Also, several industrial plants use facilities and equipment to reduce NMVOC in exhaust gases and room ventilation output. Often, this implies the feeding of air with high NMVOC

content into the burning chamber of boilers or other facilities to incinerate NMVOC. This post-combustion of NMVOC leads to additional direct CO₂ emissions. They are estimated based on industry data and expert estimates.

Emission Factors

Emission factors for NMVOC emissions base on data from the Swiss association for coating and paint applications (VSLF) and from relevant retailers (source category 3A1 Paint Applications in Households), documented in the EMIS database (EMIS 2014/3A).

For paint application in construction, which is the most important NMVOC source in 3A Paint Application, the emission factor amounts to 59 kg NMVOC per ton of paint in 2012 (EMIS 2014/3A1 Farben-Anwendung Bau).

The emission factor for indirect CO₂ emissions from decomposition of NMVOC in the atmosphere is 2.2 Gg CO₂/Gg NMVOC (carbon content fraction * molecular weight of carbon dioxide / molecular weight of carbon).

Activity Data

Activity data corresponds to the annual consumption of paints. Data on paint consumption is taken from the Swiss association for coating and paint applications (VSLF) and from relevant retailers (source category 3A1 Paint Applications in Households), documented in the EMIS database (EMIS 2014/3A).

For paint application in construction, which is the most important NMVOC source in source category 3A Paint Application, the activity data equals the consumption of 54'000 t paint in 2012 (EMIS 2014/3A1 Farben-Anwendung Bau).

5.2.3 Uncertainties and Time-Series Consistency

The uncertainty of total CO₂ emissions from the entire source category Solvent and Other Product Use is estimated to be 50% (expert estimate).

Time series is consistent.

5.2.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

5.2.5 Source-Specific Recalculations

3A1 Paint applications in households: Activity data has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.

5.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

5.3 Source Category 3B – Degreasing and Dry Cleaning

5.3.1 Source Category Description

Source category 3B Degreasing and Dry Cleaning comprises NMVOC emissions from degreasing, dry cleaning and cleaning in electronic industry. Also, it includes indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere and direct CO₂ emissions resulting from post-combustion of NMVOC in exhaust gases.

Table 5-4 Specification of source category 3B Degreasing and Dry Cleaning. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| | Source | Specification | Data Source |
|----|-----------------------------|---|-------------------------------|
| 3B | Degreasing and Dry Cleaning | Emissions of CO ₂ and NMVOC from degreasing; dry cleaning; cleaning of electronic components; cleaning of parts in metal processing; other industrial cleaning | AD, EF: EMIS 2014/3B1 and 3B2 |

5.3.2 Methodological Issues

Methodology

For determination of NMVOC emissions from degreasing and dry cleaning a country specific method based on the consumption of solvents is used. Switzerland's Informative Inventory Report 2014 contains a description of the country-specific methods used for estimating the NMVOC emissions from the most important sources within source category 3B (FOEN 2014e, section 5.3.2).

The indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere are calculated using a carbon content fraction of 0.6 according to the 2006 IPCC Guidelines (IPCC 2006).

The direct CO₂ emissions resulting from post-combustion of NMVOC in exhaust gases is estimated based on industry data and expert estimates.

Emission Factors

Emission factors for NMVOC emissions are based on data from Swiss industry and expert estimates, documented in the EMIS database (EMIS 2014/3B).

For degreasing of metal, which is the most important NMVOC source in source category 3B Degreasing and Dry Cleaning, the emission factor amounts to 550 kg NMVOC per ton of solvent in 2012 (EMIS 2014/3B1 Metallreinigung).

The emission factor for indirect CO₂ emissions from decomposition of NMVOC in the atmosphere is 2.2 Gg CO₂/Gg NMVOC (carbon content fraction * molecular weight of carbon dioxide / molecular weight of carbon).

Activity Data

Activity data corresponds to the annual consumption of solvents for degreasing and dry cleaning. Data bases on industry data and expert estimates, documented in the EMIS database (EMIS 2014/3B).

For degreasing of metal, which is the most important NMVOC source in source category 3B Degreasing and Dry Cleaning, the activity data equals to 2'272 t solvent in 2012 (EMIS 2014/3B1 Metallreinigung).

5.3.3 Uncertainties and Time-Series Consistency

The uncertainty of total CO₂ emissions from the entire source category 3 Solvent and Other Product Use is estimated to be 50% (expert estimate).

The time series is consistent.

5.3.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

5.3.5 Source-Specific Recalculations

3B2 Dry cleaning: Activity data and EF value have been updated for 2012 resulting as well in revised interpolated values for 2007-2011 and 1991-2011, respectively.

5.3.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

5.4 Source Category 3C – Chemical Products, Manufacture and Processing

5.4.1 Source Category Description

Source category 3C Chemical Products, Manufacture and Processing comprises NMVOC emissions from manufacturing and processing chemical products. Also, it includes indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere and direct CO₂ emissions resulting from post-combustion of NMVOC in exhaust gases.

Table 5-5 Specification of source category 3C Chemical Products, Manufacture and Processing. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| | Source | Specification | Data Source |
|----|---|--|----------------------|
| 3C | Chemical Products, Manufacture and Processing | Emissions of CO ₂ and NMVOC from handling and storage of solvents; fine chemical production; production of pharmaceuticals; manufacturing of paint, inks, glues, adhesive tape, rubber; processing of PVC, polystyrene foam, polyurethane and polyester | AD, EF: EMIS 2014/3C |

5.4.2 Methodological Issues

Methodology

For determination of NMVOC emissions from chemical products, manufacture and processing a country specific method is used. The emissions from fine chemical and pharmaceutical production are based on production data and expert estimates. The emissions of handling and storage of solvents are calculated based on the imported quantities. The emissions from manufacturing paint, glues, inks, adhesive tape, rubber and polyurethane as well as the processing of PVC are calculated based on production data. The emissions from processing of polystyrene foam and polyester are calculated based on consumption data. Switzerland's Informative Inventory Report 2014 contains a description of the country-specific methods used for estimating the NMVOC emissions from the most important sources within source category 3C (FOEN 2014e, section 5.4.2).

The indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere are calculated using a carbon content fraction of 0.6 according to the 2006 IPCC Guidelines (IPCC 2006).

Direct CO₂ emissions result from post-combustion of NMVOC. Those emissions are estimated based on industry data and expert estimates.

Emission Factors

Emission factors for NMVOC emissions are based on data from Swiss industry, industry associations and expert estimates, documented in the EMIS database (EMIS 2014/3C). Emission factors for handling and storage of solvents are estimated according to the solvent vapour pressure.

For fine chemical production, which is the most important NMVOC source in source category 3C Chemical Products, Manufacture and Processing, the emission factor amounts to 3.7 ton NMVOC per production index in 2012 (EMIS 2014/3C Feinchemikalien-Produktion).

The emission factor for indirect CO₂ emissions from decomposition of NMVOC in the atmosphere is 2.2 Gg CO₂/Gg NMVOC (carbon content fraction * molecular weight of carbon dioxide / molecular weight of carbon).

Activity Data

Activity data corresponds to the annual consumption of solvents and bases on data from industry, industry associations and expert estimates, documented in the EMIS database (EMIS 2014/3C).

For fine chemical production, which is the most important NMVOC source in source category 3C Chemical Products, Manufacture and Processing, the activity data equals to a production

index of 303 in 2012 (EMIS 2014/3C Feinchemikalien-Produktion). For activity data the index of production according to the Swiss Federal Office of Statistics is used.

5.4.3 Uncertainties and Time-Series Consistency

The uncertainty of total CO₂ emissions from the entire source category 3 Solvent and Other Product Use is estimated to be 50% (expert estimate).

Time series is consistent.

5.4.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-tables
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

5.4.5 Source-Specific Recalculations

3C Paint and ink manufacturing: Activity data has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.

3C Manufacturing of rubber: Activity data has been updated for 2011 and EF value has been updated for 2012 resulting as well in revised interpolated values for 2008-2010 and 1998-2011, respectively.

3C Manufacturing of polyester: Activity data has been updated for 2010 and 2011 and EF values have been updated for 2010 and 2012 resulting as well in revised interpolated values for 2008-2009 and 2008, 2009 and 2011, respectively.

3C Manufacturing of polystyrene: Activity data has been updated for 2011 resulting in revised interpolated values for 2008-2010 as well.

3C Manufacturing of PVC: Activity data has been updated for 2010 and 2011 and EF values have been updated for 2004 and 2012 resulting as well in revised interpolated values for 2009 and 1991-2003 and 2005- 2011, respectively.

5.4.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

5.5 Source Category 3D – Other

5.5.1 Source Category Description

Source category 3D Other comprises emissions from the application of N₂O in households and hospitals as well as of NMVOC from many different solvent applications. Also, it includes indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere and direct CO₂ emissions resulting from post-combustion of NMVOC in exhaust gases.

Additional emissions of CO₂, NO_x, CO and SO₂ result from the use of fireworks.

Table 5-6 Specification of source category 3D Other. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| | Source | Specification | Data Source |
|-----|---|---|-----------------------|
| 3D1 | Use of N ₂ O for Anaesthesia | Emissions of N ₂ O from the use of N ₂ O in hospitals | AD, EF: EMIS 2014/3D1 |
| 3D3 | N ₂ O from Aerosol Cans | Emissions of N ₂ O from the use of aerosol cans | AD, EF: EMIS 2014/3D3 |
| 3D5 | Other | Emissions of CO ₂ and NMVOC from use of spray cans in industry and households; domestic solvent use; print industry; application of glues and adhesives; use of concrete additives; removal of paint and lacquer; car underbody sealant; de-icing of airplanes; tanning of leather; impregnating of glass and mineral wool; use of cooling and other lubricants; extraction of oils and fats; use of pesticides; use of pharmaceutical products in households; house cleaning industry/craft/services; hairdressers; scientific laboratories; textile production; paper and paper board production; clothing production; cosmetic institutions; production and use of tobacco products; vehicles dewaxing; wood preservation; medical practitioners; other health care institutions; not attributable solvent emissions Emissions of CO ₂ , NO _x , CO and SO ₂ from use of fireworks | AD, EF: EMIS 2014/3D5 |

5.5.2 Methodological Issues

Methodology

Emissions of N₂O from source category 3D1 occur from anaesthesia use in hospitals and in source category 3D3 from the use of aerosol cans in households. For both categories a country specific method based on the production/consumption of N₂O and of the different solvent applications is used.

The emissions from source category 3D5 Domestic solvent use, which is the most important NMVOC emission source in 3D5 Other, is calculated proportional to the number of inhabitants in Switzerland. Switzerland's Informative Inventory Report 2014 contains a description of the country-specific methods used for estimating the NMVOC emissions from the most important sources within source category 3D (FOEN 2014e, section 5.5.2).

The indirect CO₂ emissions due to decomposition of NMVOC in the atmosphere are calculated using a carbon content fraction of 0.6 according to the 2006 IPCC Guidelines (IPCC 2006).

Direct CO₂ emissions result from post-combustion of NMVOC. Those emissions are estimated based on industry data and expert estimates.

Emission Factors

For source category 3D1 Use of N₂O for anaesthesia the emission factor is calculated based on the amount of N₂O sold in Switzerland divided by the number of inhabitants. The yearly amount of N₂O sold for anaesthesia purpose is provided from sales information from the companies concerned and has been updated for the NIR submission 2013 based on annual data from 2005 onwards (EMIS 2014/3D1 Lachgasanwendung Spitler).

Source 3D3 N₂O from aerosol cans include N₂O emissions from whipped cream makers using gas capsules from private households and restaurants. The emission factor is calculated based on the amount of gas capsules sold in Switzerland divided by the number of inhabitants. The emission factor has been updated for the NIR submission 2013 based on sales figures and N₂O content of aerosol cans in 2011 (EMIS 2014/3D3 Lachgasanwendung Haushalt).

In Table 5-7 emission factors for the emission of N₂O is given for source categories 3D1 Use of N₂O for anaesthesia and 3D3 N₂O from aerosol cans.

Table 5-7 Emission factors for N₂O for source category 3D1 and 3D3 in g/inhabitant in 2012 (EMIS 2014/3D1 Lachgasanwendung Spitler; EMIS 2014/3D3 Lachgasanwendung Haushalt).

| 3D1 Use of N ₂ O for anaesthesia | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|--------------|------|------|------|------|------|------|------|------|------|------|
| N ₂ O | g/inhabitant | 43 | 40.4 | 37.7 | 35.1 | 32.5 | 29.8 | 27.2 | 24.5 | 21.9 | 19.3 |
| 3D3 N ₂ O from aerosol cans | | | | | | | | | | | |
| N ₂ O | g/inhabitant | 9.3 | 9.3 | 9.4 | 9.4 | 9.5 | 9.6 | 9.6 | 9.7 | 9.7 | 9.8 |

| 3D1 Use of N ₂ O for anaesthesia | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|--------------|------|------|------|------|------|------|------|------|------|------|
| N ₂ O | g/inhabitant | 16.6 | 14 | 13 | 13 | 12 | 12 | 10 | 9 | 8 | 7 |
| 3D3 N ₂ O from aerosol cans | | | | | | | | | | | |
| N ₂ O | g/inhabitant | 9.8 | 9.9 | 9.9 | 10 | 10.3 | 10.5 | 10.8 | 11.0 | 11.3 | 11.5 |

| 3D1 Use of N ₂ O for anaesthesia | Unit | 2010 | 2011 | 2012 |
|---|--------------|------|------|------|
| N ₂ O | g/inhabitant | 7 | 6 | 5.8 |
| 3D3 N ₂ O from aerosol cans | | | | |
| N ₂ O | g/inhabitant | 11.8 | 12 | 12.2 |

For source category 3D5 Other the emission factors for NMVOC as well as the emission factors for CO₂, NO_x, CO and SO₂ from the use of fireworks emissions are based on data from Swiss industry and expert estimates, documented in the EMIS database (EMIS 2014/3D). For house cleaning, which is the most important emission source in source category 3D5 Other, the emission factor for NMVOC amounts to 892 g per inhabitant in 2012 (EMIS 2014/3D5 Reinigungs- und Lsemittel, Haushalte).

The emission factor for indirect CO₂ emissions from decomposition of NMVOC in the atmosphere is 2.2 Gg CO₂/Gg NMVOC (carbon content fraction * molecular weight of carbon dioxide / molecular weight of carbon).

Activity Data

For source categories 3D1 Use of N₂O for anaesthesia and 3D3 N₂O from aerosol cans the activity data corresponds to the number of inhabitants in Switzerland and amounts to

7'997'000 in 2012 (EMIS 2014/3D1 Lachgasanwendung Spitäler and EMIS 2014/3D3 Lachgasanwendung Haushalt).

For source category 3D5 Other the activity data corresponds to the annual production/ consumption of solvents. Data bases on industry data and expert estimates, documented in the EMIS database (EMIS 2014/3D5). For house cleaning, which is the most important emission source in source category 3D Other, the activity data is the number of inhabitants in Switzerland and amounts to 7'997'000 in 2012 (EMIS 2014/3D5 Reinigungs- und Lösemittel, Haushalte).

5.5.3 Uncertainties and Time-Series Consistency

The uncertainty of total CO₂ emissions from the entire source category 3 Solvent and Other Product Use is estimated to be 50% (expert estimate).

The uncertainty of N₂O emissions is estimated to be 80% (expert estimate, see table Table 1-14).

Time series is consistent.

5.5.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF-table
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

5.5.5 Source-Specific Recalculations

3D5 Print industry: Activity data has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.

3D5 Impregnating of glass and mineral wool: Activity data of one of the two production plants has been revised for 1991-2004 based on effective production data for 1996-2004.

3D5 Production and use of tobacco products: The estimation of consumption of tobacco products has been improved and updated for the years 2005-2011.

3D5 Use of concrete additives: Activity data has been revised and updated for 1990, 1998, 2001 and 2008-2011, respectively. This has resulted in revised interpolated values in between. EF values have been corrected to interpolated values and updated for 1999-2006 and 2008-2011, respectively.

3D5 Car underbody sealant: Activity data has been revised and updated for 1990, 1998 and 2012, respectively, resulting in revised interpolated values in between. EF values have been corrected to interpolated values and updated for 1999-2003 and 2012, respectively, resulting in revised interpolated values for 2005-2011 as well.

3D5 De-icing of airplanes and other de-icing: Activity data and EF values have been updated for 2012 resulting in revised interpolated values for 2008-2011 as well.

3D5 Use of cooling and other lubricants: Activity data and EF values have been updated for 2008-2011 and 2012, respectively, resulting as well in revised interpolated values from 2005 onwards.

3D5 Use lubricants: Activity data has been updated for 2008-2011 resulting in revised interpolated values for 2005-2007 as well. EF values have been corrected to interpolated values and updated for 1999-2003 and 2012, respectively resulting in revised interpolated values for 2008-2011 as well.

5.5.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

6 Agriculture

6.1 Overview

This chapter provides information on the estimation of the greenhouse gas emissions from the sector Agriculture. The following source categories are reported:

- 4A Enteric Fermentation, CH₄ emissions from domestic livestock,
- 4B Manure Management, emissions of CH₄ and N₂O,
- 4D Agricultural Soils, emissions of N₂O, NO_x and NMVOC,

Categories 4C Rice Cultivation and 4E Burning of Savannahs are not occurring and therefore not reported in Switzerland.

Emissions from field burning of agricultural residues, formerly reported under Source Category 4F, have been moved to Source Category 6C (Waste Incineration, Chap. 8.4), in accordance with the EMEP Guidebook 2009 (EEA 2010).

Total greenhouse gas emissions from agriculture in 2012 were 5'539 Gg CO₂ equivalents in total which is a contribution of 10.9% to the total of Swiss greenhouse gas emissions. Main agricultural sources of greenhouse gases in 2012 were enteric fermentation emitting 2'497 Gg CO₂ equivalents (45% of all agricultural greenhouse gases), followed by agricultural soils with 2'060 Gg CO₂ equivalents (37%) and Manure Management with 982 Gg CO₂ equivalents (18%).




Figure 6-1 Greenhouse gas emissions of agriculture in Gg CO₂ equivalents 1990-2012.

Main greenhouse gases are CH₄ and N₂O. There are no CO₂ emissions reported in the agricultural sector. CO₂ emissions from soils are reported under Land Use, Land-use Change and Forestry. CO₂ emissions from energy use in agriculture are reported under 1A4c Energy; Others Sectors, Agriculture/Forestry/Fisheries.

Table 6-1 Greenhouse gas emissions in Gg CO₂ equivalents from agriculture 1990-2012.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| CH ₄ | 3'307 | 3'298 | 3'236 | 3'200 | 3'213 | 3'201 | 3'171 | 3'107 | 3'092 | 3'055 |
| N ₂ O | 2'785 | 2'772 | 2'743 | 2'678 | 2'630 | 2'618 | 2'609 | 2'499 | 2'486 | 2'456 |
| Sum | 6'092 | 6'069 | 5'979 | 5'877 | 5'843 | 5'819 | 5'780 | 5'606 | 5'578 | 5'511 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| CH ₄ | 3'047 | 3'080 | 3'069 | 3'042 | 3'031 | 3'060 | 3'088 | 3'118 | 3'203 | 3'175 |
| N ₂ O | 2'449 | 2'481 | 2'466 | 2'418 | 2'417 | 2'414 | 2'405 | 2'437 | 2'442 | 2'412 |
| Sum | 5'496 | 5'561 | 5'536 | 5'461 | 5'447 | 5'474 | 5'493 | 5'556 | 5'645 | 5'587 |

| Gas | 2010 | 2011 | 2012 |
|--------------------------|-------|-------|-------|
| CO ₂ eq. (Gg) | | | |
| CO ₂ | 0 | 0 | 0 |
| CH ₄ | 3'166 | 3'155 | 3'143 |
| N ₂ O | 2'470 | 2'417 | 2'395 |
| Sum | 5'637 | 5'572 | 5'539 |

CH₄ and N₂O emissions were declining from 1990 until 2004. This general trend can be explained by a reduction of the number of cattle and a reduced input of mineral fertilisers due to the introduction of the "Proof of Ecological Performance (PEP)" (ART 2013a, Leifeld and Fuhrer 2005). From 2004 to 2008 CH₄ emissions increased again due to higher livestock numbers (mainly cattle). Since 2008 total emissions seem to be fluctuating on a rather stable level possibly due to a more or less stable cattle population due to the suspension of the milk quotation. Most emission factors did not change significantly.




Figure 6-2 Trend of the greenhouse gases of the agricultural sector 1990-2012. The base year 1990 represents 100%.

Among the key categories of the Swiss inventory, six are from the agricultural sector:




Figure 6-3 Key sources (Tier 1 and Tier 2) in Agriculture, emissions 1990 and 2012 in CO₂ equivalents (Gg).

6.2 Source Category 4A – Enteric Fermentation

6.2.1 Source Category Description

Tier 1 Key Category 4A

CH₄ emissions from Enteric Fermentation (level)

Tier 2 Key category 4A

CH₄ emissions from Enteric Fermentation (level)

The emission source is the domestic livestock population broken down into 3 cattle categories (mature dairy cattle, mature non-dairy cattle, young cattle), sheep, goats, horses, mules and asses, swine and poultry. Emissions from enteric fermentation were declining from 1990 until 2004, mainly due to a reduction of the number of cattle. However, between 2004 and 2008 cattle livestock numbers and subsequently CH₄ emissions were increasing again, whereas since 2008 they are decreasing. Emissions from cattle contribute to over 90% of the emissions from enteric fermentation.

Table 6-2 Specification of source category 4A Enteric Fermentation. (AD: Activity data; EF: Emission factor).

| 4A | Source | Specification | Data Source |
|------------|---------------------------|---|---|
| 4A1 | Cattle | Mature dairy cattle | AD: Livestock data from SBV 2013, ART/SHL 2012, SFSO 2013d; Net energy and metabolisable energy (calves) from RAP 1999; EF: Soliva 2006 |
| | | Mature non-dairy cattle | |
| | | Young cattle (fattening calves, pre-weaned calves, breeding cattle 1st year (breeding calves + breeding cattle 4-12 months), breeding cattle > 1 year, fattening cattle (fattening calves 0-4 months, fattening cattle 4-12 months) | |
| 4A3 4A4 | Sheep Goats | | AD: Livestock data from SBV 2013 and ART/SHL 2012; net energy data from Giuliani 2013; EF: Soliva 2006 |
| 4A6 4A7 | Horses Mules and asses | | AD: Livestock data from SBV 2013 and ART/SHL 2012; digestible energy data from Stricker 2012; EF: Soliva 2006 |
| 4A8 | Swine | | AD: Livestock data from SBV 2013 and ART/SHL 2012; net energy data from Giuliani 2013; EF: Soliva 2006 |
| 4A9 | Poultry | | AD: Livestock data from SBV 2013 and ART/SHL 2012; net energy data from Giuliani 2013; EF: Hadorn and Wenk 1996 cited in Soliva 2006 |

6.2.2 Methodological Issues

Methodology

The calculation is based on methods described in the IPCC Good Practice Guidance (IPCC 2000, equation 4.14). CH₄ emissions from enteric fermentation of the livestock population have been estimated using Tier 2 methodology. This means that detailed country specific data on nutrient requirements, feed intake and CH₄ conversion rates for specific feed types are required.

For calculating the gross energy intake, a country specific method based on available data on requirements of net energy (lactation, growth), digestible energy and metabolisable energy has been applied. Data on energy intake is based on RAP (1999) and SBV (2013) as well as on Stricker 2012. The method is described in detail in Soliva (2006) and is realised in Agroscope (2014).

Different energy levels (Figure 6-4) are used to express the energy conversion from energy required for maintenance and performance to gross energy intake.




Figure 6-4 Levels of feed energy conversion. Reference: Soliva 2006.

Net energy (NE) is used to express the energy required by the ruminants such as cattle, sheep and goats. NE in cattle feeding is further sub-divided into NE for lactation (NEL) and NE for growth (NEV). For some of the young cattle categories NEL is used rather than NEV what would seem natural. However, cattle raising is often coupled with dairy cattle activities and therefore the same energy unit (NEL) is used in these cases (RAP 1999). Exceptions are the fattening calves (milk-fed calves), whose requirements for energy are expressed as metabolisable energy (ME). Horses, mules, asses and swine are fed on the basis of digestible energy (DE), whereas poultry are fed according to metabolisable energy (ME).

For the cattle categories detailed estimations for NE requirements are necessary. As the Swiss Farmers Union (SBV) does not calculate the NE for detailed cattle sub-categories, NE data for each cattle source category was calculated individually according to the animal's requirements following the feeding recommendations of RAP (1999). These RAP recommendations are also used by the Swiss farmers as basis for their cattle feeding regime and for filling in application forms for subsidies for ecological services, and are therefore highly appropriate. In the calculation of the NE data, the animal's weight, daily growth rate, daily feed intake (dry matter), daily feed energy intake, and energy required for milk production and pregnancy for the respective sub-categories were considered (Soliva 2006).

For estimating the gross energy intake out of the available data on net energy, metabolisable energy and digestible energy, the following conversion factors were applied:

Table 6-3 Conversion factors used for calculation of energy requirements of individual livestock categories.
Reference: Soliva 2006: p.3. GE: Gross energy; DE: Digestible Energy; ME: Metabolisable Energy;
NEL: Net energy for lactation; NEV: Net energy for growth.

| Livestock Category | | Conversion Factors | |
|-------------------------|--------------------------------|--------------------|-------|
| Mature Dairy Cattle | | NEL to GE | 0.318 |
| Mature Non-Dairy Cattle | | NEL to GE | 0.275 |
| Young Cattle | Fattening Calves | ME to GE | 0.930 |
| | Pre-Weaned Calves | NEL to GE | 0.291 |
| | Breeding Calves | NEL to GE | 0.341 |
| | Breeding Cattle (4-12 months) | NEL to GE | 0.322 |
| | Breeding Cattle (> 1 year) | NEL to GE | 0.313 |
| | Fattening Calves (0-4 months) | NEV to GE | 0.350 |
| | Fattening Cattle (4-12 months) | NEV to GE | 0.401 |
| Milkshoop | | NEL to GE | 0.287 |
| Fattening Sheep | | NEV to GE | 0.350 |
| Goats | | NEL to GE | 0.283 |
| Horses | | DE to GE | 0.700 |
| Mules and Asses | | DE to GE | 0.700 |
| Swine | | DE to GE | 0.682 |
| Poultry | | ME to GE | 0.700 |

Emission factors

All emission factors for enteric fermentation are country specific, based on IPCC equation 4.14 IPCC 2000: p. 4.26.

$$EF = \frac{GE * Y_m * 365 \text{ days} / y}{55.65 \text{ MJ} / \text{kg} CH_4}$$

GE = Gross energy intake (MJ/head/day)

Y_m = Methane conversion rate, which is the fraction of gross energy in feed converted to methane

55.65 MJ/kg = energy content of methane.

The following input data are used:

Table 6-4 Gross energy intake per head of different livestock groups. Calculation is based on the above mentioned parameters net energy, digestible energy, metabolisable energy according to the method described in Soliva (2006). Input data on net energy, digestible energy and metabolisable energy is taken from Giuliani (2013), RAP (1999) and Stricker (2012). All sub-categories displayed in italic.

| Gross Energy Intake | | 1990-1999 | | | | | | | | | |
|---------------------------------|--------------------------------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | | MJ/head/day | | | | | | | | | |
| Mature Dairy Cattle | | 258.0 | 260.3 | 260.6 | 263.9 | 263.6 | 266.4 | 265.4 | 269.6 | 273.7 | 277.2 |
| Mature Non-Dairy Cattle | | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 |
| Young Cattle Average (weighted) | | 93.6 | 93.5 | 93.6 | 93.4 | 94.0 | 94.3 | 93.7 | 94.1 | 93.1 | 92.1 |
| | Fattening Calves | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 |
| | Pre-Weaned Calves | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 |
| | Breeding Calves | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 |
| | Breeding Cattle (4-12 months) | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 |
| | Breeding Cattle (> 1 year) | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 |
| | Fattening Calves (0-4 months) | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 |
| | Fattening Cattle (4-12 months) | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 |
| Sheep | | 21.2 | 21.7 | 22.2 | 22.4 | 23.8 | 24.0 | 22.0 | 22.1 | 21.5 | 22.9 |
| Goats | | 25.0 | 24.6 | 25.0 | 25.4 | 25.5 | 27.9 | 25.3 | 25.6 | 26.9 | 25.8 |
| Horses | | 107.3 | 107.3 | 107.3 | 107.3 | 107.1 | 106.9 | 107.1 | 107.3 | 107.3 | 107.2 |
| Mules and Asses | | 39.2 | 39.2 | 39.2 | 39.2 | 39.5 | 39.7 | 39.7 | 39.8 | 39.6 | 39.8 |
| Swine | | 28.3 | 28.9 | 29.0 | 29.1 | 28.5 | 31.9 | 29.8 | 29.9 | 27.9 | 29.0 |
| Poultry ¹⁾ | | 1.5 | 1.5 | 1.6 | 1.3 | 1.4 | 1.3 | 1.4 | 1.4 | 1.3 | 1.4 |

| Gross Energy Intake | | 2000-2009 | | | | | | | | | |
|---------------------------------|--------------------------------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| | | MJ/head/day | | | | | | | | | |
| Mature Dairy Cattle | | 280.1 | 282.0 | 285.0 | 288.6 | 294.0 | 294.1 | 295.1 | 300.5 | 304.1 | 309.2 |
| Mature Non-Dairy Cattle | | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 | 205.1 |
| Young Cattle Average (weighted) | | 93.4 | 92.3 | 91.9 | 91.6 | 91.3 | 90.8 | 90.9 | 90.7 | 90.8 | 90.4 |
| | Fattening Calves | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 | 47.6 |
| | Pre-Weaned Calves | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 | 55.7 |
| | Breeding Calves | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 | 26.9 |
| | Breeding Cattle (4-12 months) | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 | 89.2 |
| | Breeding Cattle (> 1 year) | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 | 129.1 |
| | Fattening Calves (0-4 months) | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 | 55.6 |
| | Fattening Cattle (4-12 months) | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 | 124.6 |
| Sheep | | 22.4 | 22.9 | 22.8 | 22.7 | 23.3 | 22.8 | 22.6 | 22.2 | 22.0 | 22.7 |
| Goats | | 25.7 | 26.0 | 25.2 | 25.4 | 25.2 | 25.4 | 25.3 | 25.0 | 25.0 | 25.3 |
| Horses | | 107.4 | 107.5 | 107.6 | 107.6 | 107.6 | 107.7 | 107.7 | 107.7 | 107.7 | 107.8 |
| Mules and Asses | | 39.5 | 39.6 | 39.6 | 39.6 | 39.5 | 39.4 | 39.5 | 39.3 | 39.2 | 40.0 |
| Swine | | 28.0 | 27.7 | 27.1 | 27.0 | 27.2 | 26.6 | 26.3 | 26.9 | 26.7 | 27.0 |
| Poultry ¹⁾ | | 1.4 | 1.4 | 1.3 | 1.3 | 1.3 | 1.2 | 1.2 | 1.3 | 1.3 | 1.3 |

| Gross Energy Intake | | 2010-2012 | | |
|---------------------------------|---------------------------------------|-----------|-------|-------|
| | | 2010 | 2011 | 2012 |
| | | MJ/h./d. | | |
| Mature Dairy Cattle | | 310.8 | 312.1 | 311.5 |
| Mature Non-Dairy Cattle | | 205.1 | 205.1 | 205.1 |
| Young Cattle Average (weighted) | | 90.4 | 90.1 | 89.8 |
| | <i>Fattening Calves</i> | 47.6 | 47.6 | 47.6 |
| | <i>Pre-Weaned Calves</i> | 55.7 | 55.7 | 55.7 |
| | <i>Breeding Calves</i> | 26.9 | 26.9 | 26.9 |
| | <i>Breeding Cattle (4-12 months)</i> | 89.2 | 89.2 | 89.2 |
| | <i>Breeding Cattle (> 1 year)</i> | 129.1 | 129.1 | 129.1 |
| | <i>Fattening Calves (0-4 months)</i> | 55.6 | 55.6 | 55.6 |
| | <i>Fattening Cattle (4-12 months)</i> | 124.6 | 124.6 | 124.6 |
| Sheep | | 22.6 | 22.6 | 22.6 |
| Goats | | 25.1 | 25.6 | 25.6 |
| Horses | | 107.9 | 107.9 | 107.9 |
| Mules and Asses | | 40.2 | 39.9 | 39.9 |
| Swine | | 27.2 | 26.9 | 26.9 |
| Poultry ¹⁾ | | 1.3 | 1.3 | 1.3 |

¹⁾ Poultry data is not Gross Energy intake (GE) but Metabolizable Energy intake (ME)

The **gross energy intake** per head for some animal categories revealed some fluctuations during the inventory period. The value for mature dairy cattle increased which is mainly a result of higher milk production (Table 6-5). Milk production of mature dairy cattle increased from 4'900 kg per head and year in 1990 to 6'879 kg per head and year in 2012. Statistics of

annual milk production are provided by the Swiss Farmers Union (SBV 2013). Milk production includes marketed milk, milk consumed by calves on farms and milk sold outside the commercial industry (MISTA 2013). It should be noted that daily milk yield refers to milk production during lactation (305 days) and not during the whole year (365 days). Accordingly, energy requirement for lactation is excluded from the two remaining months when the cows are dry.

Table 6-5: Annual milk production in Switzerland

| Milk Production Cattle | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|-------------------------------------|-------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Population Size Mature Dairy Cattle | head | 783'100 | 780'500 | 763'500 | 744'450 | 749'700 | 739'641 | 736'043 | 711'613 | 701'343 | 683'545 |
| Lactation Period | day | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 |
| Milk Yield Mature Dairy Cattle | kg/head/day | 16.06 | 16.35 | 16.39 | 16.78 | 16.75 | 17.09 | 16.96 | 17.48 | 17.97 | 18.40 |
| Milk Yield Mature Non-Dairy Cattle | kg/head/day | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 |

| Milk Production Cattle | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|-------------------------------------|-------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Population Size Mature Dairy Cattle | head | 669'410 | 669'410 | 657'924 | 638'288 | 621'008 | 620'708 | 618'065 | 614'795 | 628'516 | 599'361 |
| Lactation Period | day | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 | 305 |
| Milk Yield Mature Dairy Cattle | kg/head/day | 18.75 | 18.97 | 19.34 | 19.77 | 20.43 | 20.45 | 20.57 | 21.21 | 21.66 | 22.27 |
| Milk Yield Mature Non-Dairy Cattle | kg/head/day | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 | 8.20 |

| Milk Production Cattle | | 2010 | 2011 | 2012 |
|-------------------------------------|-------------|---------|---------|---------|
| Population Size Mature Dairy Cattle | head | 589'024 | 589'239 | 591'212 |
| Lactation Period | day | 305 | 305 | 305 |
| Milk Yield Mature Dairy Cattle | kg/head/day | 22.46 | 22.63 | 22.55 |
| Milk Yield Mature Non-Dairy Cattle | kg/head/day | 8.20 | 8.20 | 8.20 |

The gross energy intake for mature non-dairy cattle is significantly higher than IPCC default values, since this category only comprehends mature cows to produce offspring for meat (so called suckler cows or mother cows). Milk production of mature non-dairy cattle is 2500kg per head and year (305 days of lactation) and does not change over the inventory time period (RAP 1999).

The gross energy intake of young cattle was calculated separately for all sub-categories displayed in Table 6-4 (in italics) and subsequently averaged (weighted average). The values for all the 7 sub-categories summarized under young cattle are constant over time. Since the composition of the young cattle category is changing over time (e.g. more pre-weaned calves, less fattening calves, see Table 6-6) the average gross energy intake for young cattle is also slightly changing. To calculate an annual emission factor, the categories breeding calves and breeding cattle 4-12 months are combined in the category breeding cattle 1st year (not shown in Table 6-4 and Table 6-6). Subsequently the respective animals have two separate gross energy intake values, i.e. 26.9 MJ/head/day for the first 4 month and 89.2 MJ/head/day for the later 8 months. The same procedure is applied for fattening calves 0-4 months and fattening cattle 4-12 months summing up to the category fattening cattle.

For the **methane conversion rate** Y_m (%) only few country specific data exist. Therefore mainly default values recommended by the IPCC for developed countries in Western Europe were used (IPCC 1997b: Reference Manual: p. 4.32–4.35 and IPCC 2000: p. 4.27). For all juveniles consuming only milk (i.e. fattening calves) the CH_4 conversion rate is assumed to be zero (IPCC 2000). For poultry a country specific value ($Y_{poultry} = 0.1631\%$ of metabolisable energy) was used since no default value is given by the IPCC. This value was evaluated in an in vivo trial with broilers (Hadorn and Wenk 1996).

Activity data

The activity data input has been obtained from statistics published by the Swiss Farmers Union (SBV 2013) and the Swiss Federal Statistical Office (SFSO 2013d). All activity data has been revised and harmonized during a joint effort of the Agroscope Reckenholz Tänikon

Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012).

The following data were used:

Table 6-6 Activity data for calculating methane emissions from enteric fermentation (ART/SHL 2012, SBV 2013, SFSO 2013d).

| Population Size | | 1990-1999 | | | | | | | | | |
|-------------------------|--------------------------------|------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | | 1'000 head | | | | | | | | | |
| Total cattle | | 1'855 | 1'829 | 1'783 | 1'745 | 1'755 | 1'748 | 1'747 | 1'673 | 1'641 | 1'609 |
| Mature Dairy Cattle | | 783 | 781 | 764 | 744 | 750 | 740 | 736 | 712 | 701 | 684 |
| Mature Non-Dairy Cattle | | 12 | 14 | 17 | 18 | 20 | 23 | 28 | 32 | 36 | 41 |
| Young Cattle | | 1'060 | 1'034 | 1'002 | 983 | 986 | 986 | 983 | 929 | 904 | 884 |
| | Fattening Calves | 112 | 111 | 110 | 111 | 101 | 102 | 112 | 106 | 108 | 116 |
| | Pre-Weaned Calves | 10 | 11 | 14 | 14 | 16 | 18 | 22 | 26 | 29 | 33 |
| | Breeding Calves | 214 | 204 | 197 | 184 | 182 | 166 | 155 | 139 | 136 | 72 |
| | Breeding Cattle (4-12 months) | 132 | 133 | 127 | 125 | 124 | 129 | 131 | 121 | 118 | 147 |
| | Breeding Cattle (> 1 year) | 404 | 400 | 397 | 381 | 379 | 378 | 383 | 372 | 350 | 305 |
| | Fattening Calves (0-4 months) | 88 | 79 | 71 | 76 | 83 | 82 | 75 | 68 | 66 | 48 |
| | Fattening Cattle (4-12 months) | 100 | 96 | 87 | 92 | 101 | 110 | 105 | 97 | 97 | 162 |
| Sheep | | 395 | 409 | 415 | 424 | 405 | 387 | 419 | 420 | 422 | 424 |
| Goats | | 68 | 65 | 58 | 57 | 55 | 53 | 57 | 58 | 60 | 62 |
| Horses | | 51 | 52 | 53 | 54 | 58 | 62 | 62 | 65 | 64 | 65 |
| Mules and Asses | | 11 | 11 | 11 | 11 | 11 | 11 | 12 | 13 | 14 | 15 |
| Swine | | 1'787 | 1'723 | 1'706 | 1'692 | 1'569 | 1'446 | 1'379 | 1'395 | 1'487 | 1'453 |
| Poultry | | 5'938 | 5'647 | 5'502 | 6'410 | 6'330 | 6'251 | 6'440 | 6'553 | 6'740 | 6'908 |

| Population Size | | 2000-2009 | | | | | | | | | |
|-------------------------|--------------------------------|------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| | | 1'000 head | | | | | | | | | |
| Total cattle | | 1'588 | 1'611 | 1'594 | 1'570 | 1'545 | 1'555 | 1'567 | 1'572 | 1'604 | 1'597 |
| Mature Dairy Cattle | | 669 | 669 | 658 | 638 | 621 | 621 | 618 | 615 | 629 | 599 |
| Mature Non-Dairy Cattle | | 45 | 51 | 58 | 65 | 70 | 78 | 87 | 94 | 98 | 108 |
| Young Cattle | | 874 | 891 | 878 | 867 | 854 | 856 | 862 | 863 | 877 | 890 |
| | Fattening Calves | 103 | 115 | 114 | 114 | 111 | 106 | 101 | 100 | 95 | 101 |
| | Pre-Weaned Calves | 36 | 40 | 47 | 52 | 57 | 62 | 67 | 72 | 76 | 86 |
| | Breeding Calves | 76 | 78 | 76 | 73 | 71 | 75 | 77 | 76 | 80 | 77 |
| | Breeding Cattle (4-12 months) | 161 | 160 | 154 | 147 | 143 | 147 | 147 | 147 | 152 | 149 |
| | Breeding Cattle (> 1 year) | 352 | 350 | 345 | 337 | 326 | 318 | 320 | 320 | 322 | 331 |
| | Fattening Calves (0-4 months) | 43 | 40 | 38 | 39 | 36 | 35 | 35 | 34 | 36 | 35 |
| | Fattening Cattle (4-12 months) | 105 | 109 | 104 | 105 | 109 | 112 | 114 | 114 | 116 | 112 |
| Sheep | | 421 | 420 | 430 | 445 | 441 | 446 | 448 | 444 | 446 | 432 |
| Goats | | 62 | 63 | 66 | 67 | 71 | 74 | 76 | 79 | 81 | 81 |
| Horses | | 66 | 64 | 64 | 65 | 65 | 65 | 66 | 67 | 68 | 69 |
| Mules and Asses | | 16 | 16 | 17 | 17 | 18 | 19 | 19 | 20 | 20 | 22 |
| Swine | | 1'498 | 1'548 | 1'557 | 1'529 | 1'538 | 1'609 | 1'635 | 1'573 | 1'540 | 1'557 |
| Poultry | | 6'983 | 6'939 | 7'339 | 7'587 | 8'061 | 8'260 | 7'670 | 8'228 | 8'543 | 8'809 |

| Population Size | | 2010-2012 | | |
|-------------------------|--------------------------------|------------|-------|-------|
| | | 2010 | 2011 | 2012 |
| | | 1'000 head | | |
| Total cattle | | 1'591 | 1'577 | 1'565 |
| Mature Dairy Cattle | | 589 | 589 | 591 |
| Mature Non-Dairy Cattle | | 111 | 111 | 114 |
| Young Cattle | | 891 | 877 | 859 |
| | Fattening Calves | 99 | 101 | 99 |
| | Pre-Weaned Calves | 88 | 88 | 91 |
| | Breeding Calves | 77 | 75 | 73 |
| | Breeding Cattle (4-12 months) | 149 | 145 | 140 |
| | Breeding Cattle (> 1 year) | 332 | 324 | 311 |
| | Fattening Calves (0-4 months) | 34 | 34 | 34 |
| | Fattening Cattle (4-12 months) | 111 | 111 | 112 |
| Sheep | | 434 | 424 | 417 |
| Goats | | 83 | 83 | 85 |
| Horses | | 71 | 66 | 67 |
| Mules and Asses | | 23 | 22 | 23 |
| Swine | | 1'589 | 1'579 | 1'544 |
| Poultry | | 9'025 | 9'478 | 9'955 |




Figure 6-5 Relative development of main animal categories 1990-2012. The category with the strongest increase, mature non-dairy cattle, is not displayed, as it increases to over 950% of the 1990 value by 2012.

Emission estimation for cattle has been conducted at a more disaggregated level than the one displayed in the CRF. The category Mature non-dairy cattle only includes mature cows used to produce offspring for meat. The CRF livestock category Young cattle includes the sub-categories fattening calves, pre-weaned calves, breeding calves, breeding cattle 4-12 months, breeding cattle > 1 year, fattening calves 0-4 months and fattening cattle 4-12 months. Although not young cattle in the proper sense, bulls are contained in the categories Breeding Cattle (> 1 year) and Fattening Cattle (4-12 months) according to their purposes. This regrouping of the cattle category enhances the consistency and transparency of the emission estimation procedure from livestock activities (also refer to chapter 6.3).

The number of cattle was slightly declining until the year 2004, which is a result of an on-going process to a less intensive form of animal husbandry due to ecological and economic reasons. However, cattle livestock numbers were slightly increasing again between 2004 and 2008 mainly due to an increase of the number of young cattle. Since 2008 the cattle population is more or less stable possibly due to the suspension of the milk quotation.

After a decrease until 1996 the number of swine was increasing again until 2006 – a process that could be observed also in many other European countries (SBV 2004: p.69). Since then the number of swine has been fluctuating slightly below the level of 2006. The number of poultry shows a rapid increase between 1990 and 2012 with only a distinct dip between 2005 and 2006, a consequence of changed human consumption patterns as a result of the avian flu in 2006.

The number of sheep has been more or less constant while the number of goats is increasing after a decline between 1990 and 1995.

6.2.3 Uncertainties and Time-Series Consistency

For the uncertainty analysis the input data from ART (2008a) was used and was weighted with current activity and emission data. The arithmetic mean of the lower and upper bound uncertainty is used for activity data (6.4%) and for emission factors (17.2%), resulting in a combined uncertainty of 18.4% for Tier 1 analysis. Tier 2 analysis results in a slightly

different and asymmetric result: The uncertainty interval lies between -18.1% and +18.5% corresponding to a mean uncertainty of 18.3%. For further results see Section 1.7.

The time series 1990–2012 is generally consistent, with two issues that should be considered:

- Between 1998 and 1999 the questionnaire for the collection of livestock data was modified. In some animal categories this led to minor ruptures in the time series. Consequences for overall emissions are, however, of minor importance. While the average absolute trend for the years 1990–2011 over all animal categories excluding mature non-dairy cattle was 3.3%, the average absolute trend for the years 1998–1999 was 3.8% (ART/SHL 2012).
- For the last four inventory years cattle population statistics were not available in the usual format. Data for 2009 to 2012 is based on the animal traffic database. Aggregation has been adapted to the format necessary for the AGRAMMON and greenhouse gas inventories by the Swiss College of Agriculture SHL (SHL 2010). Data in the animal traffic database is considered more complete than the data from the survey of the SFSO because it includes also animals held outside agricultural enterprises.

6.2.4 Source-Specific QA/QC and Verification

All QA/QC activities are further described in a separate document (ART 2013a). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed. Furthermore, comparisons with data from other countries have been conducted and discussed where possible.

The documentation about the data set and calculation method assures transparency and traceability of the calculation methods (Soliva 2006). Additionally a document in German lists all the methodological differences between the former calculations and the current methodology (Soliva 2006a).

Livestock data was compared with the livestock data provided by the FAO and checked for plausibility. In all cases the new recalculated data according to ART/SHL (2012) is considered more reliable than the FAO data. Small inconsistencies (usually in the order of $\pm 2\%$) are due to updates of provisional data that are not considered by the FAO. For horses, mules and asses disagreements are due to the different accounting of agricultural and non-agricultural horses. The Swiss inventory systems accounts for all animals no matter whether they are held on agricultural or non-agricultural enterprises. Moreover, the numbers of mules and asses is higher in the Swiss GHG-Inventory because unlike the FAO, Switzerland accounts also for ponies and lesser horses. The total number for poultry also shows some minor discrepancies due to different accounting of turkeys, geese, ducks and quails. Seasonal fluctuation of the cattle population has been analysed for the years 2005–2007 based on detailed information from the Swiss Farmers Union (SBV 2007a). Fluctuations are usually in the order of $\pm 3\%$ with census data (April) always slightly above the annual mean.

Total NE-intake of the cattle population as calculated in the Swiss GHG- Inventory is in accordance with an independent calculation of the Swiss farmers union (SBV 2007). In a check during the submission 2010 the average absolute difference for the time period 1990–2004 was $\pm 1.2\%$.

IPCC tables with data for estimating emission factors for cattle (such as weight, weight gain, milk production) were filled in, checked for consistency and confidence and compared with IPCC default values (refer to Table A - 27 in Annex A3.3). Methane conversion rates (Y_m) and feed digestibilities were compared to literature values representative for Swiss conditions.

The emission factors of category 4A were compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available for submission 2013 (INFRAS 2012). Furthermore, emission factors have been calculated according to the original IPCC Tier 2 method. Implied emission factors for enteric fermentation for mature dairy and non-dairy cattle in Switzerland are generally higher than IPCC Tier 1 default due to relatively high gross energy intake (ART 2013a). This can be explained by the high performance of animal livestock in Switzerland (weight, weight gain, milk production). However, the IPCC Tier 2 analysis yields even higher energy intake levels and hence emission factors for all cattle animals. High feed quality together with high genetic standard i.e. high energy use efficiency of Swiss cattle might be a reason for these differences (ART 2013a). In general a straightforward comparison is difficult due to the country specific feeding regime and the inconsistent categorization of immature cattle.

During the years 2009-2012 the group of animal nutrition from the Swiss Federal Institute of Technology Zürich investigated the effect of different feeding and management strategies on methane and nitrous oxide emissions from enteric fermentation and manure management of cattle held under typical Swiss management conditions (Kreuzer 2012). Measured values of various parameters such as Y_m or MCF were compared to IPCC default values and values in the Swiss greenhouse gas inventory. Preliminary analysis suggests that overall emissions are neither over- nor underestimated (Zeitz et al. 2012). Further investigations have to show to what extent the preliminary estimates will be confirmed to provide a basis for implementation in Switzerland's GHG inventory after 2014.

During the past years a couple of studies have been conducted to verify methane emissions at regional scale comparing bottom up estimates with atmospherical measurements. While Hiller et al. (2014a) found that methane emissions could be underestimated by the inventory method, Stieger (2014) reported a very good accordance of bottom up estimates and flux measurements. Generally the methodological approaches of atmospherical measurements as conducted by Hiller et al. (2014a) still rely on a number of rather uncertain basic assumptions and are therefore not beyond doubts. Furthermore, it has been stated, that the differences between bottom up and top down estimates are possibly due to the limitations of the geographical emission allocation within the spatial explicit inventory (Hiller et al. 2014).

The time series of activity data and emission factors have been compared between the current and the previous submission. All activity data, implied emission factors and emissions estimates undergo the following triple check:

- the results for 2012 are compared with the results for 2011 within the current CRF,
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the submission 2013,
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission the 2013.

Additionally an independent quality control was conducted by INFRAS by a countercheck of the data and calculation sheets elaborated by Agroscope (Agroscope 2014).

6.2.5 Source-Specific Recalculations

New more precise activity data have been used for the years 1994 and 2006. Previously only rounded values have been available. The effects on overall emissions is negligible.

Preliminary estimates for energy requirements for non cattle populations for the years 2010 and 2011 have been revised. A new dataset has been received from the Swiss Farmers Union (Giuliani 2013). The estimates are based on the same method as earlier energy requirement statistics published until 2007 by the Swiss Farmers Union. The effect of the recalculation on overall greenhouse gas emissions is considered negligible.

Milk yield of mature dairy cattle in the year 2011 has been slightly revised due to an update of the provisional number from the Swiss Farmers Union.

6.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented. Other projects are eventually postponed in order to give first priority to changes related to the new reporting guidelines.

6.3 Source Category 4B – Manure Management

6.3.1 Source Category Description

| |
|---|
| Tier 1 and Tier2 Key categories 4B |
|---|

| |
|--|
| CH ₄ emissions from Manure Management (level) |
|--|

| |
|---|
| N ₂ O emissions from Manure Management (level and trend) |
|---|

CH₄ and N₂O emissions from manure management are reported. The total emissions from manure management closely follow the development of the cattle population. Emissions declined from 1990 until 2004, increased again until 2008 and remained more or less stable since then.

Table 6-7 Specification of source category 4B Manure Management (CH₄). (AD: Activity data; EF: Emission factor).

| 4B | Source | Specification | Data Source |
|------------|---------------------------|-------------------------|--|
| 4B1 | Cattle | Mature dairy cattle | AD: SBV 2013, ART/SHL 2012, SFSO 2013d; EF: RAP 1999, IPCC 2000, IPCC 1997c, Soliva 2006, Kupper et al. 2013 |
| | | Mature non-dairy cattle | |
| | | Young cattle | |
| 4B3 4B4 | Sheep Goats | | AD: SBV 2013, ART/SHL 2012, SFSO 2013d; EF: IPCC 2000, IPCC 1997c, Flisch et al. 2009, Kupper et al. 2013, Giuliani 2013, Soliva 2006 |
| 4B6 4B7 | Horses Mules and Asses | | AD: SBV 2013, ART/SHL 2012, SFSO 2013d; EF: IPCC 2000, IPCC 1997c, Flisch et al. 2009, Kupper et al. 2013, Stricker 2012, Soliva 2006 |
| 4B8 | Swine | | AD: SBV 2013, ART/SHL 2012, SFSO 2013d; EF: IPCC 2000, IPCC 1997c, Flisch et al. 2009, Kupper et al. 2013, Giuliani 2013, Soliva 2006 |
| 4B9 | Poultry | | AD: SBV 2013, ART/SHL 2012, SFSO 2013d; EF: IPCC 2000, IPCC 1997c, Flisch et al. 2009, Kupper et al. 2013, Giuliani 2013, Soliva 2006 |

Table 6-8 Specification of source category 4B Manure Management (N₂O). (AD: Activity data; EF: Emission factor).

| 4B | Source | Specification | Data Source |
|--------------|---|---------------------|--|
| 4B11 4B12 | Liquid systems Solid storage and dry lot | Mature dairy cattle | AD: SBV 2013, ART/SHL 2012, SFSO 2013d, Flisch et al. 2009, Kupper et al. 2013; EF: IPCC 1997c, IPCC 2000 |

6.3.2 Methodological Issues

For calculation of CH₄ and N₂O emissions slightly different livestock sub-categories are used. The livestock categories reported in the CRF-tables are the same, but the respective sub-categories as a basis for the calculation are different. Nevertheless, there is no inconsistency in the total number of animals as they are the same both for CH₄ and N₂O emissions. The calculation of CH₄ and N₂O emissions is realised in Agroscope (2014).

Calculation of CH₄ emissions is based on the domestic livestock populations mature dairy cattle, mature non-dairy cattle, young cattle (fattening calves, pre-weaned calves, breeding calves, breeding cattle 4-12 months, breeding cattle > 1 year, fattening calves 0-4 months, fattening cattle 4-12 months), sheep, goats, horses, mules and asses, swine and poultry as reported for enteric fermentation.

Calculation of N₂O emissions are based on a different livestock population break down:

- Cattle: Mature dairy cattle, mature non-dairy cattle and young cattle (fattening calves, pre-weaned calves, breeding cattle 1st year, breeding cattle 2nd year, breeding cattle 3rd year, fattening cattle). Although not young cattle in the proper sense, bulls are contained in the categories Breeding Cattle 3rd Year and Fattening Cattle according to their purposes.
- Sheep: fattening sheep, milk sheep
- Goats: goat places
- Horses: horses < 3 years, horses > 3 years
- Mules and asses: mules, asses
- Swine: piglets, fattening pig over 25 kg, dry sows, nursing sows, boars
- Poultry: growers, layers, broilers, turkey, other poultry (geese, ducks, ostriches, quails)

This calculation is chosen because more detailed data on parameters such as N excretion or manure management system distribution for the particular animal categories are available (Flisch et al. 2009, Kupper et al. 2013). The nitrogen excretion rates are given on a yearly basis, considering replacement of animals (young cattle, swine and poultry) and including excretions from corresponding offspring and other associated animals (sheep, goats, swine) (ART/SHL 2012).

a) CH₄ Emissions

Methodology

Calculation of CH₄ emissions from manure management is based on IPCC Tier 2 (IPCC 2000: equation 4.17).

$$EF_i = VS_i \cdot 365 \text{ days / year} \cdot Bo_i \cdot 0.67 \text{ kg / m}^3 \cdot \sum_{ijk} MCF_{jk} \cdot MS_{ijk}$$

EF_i : annual emission factor for livestock population i

VS_i : daily volatile solids (VS) excreted for an animal within population i

Bo_i : maximum CH₄ producing capacity for manure produced by an animal within population i

MCF_{jk} : CH₄ conversion factors for each manure management system j by climate region k

MS_{ijk} : fraction of animal species / category i's manure handled using manure system j in climate region k

Emission factor

Calculation of the emission factor is based on the parameters volatile solids excreted (VS), the maximum CH₄ producing capacity for manure (B_o) and the CH₄ conversion factors for each manure management system (MCF).

The **daily excretions of VS** for cattle sub-categories were estimated according to the IPCC Guidelines and GPG (2000: equation 4.16: p. 4.31). Gross energy intake is calculated according to the method described in Chapter 6.2.2. For the livestock categories swine, sheep, goats, horses, mules and asses, and poultry default values from IPCC (1997c: Reference Manual: p. 4.39 to 4.47) were taken.

The **ash content** of cattle manure is assumed to amount 8% on average (IPCC 1997c: Reference Manual: p. 4.47).

The **digestible energy** of the feed for cattle is assumed to be 60% on average, except for calves with 65% (IPCC 1997c: Reference Manual: p. 4.39).

For the Methane Producing Potential (**B₀**) default values are used (IPCC 1997c: Reference Manual: p. 4.39 to 4.47).

For the Methane Conversion Factor (**MCF**) mainly IPCC default values are used (IPCC 2000, p. 4.36 and IPCC 1997c: Reference Manual: p. 4.25). In Switzerland mainly two manure management systems exist, solid storage and liquid/slurry storage. Fattening calves, sheep and goats are mainly kept in deep litter systems and there are also specific MCF values for pasture and poultry systems: The following MCF's were used:

Table 6-9 Manure management systems and Methane conversion factors (MCFs). References: IPCC 2000, p. 4.36 and IPCC 1997b: p. 4.25 (for liquid/slurry and deep litter).

| Manure Management System | Description | MCF [%] |
|--------------------------|---|---------|
| Solid manure | Dung and urine are excreted in a barn. The solids (with and without litter) are collected and stored in bulk for a long time (months) before disposal. | 1.0% |
| Liquid/slurry | Combined storage of dung and urine under animal confinements for longer than 1 month. | 10.0% |
| Pasture | Manure is allowed to lie as it is, and is not managed (distributed, etc.). | 1.0% |
| Deep litter | Dung and urine is excreted in a barn with lots of litter and is not removed for a long time (months). This is applied for the cattle sub-categories of Fattening Calves, Fattening Calves (0-4 months) and for sheep and goats. | 10.0% |
| Poultry system | Manure is excreted on the floor with or without bedding. | 1.5% |

For the MCF for deep litter the 2000 IPCC good practice guidance suggest a value of 39%. However, this would lead to a rather large overestimation of methane emissions from deep litter manure management systems in Switzerland. Since the 2000 IPCC good practice guidance state that the MCF's for cattle and swine deep litter are similar to liquid/slurry, the respective value from the 1996 IPCC guidelines (IPCC 1997b) has been adopted. The choice of a MCF of 10% for deep litter is supported by the specific feeding and manure management regime in Switzerland (especially cold winter temperatures) and confirmed by a number of studies representative for the country specific management conditions (Amon et al. 2001, Külling et al. 2002, Külling et al. 2003, Moller et al. 2004, Hindrichsen et al. 2006, Park et al. 2006 and Sommer et al. 2007). For further details see FOEN 2011 (16.5 attachment E).

The fraction of animal manure handled using different manure management systems (**MS**) as well as the percentages of the grazing time was separately calculated for each livestock category. The fractions are based on Flisch et al. (2009) and calculated within the Swiss ammonium model AGRAMMON (Kupper et al. 2013). Input data for the AGRAMMON-model for the years 1990 and 1995 is based on expert judgement and literature whereas data for 2002, 2007 and 2010 is based on extensive farm surveys. Values in between the assessment years have been interpolated linearly (Table 6-10) while values beyond 2010 are kept constant until new survey results are available. The data clearly reflects the shift towards an increased use of pasture, range and paddocks and a decrease in solid storage. The changes of the manure management system distribution reflects the shift to a more animal friendly livestock husbandry in the course of the agricultural policy reform during the 1990th and the early 20th century. One of the most important programs in this context is called "RAUS" and implies at least 156 days of pasture per year (Schweizerischer Bundesrat, 2008). Accordingly the share of mature dairy cows (and other animals) going to pastures increased substantially and the length of stay on the pasture increased by 50%. In the year

2007 78% of the dairy cows were held on farms who participated in the RAUS program. The number of pasture days was 181 whereas it was 177 in 2010. It can be assumed, that already in the early years of the new millennium most farms accomplished the transition to RAUS and that accordingly a new management standard was reached that did not change significantly afterwards.

Emissions from deep litter and poultry systems have been calculated together with solid storage and are thus reported under solid storage in CRF-table 4.B(a)s2 and 4.B(b).

Table 6-10 Manure management system distribution.

| MS Distribution | | | | | | | | | | | | | | | |
|---|---------------------------|----------------|----------------------------|---------------------------|----------------|----------------------------|---------------------------|----------------|----------------------------|---------------------------|----------------|----------------------------|---------------------------|----------------|----------------------------|
| | 1990 | | | 1995 | | | 2002 | | | 2007 | | | 2010 | | |
| | % | | | % | | | % | | | % | | | % | | |
| | Solid manure/ Deep litter | Liquid/ Slurry | Pasture range and pad-dock | Solid manure/ Deep litter | Liquid/ Slurry | Pasture range and pad-dock | Solid manure/ Deep litter | Liquid/ Slurry | Pasture range and pad-dock | Solid manure/ Deep litter | Liquid/ Slurry | Pasture range and pad-dock | Solid manure/ Deep litter | Liquid/ Slurry | Pasture range and pad-dock |
| Mature Dairy Cattle | 27.70 | 64.04 | 8.26 | 24.53 | 65.93 | 9.54 | 16.38 | 65.66 | 17.96 | 13.94 | 68.35 | 17.72 | 14.84 | 68.22 | 16.94 |
| Mature Non-Dairy Cattle | 32.20 | 41.49 | 26.30 | 34.22 | 39.53 | 26.25 | 20.82 | 40.08 | 39.11 | 20.59 | 50.42 | 28.98 | 18.41 | 49.15 | 32.44 |
| Young Cattle Average (weighted) | 36.38 | 47.84 | 15.78 | 35.47 | 48.67 | 15.86 | 30.02 | 42.47 | 27.51 | 28.23 | 46.52 | 25.26 | 30.18 | 46.47 | 23.35 |
| Fattening Calves | 85.36 | 14.64 | 0.00 | 84.72 | 15.28 | 0.00 | 77.70 | 21.96 | 0.33 | 77.10 | 22.74 | 0.16 | 81.64 | 18.13 | 0.23 |
| Pre-Weaned Calves | 32.20 | 41.49 | 26.30 | 34.22 | 39.53 | 26.25 | 21.20 | 41.54 | 37.27 | 18.98 | 50.88 | 30.14 | 33.27 | 45.86 | 20.87 |
| Breeding Cattle 1st Year | 48.63 | 37.31 | 14.06 | 47.52 | 38.25 | 14.22 | 38.92 | 34.05 | 27.03 | 34.86 | 41.88 | 23.26 | 33.89 | 44.61 | 21.50 |
| Breeding Cattle 2nd Year | 29.00 | 45.63 | 25.37 | 26.82 | 47.54 | 25.64 | 23.49 | 38.12 | 38.38 | 21.14 | 42.32 | 36.54 | 21.25 | 44.45 | 34.30 |
| Breeding Cattle 3rd Year | 29.17 | 50.81 | 20.02 | 28.03 | 51.66 | 20.31 | 22.65 | 42.54 | 34.81 | 21.70 | 46.52 | 31.78 | 21.92 | 47.48 | 30.60 |
| Fattening Cattle | 29.65 | 70.35 | 0.00 | 33.36 | 66.64 | 0.00 | 30.13 | 67.67 | 2.20 | 32.46 | 63.21 | 4.33 | 37.14 | 58.90 | 3.96 |
| Sheep | 69.90 | 0.00 | 30.10 | 69.73 | 0.00 | 30.27 | 66.82 | 0.00 | 33.18 | 60.78 | 0.00 | 39.22 | 66.32 | 0.00 | 33.68 |
| Fattening Sheep | 69.32 | 0.00 | 30.68 | 69.32 | 0.00 | 30.68 | 66.50 | 0.00 | 33.50 | 59.84 | 0.00 | 40.16 | 65.50 | 0.00 | 34.50 |
| Milksheep | 88.57 | 0.00 | 11.43 | 88.57 | 0.00 | 11.43 | 73.94 | 0.00 | 26.06 | 75.92 | 0.00 | 24.08 | 77.15 | 0.00 | 22.85 |
| Goats | 86.39 | 0.00 | 13.61 | 86.39 | 0.00 | 13.61 | 87.82 | 0.00 | 12.18 | 92.88 | 0.00 | 7.12 | 90.00 | 0.00 | 10.00 |
| Goat Places | 86.39 | 0.00 | 13.61 | 86.39 | 0.00 | 13.61 | 87.82 | 0.00 | 12.18 | 92.88 | 0.00 | 7.12 | 90.00 | 0.00 | 10.00 |
| Horses | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 76.14 | 0.00 | 23.86 | 78.66 | 0.00 | 21.34 | 74.38 | 0.00 | 25.62 |
| Horses <3 years | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 61.77 | 0.00 | 38.23 | 61.71 | 0.00 | 38.29 | 66.37 | 0.00 | 33.63 |
| Horses >3 years | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 79.27 | 0.00 | 20.73 | 81.90 | 0.00 | 18.10 | 75.62 | 0.00 | 24.38 |
| Mules and Asses | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 76.93 | 0.00 | 23.07 | 75.21 | 0.00 | 24.79 | 79.31 | 0.00 | 20.69 |
| Mules | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 76.93 | 0.00 | 23.07 | 75.21 | 0.00 | 24.79 | 79.31 | 0.00 | 20.69 |
| Asses | 93.15 | 0.00 | 6.85 | 93.15 | 0.00 | 6.85 | 76.93 | 0.00 | 23.07 | 75.21 | 0.00 | 24.79 | 79.31 | 0.00 | 20.69 |
| Swine | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.34 | 99.54 | 0.12 | 0.14 | 98.68 | 1.18 | 0.27 | 99.61 | 0.13 |
| Piglets | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.84 | 99.16 | 0.00 | 0.67 | 98.97 | 0.36 | 2.34 | 97.66 | 0.00 |
| Fattening Pig over 25 kg | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.27 | 99.56 | 0.17 | 0.00 | 98.51 | 1.49 | 0.00 | 99.85 | 0.15 |
| Dry Sows | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.03 | 99.90 | 0.07 | 0.08 | 98.90 | 1.03 | 0.00 | 99.82 | 0.17 |
| Nursing Sows | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.70 | 99.30 | 0.00 | 0.55 | 99.11 | 0.34 | 0.17 | 99.83 | 0.00 |
| Boars | 0.00 | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.54 | 99.23 | 0.23 | 0.00 | 98.84 | 1.16 | 1.23 | 98.13 | 0.64 |
| Poultry | 100.00 | 0.00 | 0.00 | 99.50 | 0.00 | 0.50 | 97.38 | 0.00 | 2.62 | 96.33 | 0.00 | 3.67 | 97.31 | 0.00 | 2.69 |
| Growers | 100.00 | 0.00 | 0.00 | 99.41 | 0.00 | 0.59 | 99.81 | 0.00 | 0.19 | 98.54 | 0.00 | 1.46 | 98.80 | 0.00 | 1.20 |
| Layers | 100.00 | 0.00 | 0.00 | 99.41 | 0.00 | 0.59 | 94.86 | 0.00 | 5.14 | 92.69 | 0.00 | 7.31 | 93.91 | 0.00 | 6.09 |
| Broilers | 100.00 | 0.00 | 0.00 | 99.61 | 0.00 | 0.39 | 99.38 | 0.00 | 0.62 | 98.84 | 0.00 | 1.16 | 99.74 | 0.00 | 0.26 |
| Turkey | 100.00 | 0.00 | 0.00 | 99.61 | 0.00 | 0.39 | 96.94 | 0.00 | 3.06 | 96.93 | 0.00 | 3.07 | 98.09 | 0.00 | 1.91 |
| Other Poultry (Geese, Ducks, Ostriches, Quails) | 100.00 | 0.00 | 0.00 | 100.00 | 0.00 | 0.00 | 96.93 | 0.00 | 3.07 | 96.93 | 0.00 | 3.07 | 98.81 | 0.00 | 1.19 |

Activity data

Activity data on all livestock categories is taken from SBV (2013) and the Swiss Federal Statistical Office (SFSO 2013d). All activity data has been revised and harmonized during a joint effort of the Agroscope Reckenholz Tänikon Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012) (refer to chapter 6.2.2 for details).

b) N₂O Emissions

Methodology

For the calculation of N₂O emissions from manure management a country specific method based on the Swiss ammonia model AGRAMMON is applied (Kupper et al. 2013). Basically the IPCC emission factors are used, but activity data is adjusted to the particular situation of Switzerland.

For calculation of emissions from manure management AGRAMMON applies other values for the nitrogen excretion per animal category than IPCC and differentiates the animal waste management systems Liquid systems and Solid storage. N₂O emissions from pasture, range

and paddock appear under the category 4D Agricultural Soils, source category 2 Animal Production. IPCC categories Daily Spread and Other Systems are not occurring. The basic animal waste management systems are defined in Flisch et al. (2009) and Menzi et al. (1997).

Emission factors

IPCC default emission factors are used for the two animal waste management systems (IPCC 1997c: Reference Manual: p. 4.104).

Table 6-11 Emission factors for calculating N₂O emissions from manure management (IPCC 1997c: p. 4.104).

| Source | Emission factor per animal waste management system (kg N ₂ O-N / kg N) |
|----------------|---|
| Liquid systems | 0.001 |
| Solid storage | 0.020 |

Activity data

Livestock population data of all categories are taken from the Swiss Farmers Union (SBV 2013) and the Swiss Federal Statistical Office (SFSO 2013d). All activity data has been revised and harmonized during a joint effort of the Agroscope Reckenholz Tänikon Research Station (ART) and the Swiss College of Agriculture (SHL) in 2011 (ART/SHL 2012). Input data is subdivided into the following livestock categories:

Table 6-12 Activity data for calculating N₂O emissions from manure management (ART/SHL 2012, SBV 2013).

| Population Size | 1990-1999 | | | | | | | | | |
|---|------------------|------|------|------|------|------|------|------|------|------|
| | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | 1000 head/places | | | | | | | | | |
| Cattle | 1855 | 1829 | 1783 | 1745 | 1755 | 1748 | 1747 | 1673 | 1641 | 1609 |
| Mature Dairy Cattle | 783 | 781 | 764 | 744 | 750 | 740 | 736 | 712 | 701 | 684 |
| Mature Non-Dairy Cattle | 12 | 14 | 17 | 18 | 20 | 23 | 28 | 32 | 36 | 41 |
| Young Cattle | 1060 | 1034 | 1002 | 983 | 986 | 986 | 983 | 929 | 904 | 884 |
| Fattening Calves | 112 | 111 | 110 | 111 | 101 | 102 | 112 | 106 | 108 | 116 |
| Pre-Weaned Calves | 10 | 11 | 14 | 14 | 16 | 18 | 22 | 26 | 29 | 33 |
| Breeding Cattle 1st Year | 346 | 337 | 324 | 308 | 306 | 295 | 286 | 260 | 254 | 219 |
| Breeding Cattle 2nd Year | 253 | 252 | 251 | 239 | 237 | 239 | 243 | 233 | 217 | 188 |
| Breeding Cattle 3rd Year | 151 | 148 | 147 | 142 | 141 | 139 | 140 | 139 | 133 | 118 |
| Fattening Cattle | 188 | 175 | 158 | 168 | 184 | 193 | 180 | 165 | 163 | 210 |
| Sheep | 395 | 409 | 415 | 424 | 405 | 387 | 419 | 420 | 422 | 424 |
| Fattening Sheep | 191 | 201 | 201 | 211 | 201 | 191 | 208 | 208 | 209 | 222 |
| Milkshew | 4 | 4 | 4 | 4 | 3 | 3 | 3 | 3 | 4 | 6 |
| Goats | 68 | 65 | 58 | 57 | 55 | 53 | 57 | 58 | 60 | 62 |
| Goat Places | 45 | 43 | 38 | 37 | 36 | 35 | 37 | 38 | 40 | 41 |
| Horses | 51 | 52 | 53 | 54 | 58 | 62 | 62 | 65 | 64 | 65 |
| Horses <3 years | 11 | 11 | 11 | 12 | 14 | 16 | 16 | 14 | 14 | 15 |
| Horses >3 years | 40 | 41 | 42 | 43 | 44 | 45 | 47 | 51 | 50 | 51 |
| Mules and Asses | 11 | 11 | 11 | 11 | 11 | 11 | 12 | 13 | 14 | 15 |
| Mules | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| Asses | 10 | 11 | 11 | 11 | 11 | 11 | 12 | 13 | 13 | 15 |
| Swine | 1787 | 1723 | 1706 | 1692 | 1569 | 1446 | 1379 | 1395 | 1487 | 1453 |
| Piglets | 299 | 283 | 291 | 300 | 287 | 275 | 241 | 252 | 262 | 281 |
| Fattening Pig over 25 kg | 1025 | 990 | 973 | 943 | 855 | 768 | 779 | 780 | 837 | 734 |
| Dry Sows | 129 | 126 | 125 | 125 | 117 | 109 | 99 | 104 | 111 | 101 |
| Nursing Sows | 37 | 37 | 37 | 37 | 35 | 33 | 30 | 30 | 31 | 35 |
| Boars | 8 | 8 | 8 | 8 | 8 | 7 | 6 | 6 | 6 | 6 |
| Poultry | 5938 | 5647 | 5502 | 6410 | 6330 | 6251 | 6440 | 6553 | 6740 | 6908 |
| Growers | 719 | 664 | 710 | 719 | 717 | 714 | 732 | 733 | 793 | 761 |
| Layers | 3083 | 2645 | 2536 | 2518 | 2318 | 2118 | 2226 | 2278 | 2270 | 2223 |
| Broilers | 2020 | 2199 | 2096 | 2990 | 3111 | 3231 | 3293 | 3342 | 3502 | 3747 |
| Turkey | 95 | 117 | 140 | 163 | 166 | 170 | 174 | 184 | 158 | 155 |
| Other Poultry (Geese, Ducks, Ostriches, Quails) | 22 | 21 | 21 | 20 | 18 | 17 | 15 | 16 | 16 | 22 |

Table continued from last page:

| Population Size | 2000-2009 | | | | | | | | | |
|---|------------------|------|------|------|------|------|------|------|------|------|
| | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| | 1000 head/places | | | | | | | | | |
| Cattle | 1588 | 1611 | 1594 | 1570 | 1545 | 1555 | 1567 | 1572 | 1604 | 1597 |
| Mature Dairy Cattle | 669 | 669 | 658 | 638 | 621 | 621 | 618 | 615 | 629 | 599 |
| Mature Non-Dairy Cattle | 45 | 51 | 58 | 65 | 70 | 78 | 87 | 94 | 98 | 108 |
| Young Cattle | 874 | 891 | 878 | 867 | 854 | 856 | 862 | 863 | 877 | 890 |
| Fattening Calves | 103 | 115 | 114 | 114 | 111 | 106 | 101 | 100 | 95 | 101 |
| Pre-Weaned Calves | 36 | 40 | 47 | 52 | 57 | 62 | 67 | 72 | 76 | 86 |
| Breeding Cattle 1st Year | 236 | 238 | 230 | 220 | 215 | 222 | 223 | 223 | 232 | 226 |
| Breeding Cattle 2nd Year | 222 | 219 | 219 | 213 | 205 | 205 | 210 | 210 | 213 | 212 |
| Breeding Cattle 3rd Year | 130 | 130 | 126 | 124 | 121 | 113 | 110 | 109 | 110 | 119 |
| Fattening Cattle | 147 | 148 | 142 | 144 | 145 | 147 | 149 | 148 | 152 | 147 |
| Sheep | 421 | 420 | 430 | 445 | 441 | 446 | 448 | 444 | 446 | 432 |
| Fattening Sheep | 217 | 217 | 220 | 229 | 227 | 229 | 231 | 230 | 229 | 227 |
| Milksheep | 7 | 7 | 7 | 8 | 8 | 9 | 10 | 10 | 11 | 12 |
| Goats | 62 | 63 | 66 | 67 | 71 | 74 | 76 | 79 | 81 | 81 |
| Goat Places | 41 | 42 | 43 | 45 | 46 | 48 | 51 | 52 | 53 | 54 |
| Horses | 66 | 64 | 64 | 65 | 65 | 65 | 66 | 67 | 68 | 69 |
| Horses <3 years | 13 | 12 | 12 | 11 | 11 | 11 | 11 | 11 | 11 | 10 |
| Horses >3 years | 53 | 52 | 52 | 53 | 53 | 54 | 55 | 56 | 57 | 59 |
| Mules and Asses | 16 | 16 | 17 | 17 | 18 | 19 | 19 | 20 | 20 | 22 |
| Mules | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Asses | 15 | 15 | 16 | 17 | 17 | 18 | 19 | 19 | 20 | 21 |
| Swine | 1498 | 1548 | 1557 | 1529 | 1538 | 1609 | 1635 | 1573 | 1540 | 1557 |
| Piglets | 297 | 319 | 327 | 323 | 328 | 338 | 367 | 345 | 336 | 338 |
| Fattening Pig over 25 kg | 751 | 763 | 768 | 752 | 753 | 797 | 786 | 767 | 763 | 779 |
| Dry Sows | 105 | 108 | 109 | 105 | 108 | 113 | 115 | 106 | 105 | 105 |
| Nursing Sows | 37 | 38 | 36 | 36 | 35 | 36 | 37 | 35 | 33 | 33 |
| Boars | 6 | 6 | 6 | 5 | 5 | 5 | 5 | 4 | 4 | 4 |
| Poultry | 6983 | 6939 | 7339 | 7587 | 8061 | 8260 | 7670 | 8228 | 8543 | 8809 |
| Growers | 832 | 745 | 754 | 809 | 853 | 868 | 888 | 902 | 919 | 967 |
| Layers | 2150 | 2069 | 2154 | 2117 | 2089 | 2189 | 2147 | 2198 | 2255 | 2318 |
| Broilers | 3808 | 3993 | 4298 | 4518 | 4971 | 5060 | 4481 | 5002 | 5300 | 5456 |
| Turkey | 173 | 123 | 124 | 134 | 139 | 132 | 137 | 112 | 54 | 52 |
| Other Poultry (Geese, Ducks, Ostriches, Quails) | 21 | 9 | 8 | 9 | 9 | 11 | 16 | 14 | 15 | 16 |

| Population Size | 2010-2012 | | |
|---|------------------|------|------|
| | 2010 | 2011 | 2012 |
| | 1000 head/places | | |
| Cattle | 1591 | 1577 | 1565 |
| Mature Dairy Cattle | 589 | 589 | 591 |
| Mature Non-Dairy Cattle | 111 | 111 | 114 |
| Young Cattle | 891 | 877 | 859 |
| Fattening Calves | 99 | 101 | 99 |
| Pre-Weaned Calves | 88 | 88 | 91 |
| Breeding Cattle 1st Year | 226 | 221 | 212 |
| Breeding Cattle 2nd Year | 213 | 207 | 200 |
| Breeding Cattle 3rd Year | 119 | 116 | 112 |
| Fattening Cattle | 145 | 145 | 146 |
| Sheep | 434 | 424 | 417 |
| Fattening Sheep | 228 | 222 | 219 |
| Milksheep | 12 | 12 | 13 |
| Goats | 83 | 83 | 85 |
| Goat Places | 55 | 56 | 57 |
| Horses | 71 | 66 | 67 |
| Horses <3 years | 10 | 10 | 9 |
| Horses >3 years | 61 | 56 | 58 |
| Mules and Asses | 23 | 22 | 23 |
| Mules | 1 | 1 | 1 |
| Asses | 22 | 21 | 22 |
| Swine | 1589 | 1579 | 1544 |
| Piglets | 351 | 353 | 345 |
| Fattening Pig over 25 kg | 788 | 787 | 776 |
| Dry Sows | 106 | 103 | 97 |
| Nursing Sows | 34 | 32 | 31 |
| Boars | 4 | 3 | 3 |
| Poultry | 9025 | 9478 | 9955 |
| Growers | 926 | 970 | 1076 |
| Layers | 2438 | 2437 | 2521 |
| Broilers | 5580 | 5984 | 6282 |
| Turkey | 58 | 58 | 51 |
| Other Poultry (Geese, Ducks, Ostriches, Quails) | 23 | 29 | 25 |

Data on nitrogen excretion per animal category (kg N/head/year) is taken from Kupper et al. (2013) (see Table 6-13). These values are based on Flisch et al. (2009) and adjusted according to the Swiss ammonia model AGRAMMON. Unlike IPCC, the age structure of the animals and the different use of the animals (e.g. fattening and breeding) are considered. Standard nitrogen excretion rates are modified within the AGRAMMON model to account for changing agricultural structures and production techniques along the years (e.g. milk yield, protein reduced animal feed, use of feed concentrates etc.). Calculation of nitrogen excretion of mature dairy cattle is dependent on milk production and is therefore increasing from 1990 to 2007. In the year 2007 milk yield reaches 6500 liter per head and year. To reach higher milk yields farmers usually apply higher shares of energy rich feed concentrates. Consequently increases in nitrogen excretion rates are lower or nonexistent beyond a milk yield of 6500 liter per head and year (Flisch et al. 2009). In accordance with the AGRAMMON model the same nitrogen excretion rates as in 2010 have been used for the years 2011 and 2012 due to the lack of further survey results. Sheep in Switzerland are fed mainly according to a regime based on roughage from extensive pasture and meadows (Flisch et al. 2009) and are estimated to excrete approximately 8.0 kg N per head and year. This is considerably lower than IPCC default. However, nitrogen excretion is averaged over the whole population of which roughly 50% are lambs and other immature animals. Swine show a significant decrease in nitrogen excretion per head over almost the whole inventory time period which can be explained by the increasing use of protein reduced fodder.

The consideration of adopted nitrogen excretion values is one of the major advantages of the country specific method in Switzerland. The more disaggregated approach leads to considerable lower calculated nitrogen excretion rates compared to IPCC, which therefore also implies lower total N₂O emissions from manure management.

Table 6-13 Nitrogen excretion per animal category, 1990-2012 (Kupper et al. 2013).

| Nitrogen Excretion | Unit | 1990-1999 | | | | | | | | | |
|---------------------------------|---|--------------------|-------|-------|-------|-------|-------|-------|--------|--------|--------|
| | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | | kg N / unit / year | | | | | | | | | |
| Mature Dairy Cattle | head | 96.06 | 96.57 | 97.09 | 97.61 | 98.13 | 98.65 | 99.35 | 100.05 | 100.75 | 101.45 |
| Mature Non-Dairy Cattle | head | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 |
| Young Cattle Average (weighted) | head | 33.08 | 33.11 | 33.21 | 33.13 | 33.31 | 33.37 | 33.28 | 33.56 | 33.31 | 32.85 |
| | Fattening Calves | place | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 |
| | Pre-Weaned Calves | head | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 |
| | Breeding Cattle 1st Year | head | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 |
| | Breeding Cattle 2nd Year | head | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 |
| | Breeding Cattle 3rd Year | head | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 |
| | Fattening Cattle | place | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 |
| Sheep | head | 7.46 | 7.56 | 7.46 | 7.64 | 7.62 | 7.59 | 7.58 | 7.58 | 7.63 | 8.14 |
| | Fattening Sheep | place | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 |
| | Milksheep | place | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 |
| Goats | head | 10.49 | 10.58 | 10.56 | 10.53 | 10.47 | 10.41 | 10.43 | 10.42 | 10.58 | 10.59 |
| | Goat Places | place | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 |
| Horses | head | 43.57 | 43.57 | 43.57 | 43.57 | 43.51 | 43.47 | 43.50 | 43.56 | 43.57 | 43.55 |
| | Horses <3 years | head | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 |
| | Horses >3 years | head | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 |
| Mules and Asses | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| | Mules | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| | Asses | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| Swine | head | 13.37 | 13.38 | 13.31 | 13.13 | 12.95 | 12.75 | 12.72 | 12.35 | 11.99 | 10.88 |
| | Piglets | place | 4.60 | 4.60 | 4.60 | 4.60 | 4.60 | 4.60 | 4.60 | 4.60 | 4.60 |
| | Fattening Pig over 25 kg | place | 17.01 | 16.95 | 16.88 | 16.81 | 16.75 | 16.68 | 16.15 | 15.63 | 14.58 |
| | Dry Sows | place | 24.28 | 24.28 | 24.28 | 24.28 | 24.28 | 23.53 | 22.77 | 22.02 | 21.26 |
| | Nursing Sows | place | 47.57 | 47.57 | 47.57 | 47.57 | 47.57 | 46.77 | 45.98 | 45.18 | 44.39 |
| | Boars | head | 20.50 | 20.50 | 20.50 | 20.50 | 20.50 | 20.02 | 19.53 | 19.04 | 18.56 |
| Poultry | head | 0.57 | 0.56 | 0.56 | 0.54 | 0.53 | 0.53 | 0.54 | 0.54 | 0.54 | 0.55 |
| | Growers | place | 0.34 | 0.34 | 0.34 | 0.34 | 0.34 | 0.34 | 0.33 | 0.33 | 0.32 |
| | Layers | place | 0.71 | 0.71 | 0.71 | 0.71 | 0.71 | 0.72 | 0.74 | 0.75 | 0.76 |
| | Broilers | place | 0.40 | 0.40 | 0.40 | 0.40 | 0.40 | 0.41 | 0.41 | 0.42 | 0.43 |
| | Turkey | place | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 |
| | Other Poultry (Geese, Ducks, Ostriches, Quails) | place | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 |

| Nitrogen Excretion | Unit | 2000-2009 | | | | | | | | | |
|---------------------------------|---|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| | | kg N / unit / year | | | | | | | | | |
| Mature Dairy Cattle | head | 102.15 | 102.85 | 103.55 | 104.49 | 105.42 | 106.35 | 107.28 | 108.21 | 108.20 | 108.18 |
| Mature Non-Dairy Cattle | head | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 | 80.00 |
| Young Cattle Average (weighted) | head | 33.56 | 33.27 | 33.26 | 33.27 | 33.25 | 33.12 | 33.18 | 33.17 | 33.25 | 33.42 |
| | Fattening Calves | place | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 | 13.00 |
| | Pre-Weaned Calves | head | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 | 34.00 |
| | Breeding Cattle 1st Year | head | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 | 25.00 |
| | Breeding Cattle 2nd Year | head | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 | 40.00 |
| | Breeding Cattle 3rd Year | head | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 | 55.00 |
| | Fattening Cattle | place | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 | 33.00 |
| Sheep | head | 8.06 | 8.08 | 8.03 | 8.09 | 8.13 | 8.13 | 8.18 | 8.26 | 8.22 | 8.47 |
| | Fattening Sheep | place | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 |
| | Milksheep | place | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 | 21.00 |
| Goats | head | 10.60 | 10.69 | 10.43 | 10.66 | 10.47 | 10.49 | 10.59 | 10.50 | 10.49 | 10.70 |
| | Goat Places | place | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 | 16.00 |
| Horses | head | 43.60 | 43.61 | 43.63 | 43.64 | 43.65 | 43.66 | 43.66 | 43.67 | 43.67 | 43.70 |
| | Horses <3 years | head | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 | 42.00 |
| | Horses >3 years | head | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 | 44.00 |
| Mules and Asses | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| | Mules | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| | Asses | head | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 | 15.70 |
| Swine | head | 10.53 | 10.10 | 9.75 | 9.61 | 9.50 | 9.42 | 9.21 | 9.09 | 9.18 | 9.23 |
| | Piglets | place | 4.60 | 4.60 | 4.60 | 4.56 | 4.53 | 4.49 | 4.45 | 4.42 | 4.38 |
| | Fattening Pig over 25 kg | place | 14.05 | 13.53 | 13.00 | 12.78 | 12.55 | 12.33 | 12.11 | 11.89 | 11.95 |
| | Dry Sows | place | 20.51 | 19.75 | 19.00 | 19.08 | 19.16 | 19.24 | 19.33 | 19.41 | 19.51 |
| | Nursing Sows | place | 43.59 | 42.80 | 42.00 | 42.43 | 42.86 | 43.30 | 43.73 | 44.16 | 43.17 |
| | Boars | head | 18.07 | 17.58 | 17.10 | 17.18 | 17.27 | 17.35 | 17.43 | 17.52 | 17.73 |
| Poultry | head | 0.55 | 0.55 | 0.55 | 0.55 | 0.54 | 0.54 | 0.55 | 0.54 | 0.53 | 0.53 |
| | Growers | place | 0.32 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 | 0.31 |
| | Layers | place | 0.77 | 0.79 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 |
| | Broilers | place | 0.44 | 0.44 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 |
| | Turkey | place | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 | 1.40 |
| | Other Poultry (Geese, Ducks, Ostriches, Quails) | place | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 | 0.56 |

Table continued from last page:

| Nitrogen Excretion | Unit | 2010-2012 | | |
|---|-------|--------------------|--------|--------|
| | | 2010 | 2011 | 2012 |
| | | kg N / unit / year | | |
| Mature Dairy Cattle | head | 108.17 | 108.17 | 108.17 |
| Mature Non-Dairy Cattle | head | 80.00 | 80.00 | 80.00 |
| Young Cattle Average (weighted) | head | 33.45 | 33.36 | 33.32 |
| Fattening Calves | place | 13.00 | 13.00 | 13.00 |
| Pre-Weaned Calves | head | 34.00 | 34.00 | 34.00 |
| Breeding Cattle 1st Year | head | 25.00 | 25.00 | 25.00 |
| Breeding Cattle 2nd Year | head | 40.00 | 40.00 | 40.00 |
| Breeding Cattle 3rd Year | head | 55.00 | 55.00 | 55.00 |
| Fattening Cattle | place | 33.00 | 33.00 | 33.00 |
| Sheep | head | 8.48 | 8.46 | 8.53 |
| Fattening Sheep | place | 15.00 | 15.00 | 15.00 |
| Milksheep | place | 21.00 | 21.00 | 21.00 |
| Goats | head | 10.57 | 10.76 | 10.83 |
| Goat Places | place | 16.00 | 16.00 | 16.00 |
| Horses | head | 43.72 | 43.71 | 43.73 |
| Horses <3 years | head | 42.00 | 42.00 | 42.00 |
| Horses >3 years | head | 44.00 | 44.00 | 44.00 |
| Mules and Asses | head | 15.70 | 15.70 | 15.70 |
| Mules | head | 15.70 | 15.70 | 15.70 |
| Asses | head | 15.70 | 15.70 | 15.70 |
| Swine | head | 9.18 | 9.17 | 9.15 |
| Piglets | place | 4.36 | 4.36 | 4.36 |
| Fattening Pig over 25 kg | place | 12.09 | 12.09 | 12.09 |
| Dry Sows | place | 19.73 | 19.73 | 19.73 |
| Nursing Sows | place | 41.19 | 41.19 | 41.19 |
| Boars | head | 18.16 | 18.16 | 18.16 |
| Poultry | head | 0.54 | 0.53 | 0.53 |
| Growers | place | 0.31 | 0.31 | 0.31 |
| Layers | place | 0.80 | 0.80 | 0.80 |
| Broilers | place | 0.45 | 0.45 | 0.45 |
| Turkey | place | 1.40 | 1.40 | 1.40 |
| Other Poultry (Geese, Ducks, Ostriches, Quails) | place | 0.56 | 0.56 | 0.56 |

The split of nitrogen flows into the different animal waste management systems and its temporal dynamic is based on Kupper et al. (2013). The distribution is consistent with the allocation of volatile solids used for the calculation of CH₄ emissions (for further information refer to the previous section on CH₄ emissions).

6.3.3 Uncertainties and Time-Series Consistency

For the uncertainty analysis the input data from ART (2008a) was used and was weighted with current activity and emission data. The arithmetic mean of the lower and upper bound is used for activity data and for emission factors resulting in a combined uncertainty of 55% for CH₄ and 64% for N₂O in Tier 1 analysis, where N₂O includes the sum of liquid and solid storage. For Tier 2 analysis, liquid and solid storage are treated separately giving the following results for the uncertainty intervals and the mean uncertainties:

- 4B CH₄ [-54.5%; 55.1%], mean uncertainty 54.7%
- 4B N₂O liquid: [-86.4%; 69.3%], mean uncertainty 77.9%
- 4B N₂O solid: [-60.7%; 58.0%], mean uncertainty 59.4%

Further results of Tier 2 uncertainty analysis are shown in Table 1-18.

Time series consistency of livestock population data and gross energy intake: See Chapter 6.2.3.)

Input data from the AGRAMMON-model are available for the years 1990 and 1995 (expert judgement and literature) as well as for 2002, 2007 and 2010 (extensive farm surveys).

Values in between the assessment years were interpolated linearly while values beyond 2010 are kept constant until new survey results are available.

6.3.4 Source-Specific QA/QC and Verification

All QA/QC activities are further described in a separate document (ART 2013a). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed.

For quality of livestock population data and animal energy intake please consult Chapter 6.2.4.

The time series of activity data and emission factors have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the following triple check:

- the results for 2012 are compared with the results for 2011 within the current CRF
- the results for 2011 are compared between the current CRF-tables and the CRF-tables of the submission 2013
- the results for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the submission 2013.

Additionally an independent quality control was conducted by INFRAS by a countercheck of the data and calculation sheets elaborated by Agroscope (Agroscope 2014).

a) CH₄

For CH₄ the documentation about the data set and calculation method assures transparency and traceability of the calculation methods (Soliva 2006). Additionally a document in German lists all the methodological differences between the former calculations and the current methodology regarding CH₄ estimations (Soliva 2006a).

IPCC tables with data for estimating emission factors for all livestock categories (such as weight, feed digestibility, maximum CH₄ producing capacity (B₀) or daily excretion of volatile solids) were filled in, checked for consistency and confidence and compared with IPCC default values (refer to Table A - 28 in Annex A3.3). Factors for methane conversion (MCF) and manure management distribution (MS) were analysed considering the Swiss national agricultural context.

The emission factors of 4B CH₄ were compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available for submission 2013 (ART 2013a, INFRAS 2012). Most implied emission factors for CH₄ emissions from manure management in Switzerland are considerably above IPCC default. Differences are mainly due to different allocations to manure management systems i.e., a higher share of manure stored in liquid systems and as deep litter.

During the years 2009-2012 the group of animal nutrition from the Swiss Federal Institute of Technology Zürich investigated the effect of different feeding and management strategies on methane and nitrous oxide emissions from enteric fermentation and manure management of cattle held under typical Swiss management conditions (Kreuzer 2012). Measured values of various parameters such as digestible energy, B₀ or MCF have been compared to IPCC default values and values in the Swiss greenhouse gas inventory. Preliminary analysis suggests that overall emissions are neither over- nor underestimated (Zeitz et al. 2012). Further investigations have to show to what extent the preliminary estimates will be confirmed to provide a basis for implementation in Switzerland's GHG inventory.

During the past years a couple of studies have been conducted to verify methane emissions at regional scale comparing bottom up estimates with atmospherical measurements (Hiller et al. 2014, Hiller et al. 2014a, Stieger 2014). For further information see section 6.2.4.

b) N₂O

N₂O estimation is based on the Swiss ammonium emission model AGRAMMON that is documented in Kupper et al. (2013).

All relevant data needed for the calculation of N₂O emissions such as nitrogen excretion, manure management system distribution and N₂O emission factors have been checked for consistency and have been compared to the corresponding values of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) and to the IPCC default value if available (ART 2013a). As one of the most important parameters, nitrogen excretion has been analysed in more detail. A comparison in 2011 revealed that bottom up calculations of total nitrogen excretion in the Swiss GHG inventory are only 5-8% below the values of an independent top down approach subtracting all nitrogen contained in animal products from the total amount of nitrogen in animal feedstuff produced in or imported to the country (Peter et al. 2006, Spiess 2005). Furthermore N_{ex} values for the most important animal categories (mature dairy cattle, mature non-dairy cattle and swine), being responsible for almost 70% of total nitrogen excretion, are very well in line with the alternative gross energy approach suggested in the 2006 IPCC guidelines.

6.3.5 Source-Specific Recalculations

For recalculation of livestock numbers, energy requirements and milk yield see chapter 6.2.5.

The nitrogen excretion rate of mature dairy cattle of the year 2011 has been revised in order to be consistent with the AGRAMMON model (Kupper et al. 2013). The reduced nitrogen excretion rate resulted in an overall emission reduction (including source category 4B and 4D) of less than 10 Gg CO₂ equivalent.

6.3.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented.

Currently manure used for biogas production as reported under 1A1a and 6D is not subtracted from animal manure in sector 4B. It is planned to improve the respective cross sectoral reporting in future submissions to avoid double counting of emissions. The regulations for national compensation projects in the field of anaerobic manure treatment are still under revision. These regulations will be an important basis for the respective estimates in the national inventory as the issued emission reduction certificates should correspond to emission reductions in the inventory. The regulations should be finalized during 2014, so that the respective recalculations will be included in the GHG-Inventory submission 2015. Currently the agriculture expert is in contact with Ökostrom Schweiz, the association of approximately 100 energy-producing farmers in order to get the required data on manure processed in the digesters.

Other projects are eventually postponed in order to give first priority to changes related to the new reporting guidelines.

6.4 Source Category 4C – Rice Cultivation

Rice Cultivation is of minor importance in Switzerland. The agricultural land used for rice cultivation and the annual yield of rice are not estimated by the Swiss Farmers Union (SBV 2013). There is only some insignificant upland rice cultivation. CH₄ Emissions are assumed to be zero. They are therefore not considered in the emission calculation.

6.5 Source Category 4D – Agricultural Soils

6.5.1 Source Category Description

Tier 1 and Tier 2 Key category 4D:

4D1: N₂O emissions from Agricultural Soils; Direct Soil Emissions (level and trend)

4D2: N₂O emissions from Agric.Soils; Pasture, Range and Paddock Manure (level and trend)

4D3: N₂O emissions from Agricultural Soils; Indirect Soil Emissions (level and trend)

The source category 4D includes the following emissions: Direct N₂O emissions from soils and from animal production (emission from pasture, range and paddock), indirect N₂O emissions, other N₂O emissions from agricultural soils (application of sewage sludge and compost), NO_x emissions from soils and NMVOC emissions.

Direct and indirect N₂O emissions are decreasing since 1990 in almost all sub-categories. Contrarily N₂O emissions from animal production have been increasing due to a higher share of manure excreted on pasture, range and paddock. NO_x emissions declined by more than 18% since 1990.

The general trend can be explained by a reduction of the number of cattle and a reduced input of mineral fertilisers due to the introduction of the “Proof of Ecological Performance (PEP)” (ART 2013a, Leifeld and Fuhrer 2005). From 2004 on the cattle population increased again which lead to higher total animal manure nitrogen excretion.

Table 6-14 Specification of source category 4D Agricultural Soils. (AD: Activity data; EF: Emission factor).

| 4D | Source | Specification | Data Source |
|-----|--|--|--|
| 4D1 | Direct soil emissions | Includes emissions from synthetic fertilizer, animal manure and crop residues, N-fixing crops, organic soils, residues from meadows and pasture, N-fixation on meadows and pasture | AD: SBV 2013, ART/SHL 2012, Agricura 2012, Flisch et al. 2009, FAL/RAC 2001, Kupper et al. 2013, Leifeld et al. 2003, Schmid et al. 2000, Walther et al. 1994; EF: IPCC 1997c (N ₂ O), IPCC 2000 |
| 4D2 | Pasture, range and paddock manure | Emissions from pasture, range and paddock | AD: SBV 2013, ART/SHL 2012, Flisch et al. 2009, Kupper et al. 2013; EF: IPCC 1997c |
| 4D3 | Indirect emissions | Leaching and runoff, N deposition air to soil | AD: SBV 2013, ART/SHL 2012, Flisch et al. 2009, Kupper et al. 2013, Prasuhn and Braun 1994, Braun et al. 1994, Schmid et al. 2000, EEA 2007; EF: IPCC 2000 |
| 4D4 | Other (sewage sludge and compost used for fertilizing) | | AD: SBV 2013, Kupper et al. 2013; EF: IPCC 1997c |

6.5.2 Methodological Issues

Methodology

For calculation of N₂O emissions from agricultural soils the national method IULIA is applied. IULIA is an IPCC-derived method for the calculation of N₂O emissions from agriculture that basically uses the same emission factors, but adjusts the activity data to the particular situation of Switzerland (Schmid et al. 2000). According to Schmid et al. (2000) IULIA is better adapted to the conditions of Swiss agriculture, compared to the IPCC method.

IULIA has been updated with new parameters derived from the Swiss ammonium model AGRAMMON (Kupper et al. 2013). New values for nitrogen excretion, manure system distribution and ammonium emission factors have been adopted. Furthermore the updated version of the "Principles of Fertilization in Arable and Forage Crop Production" (GruDAF; Flisch et al. 2009) has been used instead of obsolete data from FAL/RAC 2001 and Walther et al. 1994.

The modelling of the N₂O emissions is realised in Agroscope (2014). The model structure is displayed in the following figure.




Figure 6-6 Diagram depicting the methodology of the approach to calculate the N₂O emissions in Agriculture. Note that the figure shows explicitly the methodology of the approach and not the physical nitrogen flow.

Main differences between the IULIA/AGRAMMON method and IPCC are (Schmid et al. 2000: p. 74):

- IULIA/AGRAMMON estimates lower nitrogen excretion per animal category, especially due to the lower excretions of young cattle (refer to chapter 6.3.2.b).
- The amount of losses to the atmosphere from the excreted nitrogen is almost 50% higher compared to IPCC.
- The amount of leaching (of manure nitrogen and of synthetic fertilizers) is lower by 1/3 compared to IPCC.

- Compared to the IPCC default method more manure is managed in liquid systems and less manure is excreted on pasture, range and paddock. Furthermore the manure management system distribution is not constant over the time series.
- The nitrogen inputs from biological fixation are higher by more than a factor of 30 since fixation on meadows and pastures are also considered. The consideration of nitrogen fixation from grassland is one of the major advantages of the method IULIA as the grassland accounts for the majority of nitrogen fixed in Swiss Agriculture.
- The nitrogen inputs from crop residues are only 25% higher although emissions from plant residue on grasslands are considered. This is explained by the fact that the emissions from plant residues returned to soils on cropland are estimated 50% below the IPCC defaults (see Schmid et al. 2000 p. 68).

Despite the different assumptions of the two methods, differences at the level of the N₂O emissions are quite moderate. In a comparison of the 1996 N₂O inventory, IULIA estimations of the N₂O emissions from agriculture were approximately 15% lower than the IPCC estimations (Schmid et al. 2000: p. 75). This comparison has been made with the original IULIA model in the year 2000. Since then the model has been developed further (e.g. implementation of the AGRAMMON model). A comprehensive comparison as conducted by Schmid et al. 2000 has not been made since.

Direct emissions from soil (4D1)

Calculation of direct N₂O emissions from soil is based on IPCC 2000 Tier 1b.

- Emissions from **synthetic fertilizer** include urea and other mineral fertilizers (mainly ammonium-nitrate). The amount of nitrogen input due to these fertilizers is taken from SBV (2013), Agricura (2012) and Kupper et al. (2013). Fertilizer statistics is based on sales statistics by the compulsory storekeepers of fertilizers (Pflichtlagerhalter) and small importers. Agricura conducts plausibility checks with import-data received by the Directorate General of Customs (Oberzolldirektion). From the amount of nitrogen in fertilizer, losses to the atmosphere in form of NH₃ are subtracted and the rest is multiplied with the corresponding N₂O emission factor. According to AGRAMMON NH₃ losses to the atmosphere are 15% for urea and 2% for other synthetic fertilizers (van der Weerden and Jarvis 1997) instead of the IPCC value of 10% for NH₃ and NO_x (see Table 6-16). For more information on ammonia volatilization from synthetic fertilizers see the paragraph on Indirect emissions (4D3). NO_x emissions are not subtracted since they occur mainly after the fertilizer application. Thus, the basis for N₂O-emissions is the synthetic fertilizer including the nitrogen that will be lost as NO_x later (Berthoud 2004).
- To model the emissions of **animal manure applied to soils**, nitrogen input from manure is calculated as the total N excretion minus N excreted on pasture, range and paddock minus ammonia volatilization from solid and liquid manure. The losses (to the atmosphere) as ammonia are specified for each animal category separately instead of using a fixed ratio of 20% (Kupper et al. 2013). For more information on ammonia volatilization from animal manure see the paragraph on indirect emissions (4D3). NO_x emissions are not subtracted since they occur after the application of animal wastes (Berthoud 2004). $Frac_{GASM}$ in CRF-table 4.Ds2 represents the amount of nitrogen volatilized as NH₃ from housing, manure storage and manure application divided by the manure excreted in the stable. The nitrogen input from manure applied to soils in CRF-table 4.Ds1 can thus be calculated with the numbers given in CRF-table 4.B(b) and 4.Ds2. For further details regarding the volatilized N refer to Table 6-16.
- Emissions from **crop residues** are based on the amount of nitrogen in crop residues returned to soil. According to IULIA (Schmid et al. 2000: p. 68 and p. 100) the calculation of nitrogen in crop residues is based on data reported on crop yields (SBV

2013), the standard values for arable crop yields (FAL/RAC 2001 and Flisch et al. 2009) and standard amounts of nitrogen in crop residues returned to soils (FAL/RAC 2001 and Flisch et al. 2009). The calculation of the amount of nitrogen in crop residues returned to soil according to IULIA is as follows (Schmid et al. 2000: p. 101):

$$F_{CR} = \sum_{Cr} (E_{Cr} * \frac{NR_{Cr}}{Y_{Cr}})$$

F_{CR} : Amount of nitrogen in crop residues returned to soils (t N)

E_{Cr} : Amount of crop yields for culture Cr (t)

Y_{Cr} : Standard values for arable crop yields for culture Cr (t/ha)

NR_{Cr} : Standard amount of nitrogen in crop residues returned to soils (t/ha)

In addition to the N transfer from crop residues, IULIA also takes into account the plant residue returned to soils on meadows and pastures (Schmid et al. 2000). Three quarters of the agricultural land use consists of grassland which underscores the importance of this source for Switzerland. Input data on the managed area of meadows and pastures are taken from SBV (2013) and the Swiss Federal Statistical Office (SFSO 2013d). Estimated values of total crop production, nitrogen incorporated with crop residues $F_{(CR)}$, residue/crop ratio, dry matter (dm) fraction of residues and nitrogen content of residues are provided in Annex A3.3.

- For calculation of emissions from **N-fixing crops**, IULIA assumes that 60% of the nitrogen in leguminous crops originates from biological nitrogen fixation (Schmid et al. 2000: p. 70). This is in line with the IPCC Guidelines that state that biological nitrogen fixation supplies 50-60 per cent of the nitrogen in grain legumes (IPCC 1997c, p. 4.89). The total amount of nitrogen is calculated according to the calculation of nitrogen in crop residues, additionally taking into account the nitrogen contained in the crop product. In addition, IULIA takes biological nitrogen fixation on meadows and pastures into account, assuming a nitrogen concentration of 3.5% in the dry matter from which 80% derives from biological nitrogen fixation. For the dry matter production of clover on pastures and meadows statistical data were used (Schmid et al. 2000: p. 70). The following table gives an overview of the calculation of emissions from N-fixing crops.

Table 6-15 Input values for calculation of emissions from N-fixing crops according to IULIA (Schmid et al. 2000: p. 70).

| Fixation | Share of N caused by fixation | Share of N in dry matter |
|---------------------------------------|-------------------------------|--------------------------|
| Leguminous (N-fixing crops) | 0.6 | crop-specific |
| Clover (Fixation meadows and pasture) | 0.8 | 0.035 |

Estimates of total crop production and nitrogen fixed per kg crop dry matter are provided in Annex A3.3.

- Emissions from **cultivated organic soils** are based on estimations on the area of cultivated organic soils and the IPCC default emission factor for N_2O emissions from cultivated organic soils (IPCC 1997b). The area of cultivated organic soils corresponds to the total area of organic soils under Cropland and Grassland as reported in CRF-table 5.B and 5.C (see also chapter 7.2.3).

Emissions from animal production (4D2)

Calculation of emissions from animal production is based on AGRAMMON (Kupper et al. 2013). IPCC equation 4.18, IPCC 2000: p. 4.42 is used, but country specific N excretion rates and manure management system distribution fractions (MS) are used (refer to chapter

6.3.2). The relevant input data are based on Flisch et al. (2009) and calculated within the Swiss ammonium model AGRAMMON.

Only emissions of pasture, range and paddock are to be reported under agricultural soils. Other emissions from animal production are reported under Manure Management.

Indirect emissions (4D3)

Calculation of the indirect emissions is based on IPCC 2000 Tier 1b.

- For calculation of N_2O emissions from **leaching and run-off**, N-leaching from commercial fertilizers (including synthetic fertilizers, sewage sludge and compost) and animal manure has to be estimated. The relevant input data is based on Flisch et al. 2009, Prasuhn and Braun (1994), Braun et al. (1994) and Prasuhn and Mohni (2003). $Frac_{Leach}$ is set as 0.2 instead of the IPCC default of 0.3. This country specific value is extrapolated from long-term monitoring and modelling studies from the canton of Berne (Prasuhn and Mohni 2003) while the default value is based on a global model which assumes that 30% of nitrogen from synthetic fertilizer and atmospheric deposition is reaching water bodies. According to Schmid et al. (2000: p.71) this later amount is not representative for N-excretion of livestock animals in Switzerland and would lead to a significant overestimation.
- N_2O emissions from **deposition** are based on NH_3 and NO_x emissions. NH_3 -losses to the atmosphere are calculated according to the Swiss ammonium emission model AGRAMMON (Kupper et al. 2013). Input data for AGRAMMON for the years 1990 and 1995 are mainly based on expert judgements and literature studies whereas data for 2002, 2007 and 2010 include extensive farm surveys. Values in between the assessment years have been interpolated linearly while values beyond 2010 are kept constant until new survey results are available. For the calculation of NH_3 emissions changes of agricultural structures (changes to more animal friendly housing systems) and techniques (manure management, measures to reduce NH_3 emissions) are considered and explain temporal dynamics. Specific losses for all livestock categories are estimated. Ammonium volatilization of nitrogen in synthetic fertilizers is 15% for urea and 2% for other synthetic fertilizers. These estimates are based on a literature review by van der Weerden and Jarvis (1997) who examined ammonia emission factors for ammonium nitrate and urea for grassland and cropland soils. The emission factors for all other applied synthetic nitrogen (as straight and compound fertilizers) were assumed to be similar to that for ammonium nitrate. Ammonia emission factors for recycling fertilizers (sewage sludge and compost) are between 10 and 20% depending on the relative share of the individual fertilizer types (Kupper et al. 2013). Total $Frac_{GASF}^{14}$ has declined considerably due to a change in the shares of the different components that contribute to $Frac_{GASF}$ (weighted mean): the use of urea and sewage sludge (which both have high NH_3 emission factors) has been declining since 1990. Furthermore, volatilization of 2.0 kg NH_3 -N/ha agricultural soil is assumed due to processes in the vegetation cover (Kupper et al. 2013). Details about the amount of volatilized NH_3 are provided in the following table.

¹⁴ The concepts of $Frac_{GASF}$ and $Frac_{GASM}$ in the Swiss GHG-Inventory are not used as straight-forward as in the IPCC Guidelines and GPG. The Swiss Model applies two respective data sets: one for the estimation of nitrogen applied to agricultural soils and another used for the estimation of atmospheric nitrogen deposition. $Frac_{GASM}$ as reported in CRF-table 4.Ds2 represents the amount of nitrogen volatilized as NH_3 from housing, manure storage and manure application divided by the manure excreted in the stable. The nitrogen input from manure applied to soils in CRF-table 4.Ds1 can thus be calculated with the numbers given in CRF-table 4.B(b) and 4.Ds2. For the calculation of indirect N_2O emissions from atmospheric deposition all nitrogen volatilizations are considered including NO_x emissions and ammonia volatilization on pasture, range and paddock.

Table 6-16 Overview of Ammonia emission factors 1990–2010. Data source: Kupper et al. (2013)

| Ammonia Emission Factor | | 1990 | 1995 | 2002 | 2007 | 2010 |
|--------------------------------------|--|-------|-------|-------|-------|-------|
| | | % | | | | |
| Mature Dairy Cattle | | 35.12 | 35.25 | 32.74 | 33.69 | 33.33 |
| Mature Non-Dairy Cattle | | 31.47 | 32.08 | 28.17 | 33.63 | 31.24 |
| Young Cattle Average (weighted) | | 33.94 | 34.43 | 30.99 | 33.21 | 33.07 |
| | <i>Fattening Calves</i> | 39.58 | 39.97 | 38.66 | 40.83 | 40.54 |
| | <i>Pre-Weaned Calves</i> | 31.47 | 32.08 | 29.04 | 33.24 | 35.01 |
| | <i>Breeding Cattle 1st Year</i> | 33.94 | 34.45 | 31.05 | 33.86 | 33.80 |
| | <i>Breeding Cattle 2nd Year</i> | 30.23 | 30.63 | 27.14 | 29.33 | 29.47 |
| | <i>Breeding Cattle 3rd Year</i> | 32.09 | 32.55 | 28.46 | 31.17 | 31.10 |
| | <i>Fattening Cattle</i> | 41.31 | 41.46 | 40.14 | 39.61 | 38.11 |
| Sheep | | 21.20 | 21.21 | 20.32 | 19.14 | 20.51 |
| | <i>Fattening Sheep</i> | 21.08 | 21.13 | 20.31 | 18.91 | 20.35 |
| | <i>Milksheep</i> | 24.86 | 24.92 | 20.53 | 22.88 | 22.65 |
| Goats | | 24.64 | 24.71 | 24.32 | 24.88 | 24.25 |
| | <i>Goat Places</i> | 24.64 | 24.71 | 24.32 | 24.88 | 24.25 |
| Horses | | 25.77 | 25.84 | 22.12 | 23.36 | 21.90 |
| | <i>Horses <3 years</i> | 25.77 | 25.84 | 19.14 | 18.83 | 19.84 |
| | <i>Horses >3 years</i> | 25.77 | 25.84 | 22.77 | 24.22 | 22.22 |
| Mules and Asses | | 25.77 | 25.84 | 21.86 | 22.18 | 23.24 |
| | <i>Mules</i> | 25.77 | 25.84 | 21.86 | 22.18 | 23.24 |
| | <i>Asses</i> | 25.77 | 25.84 | 21.86 | 22.18 | 23.24 |
| Swine | | 38.26 | 38.87 | 45.01 | 45.34 | 44.40 |
| | <i>Piglets</i> | 38.26 | 38.39 | 38.24 | 39.94 | 43.37 |
| | <i>Fattening Pig over 25 kg</i> | 38.26 | 38.96 | 46.68 | 46.60 | 44.07 |
| | <i>Dry Sows</i> | 38.26 | 38.99 | 46.30 | 47.47 | 48.36 |
| | <i>Nursing Sows</i> | 38.26 | 38.39 | 38.94 | 40.27 | 41.69 |
| | <i>Boars</i> | 38.26 | 38.39 | 46.53 | 47.78 | 47.28 |
| Poultry | | 41.77 | 38.93 | 32.48 | 30.65 | 30.86 |
| | <i>Growers</i> | 48.76 | 46.29 | 44.96 | 35.39 | 31.95 |
| | <i>Layers</i> | 44.93 | 44.40 | 36.55 | 35.54 | 35.62 |
| | <i>Broilers</i> | 32.72 | 32.32 | 27.59 | 25.96 | 27.12 |
| | <i>Turkey</i> | 32.72 | 33.07 | 29.66 | 34.40 | 27.11 |
| | <i>Other Poultry (Geese, Ducks, Ostriches, Quails)</i> | 32.72 | 32.99 | 32.53 | 33.34 | 36.70 |
| Fertilizer | | 6.10 | 5.83 | 4.99 | 4.61 | 4.11 |
| | <i>Urea</i> | 15.00 | 15.00 | 15.00 | 15.00 | 15.00 |
| | <i>Other Mineral Fertilizers</i> | 2.00 | 2.00 | 2.00 | 2.00 | 2.00 |
| | <i>Recycling Fertilizers</i> | 17.58 | 19.74 | 18.03 | 12.32 | 9.58 |
| Other: Vegetation cover (kg/ha/year) | | 2.00 | 2.00 | 2.00 | 2.00 | 2.00 |

Other (sewage sludge and compost used for fertilizing) (4D4)

This source category covers N₂O emissions from sewage sludge and from compost used for fertilization. The calculation of the emissions corresponds to the one for synthetic fertilizer. Both direct soil emissions as well as indirect soil emissions (atmospherical deposition and leaching and runoff) are considered. Since 2003 the use of sewage sludge as fertilizer is prohibited in Switzerland. However, a transition period applies for some areas. Accordingly the cantons could prolong this period until 2008 in individual cases (UVEK 2003).

Activity data is based on Kupper et al. (2013) and has been consolidated by the responsible persons at the School of Agricultural, Forest and Food Science (HAFL, Kupper et al. 2013). Estimates are available for the years 1990, 1995, 2000, 2005, 2007 and 2010 and years in between have been interpolated linearly. Beyond 2010 constant values as in 2010 are used until further survey results are available.

NO_x emissions

NO_x emissions are estimated to be 0.7% of total nitrogen from animal manure and synthetic fertilizer, sewage sludge and compost. This factor is based on the CORINAIR Emission Inventory Guidebook 2003 (EEA 2007). Data on N-excretion (kg N/head/yr) is based on

Flisch et al. (2009) and calculated within AGRAMMON (Kupper et al. 2013). The amount of nitrogen from synthetic fertilizer, sewage sludge and compost is taken from Agricura (2012), Kupper et al. (2013) and SBV (2013).

NM VOC emissions

Estimation of NM VOC emissions of meadows and arable land is based on Spirig and Neftel (2002). VOC flows are estimated in Warneke et al. (2002) (for meadows) and König et al. (1995) (for arable land). Emissions were measured in a field trial in Austria (Karl et al. 2001).

Emission factors

The following IPCC default emission factors for calculating N₂O emissions from agricultural soils are used.

Table 6-17 Emission factors for calculating N₂O emissions from agricultural soils (IPCC 1997c: tables 4.18 (direct emissions), 4.22 (pasture, range and paddock) and 4.23 (indirect emissions); IPCC 2000: table 4.17 (organic soils).

| Emission Source | Emission factor |
|---|-----------------|
| Direct Emissions | |
| Synthetic fertilizer (kg N ₂ O-N/kg) | 0.0125 |
| Animal manure nitrogen used as fertilizer (kg N ₂ O-N/kg) | 0.0125 |
| Crop residue (kg N ₂ O-N/kg) | 0.0125 |
| N-fixing crops (kg N ₂ O-N/kg) | 0.0125 |
| Organic soils (kg N ₂ O-N/ha) | 8 |
| Residues meadows and pasture (kg N ₂ O-N/kg) | 0.0125 |
| N-fixing meadows and pasture (kg N ₂ O-N/kg) | 0.0125 |
| Animal production | |
| Pasture, range and paddock (kg N ₂ O-N/kg) | 0.0200 |
| Indirect emissions | |
| Leaching and run-off (kg N ₂ O-N/kg) | 0.0250 |
| Deposition (kg N ₂ O-N/kg) | 0.01 |
| Other | |
| Other (sewage sludge and compost used for fertilizing) (kg N ₂ O-N/kg) | 0.0125 |

Activity data

Activity data for calculation of direct soil emissions has been provided by ART/SHL (2012; animal livestock population), SBV (2013; use of synthetic fertilizer, crop yields, area of meadows and pasture), Agricura (2012; use of synthetic fertilizer), FAL/RAC (2001: p. 48/49), Schmid et al. (2000), Walther et al. (1994), Flisch et al. (2009) and Kupper et al. (2013).

Use of synthetic fertilizers in public green areas, sports grounds and home gardens (domestic synthetic fertilizer use) is 4% of all synthetic fertilizers (Kupper et al. 2013) and reported under 4D1.6 Other Direct Emissions.

The relevant activity data for calculating N₂O emissions from soils is displayed in the following table. Additional information is given in Table A - 29 and Table A - 30 in Annex A3.3.

Table 6-18 Activity data for calculating N₂O emissions from agricultural soils.

| Related Activity Data | | 1990-1999 | | | | | | | | | |
|----------------------------|--|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
| | | Value | | | | | | | | | |
| Direct Emissions | | | | | | | | | | | |
| Fertilizers (t N/yr) | Total commercial fertilizers | 70'744 | 71'611 | 71'543 | 66'869 | 63'272 | 63'399 | 61'050 | 53'460 | 53'581 | 55'667 |
| | Mineral fertilizer (t N/yr) | 66'096 | 66'877 | 66'724 | 61'964 | 58'283 | 58'326 | 56'213 | 48'855 | 49'207 | 51'521 |
| | Sewage sludge (t N/yr) | 3'852 | 3'834 | 3'816 | 3'797 | 3'778 | 3'758 | 3'497 | 3'239 | 2'984 | 2'731 |
| | Compost (t N/yr) | 796 | 900 | 1'003 | 1'107 | 1'211 | 1'315 | 1'340 | 1'365 | 1'390 | 1'416 |
| Animal manure | Nitrogen input from manure applied to soils (t N/yr) | 81'496 | 80'382 | 78'701 | 77'379 | 76'677 | 75'041 | 73'762 | 70'902 | 69'587 | 66'873 |
| N-fixation | N fixation peas, dry beans, soybeans and leguminous vegetables (t N/yr) | 681 | 774 | 909 | 796 | 811 | 861 | 931 | 1'131 | 1'107 | 1'049 |
| | N fixation meadows and pasture (t N/yr) | 29'027 | 29'325 | 29'728 | 29'602 | 28'913 | 30'270 | 30'645 | 30'862 | 30'868 | 30'852 |
| Crop residue | N from crop residues (t N/yr) | 11'335 | 11'170 | 11'053 | 11'249 | 10'634 | 10'838 | 12'145 | 11'742 | 11'803 | 10'555 |
| | N from residues meadows and pasture (t N/yr) | 21'473 | 21'574 | 21'713 | 21'677 | 21'461 | 21'903 | 22'032 | 22'080 | 22'156 | 22'069 |
| | Area of meadows and pasture (ha) | 784'867 | 788'089 | 792'338 | 791'387 | 785'006 | 798'550 | 802'514 | 803'722 | 807'945 | 805'131 |
| Organic soils | Area of cultivated organic soils (ha) | 18'493 | 18'458 | 18'423 | 18'391 | 18'353 | 18'314 | 18'276 | 18'241 | 18'205 | 18'170 |
| Animal production | | | | | | | | | | | |
| Pasture, range and paddock | N excretion on pasture range and paddock (t N/yr) | 13'148 | 13'399 | 13'498 | 13'421 | 13'691 | 13'833 | 15'615 | 16'805 | 18'040 | 18'800 |
| Indirect emissions | | | | | | | | | | | |
| Leaching and run-off | N excretion of all animals (t N/yr) | 144'593 | 143'184 | 140'768 | 138'724 | 138'066 | 135'756 | 135'750 | 132'553 | 131'987 | 128'452 |
| | Fertilizer (t N/yr) | 75'339 | 75'675 | 75'612 | 70'848 | 67'185 | 67'321 | 64'841 | 56'660 | 56'580 | 58'699 |
| Deposition | N from fertilizers and animal manure that is lost through leaching and run off (t N/yr) | 43'986 | 43'772 | 43'276 | 41'914 | 41'050 | 40'615 | 40'118 | 37'843 | 37'713 | 37'430 |
| | Emissions NH ₃ from fertilizers, animal manure and agricultural soils (tN/yr) | 57'331 | 56'274 | 55'455 | 54'723 | 54'449 | 53'655 | 53'113 | 51'045 | 50'426 | 48'908 |
| | Emissions NO _x from fertilizers and animal manure (t N/yr) | 1'540 | 1'532 | 1'515 | 1'467 | 1'437 | 1'422 | 1'404 | 1'324 | 1'320 | 1'310 |
| | Area of agricultural soils (ha) | 1'066'981 | 1'069'630 | 1'072'279 | 1'074'928 | 1'077'577 | 1'080'226 | 1'082'875 | 1'075'727 | 1'078'405 | 1'071'899 |

| Related Activity Data | | 2000-2009 | | | | | | | | | |
|----------------------------|--|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
| | | Value | | | | | | | | | |
| Direct Emissions | | | | | | | | | | | |
| Fertilizers (t N/yr) | Total commercial fertilizers | 54'824 | 58'660 | 57'102 | 54'666 | 54'749 | 53'587 | 52'708 | 54'859 | 52'121 | 49'542 |
| | Mineral fertilizer (t N/yr) | 50'903 | 54'896 | 53'496 | 51'217 | 51'458 | 50'454 | 49'559 | 51'694 | 48'886 | 46'220 |
| | Sewage sludge (t N/yr) | 2'481 | 2'169 | 1'858 | 1'546 | 1'235 | 923 | 779 | 635 | 423 | 212 |
| | Compost (t N/yr) | 1'441 | 1'595 | 1'749 | 1'903 | 2'056 | 2'210 | 2'370 | 2'530 | 2'812 | 3'110 |
| Animal manure | Nitrogen input from manure applied to soils (t N/yr) | 65'209 | 64'609 | 63'125 | 62'268 | 61'606 | 62'443 | 62'667 | 62'702 | 64'371 | 64'117 |
| N-fixation | N fixation peas, dry beans, soybeans and leguminous vegetables (t N/yr) | 840 | 734 | 1'144 | 1'256 | 1'322 | 1'179 | 1'107 | 1'093 | 1'069 | 1'028 |
| | N fixation meadows and pasture (t N/yr) | 30'817 | 31'120 | 31'143 | 31'485 | 31'623 | 31'089 | 31'204 | 31'639 | 31'671 | 31'872 |
| Crop residue | N from crop residues (t N/yr) | 11'887 | 10'386 | 11'446 | 9'737 | 11'814 | 11'513 | 10'529 | 11'508 | 11'492 | 11'874 |
| | N from residues meadows and pasture (t N/yr) | 22'055 | 22'217 | 22'220 | 22'321 | 22'334 | 22'174 | 22'199 | 22'267 | 22'249 | 22'269 |
| | Area of meadows and pasture (ha) | 806'369 | 809'441 | 809'597 | 812'624 | 812'370 | 807'793 | 808'416 | 809'187 | 808'300 | 807'927 |
| Organic soils | Area of cultivated organic soils (ha) | 18'134 | 18'098 | 18'063 | 18'027 | 17'991 | 17'970 | 17'937 | 17'904 | 17'883 | 17'853 |
| Animal production | | | | | | | | | | | |
| Pasture, range and paddock | N excretion on pasture range and paddock (t N/yr) | 20'921 | 22'643 | 24'166 | 23'858 | 23'432 | 23'555 | 23'719 | 23'718 | 23'868 | 23'402 |
| Indirect emissions | | | | | | | | | | | |
| Leaching and run-off | N excretion of all animals (t N/yr) | 128'095 | 129'120 | 128'423 | 127'016 | 125'842 | 127'797 | 128'785 | 129'136 | 131'539 | 130'195 |
| | Fertilizer (t N/yr) | 57'919 | 61'759 | 60'100 | 57'340 | 57'481 | 56'021 | 55'015 | 57'510 | 54'469 | 51'628 |
| Deposition | N from fertilizers and animal manure that is lost through leaching and run off (t N/yr) | 37'203 | 38'176 | 37'705 | 36'871 | 36'664 | 36'764 | 36'760 | 37'329 | 37'202 | 36'365 |
| | Emissions NH ₃ from fertilizers, animal manure and agricultural soils (tN/yr) | 48'265 | 48'259 | 47'499 | 46'918 | 46'869 | 47'582 | 48'067 | 48'724 | 48'999 | 48'077 |
| | Emissions NO _x from fertilizers and animal manure (t N/yr) | 1'302 | 1'336 | 1'320 | 1'290 | 1'283 | 1'287 | 1'287 | 1'307 | 1'302 | 1'273 |
| | Area of agricultural soils (ha) | 1'072'492 | 1'071'130 | 1'069'770 | 1'067'055 | 1'064'573 | 1'065'118 | 1'065'199 | 1'060'242 | 1'058'100 | 1'055'648 |

| Related Activity Data | | 2010-2012 | | |
|---------------------------------|--|-----------|-----------|-----------|
| | | 2010 | 2011 | 2012 |
| | | Value | | |
| Direct Emissions | | | | |
| Fertilizers (t N/yr) | Total commercial fertilizers | 56'849 | 50'464 | 48'954 |
| | Mineral fertilizer (t N/yr) | 53'425 | 47'040 | 45'529 |
| | Sewage sludge (t N/yr) | 0 | 0 | 0 |
| | Compost (t N/yr) | 3'424 | 3'424 | 3'424 |
| Animal manure | Nitrogen input from manure applied to soils (t N/yr) | 64'504 | 64'187 | 64'145 |
| N-fixation | N fixation peas, dry beans, soybeans and leguminous vegetables (t N/yr) | 1'037 | 1'096 | 1'014 |
| | N fixation meadows and pasture (t N/yr) | 31'983 | 32'164 | 32'117 |
| Crop residue | N from crop residues (t N/yr) | 10'442 | 12'178 | 11'251 |
| | N from residues meadows and pasture (t N/yr) | 22'266 | 22'332 | 22'299 |
| | Area of meadows and pasture (ha) | 807'226 | 809'513 | 808'355 |
| Organic soils | Area of cultivated organic soils (ha) | 17'822 | 17'790 | 17'759 |
| Animal production | | | | |
| Pasture, range and paddock | N excretion on pasture range and paddock (t N/yr) | 22'992 | 22'726 | 22'662 |
| Indirect emissions | | | | |
| Leaching and run-off | N excretion of all animals (t N/yr) | 129'900 | 129'089 | 128'871 |
| | Fertilizer (t N/yr) | 59'287 | 52'687 | 50'987 |
| Deposition | N from fertilizers and animal manure that is lost through leaching and run off (t N/yr) | 37'838 | 36'355 | 35'972 |
| | Emissions NH ₃ from fertilizers, animal manure and agricultural soils (tN/yr) | 48'122 | 47'666 | 47'361 |
| | Emissions NO _x from fertilizers and animal manure (t N/yr) | 1'324 | 1'272 | 1'259 |
| Area of agricultural soils (ha) | | 1'051'748 | 1'051'866 | 1'051'063 |

6.5.3 Uncertainties and Time-Series Consistency

For the uncertainty analysis the input data from ART (2008a) was used and was weighted with current activity and emission data. The arithmetic mean of the lower and upper bound uncertainty is used for activity data and for emission factors, resulting in the following

combined uncertainties for Tier 1 analysis: 4D1: 83%, 4D2: 85%, 4D3: 166% and 4D4 80%. To aggregate fertilizer and organic soils to a single category 4D1 and atmospheric deposition, leaching and run-off to 4D3 (as required for input into Tier 1 analysis), the combined uncertainty of the emissions is determined by using Tier 1 error propagation for the sub-systems.

For Tier 2 analysis the uncertainties are shown in Table 6-19. For further results see Section 1.7 and Annex 7.

Table 6-19 Combined uncertainties for N₂O emissions from 4D Agricultural Soils.

| Source categories 4D | | | gas | lower bound 2.5% (in %) | upper bound 97.5% (in %) | mean uncertainty % |
|----------------------|--|---------------------|------------------|----------------------------|-----------------------------|-----------------------|
| 4D1 | D. Agricultural Soils; Direct Soil Emissions | fertilizer | N ₂ O | -6 | 242 | 124 |
| 4D1 | | organic soils | N ₂ O | 41 | 171 | 65 |
| 4D2 | D. Agricultural Soils; Pasture, Range and Paddock Manure | | N ₂ O | 40 | 151 | 55 |
| 4D3 | D. Agricultural Soils; Indirect Emissions | deposition | N ₂ O | 37 | 182 | 72 |
| 4D3 | | leaching and runoff | N ₂ O | 39 | 224 | 92 |
| 4D4 | D. Agricultural Soils; Sewage sludge and compost | | N ₂ O | 20 | 180 | 80 |

For details on time-series consistency see Chapter 6.2.3 and 6.3.3.

6.5.4 Source-Specific QA/QC and Verification

All QA/QC activities are further described in a separate document (ART 2013a). General information on agricultural structures and policies is provided and eventual differences between national and (IPCC) standard values are being analysed and discussed.

An internal documentation of the Agroscope Research Station about the calculation of the greenhouse gas emissions in agriculture assures transparency and traceability of the calculation methods (Berthoud 2004). IULIA is described in Schmid et al. (2000) and the Swiss ammonium emission model AGRAMMON is documented in Kupper et al. (2013) and Agrammon 2010.

All relevant data needed for the calculation of direct and indirect nitrogen inputs to agricultural soils (e.g. F_{SN} , MS-distribution, $Frac_{GASF}$, N_{ex} , $Frac_{GASM}$, F_{BN} , F_{CR}) have been checked for consistency and confidence and have been compared (where possible) to IPCC default values, values of other countries as well as literature values. As one of the most important parameters, nitrogen excretion has been analysed in more detail as described in Chapter 6.3.4.

For quality of livestock population data consult Chapter 6.2.4.

The implied emission factors have been compared to the corresponding emission factors of other countries (UNFCCC <http://unfccc.int/di/FlexibleQueries.do>) (INFRAS 2012). Additionally, N₂O emission factors have been compared to literature values to assure plausibility. Implied emission factors are generally in line with measurements representative for Swiss conditions (ART 2013a).

The estimate for cultivated histosols in the agricultural sector is consistent with the estimates reported under Cropland and Grassland in the LULUCF sector. A literature study conducted by Leifeld et al. (2003) comes up with an estimate of 17'000 ± 5'000 ha which is close to the numbers reported in the LULUCF sector (18'100 ha on average).

The time series of activity data and emission factors have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the following triple check:

- the results for 2012 are compared with the results for 2011 within the current CRF

- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of the submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of the submission 2013

Additionally an independent quality control was conducted by INFRAS by a countercheck of the data and calculation sheets elaborated by Agroscope (Agroscope 2014).

6.5.5 Source-Specific Recalculations

For recalculation of livestock numbers see chapter 6.2.5.

For recalculation of the nitrogen excretion rate of mature dairy cattle of the year 2011 see chapter 6.3.5.

The area of cultivated organic soils has been revised due to new projections in the LULUCF-sector. Average absolute difference in area of organic soils are only 0.03% and the mean change of overall greenhouse gas emissions is below 0.02 Gg CO₂ equivalent.

Activity data for compost for the years 2008-2011 has been revised due to an error correction. New emission estimates are lower by approximately 7-8 Gg CO₂ equivalent.

The ammonia emission factor for recycling fertilizers (liquid digestate) has been adjusted for the years 2008 - 2011 due to the increasing use of trailing hoses during land application as fertilizer. Due to the low quantities of applied nitrogen from this source, impact on Frac_{GASF} and on overall emissions is negligible.

A general recalculation for the year 2011 has been carried out due to some updates of crop yield data from the Swiss Farmers Union (SBV 2012). The respective changes are only of minor importance for total emission estimates.

6.5.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented. Other projects are eventually postponed in order to give first priority to changes related to the new reporting guidelines.

6.6 Source Category 4E – Burning of savannahs

Burning of savannahs does not occur (NO) in Switzerland.

6.7 Source Category 4F – Field Burning of Agricultural Residues

Source category 4F Field Burning of Agricultural Residues has been moved to sector 6 Waste. Emissions from open burning of branches in agriculture and forestry have been reported here in the past. However, since branches in agriculture and forestry are burned only after they have been translocated from their place of origin, they should be reported under sector 6 Waste in accordance to the EMEP guidebook 2009 (EEA 2010). Accordingly the source category has been moved to 6C1 and is now reported under 6.C Waste Incineration, a. biogenic.

7 LULUCF

7.1 Overview of LULUCF

7.1.1 Methodology

Chapter 7 presents estimates of greenhouse gas emissions by sources and removals by sinks from land use, land-use change and forestry (LULUCF). Data acquisition and calculations are based on the Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC 2003) and are completed by country specific methodologies.

The land areas in the period 1990-2012 are represented by geographically explicit land-use data with a resolution of one hectare (following approach 3 for representing land areas; IPCC 2003). Direct and repeated assessment of land use with full spatial coverage also enables to calculate spatially explicit land-use change matrices. In 2004, the Swiss Land Use Statistics AREA was launched. Simultaneously, aerial photos from two earlier Swiss Land Use Statistics (1979/85 and 1992/97) were re-evaluated, applying the same approach. At the editorial deadline AREA had been completed and the interpretation of the entire Swiss territory was available for three time slices.

The six main land-use categories required by IPCC (2003) are: A. Forest Land, B. Cropland, C. Grassland, D. Wetlands, E. Settlements and F. Other Land. These categories were divided in 18 sub-divisions of land use. A further spatial stratification reflects the criteria "altitude" (3 zones), "geomorphologic and climatic conditions" (adopting the five production regions of the National Forest Inventory; NFI) and "soil type" (mineral, organic).

Country specific emission factors and carbon stocks for Forest Land were derived from four Swiss National Forest Inventories (NFI 1 – NFI 4a+), which had been finalised in 1985, 1995, 2006 and 2012, respectively. The inventories comprehended ca. 2'600 (2012), 6'000 (1995, 2006) and 12'000 (1985) terrestrial sampling plots (see Table 7-11), where biomass stock, growth, harvesting and mortality had been measured.

For the remaining land-use categories, carbon stocks and GHG emissions/removals were derived from particular research activities, domestic surveys and measurements in the fields of agriculture (cropland, grassland) and nature conservation (wetlands). Partially, also IPCC default values and expert estimates were used.

7.1.2 Emissions and Removals

Table 7-1 and Figure 7-1 summarize the CO₂ emissions and removals as a result of carbon losses and gains for the years 1990-2012. The total net emissions and removals of CO₂ from 1990 to 2012 vary between -3'331 Gg (1993) and 1'266 Gg (2001).

In Table 7-1 and Figure 7-1, four components of the CO₂ balance are differentiated:

- Gains in carbon stock of living biomass on all land uses and due to land-use changes; it represents the largest sink of carbon.
- Losses in carbon stock of living biomass on all land uses and due to land-use changes; it represents the largest source of carbon. The highest losses are observed in the years following a heavy storm with windfall in December 1999.
- Net carbon stock changes in dead organic matter (DOM) on forest land remaining forest land as well as on forest land converted to non-forest land: it represents a sink of carbon in most years from 1990-2006. Since 2007 this item is a source.
- Balance of carbon emissions and removals (1) in soils due to the use of soils (especially of organic soils) and due to land-use changes, (2) by agricultural lime application, and (3)

by wildfires. In the period under investigation this accumulative component persistently represents a source of carbon.

In forests, growth of biomass exceeds the harvesting and mortality rate except for the years 2000-2002. Compared to CO₂ fluxes involved in forest biomass dynamics, the net CO₂ emissions arising from the use of soils, from agricultural lime application, wildfires, and from all land-use changes are relatively small. Overall, the LULUCF sector was a sink of -1'702 Gg CO₂ on the average between 1990 and 2012 (see Table 7-1 and Figure 7-2).

Table 7-1 Switzerland's CO₂ emissions and removals (Gg) of category 5 Land Use, Land-Use Change and Forestry 1990-2012. Positive values refer to emissions; negative values refer to removals. In this data set, emissions of CH₄ and N₂O are not included. Land-use changes include Afforestation and Deforestation.

| LULUCF | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------------|--------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| | Gg CO ₂ | | | | | | | | | |
| Gains of living biomass | -12'690 | -12'627 | -12'772 | -12'725 | -12'665 | -12'913 | -12'962 | -12'906 | -12'552 | -12'562 |
| Losses of living biomass | 9'886 | 10'054 | 10'018 | 8'903 | 9'392 | 9'032 | 9'829 | 9'703 | 10'008 | 10'777 |
| Net change in dead organic matter | 136 | -123 | -74 | -193 | 141 | 9 | -326 | -787 | -1'154 | -1'098 |
| Net change in soil, liming, wildfires | 706 | 683 | 680 | 683 | 701 | 708 | 707 | 733 | 719 | 713 |
| Total Sector 5: LULUCF | -1'962 | -2'013 | -2'147 | -3'331 | -2'432 | -3'164 | -2'752 | -3'257 | -2'979 | -2'169 |

| LULUCF | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------------|--------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| | Gg CO ₂ | | | | | | | | | |
| Gains of living biomass | -12'981 | -12'579 | -12'635 | -12'597 | -13'303 | -12'718 | -12'846 | -13'599 | -13'383 | -13'419 |
| Losses of living biomass | 13'306 | 14'002 | 13'543 | 11'272 | 10'743 | 11'166 | 11'105 | 11'071 | 11'057 | 10'783 |
| Net change in dead organic matter | -1'041 | -868 | -667 | -838 | -832 | -1'131 | -662 | 22 | 856 | 1'143 |
| Net change in soil, liming, wildfires | 712 | 711 | 718 | 721 | 709 | 730 | 730 | 720 | 681 | 667 |
| Total Sector 5: LULUCF | -3 | 1'266 | 961 | -1'441 | -2'683 | -1'953 | -1'672 | -1'785 | -789 | -827 |

| LULUCF | 2010 | 2011 | 2012 | Mean |
|---------------------------------------|--------------------|---------|---------|---------|
| | Gg CO ₂ | | | |
| Gains of living biomass | -13'348 | -13'995 | -13'361 | -12'962 |
| Losses of living biomass | 10'865 | 10'539 | 10'733 | 10'773 |
| Net change in dead organic matter | 859 | 863 | 809 | -215 |
| Net change in soil, liming, wildfires | 677 | 680 | 677 | 703 |
| Total Sector 5: LULUCF | -948 | -1'914 | -1'142 | -1'702 |




Figure 7-1 (i) CO₂ removals due to the gain (growth) of living biomass, (ii) CO₂ emissions due to the loss (harvest and mortality) of living biomass, (iii) net CO₂ emissions and removals due to changes in dead organic matter, and (iv) net CO₂ emissions from soils, due to liming and wildfires, 1990–2012.

The non-CO₂ emissions associated with land use, land-use change and forestry are very small. Between 1990 and 2012 annual CH₄ emissions add up to less than 1.81 Gg, and annual N₂O emissions equal at maximum 0.04 Gg. Those emissions arise from flooded land/reservoirs (CH₄; CRF-table 5(II)), soil disturbance associated with land-conversion to cropland (N₂O; CRF-table 5(III)), and wildfires on forest land (CH₄ and N₂O; CRF-table 5(V)). The calculation methods are based on default procedures of IPCC (2003; Chapter 3) and are summarized in Chapters 7.3.4.13, 7.4.4.4 and 7.6.4.4, respectively.

Figure 7-2 shows the resulting net GHG balances of LULUCF 1990–2012 including all CO₂ and non-CO₂ fluxes. Further representations of LULUCF CO₂ eq data can be found in Chapter 2 “Trends in Greenhouse Gas Emissions and Removals”.




Figure 7-2 Switzerland's net GHG balance of category 5 Land Use, Land-Use Change and Forestry 1990–2012 (in Gg CO₂ eq). Positive values refer to emissions, negative values refer to removals.

7.1.3 Approach for Calculating Carbon Emissions and Removals

7.1.3.1 Work Steps

The selected procedure for calculating carbon emissions and removals in the LULUCF sector corresponds to a Tier 2 approach as described in IPCC (2003; Chapter 3). It can be summarised as follows:

- Define land use categories and sub-divisions with respect to available land-use data (see Table 7-2). For the present study, so-called combination categories (CC) were defined on the basis of the AREA land-use and land-cover categories (Table 7-6; SFSO 2006a).
- Define criteria and collect data for the spatial stratification of the land-use categories.
- Measure or estimate the carbon stocks and carbon stock changes for each spatial stratum of the land-use categories.
- Calculate the land use and the land-use change matrix in each spatial stratum.
- Calculate the carbon stock changes in living biomass (ΔC_l), in dead organic matter (ΔC_d) and in soil (ΔC_s) for all cells of the land-use change matrix.
- Finally, aggregate the results by summarising the carbon stock changes over land-use categories and strata according to the level of disaggregation displayed in the CRF-tables.

Table 7-2 Land-use categories used in this report (so-called combination categories CC): 6 main land-use categories (identical to the UNFCCC land-use categories) and 18 sub-divisions. Additionally, descriptive remarks, abbreviations used in the CRF-tables, and CC codes are given. For a detailed definition of the combination categories see Table 7-6 (FOEN 2007f) and SFSO (2006a).

| CC Main category | CC Sub-division | Remarks | Terminology in CRF tables | CC code |
|------------------|--|--|---------------------------|---------|
| A. Forest Land | Afforestations | areas converted to forest by active measures, e.g. planting | afforestation | 11 |
| | Productive Forest | dense and open forest meeting the criteria of forest land | productive | 12 |
| | Unproductive Forest | brush forest and forest on unproductive areas meeting the criteria of forest land | unproductive | 13 |
| B. Cropland | | arable and tillage land (annual crops and leys in arable rotations) | | 21 |
| C. Grassland | Permanent Grassland | meadows, pastures (low-land and alpine) | permanent | 31 |
| | Shrub Vegetation | agricultural and unproductive areas predominantly covered by shrubs | woody, shrub | 32 |
| | Vineyards, Low-Stem Orchards, Tree Nurseries | perennial agricultural plants with woody biomass (no trees) | woody, vine | 33 |
| | Copse | agricultural and unproductive areas covered by perennial woody biomass including trees | woody, copse | 34 |
| | Orchards | permanent grassland with fruit trees | woody, orchard | 35 |
| | Stony Grassland | grass, herbs and shrubs on stony surfaces | unproductive, stony | 36 |
| | Unproductive Grassland | unmanaged grass vegetation | unproductive | 37 |
| D. Wetlands | Surface Waters | lakes and rivers | surface | 41 |
| | Unproductive Wetland | reed, unmanaged wetland | unproductive | 42 |
| E. Settlements | Buildings and Constructions | areas without vegetation such as houses, roads, construction sites, dumps | building | 51 |
| | Herbaceous Biomass in Settlements | areas with low vegetation, e.g. lawns | herb | 52 |
| | Shrubs in Settlements | areas with perennial woody biomass (no trees) | shrub | 53 |
| | Trees in Settlements | areas with perennial woody biomass including trees | tree | 54 |
| F. Other Land | | areas without soil and vegetation: rocks, sand, scree, glaciers | | 61 |

The sub-categories listed in Table 7-2 were defined with respect to optimal distinction of biomass and soil carbon contents and dynamics (see Chapter 7.2.2.2, Table 7-6). The underlying criteria to include land-use sub-categories such as Shrub vegetation, Vineyards, Low-stem Orchards, Tree Nurseries, Copse and Orchards under Grassland with woody biomass are: (1) They do not fulfil the criteria for forests; (2) There is an agricultural management in general; (3) They all have woody biomass (i.e. perennial vegetation) with grass understory. Under Cropland there are no perennial crops, but only annual crops and leys in arable rotations. All perennial crops are included in the grassland sub-categories.

7.1.3.2 Calculating Carbon Stock Changes

For calculating carbon stock changes, the following input parameters (mean values per hectare) must be quantified for all combination categories (CC) and spatial strata (i):

| | |
|--------------------|---|
| stock $C_{l,i,CC}$ | carbon stock in living biomass ($t\ C\ ha^{-1}$) |
| stock $C_{d,i,CC}$ | carbon stock in dead organic matter ($t\ C\ ha^{-1}$) |
| stock $C_{s,i,CC}$ | carbon stock in soil ($t\ C\ ha^{-1}$) |
| gain $C_{l,i,CC}$ | annual gain (growth) of carbon in living biomass ($t\ C\ ha^{-1}\ yr^{-1}$) |
| loss $C_{l,i,CC}$ | annual loss (harvesting and mortality) of carbon in living |

| | |
|---------------------------|--|
| | biomass (t C ha ⁻¹ yr ⁻¹) |
| changeC _{d,i,CC} | annual net carbon stock change in dead organic matter (t C ha ⁻¹ yr ⁻¹) |
| changeC _{s,i,CC} | annual net carbon stock change in soil (t C ha ⁻¹ yr ⁻¹) |

For the reporting under the UNFCCC, the carbon content of litter (organic soil horizons) is included in stockC_{d,i,CC}. For the reporting under the Kyoto Protocol, litter is calculated and reported separately (see Chapter 11.3.1.1).

On this basis, the total carbon fluxes (t C yr⁻¹) in living biomass (deltaC_l), in dead organic matter (deltaC_d) and in soil (deltaC_s) are calculated for all cells of the land-use change matrix. Each cell is characterized by a land-use category before the reporting year (b), a land-use category at the end of the reporting year (a), and the area of converted land within the spatial stratum (i). It includes the case without any land-use change (a = b).

Equations 7.1.-7.6 show, according to the IPCC good practice guidance (GPG) for LULUCF (IPCC 2003), two approaches and their application for calculating carbon emissions and removals: (1) the gain-loss approach (GPG Equation 3.1.1) and (2) the stock-change approach (GPG Equation 3.1.2).

The gain-loss approach takes into account the net carbon stock changes in living biomass (l), dead organic matter (d) and soils (s) of solely the land-use category after the conversion. The gain-loss approach is also used in cases of no land-use change. The stock-change approach takes into account the stock changes due to conversion of land use (difference of the stocks before and after the conversion).

The gain-loss approach is defined as:

$$\text{deltaC}_{l,i,ba} = (\text{gainC}_{l,i,a} - \text{lossC}_{l,i,a}) * A_{i,ba} \quad (7.1)$$

$$\text{deltaC}_{d,i,ba} = \text{changeC}_{d,i,a} * A_{i,ba} \quad (7.2)$$

$$\text{deltaC}_{s,i,ba} = \text{changeC}_{s,i,a} * A_{i,ba} \quad (7.3)$$

The formulation of the stock-change approach is:

$$\text{deltaC}_{l,i,ba} = [(\text{stockC}_{l,i,a} - \text{stockC}_{l,i,b}) / \text{CT}] * A_{i,ba} \quad (7.4)$$

$$\text{deltaC}_{d,i,ba} = [(\text{stockC}_{d,i,a} - \text{stockC}_{d,i,b}) / \text{CT}] * A_{i,ba} \quad (7.5)$$

$$\text{deltaC}_{s,i,ba} = [(\text{stockC}_{s,i,a} - \text{stockC}_{s,i,b}) / \text{CT}] * A_{i,ba} \quad (7.6)$$

where:

| | |
|-------------------|---|
| a | land-use category after conversion (CC = a) |
| b | land-use category before conversion (CC = b) |
| ba | land use conversion from b to a |
| i | spatial stratum |
| A _{i,ba} | area of land (ha) converted from b to a in the spatial stratum i (activity data from the land-use change matrix) |
| CT | conversion time (yr), see Chapter 7.1.3.3 |

Table 7-3 pinpoints which approach is used for calculating the carbon fluxes for the various types of land-use conversion and carbon pools (living biomass, dead organic matter and soil): The gain-loss approach is generally used for smooth transitions, e.g. the growth of living biomass on land converted to forest land; the stock-change approach is used for quick changes (e.g. loss of biomass by deforestation, CT = 1 year) as well as slow processes such as the change in soil carbon content (CT = 20 years, see Chapter 7.1.3.3).

For the conversions between different forest categories (e.g. CC12 to CC13 and CC13 to CC12) the method is chosen in such a way that no potential carbon losses are underestimated.

In case of land-use changes involving "Buildings and Constructions" (CC51), 50% of the difference between carbon stocks before and after the conversion is reported as a source or sink, respectively; for a detailed documentation see Chapter 7.7.4.2 and Chapter 11.3.1.1.

Table 7-3 Calculation approach (gain-loss or stock-change) and conversion time periods (CT, years) applied for different land-use transitions and carbon pools. KP = corresponding activity under the Kyoto Protocol; NF = non-forest category. Combination categories CC11 to CC61 are introduced in Table 7-2.

| Change in main land-use category or sub-division | Living biomass | UNFCCC: Dead organic matter KP: Deadwood, Litter | Soil | Remarks |
|--|------------------|--|------------------|--|
| No change in category KP and UNFCCC | gain-loss | gain-loss | gain-loss | |
| CC13 to CC12 UNFCCC: 5A1 KP: forest management | gain-loss | stock-change, 20 | stock-change, 20 | |
| CC12 to CC13 UNFCCC: 5A1 KP: forest management | stock-change, 20 | stock-change, 20 | stock-change, 20 | |
| CC11 to CC12 UNFCCC: 5A1 KP: afforestation >20 years | gain-loss | gain-loss | gain-loss | |
| Change to CC11 UNFCCC: 5A2 KP: afforestation ≤20 years | gain-loss | stock-change, 20 | stock-change, 20 | Dead organic matter is 0 in CC11 and in NF; directly human-induced |
| NF to CC12/CC13 UNFCCC: 5A2 KP: forest management | gain-loss | stock-change, 20 | stock-change, 20 | |
| Change to CC51 UNFCCC: 5E2 KP: deforestation | stock-change, 1 | stock-change, 1 | stock-change, 20 | Buildings/constructions; soil: carbon stock reduced by 50% |
| Change to CC52-54 UNFCCC: 5E2 | stock-change, 1 | stock-change, 1 | stock-change, 20 | Green settlement areas |
| Change to CC21 UNFCCC: 5B2 | stock-change, 1 | stock-change, 1 | stock-change, 20 | Cropland |
| Change to CC31-37 UNFCCC: 5C2 | stock-change, 1 | stock-change, 1 | stock-change, 20 | Grassland |
| Change to CC41 UNFCCC: 5D2 | stock-change, 1 | stock-change, 1 | stock-change, 1 | Surface water |
| Change to CC42 UNFCCC: 5D2 | stock-change, 1 | stock-change, 1 | stock-change, 20 | Unproductive wetland |
| Change to CC61 UNFCCC: 5F2 | stock-change, 1 | stock-change, 1 | stock-change, 20 | Other land |

7.1.3.3 Considering the Conversion Time (CT)

Changes in the soil carbon stock – this is also true for the increase of woody biomass – as a result of land-use changes are slow processes that might take decades. Therefore, IPCC (2003) suggests implementing a conversion time (CT). Following the IPCC default value (CT = 20 years), the carbon emission or removal due to a soil carbon stock difference ($\text{stockC}_{s,i,a} - \text{stockC}_{s,i,b}$) does not occur in one year but is distributed evenly over the 20 years following the land-use conversion.

A conversion time of 20 years has been applied to all soil carbon stock changes (except land converted to surface water). Accordingly, the CRF-tables 5A2, 5B2, 5C2, 5D2, 5E2 and 5F2 contain the cumulative area remaining in the respective category in the reporting year.

In addition, the default conversion time of 20 years has been assumed for carbon stock changes in biomass (living and dead) for land converted from productive forest to unproductive forest (CC12 to CC13).

The land-use category Afforestations (CC11) is inherently a transitional category by definition in the land-use survey. Areas converted to afforestations are reported in the CRF-table 5A2 with the same conversion time as for other forest sub-categories (20 years). However, afforestations remaining afforestations (according to the land-use survey) are reported in CRF-table 5A1 and are merged with source category Productive Forest (CC12) after having been reported 20 years under land converted to forest land.

Table 7-3 summarises the conversion times applied to carbon stock changes in living biomass, in dead organic matter, and in soils for all types of land-use transitions.

There is no consistent data on land-use changes before 1990, but it is well known (ARE/SAEFL 2001, FOEN 2013g) that the main trends of the Swiss land-use dynamics (e.g. increase of forests and settlements) did arise before 1970. Therefore, it was assumed that between 1971 and 1989 the annual rate of all land-use changes was the same as in 1990. Based on this assumption it has been possible to produce the land-use data required for the consideration of the conversion time in that period.

7.1.3.4 Displaying Results in the Common Reporting Format (CRF)

In the CRF-tables 5A to 5F, a part of the combination categories (CC) and associated spatial strata are shown at an aggregated level for optimal documentation and overview. The values of ΔC are accordingly summarised. Positive values of $\Delta C_{l,i,ba}$ are inserted in the column "Gains" and negative values in the column "Losses", respectively. The values of $\Delta C_{d,i,ba}$ and $\Delta C_{s,i,ba}$ are inserted in column "Net carbon stock change in dead organic matter" and "Net carbon stock change in soils", respectively.

The CRF-tables 5A to 5F are subdivided in two parts: (1) X Land remaining X Land and (2) Land converted to X Land. Unchanged areas as well as changes occurring from one combination category to another belonging to the same main land-use category are reported in the first part of the CRF. For example, the area of "shrub vegetation" (CC32) converted to "permanent grassland" (CC31) is reported in CRF-table 5C1 in the sub-division "permanent" in the respective altitude zone. As CC31 and CC32 do have different carbon stocks in biomass, a carbon stock change is calculated according to the equations presented in Chapter 7.1.3.2.

7.1.4 Carbon Stocks, Emission Factors, and Net Changes at a Glance

Table 7-4 lists all values of carbon stocks, gains, losses and net changes of carbon specified for combination category (CC) and associated spatial strata for the year 1990. These values remain constant during the period 1990-2012 with the following exceptions (highlighted cells):

- Carbon stock, gain and loss of living biomass, carbon stock and net change in dead organic matter as well as net change in mineral soils of productive forest (CC12): Deduction and values of the annually changing data of CC12 are described in Chapters 7.3.4.6, 7.3.4.7 and 7.3.4.8.
- Carbon stock, gain and loss of living biomass of cropland (CC21): Annual data of CC21 are listed in Chapter 7.4.4.

The deduction of the individual carbon stocks and emission factors is explained in detail in the Chapters 7.3 to 7.8.

Table 7-4 Carbon stocks and changes in living biomass, in dead organic matter and in soils for the combination categories (CC), disaggregated for altitude, NFI region, and soil type. The values are valid for the period 1990-2012 with the exception of the values in the highlighted cells, which change annually (numbers given here are for the year 1990); cf. main text.

| land-use code CC | NFI region | altitude zone z | carbon stock in living biomass (stockCl,i) | carbon stock in dead org. matter (stockCd,i) | carbon stock in mineral soil (stockCs,i) | carbon stock in organic soil (stockCs,i) | gain of living biomass (gainCl,i) | loss of living biomass (lossCl,i) | net change in dead org. matter (changeCd,i) | net change in mineral soil (changeCs,i) | net change in organic soil (changeCs,i) |
|------------------------|------------|-----------------|--|--|--|--|--|-----------------------------------|---|---|---|
| | Strata | | [t C ha ⁻¹] | | | | [t C ha ⁻¹ yr ⁻¹] | | | | |
| 11 Afforestations | 1 | 1 | 7.84 | 0 | 82.65 | 240 | 1.63 | 0 | 0 | 0 | -0.68 |
| | 1 | 2 | 4.30 | 0 | 102.03 | 240 | 1.09 | 0 | 0 | 0 | -0.68 |
| | 1 | 3 | 1.61 | 0 | 121.34 | 240 | 0.57 | 0 | 0 | 0 | -0.68 |
| | 2 | 1 | 7.84 | 0 | 55.40 | 240 | 1.63 | 0 | 0 | 0 | -0.68 |
| | 2 | 2 | 4.30 | 0 | 62.12 | 240 | 1.09 | 0 | 0 | 0 | -0.68 |
| | 2 | 3 | 1.61 | 0 | 122.00 | 240 | 0.57 | 0 | 0 | 0 | -0.68 |
| | 3 | 1 | 7.84 | 0 | 66.10 | 240 | 1.63 | 0 | 0 | 0 | -0.68 |
| | 3 | 2 | 4.30 | 0 | 75.91 | 240 | 1.09 | 0 | 0 | 0 | -0.68 |
| | 3 | 3 | 1.61 | 0 | 95.78 | 240 | 0.57 | 0 | 0 | 0 | -0.68 |
| | 4 | 1 | 7.84 | 0 | 66.47 | 240 | 1.63 | 0 | 0 | 0 | -0.68 |
| | 4 | 2 | 4.30 | 0 | 74.39 | 240 | 1.09 | 0 | 0 | 0 | -0.68 |
| | 4 | 3 | 1.61 | 0 | 69.48 | 240 | 0.57 | 0 | 0 | 0 | -0.68 |
| | 5 | 1 | 7.84 | 0 | 102.37 | 240 | 1.63 | 0 | 0 | 0 | -0.68 |
| | 5 | 2 | 4.30 | 0 | 108.99 | 240 | 1.09 | 0 | 0 | 0 | -0.68 |
| | 5 | 3 | 1.61 | 0 | 107.08 | 240 | 0.57 | 0 | 0 | 0 | -0.68 |
| 12 Productive forest | 1 | 1 | 126.87 | 14.95 | 82.65 | 240 | 3.60 | -2.41 | -0.01 | 0.00 | -0.68 |
| | 1 | 2 | 124.88 | 14.32 | 102.03 | 240 | 3.21 | -2.27 | -0.15 | 0.00 | -0.68 |
| | 1 | 3 | 84.73 | 13.66 | 121.34 | 240 | 1.95 | -1.34 | -0.10 | 0.00 | -0.68 |
| | 2 | 1 | 134.18 | 17.82 | 55.40 | 240 | 4.63 | -4.13 | -0.16 | 0.00 | -0.68 |
| | 2 | 2 | 146.77 | 20.33 | 62.12 | 240 | 4.63 | -3.93 | -0.04 | 0.00 | -0.68 |
| | 2 | 3 | 101.21 | 20.33 | 122.00 | 240 | 1.60 | -0.86 | -0.04 | 0.00 | -0.68 |
| | 3 | 1 | 135.06 | 18.29 | 66.10 | 240 | 4.56 | -3.04 | 0.83 | 0.00 | -0.68 |
| | 3 | 2 | 147.43 | 24.23 | 75.91 | 240 | 4.15 | -3.06 | -0.09 | 0.00 | -0.68 |
| | 3 | 3 | 119.32 | 35.36 | 95.78 | 240 | 2.48 | -2.11 | -0.38 | 0.00 | -0.68 |
| | 4 | 1 | 94.81 | 11.46 | 66.47 | 240 | 3.24 | -2.71 | 0.59 | 0.00 | -0.68 |
| | 4 | 2 | 104.42 | 27.27 | 74.39 | 240 | 2.49 | -1.81 | -0.07 | 0.00 | -0.68 |
| | 4 | 3 | 96.41 | 41.36 | 69.48 | 240 | 1.81 | -1.62 | -0.05 | 0.00 | -0.68 |
| | 5 | 1 | 70.67 | 11.05 | 102.37 | 240 | 2.74 | -1.01 | -0.39 | 0.00 | -0.68 |
| | 5 | 2 | 76.70 | 13.95 | 108.99 | 240 | 2.20 | -0.71 | -0.33 | 0.00 | -0.68 |
| | 5 | 3 | 76.70 | 33.60 | 107.08 | 240 | 1.61 | -0.48 | -0.03 | 0.00 | -0.68 |
| 13 Unproductive forest | 1 | 1 | 45.90 | 9.51 | 82.65 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 1 | 2 | 48.20 | 7.53 | 102.03 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 1 | 3 | 48.03 | 7.76 | 121.34 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 2 | 1 | 46.64 | 8.70 | 55.40 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 2 | 2 | 45.90 | 11.42 | 62.12 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 2 | 3 | 12.86 | 11.42 | 122.00 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 3 | 1 | 45.90 | 7.51 | 66.10 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 3 | 2 | 47.68 | 16.29 | 75.91 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 3 | 3 | 29.08 | 26.21 | 95.78 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 4 | 1 | 40.47 | 3.15 | 66.47 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 4 | 2 | 38.37 | 19.99 | 74.39 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 4 | 3 | 18.58 | 33.37 | 69.48 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 5 | 1 | 38.59 | 8.22 | 102.37 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 5 | 2 | 33.46 | 11.03 | 108.99 | 240 | 0 | 0 | 0 | 0 | -0.68 |
| | 5 | 3 | 21.14 | 30.77 | 107.08 | 240 | 0 | 0 | 0 | 0 | -0.68 |

(Table 7-4 continued)

| | | | | | | | | | | | |
|-----------------------------|------|------|-------|---|-------|-----|------|------|---|---|-------|
| 21 Cropland | n.s. | n.s. | 4.34 | 0 | 53.40 | 240 | 0.05 | 0.00 | 0 | 0 | -9.52 |
| 31 Permanent Grassland | n.s. | 1 | 7.08 | 0 | 62.02 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| | n.s. | 2 | 6.00 | 0 | 67.50 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| | n.s. | 3 | 7.95 | 0 | 75.18 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| 32 Shrub Vegetation | n.s. | 1 | 12.90 | 0 | 62.02 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| | n.s. | 2 | 12.90 | 0 | 67.50 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| | n.s. | 3 | 12.90 | 0 | 75.18 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 33 Vineyards et al. | n.s. | n.s. | 3.74 | 0 | 53.40 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| 34 Copse | n.s. | 1 | 12.90 | 0 | 62.02 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| | n.s. | 2 | 12.90 | 0 | 67.50 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| | n.s. | 3 | 12.90 | 0 | 75.18 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 35 Orchards | n.s. | n.s. | 24.32 | 0 | 64.76 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| 36 Stony Grassland | n.s. | n.s. | 4.52 | 0 | 26.31 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 37 Unproductive Grassland | n.s. | n.s. | 7.01 | 0 | 68.23 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 41 Surface Waters | n.s. | n.s. | 0 | 0 | 0 | 240 | 0 | 0 | 0 | 0 | 0 |
| 42 Unproductive Wetland | n.s. | n.s. | 6.50 | 0 | 68.23 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 51 Buildings, Constructions | n.s. | n.s. | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 52 Herbaceous Biomass in S. | n.s. | n.s. | 9.54 | 0 | 53.40 | 240 | 0 | 0 | 0 | 0 | -9.52 |
| 53 Shrubs in Settlements | n.s. | n.s. | 15.43 | 0 | 53.40 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 54 Trees in Settlements | n.s. | n.s. | 20.72 | 0 | 53.40 | 240 | 0 | 0 | 0 | 0 | -5.3 |
| 61 Other Land | n.s. | n.s. | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Legend*altitude zones:*

- 1 < 601 m
- 2 601 - 1200 m
- 3 > 1200 m

NFI-regions:

- 1 Jura
- 2 Central Plateau
- 3 Pre-Alps
- 4 Alps
- 5 Southern Alps

n.s. = no stratification

annually changing data

7.1.5 Uncertainty Estimates

Table 7-5 gives an overview of uncertainty estimates of activity data (AD) and of emission factors (EF). In most cases (highlighted in yellow; reasons for exceptions are indicated in column "Remark"), the uncertainty of AD depends on the quality of the AREA survey data.

In general, AD uncertainty is lower than EF uncertainty, because AD are based on a systematic survey with high spatial resolution (AREA, see Chapter 7.2), while EF include parameters that are difficult to measure or model such as carbon stocks in biomass, growth rates and other biological processes.

Uncertainty estimates of AD are presented in Chapter 7.2.5, while uncertainty estimates of EF are presented in detail in the respective chapters (7.X.5) of the LULUCF source categories.

Table 7-5 Uncertainty estimates in the LULUCF sector, expressed as half of the 95% confidence intervals.

| IPCC category | | Gas | Activity data uncertainty | Emission factor uncertainty | Remark |
|---------------|--------------------------------------|------------------|---------------------------|-----------------------------|--------------|
| | | | % | % | |
| 5A1 | 1. Forest Land remaining Forest Land | CO ₂ | 2 | 63 | |
| 5A2 | 2. Land converted to Forest Land | CO ₂ | 2 | 63 | |
| 5B1 | 1. Cropland remaining Cropland | CO ₂ | 5 | 110 | |
| 5B2 | 2. Land converted to Cropland | CO ₂ | 6 | 143 | |
| 5B2 | 2. Land converted to Cropland | N ₂ O | 6 | 90 | |
| 5C1 | 1. Grassland remaining Grassland | CO ₂ | 6 | 2084 | |
| 5C2 | 2. Land converted to Grassland | CO ₂ | 6 | 67 | |
| 5D1 | 1. Wetlands remaining Wetlands | CO ₂ | 30 | 100 | organic soil |
| 5D2 | 2. Land converted to Wetlands | CO ₂ | 5 | 50 | |
| 5E1 | 1. Settlements remaining Settlements | CO ₂ | 5 | 50 | |
| 5E2 | 2. Land converted to Settlements | CO ₂ | 5 | 50 | |
| 5F2 | 2. Land converted to Other Land | CO ₂ | 4 | 50 | |
| 5(IV) | Agricultural lime application | CO ₂ | 40 | 25 | |
| 5(V) | Forest Land | CO ₂ | 10 | 70 | wildfire |
| 5(V) | Forest Land and Grasland | CH ₄ | 10 | 70 | wildfire |
| 5(V) | Forest Land and Grasland | N ₂ O | 10 | 70 | wildfire |

7.2 Activity Data – Land Areas

7.2.1 Description

Chapter 7.2 presents information related to activity data that is valid for all LULUCF categories, including information on land-use databases, approaches used for representing land areas, classification systems, uncertainties of land-use data as well as land-use related QA/QC, recalculations and planned improvements. The chapter, hence, is structured in a similar way as the subsequent category-specific Chapters 7.3 – 7.8.

7.2.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for Inventory Preparation

7.2.2.1 Swiss Land Use Statistics (AREA)

Data of the Swiss Land Use Statistics (AREA) evaluated by the Swiss Federal Statistical Office (SFSO 2013) are the basis of activity data. In the course of the AREA surveys, every hectare of Switzerland's territory (4'128 kha) was assigned to one of 46 land-use categories and to one of 27 land-cover categories by means of stereographic interpretation of aerial photos (SFSO 2006a). The AREA surveys were launched in 2004 and completed in 2013.

For the reconstruction of the land use conditions in Switzerland during the period 1990-2012 three datasets are used:

- Land Use Statistics "1979/85" (AREA1)
- Land Use Statistics "1992/97" (AREA2)
- Land Use Statistics "2004/09" (AREA3)

The aerial photos for AREA1, AREA2 and AREA3 were taken 1977-1986, 1990-1998 and 2004-2009, respectively. They were simultaneously evaluated according to the newly designed AREA set of land-use and land-cover categories based on the nomenclature 'NOAS04' (SFSO 2006a).

The inter-survey period is not identical throughout the Swiss territory, but varies regionally. It averages approximately 12 years. This methodical characteristic needs to be considered when reconstructing the annual country-wide status of land use or when calculating annual rates of land-use change.

7.2.2.2 Combination Categories (CC) as derived from AREA Land Use Statistics

The 46 land-use categories and 27 land-cover categories of AREA were aggregated to 18 combination categories (CC) as shown in Table 7-6 (FOEN 2007f), thus implementing the main categories proposed by IPCC as well as country specific sub-divisions (see Table 7-2). The sub-divisions were defined with respect to optimal distinction of biomass densities, carbon turnover, and soil carbon contents.

The first digit of the CC code represents the land-use category according to IPCC, whereas the second digit stands for sub-divisions of the land-use categories.

15 April 2014

[illegible]

7.2.2.3 Interpolation of the Status for each Year

The exact dates of aerial photo shootings are known for each hectare. However, the exact occurrence date (year) of a land-use change on a specific hectare is unknown. The actual change can have taken place in any year between two AREA surveys. In this study, it is assumed that the probability of a land-use change from AREA1 to AREA2 and from AREA2 to AREA3 is uniformly distributed over the respective interim period between two surveys. Therefore, the land-use change of each hectare has to be equally distributed over its specific interim period.

Thus, the land-use status for the years between two data collection dates can be calculated by linear interpolation. Dates of aerial photo shootings (i.e. starting and ending year of the inter-survey period) and the land-use categories of AREA1, AREA2 and AREA3 for every hectare are used for these calculations. An example is shown in Figure 7-3: A hectare has been assigned to the land-use category Cropland in AREA1 (aerial photo in 1980). A land-use change to Surrounding of Buildings has been discovered 10 years later (1990) in AREA2.




Figure 7-3 Hypothetical linear development of land-use changes between AREA1, AREA2 and AREA3 considering as example a hectare changing from “Cropland” to “Surrounding of Buildings” and then from “Surrounding of Buildings” to “Buildings”. For 2009, a linear interpolation has been carried out between AREA3 and a virtual fourth survey modelled for the year 2016 (here resulting in no change of land use).

The “state 1988” of that hectare is determined by calculating the fractions of the two land-use categories for the year 1988. A linear development from “Cropland” to “Surrounding of Buildings” during the whole interim period is assumed. Thus, in 1988 the hectare is split up in two fractions: 80% is “Surrounding of Buildings” and 20% is “Cropland”. The same procedure can be applied for two survey dates between AREA2 and AREA3 (here exemplarily shown for the period 1990-2004, highlighting “state 2003”).

AREA3 comprehends aerial photos from six years (2004-2009). Therefore, the land-use changes occurring after AREA3 are calculated from the linear development detected between AREA3 and a virtual fourth survey, AREA4 (see Figure 7-3: example “state 2009”). AREA4 was modeled for each sample plot using a Markov-chain approach, where transition probabilities between AREA3 and AREA4 were assessed based on transition distribution

between AREA2 and AREA3 within each spatial stratum (Sigmaplan 2014). This approach was evaluated successfully by modeling a virtual AREA3 from transition probabilities between AREA1 and AREA2 and comparing the results to the actual interpretation of AREA3.

The status for each individual year in the period 1990-2012 for the whole Swiss territory results from the summation of the fractions of all hectares per combination category CC, additionally considering the spatial strata where appropriate.

7.2.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

7.2.3.1 Spatial Stratification

In order to quantify carbon stocks and GHG emissions by sources and removals by sinks in the LULUCF sector as accurately as possible, Switzerland's territory was stratified by means of three site criteria: soil type (mineral or organic), altitude and forest production region.

Most soils in Switzerland are mineral soil types. For mapping the occurrence of organic soils, two datasets were used: (i) the digital soil map "BEK" (SFSO 2000a) and (ii) the Inventory of Raised Bogs of National Importance (Appendix to Swiss Confederation 1991a).

Two units of the digital soil map contain mainly organic soils (Figure 7-4): The codes F1 and Q3, representing Histosols in the Central Plateau and in Alpine valleys, respectively, are good indicators for organic soils in the lowlands. As the soil map has no appropriate unit for organic soils in mountainous areas the maps of the Inventory of Raised Bogs (with a scale of 1:25'000) were used in addition. All areas covered by this inventory were assumed to have organic soils (see Figure 7-4).

For Forest Land and – in part – Grassland, three altitudinal belts were differentiated: <601 m a.s.l. (meters above sea level), 601-1200 m a.s.l., and >1200 m a.s.l. (Figure 7-4). Altitude data are available on a hectare-grid from the Swiss Federal Statistical Office (SFSO 1997).

Forest Land was furthermore differentiated into the five production regions of the National Forest Inventory NFI (EAFV/BFL 1988; Brassel and Brändli 1999; Brändli 2010). The NFI regions were adopted from EAFV/BFL (1988) as shown in Figure 7-4:

1. Jura
2. Central Plateau
3. Pre-Alps
4. Alps
5. Southern Alps.

Applying all spatial stratifications, 30 different strata (referred to as subscript *i* in Chapter 7.1.3.2) would be theoretically possible. Not all of them, but altogether 28 have been actually realised and applied for the calculation of LULUCF-associated carbon emissions and removals.




Figure 7-4 Map showing the spatial stratification according to NFI region, altitude, and soil type.

7.2.3.2 The Land-use Tables and Change Matrices

In Table 7-7 the land-use statistics resulting from spatial stratification (Chapter 7.2.3.1) and interpolation in time (Chapter 7.2.2.3) are exemplarily shown for the year 1990. This table gives also an overview of the size of the individual spatial strata. The combination categories (CC) have been introduced in Table 7-2.

Table 7-7 Land use (in terms of combination categories CC) projection by the end of 1990, stratified separately for altitude (3 zones), soil type (mineral or organic) and NFI region (1-5), in kha.

| CC: | 11 | 12 | 13 | 21 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 41 | 42 | 51 | 52 | 53 | 54 | 61 | Sum |
|-------------------|-----|--------|------|-------|-------|-------|------|-------|-----|-------|------|-------|------|-------|------|-----|------|-------|--------|
| Altitude | | | | | | | | | | | | | | | | | | | |
| <601 | 1.1 | 222.7 | 0.5 | 299.5 | 152.0 | 2.6 | 22.7 | 37.8 | 1.2 | 0.5 | 2.9 | 137.5 | 5.2 | 116.2 | 47.4 | 2.8 | 18.6 | 1.9 | 1072.6 |
| 601-1200 | 1.5 | 501.4 | 8.0 | 132.3 | 360.0 | 8.6 | 3.8 | 40.1 | 0.4 | 2.4 | 1.5 | 10.0 | 5.6 | 47.0 | 17.1 | 1.0 | 5.4 | 7.7 | 1153.8 |
| >1200 | 1.4 | 382.8 | 77.1 | 0.4 | 425.4 | 144.5 | 0.0 | 30.0 | 0.0 | 148.8 | 61.9 | 13.5 | 14.4 | 11.4 | 3.7 | 0.2 | 1.0 | 585.6 | 1902.0 |
| | 4.0 | 1106.8 | 85.6 | 432.2 | 937.4 | 155.7 | 26.5 | 107.8 | 1.6 | 151.6 | 66.3 | 160.9 | 25.1 | 174.5 | 68.2 | 3.9 | 24.9 | 595.2 | 4128.4 |
| Soil | | | | | | | | | | | | | | | | | | | |
| mineral | 4.0 | 1103.5 | 85.6 | 420.4 | 931.8 | 155.6 | 26.4 | 107.3 | 1.6 | 151.6 | 66.0 | 160.4 | 21.9 | 172.7 | 67.5 | 3.9 | 24.8 | 595.2 | 4100.1 |
| organic | 0.0 | 3.3 | 0.1 | 11.8 | 5.6 | 0.1 | 0.1 | 0.5 | 0.0 | 0.0 | 0.3 | 0.5 | 3.2 | 1.8 | 0.7 | 0.1 | 0.1 | 0.027 | 28.3 |
| | 4.0 | 1106.8 | 85.6 | 432.2 | 937.4 | 155.7 | 26.5 | 107.8 | 1.6 | 151.6 | 66.3 | 160.9 | 25.1 | 174.5 | 68.2 | 3.9 | 24.9 | 595.2 | 4128.4 |
| NFI-region | | | | | | | | | | | | | | | | | | | |
| 1 | 0.7 | 197.2 | 5.3 | 78.0 | 122.6 | 0.9 | 4.7 | 14.8 | 0.3 | 0.2 | 0.6 | 23.6 | 1.2 | 26.8 | 10.9 | 0.5 | 4.7 | 0.5 | 493.5 |
| 2 | 0.8 | 227.1 | 0.4 | 306.9 | 152.4 | 0.9 | 9.9 | 31.1 | 1.0 | 0.2 | 1.6 | 70.3 | 4.1 | 84.9 | 34.7 | 1.6 | 12.6 | 0.7 | 941.2 |
| 3 | 1.0 | 214.3 | 9.2 | 30.4 | 261.3 | 10.4 | 0.8 | 21.7 | 0.1 | 8.5 | 6.8 | 30.2 | 12.0 | 26.8 | 9.2 | 0.5 | 2.9 | 15.0 | 661.0 |
| 4 | 1.2 | 331.6 | 49.5 | 13.8 | 365.4 | 110.2 | 9.5 | 31.0 | 0.2 | 118.0 | 49.2 | 26.2 | 7.2 | 26.9 | 9.8 | 0.8 | 3.0 | 524.7 | 1678.1 |
| 5 | 0.3 | 136.6 | 21.2 | 3.0 | 35.7 | 33.3 | 1.5 | 9.2 | 0.0 | 24.6 | 8.1 | 10.7 | 0.7 | 9.2 | 3.7 | 0.6 | 1.9 | 54.2 | 354.6 |
| | 4.0 | 1106.8 | 85.6 | 432.2 | 937.4 | 155.7 | 26.5 | 107.8 | 1.6 | 151.6 | 66.3 | 160.9 | 25.1 | 174.5 | 68.2 | 3.9 | 24.9 | 595.2 | 4128.4 |

Table 7-8 shows the overall trends of land-use changes between 1990 and 2012. For example, the area of afforestations (CC11) decreased by 76% during this period, while the area of unproductive forests (CC13) increased by 7%. CC11 is decreasing because the area

of new afforestations has been decreasing during this period and because most of the afforestation areas develop to productive forests after a certain time period.

Table 7-8 Statistics of land use (in terms of combination categories CC) and relative change (%) between 1990 and 2012, in kha.

| CC: | 11 | 12 | 13 | 21 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 41 | 42 | 51 | 52 | 53 | 54 | 61 | Sum |
|---------|-----|--------|------|-------|-------|-------|------|-------|-----|-------|------|-------|------|-------|------|-----|------|-------|--------|
| Year: | | | | | | | | | | | | | | | | | | | |
| 1990 | 4.0 | 1106.8 | 85.6 | 432.2 | 937.4 | 155.7 | 26.5 | 107.8 | 1.6 | 151.6 | 66.3 | 160.9 | 25.1 | 174.5 | 68.2 | 3.9 | 24.9 | 595.2 | 4128.4 |
| 1991 | 3.9 | 1109.0 | 86.0 | 431.4 | 935.5 | 155.2 | 26.5 | 106.6 | 1.5 | 151.4 | 66.1 | 160.9 | 25.1 | 176.2 | 68.7 | 4.0 | 25.3 | 594.9 | 4128.4 |
| 1992 | 3.8 | 1111.2 | 86.4 | 430.6 | 933.7 | 154.7 | 26.6 | 105.4 | 1.4 | 151.1 | 66.0 | 160.9 | 25.1 | 177.9 | 69.3 | 4.0 | 25.6 | 594.5 | 4128.4 |
| 1993 | 3.7 | 1113.3 | 86.8 | 429.8 | 932.1 | 154.2 | 26.5 | 104.2 | 1.4 | 150.9 | 65.8 | 160.9 | 25.1 | 179.5 | 69.8 | 4.1 | 25.9 | 594.2 | 4128.4 |
| 1994 | 3.5 | 1115.3 | 87.1 | 428.6 | 931.1 | 153.6 | 26.5 | 103.1 | 1.3 | 150.7 | 65.7 | 161.0 | 25.1 | 181.1 | 70.4 | 4.1 | 26.1 | 594.0 | 4128.4 |
| 1995 | 3.3 | 1117.1 | 87.5 | 427.0 | 930.7 | 153.0 | 26.5 | 102.0 | 1.3 | 150.5 | 65.6 | 161.0 | 25.2 | 182.7 | 71.1 | 4.2 | 26.1 | 593.7 | 4128.4 |
| 1996 | 3.1 | 1118.6 | 87.8 | 425.4 | 930.5 | 152.5 | 26.4 | 101.0 | 1.3 | 150.4 | 65.5 | 161.0 | 25.2 | 184.2 | 71.8 | 4.2 | 26.1 | 593.4 | 4128.4 |
| 1997 | 2.9 | 1120.1 | 88.1 | 423.7 | 930.5 | 152.0 | 26.3 | 99.9 | 1.3 | 150.3 | 65.4 | 161.0 | 25.2 | 185.7 | 72.6 | 4.2 | 25.9 | 593.1 | 4128.4 |
| 1998 | 2.7 | 1121.4 | 88.4 | 421.9 | 930.5 | 151.8 | 26.2 | 98.9 | 1.2 | 150.2 | 65.3 | 161.1 | 25.2 | 187.3 | 73.4 | 4.2 | 25.8 | 592.9 | 4128.4 |
| 1999 | 2.5 | 1122.7 | 88.6 | 420.2 | 930.5 | 151.5 | 26.1 | 97.8 | 1.2 | 150.3 | 65.2 | 161.1 | 25.2 | 188.8 | 74.3 | 4.2 | 25.6 | 592.6 | 4128.4 |
| 2000 | 2.2 | 1123.9 | 88.9 | 418.5 | 930.4 | 151.3 | 26.1 | 96.7 | 1.2 | 150.3 | 65.1 | 161.1 | 25.2 | 190.3 | 75.1 | 4.2 | 25.5 | 592.4 | 4128.4 |
| 2001 | 2.0 | 1125.2 | 89.2 | 416.8 | 930.4 | 151.0 | 26.0 | 95.6 | 1.1 | 150.3 | 65.0 | 161.2 | 25.2 | 191.9 | 75.9 | 4.2 | 25.3 | 592.2 | 4128.4 |
| 2002 | 1.8 | 1126.5 | 89.4 | 415.0 | 930.4 | 150.8 | 25.9 | 94.6 | 1.1 | 150.3 | 64.9 | 161.2 | 25.3 | 193.4 | 76.7 | 4.2 | 25.2 | 591.9 | 4128.4 |
| 2003 | 1.6 | 1127.8 | 89.7 | 413.3 | 930.3 | 150.6 | 25.8 | 93.5 | 1.1 | 150.3 | 64.8 | 161.2 | 25.3 | 194.9 | 77.5 | 4.2 | 25.0 | 591.7 | 4128.4 |
| 2004 | 1.3 | 1129.1 | 89.9 | 411.6 | 930.3 | 150.3 | 25.7 | 92.4 | 1.0 | 150.3 | 64.6 | 161.3 | 25.3 | 196.5 | 78.3 | 4.2 | 24.8 | 591.4 | 4128.4 |
| 2005 | 1.2 | 1131.1 | 90.2 | 411.1 | 930.4 | 149.9 | 25.5 | 90.7 | 1.0 | 150.2 | 64.4 | 161.3 | 25.3 | 197.9 | 78.8 | 4.1 | 24.3 | 591.0 | 4128.4 |
| 2006 | 1.1 | 1133.1 | 90.5 | 411.0 | 930.5 | 149.6 | 25.2 | 88.9 | 1.0 | 150.2 | 64.2 | 161.4 | 25.3 | 199.1 | 79.1 | 4.0 | 23.8 | 590.5 | 4128.4 |
| 2007 | 1.1 | 1135.0 | 90.6 | 411.2 | 930.5 | 149.3 | 25.0 | 87.1 | 0.9 | 150.1 | 63.9 | 161.5 | 25.2 | 200.1 | 79.4 | 3.9 | 23.3 | 590.0 | 4128.4 |
| 2008 | 1.0 | 1137.2 | 90.9 | 411.4 | 930.9 | 148.7 | 24.8 | 85.6 | 0.9 | 150.1 | 63.7 | 161.6 | 25.2 | 200.8 | 79.5 | 3.8 | 22.9 | 589.3 | 4128.4 |
| 2009 | 1.0 | 1138.8 | 91.2 | 409.9 | 930.9 | 148.3 | 24.7 | 84.5 | 0.9 | 150.1 | 63.5 | 161.7 | 25.2 | 202.2 | 80.2 | 3.9 | 22.7 | 588.9 | 4128.4 |
| 2010 | 1.0 | 1139.8 | 91.4 | 408.4 | 930.6 | 148.1 | 24.6 | 83.6 | 0.9 | 150.1 | 63.4 | 161.7 | 25.2 | 203.7 | 80.9 | 3.9 | 22.6 | 588.7 | 4128.4 |
| 2011 | 1.0 | 1140.7 | 91.6 | 406.8 | 930.3 | 147.9 | 24.5 | 82.8 | 0.8 | 150.1 | 63.3 | 161.7 | 25.3 | 205.1 | 81.5 | 3.9 | 22.5 | 588.4 | 4128.4 |
| 2012 | 1.0 | 1141.7 | 91.8 | 405.2 | 930.0 | 147.7 | 24.4 | 82.0 | 0.8 | 150.1 | 63.2 | 161.8 | 25.3 | 206.6 | 82.2 | 3.9 | 22.4 | 588.2 | 4128.4 |
| Change: | -76 | 3 | 7 | -6 | -1 | -5 | -8 | -24 | -48 | -1 | -5 | 0 | 1 | 18 | 21 | -1 | -10 | -1 | 0 |

The annual rates of change in the entire territory of Switzerland (change-matrices, Table 7-9) are achieved by adding up the annual change rates of all hectares per combination category (CC). For calculating the carbon stock changes, fully stratified (up to 28 strata, cf. Chapter 7.2.3.1) land-use change tables are used for each year (Meteotest 2014).

It is worth noting that in general the numbers given in the tables above cannot be directly compared with the numbers reported in the CRF-tables: The CRF-tables 5A2–5F2 contain the cumulative area remaining in the respective category in the reporting year. As described in Chapter 7.1.3.3, a conversion time of 20 years is applied to those land-use transitions and during the conversion time, the converted areas are reported under CRF-tables 5X2. In contrast, the change matrices present the land-use changes occurring in the specified year alone.

Table 7-9 Annual rates of land-use change in 1990 and in 2012 (change matrices). Units: ha/year, rounded values. Empty cells indicate that no change has occurred.

| 1990 | | change to CC | | | | | | | | | | | | | | | | | | | |
|----------------|----------|--------------|------|------|------|------|------|-----|------|----|-----|-----|-----|----|------|------|-----|-----|-----|----------|--|
| | | 11 | 12 | 13 | 21 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 41 | 42 | 51 | 52 | 53 | 54 | 61 | decrease | |
| change from CC | 11 | | 369 | 1 | 0 | 0 | 0 | | 0 | | | | 41 | | 0 | 0 | | | 0 | 372 | |
| | 12 | | | 135 | 5 | 125 | 86 | 6 | 82 | | 12 | 19 | 11 | 7 | 117 | 27 | 11 | 17 | 49 | 709 | |
| | 13 | | 534 | | | 128 | 37 | 0 | 45 | | 3 | 2 | 1 | 1 | 5 | 0 | 0 | 1 | 9 | 767 | |
| | 21 | 8 | 1 | | | 663 | 6 | 181 | 40 | 1 | 4 | 4 | 4 | 4 | 632 | 317 | 21 | 18 | 22 | 1926 | |
| | 31 | 136 | 166 | 231 | 718 | | 1007 | 123 | 560 | 4 | 46 | 43 | 9 | 11 | 870 | 490 | 27 | 44 | 68 | 4554 | |
| | 32 | 24 | 1022 | 687 | 2 | 126 | | 9 | 337 | | 14 | 14 | 6 | 0 | 24 | 8 | 5 | 3 | 30 | 2312 | |
| | 33 | 1 | 2 | | 126 | 65 | 4 | | 33 | 2 | 0 | 1 | 0 | | 50 | 26 | 4 | 3 | 5 | 323 | |
| | 34 | 30 | 680 | 33 | 151 | 1091 | 60 | 40 | | 11 | 10 | 24 | 4 | 4 | 207 | 114 | 8 | 54 | 15 | 2537 | |
| | 35 | | 0 | | 8 | 13 | 0 | 4 | 47 | | | | | | 4 | 2 | 0 | 0 | 0 | 80 | |
| | 36 | 3 | 27 | 25 | 2 | 162 | 243 | 1 | 41 | | | 89 | 4 | 0 | 8 | 1 | 0 | | 45 | 652 | |
| | 37 | 7 | 26 | 6 | 1 | 8 | 234 | 1 | 68 | | 10 | | 3 | 0 | 6 | 1 | | 0 | 13 | 384 | |
| | 41 | 0 | 4 | 1 | 2 | 2 | 6 | 0 | 4 | | 4 | 1 | | 17 | 11 | 2 | 1 | 0 | 99 | 156 | |
| | 42 | 5 | 27 | 5 | 1 | 3 | 2 | 0 | 3 | | 0 | 0 | 6 | | 4 | 1 | 0 | 0 | 1 | 59 | |
| | 51 | 38 | 18 | 1 | 86 | 158 | 11 | 5 | 11 | | 3 | 5 | 6 | 4 | | 271 | 58 | 46 | 5 | 726 | |
| | 52 | 7 | 4 | | 16 | 32 | 3 | 1 | 2 | | 0 | 1 | 1 | 2 | 349 | | 68 | 387 | 0 | 874 | |
| | 53 | 5 | 9 | 0 | 6 | 7 | 2 | 0 | 2 | | | | 0 | 2 | 45 | 28 | | 46 | 0 | 150 | |
| | 54 | 2 | 6 | | 1 | 2 | 0 | 0 | 3 | | | 0 | 0 | 1 | 78 | 152 | 8 | | 0 | 253 | |
| | 61 | 4 | 41 | 16 | 16 | 67 | 93 | 8 | 32 | | 287 | 33 | 96 | 2 | 13 | 1 | 0 | 1 | | 709 | |
| | increase | 271 | 2936 | 1141 | 1140 | 2652 | 1794 | 381 | 1310 | 18 | 394 | 236 | 152 | 55 | 2425 | 1443 | 211 | 621 | 362 | 17543 | |

| 2012 | | change to CC | | | | | | | | | | | | | | | | | | | |
|----------------|----------|--------------|------|-----|-----|------|------|-----|-----|----|-----|-----|-----|----|------|------|-----|-----|-----|----------|----|
| | | 11 | 12 | 13 | 21 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 41 | 42 | 51 | 52 | 53 | 54 | 61 | decrease | |
| change from CC | 11 | | 71 | 0 | | | 0 | | | | | | | | 0 | | | | | 72 | |
| | 12 | | | 224 | 1 | 178 | 140 | 2 | 107 | | 31 | 21 | 15 | 12 | 90 | 26 | 12 | 9 | 73 | 943 | |
| | 13 | | 515 | | | 158 | 58 | | 27 | | 4 | 2 | 1 | 1 | 3 | 0 | 0 | 0 | 10 | 779 | |
| | 21 | 2 | 0 | | | 1287 | 5 | 140 | 17 | 0 | 4 | 11 | 8 | 9 | 496 | 286 | 13 | 5 | 12 | 2294 | |
| | 31 | 17 | 77 | 176 | 444 | | 725 | 73 | 319 | 2 | 71 | 30 | 7 | 9 | 756 | 403 | 14 | 9 | 81 | 3215 | |
| | 32 | 3 | 646 | 510 | 2 | 127 | | 3 | 288 | | 18 | 11 | 5 | 0 | 13 | 4 | 2 | 1 | 32 | 1664 | |
| | 33 | 0 | 1 | | 135 | 95 | 5 | | 20 | 1 | 1 | 0 | | 0 | 35 | 23 | 1 | 2 | 6 | 326 | |
| | 34 | 3 | 523 | 31 | 53 | 770 | 67 | 15 | | 4 | 10 | 23 | 6 | 1 | 126 | 70 | 3 | 22 | 18 | 1745 | |
| | 35 | | | | 1 | 6 | | 2 | 16 | | | | | | 1 | 0 | | | | 26 | |
| | 36 | 0 | 17 | 20 | 3 | 80 | 195 | 1 | 44 | | | 51 | 4 | | 3 | 0 | | | 40 | 460 | |
| | 37 | 2 | 14 | 3 | 1 | 2 | 181 | | 47 | | 13 | | 3 | 0 | 4 | 1 | | | 12 | 283 | |
| | 41 | 0 | 2 | 0 | 0 | 1 | 5 | | 2 | | 3 | 3 | | 9 | 6 | 1 | 0 | | 100 | 133 | |
| | 42 | 0 | 18 | 4 | | | 0 | 0 | | 1 | | 1 | | 7 | | 2 | 0 | | | 1 | 36 |
| | 51 | 17 | 9 | 0 | 63 | 144 | 8 | 3 | 6 | | 6 | 6 | 6 | 2 | | 287 | 51 | 21 | 5 | 635 | |
| | 52 | 6 | 3 | 0 | 16 | 41 | 3 | 1 | 3 | | 1 | 2 | 1 | 2 | 411 | | 51 | 216 | 0 | 757 | |
| | 53 | 2 | 10 | | 3 | 11 | 2 | 0 | 1 | | 0 | 1 | 0 | 0 | 49 | 39 | | 38 | 0 | 156 | |
| | 54 | 1 | 3 | | 0 | 2 | 0 | | 2 | | | 0 | | 0 | 96 | 302 | 18 | | | 425 | |
| | 61 | 1 | 28 | 11 | 16 | 46 | 73 | 4 | 28 | | 285 | 16 | 101 | 1 | 6 | 1 | 0 | | | 618 | |
| | increase | 55 | 1938 | 981 | 740 | 2947 | 1467 | 244 | 928 | 6 | 449 | 177 | 164 | 49 | 2098 | 1445 | 165 | 323 | 391 | 14568 | |

7.2.4 Methodological Issues

No further remarks.

7.2.5 Uncertainties and Time-series Consistency of Activity Data

An overview of uncertainty estimates for activity data (AD) and emission factors (or biomass parameters) is shown in Table 7-5. Details related to uncertainties of AD are presented in this chapter, while uncertainties of the emission factors are presented in the respective chapters (7.X.5) of the LULUCF source categories.

In most cases (as highlighted in yellow in Table 7-5), the uncertainty of AD depends on the quality of the AREA survey data. However, in the following cases the uncertainty is determined mainly by other parameters:

- CO₂ emissions of category 5D1 (Wetland remaining wetland) are due to net carbon stock losses in organic soils. The uncertainty of the area of organic soils is around 30% according to Leifeld et al. (2003: 61).

- CO₂ emissions of category 5(IV): Agricultural lime application. The uncertainty of the amount of lime (40%) was estimated based on a poll among the main producers in Switzerland (Agroscope 2014a; see Chapter 7.4.4.5).
- CO₂, CH₄ and N₂O emissions from category 5(V) are due to wildfires on Forest Land and Grassland. The burnt area is surveyed by cantonal authorities. An uncertainty of 10% is assumed as it is a complete survey and not a sampling approach.

The uncertainty of AREA-based activity data has two main sources (Table 7-10). They have been quantified on the basis of the AREA data (SFSO 2013) as follows:

1) Interpretation error: In the AREA survey, the interpretation of the aerial photos is checked by a second independent interpreter. The portion of sampling points with a mismatch of the first and the second interpretation is used as the uncertainty of the interpretation. This uncertainty of interpretation integrates all errors related to the manual interpretation of land-use and land-cover classes on aerial photographs. While it is clear that this is rather an estimate of the maximum potential interpretation error than of the actual interpretation error, it is reported hereafter unless more accurate information is available.

2) Statistical sampling error: In the AREA survey, the land-use types are interpreted on points situated on a regular 100x100 m grid. Thus, the uncertainty of the surface area covered by a certain land-use type or land-use change decreases with increasing numbers of sampling points. Assuming a binomial distribution of the errors, this uncertainty is calculated as

$$U_{\text{sampling}} = 100 * 1.96 * (\text{number of points})^{-0.5}$$

The number of sampling points lies between 2'472 (for 5D2) and 1'374'367 (for 5C1) leading to values of U_{sampling} between 3.9% and 0.2%.

The overall uncertainty was calculated as:

$$U_{\text{overall}} = (U_{\text{interpret}}^2 + U_{\text{sampling}}^2)^{0.5}$$

Finally, conservatively rounded values of the calculated overall uncertainties were chosen for further processing in the uncertainty analysis.

Table 7-10 Sources of AD uncertainty and overall uncertainties in the area calculations, expressed as half of the 95% confidence intervals. Exception for source category 5D1 is mentioned in the main text above. Calculations are based on AREA data from SFSO (2013).

| IPCC Description | | Interpretation uncertainty | Sampling uncertainty | Overall uncertainty, calculated value | Overall uncertainty, rounded value |
|------------------|-----------------------------------|----------------------------|----------------------|---------------------------------------|------------------------------------|
| 5A1 | Forest Land remaining Forest Land | 1.1 | 0.2 | 1.09 | 2 |
| 5A2 | Land converted to Forest Land | 1.1 | 1.2 | 1.60 | 2 |
| 5B1 | Cropland remaining Cropland | 4.9 | 0.3 | 4.89 | 5 |
| 5B2 | Land converted to Cropland | 4.9 | 2.1 | 5.31 | 6 |
| 5C1 | Grassland remaining Grassland | 5.2 | 0.2 | 5.23 | 6 |
| 5C2 | Land converted to Grassland | 5.2 | 1.0 | 5.33 | 6 |
| 5D1 | Wetlands remaining Wetlands | 0.9 | 0.5 | 1.02 | 2 |
| 5D2 | Land converted to Wetlands | 0.9 | 3.9 | 4.05 | 5 |
| 5E1 | Settlements remaining Settlements | 4.4 | 0.4 | 4.41 | 5 |
| 5E2 | Land converted to Settlements | 4.4 | 1.1 | 4.54 | 5 |
| 5F1 | Other Land remaining Other Land | 1.4 | 0.3 | 1.40 | NA |
| 5F2 | Land converted to Other Land | 1.4 | 2.8 | 3.16 | 4 |

7.2.6 QA/QC and Verification of Activity Data

The AREA survey is a well-defined and controlled, long-term process in the responsibility of the Swiss Federal Statistical Office (SFSO 2006a). The data supplied by SFSO (2013) have been checked for suitability and consistency (Sigmaplan 2014).

The temporal interpolation and extrapolation of the AREA sample is quite a complex procedure, whose internal consistency is checked systematically as described in Sigmaplan (2014). Further checks (interannual comparisons, plausibility) are carried out after producing the land-use change tables presented in Chapter 7.2.3.2.

In response to UNFCCC (2012: §115) a systematic cross-check between the activity data reported under LULUCF category 5A and under the KP activity Forest management was carried out (Meteotest 2013a). It revealed that the difference between activity data used for emission and removal estimates under the Convention and under the KP can be consistently explained. The cross-check was updated for the present submission (see Chapter 11.2.3).

7.2.7 Recalculations of Activity Data

In the previous submission (FOEN 2013), AREA coverage has been restricted to 83%. The completion of the AREA surveys in 2013 led to a recalculation in the LULUCF sector for the period 1990-2011.

The borders of the NFI regions used to define the spatial strata (see Chapter 7.2.3.1) have been updated using new, more precise data (Meteotest 2013b). As a consequence, the size of the strata changed slightly.

7.2.8 Planned Improvements for Activity Data

No improvements are planned for the compilation and deduction of activity data.

7.3 Category 5A – Forest Land

7.3.1 Description

Tier 2 Key category 5A1

CO₂ from Forest Land remaining Forest Land
(2012: level and trend)

Tier 2 Key category 5A2

CO₂ from Land converted to Forest Land
(2012: level and trend)

Only temperate forests are occurring in Switzerland. Forest is defined as a minimum area of land of 0.0625 ha with crown cover of at least 20% and a minimum width of 25 m. The minimum height of the dominant trees must be 3 m or have the potential to reach 3 m at maturity in situ (FOEN 2006h). The following forest areas are not subject of the criteria of minimum stand height and minimum crown cover, but must have the potential to achieve it: afforested, regenerated, as well as burned, cut or damaged areas. Although orchards, parks, camping grounds, open tree formations in settlements, gardens, cemeteries, sports and parking fields may fulfil the (quantitative) forest definition, they are not considered as forests (FOEN 2006h).

For reporting in the CRF-tables, the different forest types are allocated to afforestations (CC11), productive forest (CC12) and unproductive forest (CC13) based on AREA categories (see Table 7-2; Table 7-6; SFSO 2006a). A detailed description of the category unproductive forest can be found in Chapter 7.3.4.9.

7.3.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.3.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.3.4 Methodological Issues

7.3.4.1 Choice of Method and National Forest Inventories

For calculating annual changes in carbon stocks changes, the general approach was used (see IPCC 2003 Eq. 3.1.1).

Data for growing stock, gross growth, cut (harvesting) and mortality were derived from the first, second, third and first phase a of the fourth Swiss National Forest Inventories (NFI, see Table 7-11). A description of NFI 1 and NFI 2 methodologies can be found in EAFV/BFL (1988) and in Brassel and Brändli (1999). Data and methodology of NFI 3 are described in Brändli (2010). Data of NFI 4a are described in Abegg et al. (2012). The methodology remained identical to Brändli (2010).

The inventories NFI 1, 2 and 3 are based on full surveys that were repeated in intervals of approximately 10 years. Since 2009, the inventory interval has been changed: a continuous survey is being conducted (NFI 4, 2009-2017). This means that a rotating subsample of approximately 12% will be surveyed and evaluated every year. NFI 4 data for the years 2009-2012 are implemented in this submission. Abegg et al. (2012) with NFI4a data covering 2009-2011 is an official NFI release. On request, NFI4a+ data for 2009-2012 have been provided exclusively for this submission by Thürig (2014).

Table 7-11: Characteristics of the National Forest Inventories 1, 2, 3 and 4a+.

| | NFI 1 | NFI 2 | NFI 3 | NFI 4a+ |
|--------------------------|-----------|--------------|--------------|--------------|
| Inventory cycle | 1983-1985 | 1993-1995 | 2004-2006 | 2009-2012 |
| Grid size | 1 x 1 km | 1.4 x 1.4 km | 1.4 x 1.4 km | 1.4 x 1.4 km |
| Terrestrial sample plots | ~12'000 | ~6'000 | ~6'000 | ~2'600 |
| Measured single trees | ~130'000 | ~70'000 | ~70'000 | ~30'000 |

7.3.4.2 Three-year Averaging of Forest Carbon Pools

The Revised 1996 IPCC Guidelines (IPCC 1997a) recommend working with three-year averages to report carbon changes in “Forest and Other Woody Biomass Stocks”. Further, the 2003 IPCC GPG (IPCC 2003) describes how to deal with interannual variability and states that “it is good practice to consistently report emissions using longer-term averages of environmental conditions or actual annual estimates of emissions when estimating stock changes”.

Changes in the carbon pools reported for the Swiss forest sector reflect annual fluctuations in management, weather conditions and natural disturbances. Therefore, three-year moving averages are calculated for all changes in forest carbon pools in order to smooth out high interannual fluctuations.

Three-year moving averages for the inventory year X are calculated as the average of the years X, X-1 and X-2. For example, the value for the inventory year 2004 is the average value of the years 2002-2004. This “backward-averaging” was used instead of calculating the arithmetic mean (mean of the years X-1, X, X+1), because

- if X is the most recent inventory year, X+1 data generally are not available in time (for submission in year X+2);
- we argue, that growth of living biomass, cut and mortality and the amount of dead wood is more influenced by the previous years than by the following year.

This “backward-averaging” introduces a certain time-lag in the calculated values and can complicate the interpretation of the resulting CO₂ emissions and removals.

7.3.4.3 Stratification

Spatial Strata

Forests in Switzerland reveal a high heterogeneity in terms of elevation, growth conditions, tree species composition, and inter-annual growth variability.

To combine the activity data of the Swiss land use statistics (see Chapter 7.2) with the emission factors from the Swiss forest inventory, Switzerland was divided into different strata. To find explanatory variables that significantly reduce the variance of gross growth an analysis of variance was done (Table 7-12).

Table 7-12 Analysis of variance of gross growth. Explanatory variables: Tree species, NFI production region, and altitude.

| | Gross growth | |
|------------------------|--------------|---------|
| | F-value | p-value |
| Coniferous / Deciduous | 421 | <0.0001 |
| Production region | 45 | <0.0001 |
| Altitude | 34 | <0.0001 |

The analysis of variance indicated that production region, elevation, and tree species all significantly explain differences in gross growth. Therefore, the explanatory variables considered in this study are:

- tree species (coniferous and deciduous species).
- the five NFI production regions
(1. Jura, 2. Central Plateau, 3. Pre-Alps, 4. Alps, 5. Southern Alps)
- altitude (<601 m, 601-1200 m, >1200 m)

Values for growing stock, gross growth, harvesting and mortality were calculated and applied for each of these 30 strata.

Separating Mixed Forests into Coniferous and Deciduous Sites

In Switzerland, most forests are mixed stands. However, the forest area derived by the Swiss land use statistics does not allow separating coniferous and deciduous sites.

To derive species specific measures for growing stock, gross growth, harvesting and mortality, the total forest area has to be divided according to the species mixture. The emission factor per stratum is then calculated as the weighted mean of both species. The required ratio of coniferous forest area (R_c) per spatial stratum was calculated by dividing the sum of the biomass of the conifers (B_c) over the sum of the biomass of all trees (B).

$$R_{ci} = B_{ci} / B_i \quad i = \text{spatial strata}$$

As both species add up to 1 (or 100%) the rate of deciduous forest area (R_d) is:

$$R_{di} = 1 - R_{ci} \quad i = \text{spatial strata}$$

The weights for each spatial stratum are displayed in Table 7-13.

Table 7-13 Ratio of coniferous and deciduous species for 1985-1994 (derived from NFI 1 and NFI 2; source: Brassel and Brändli 1999), for 1995-2005 (derived from NFI 2 and NFI 3 data; source: Brändli 2010) and for 2006-2012 (derived from NFI 3 and NFI 4a+ data; source: Abegg et al. 2012 and Thürig 2014).

| | | 1985 - 1994 | | 1995 – 2005 | | 2006-2012 | |
|------------|--------------|-------------|-----------|-------------|-----------|------------|-----------|
| NFI region | Altitude [m] | Coniferous | Deciduous | Coniferous | Deciduous | Coniferous | Deciduous |
| 1 | <601 | 0.31 | 0.69 | 0.31 | 0.69 | 0.30 | 0.70 |
| | 601-1200 | 0.54 | 0.46 | 0.52 | 0.48 | 0.52 | 0.48 |
| | >1200 | 0.74 | 0.26 | 0.72 | 0.28 | 0.74 | 0.26 |
| 2 | <601 | 0.56 | 0.47 | 0.50 | 0.50 | 0.40 | 0.60 |
| | 601-1200 | 0.60 | 0.40 | 0.58 | 0.42 | 0.48 | 0.52 |
| | >1200 | 0.90 | 0.10 | 0.90 | 0.10 | 0.96 | 0.40 |
| 3 | <601 | 0.40 | 0.60 | 0.40 | 0.60 | 0.26 | 0.74 |
| | 601-1200 | 0.70 | 0.30 | 0.69 | 0.31 | 0.63 | 0.37 |
| | >1200 | 0.92 | 0.08 | 0.91 | 0.09 | 0.87 | 0.13 |
| 4 | <601 | 0.33 | 0.67 | 0.33 | 0.67 | 0.17 | 0.83 |
| | 601-1200 | 0.64 | 0.36 | 0.63 | 0.37 | 0.54 | 0.46 |
| | >1200 | 0.97 | 0.03 | 0.96 | 0.04 | 0.95 | 0.05 |
| 5 | <601 | 0.07 | 0.93 | 0.06 | 0.94 | 0.03 | 0.97 |
| | 601-1200 | 0.18 | 0.82 | 0.17 | 0.83 | 0.16 | 0.84 |
| | >1200 | 0.84 | 0.16 | 0.83 | 0.17 | 0.84 | 0.16 |

Additional Stratification: Eastern and Western Alps

In the Swiss Alps (NFI region 4) below an altitude of 1200 m, climate between the eastern and the western part differs substantially. We therefore included an additional stratification for the eastern and the western part of the Alps below 1200 m (Alps < 601 m east, Alps < 601 m west, Alps 601-1200 m east, Alps 601-1200 m west; see Thürig et al. 2005a for details). This additional stratification resulted in very small datasets per stratum.

Gains and losses of living biomass were estimated for the eastern and western Alps separately. The emission factors for the Alps below 1200 m were then calculated as a weighted mean of the percentage of forest biomass situated in the western and in the eastern Alps. The weights for the pooled emission factors derived from the NFI 1, NFI 2, NFI 3 and NFI 4 are listed in Table 7-14.

Table 7-14 Ratio of biomass in the eastern and western Alps (NFI production region 4) for 1985-1994 (derived from NFI 1 and NFI 2; source: Brassel and Brändli 1999), for 1995-2005 (derived from NFI 2 and NFI 3 data; source: Brändli 2010) and for 2006-2012 (derived from NFI 3 and NFI 4a+ data; source: Abegg et al. 2012, Thürig 2014).

| | | 1985 - 1994 | | 1995 – 2005 | | 2006-2012 | |
|--------------|--|------------------|------------------|------------------|------------------|--------------------|--------------------|
| Altitude [m] | | NFI 2 Eastern | NFI 2 Western | NFI 3 Eastern | NFI 3 Western | NFI 4a+ Eastern | NFI 4a+ Western |
| <601 | | 0.56 | 0.44 | 0.53 | 0.47 | 0.60 | 0.40 |
| 601-1200 | | 0.62 | 0.38 | 0.61 | 0.39 | 0.62 | 0.38 |

7.3.4.4 Estimation of Growing Stock in Biomass

The biomass of all tree compartments (stem-wood over bark including stock, coarse and small branches, needles/leaves, and roots) were estimated based on established allometries to tree-dimensions (Table 7-15; Thürig and Herold 2013). Estimates for branches, foliage and roots were derived from tree diameter at breast height (DBH). For stem-wood over bark including stock, additionally, diameter at tree height 7 m (D7) and total tree height were required. Except for roots, the biomass functions were empirically derived from a large number of single-tree data from Swiss forest sites (see references in Table 7-15).

Table 7-15 Applied allometric biomass functions, dependencies and references. DBH: tree diameter at breast height; D7: diameter at tree height 7 m.

| Tree parts | Input parameter | Nr. of trees | References |
|---------------------------------|-----------------|--------------|------------------------|
| Stem-wood over bark incl. stock | DBH, D7, height | 12'000 | Kaufmann et al. 2001 |
| Coarse branches (≥ 7 cm) | DBH | 40'000 | Kaufmann et al. 2001 |
| Small branches (< 7 cm) | DBH | 40'000 | Kaufmann et al. 2001 |
| Needles, Leaves | DBH | 400 | Perruchoud et al. 1999 |
| Broadleaved Roots | DBH | 443 | Wutzler et al. 2008 |
| Coniferous Roots | DBH | 80 | Zell and Thürig 2012 |

The biomass of all individual trees was calculated and, in a second step, single-tree estimates of gains and losses were obtained as the difference in biomass between subsequent NFIs (Thürig and Herold 2013).

7.3.4.5 Carbon Content

The IPCC default carbon content of solid wood of 50% was applied (IPCC 2003; p. 3.25).

7.3.4.6 Productive Forests (CC12): Growing Stock, Gross Growth and Cut and Mortality

Values for growing stock, gross growth, cut and mortality for productive forests (CC12, without afforestations) were derived from 5'425 common sample plots measured during NFI 1 and NFI 2 (Kaufmann 2001), 5'581 samples measured during NFI 2 and NFI 3 (Brändli 2010) and 2613 samples measured during NFI 3 and NFI 4 2009-2012 (Abegg et al. 2012; Thürig 2014). All values derived from the national forest inventories are related to above- and below-ground biomass in mass units (t C ha^{-1}) per spatial stratum. Annual values for growing stock are shown in Table 7-19 as "carbon stock in living biomass". Table 7-16 and Table 7-17 show gross growth and cut and mortality (in Table 7-19 marked as "gain of living biomass" and "loss of living biomass") for the four NFIs for coniferous and deciduous trees, respectively.

Table 7-16 Gross growth and cut and mortality for coniferous trees (related to coniferous forest biomass). In the Alps (NFI production region 4) below 1200 m, data are additionally stratified for eastern and western Alps. Data sources: Brassel and Brändli (1999), Brändli (2010), Abegg et al. (2012) and Thürig (2014).

| NFI region | Altitude [m] | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 1-2 | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 1-2 | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 2-3 | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 2-3 | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 3-4a+ | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 3-4a+ |
|------------|--------------|---|--|---|--|---|--|
| 1 | <601 | 2.13 | 1.42 | 2.26 | 2.82 | 2.07 | 2.95 |
| | 601-1200 | 3.15 | 2.52 | 3.16 | 2.80 | 3.60 | 3.05 |
| | >1200 | 2.68 | 2.16 | 2.67 | 1.71 | 3.34 | 1.19 |
| 2 | <601 | 4.51 | 4.28 | 4.21 | 6.28 | 3.93 | 5.68 |
| | 601-1200 | 5.29 | 4.60 | 4.85 | 7.21 | 5.14 | 6.54 |
| | >1200 | 2.40 | 1.40 | 1.49 | 2.20 | 5.14 | 6.54 |
| 3 | <601 | 3.27 | 1.91 | 3.01 | 3.01 | 4.92 | 4.47 |
| | 601-1200 | 5.52 | 4.10 | 5.39 | 6.38 | 4.92 | 4.47 |
| | >1200 | 4.50 | 3.57 | 4.52 | 4.64 | 5.23 | 3.61 |
| 4 east | <601 | 2.75 | 1.29 | 2.90 | 1.59 | 3.22 | 2.65 |
| 4 west | <601 | 0.72 | 0.84 | 1.23 | 0.92 | 2.22 | 1.27 |
| 4 east | 601-1200 | 3.44 | 2.86 | 3.44 | 2.31 | 3.22 | 2.65 |
| 4 west | 601-1200 | 2.40 | 2.02 | 2.17 | 1.76 | 2.22 | 1.27 |
| 4 | >1200 | 3.36 | 5.59 | 3.50 | 2.43 | 3.47 | 1.71 |
| 5 | <601 | 0.08 | 0.06 | 0.12 | 0.02 | 0.78 | 0.00 |
| | 601-1200 | 0.43 | 0.23 | 0.56 | 0.15 | 0.34 | 0.84 |
| | >1200 | 2.38 | 0.75 | 2.46 | 0.78 | 3.28 | 1.15 |

Table 7-17 Gross growth, cut and mortality for deciduous trees (related to deciduous forest biomass). In the Alps (NFI production region 4) below 1200 m, data are additionally stratified for eastern and western Alps. Data sources: Brassel and Brändli (1999); Brändli (2010), Abegg et al. (2012) and Thürig (2014).

| NFI region | Altitude [m] | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 1-2 | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 1-2 | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 2-3 | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 2-3 | Gross growth [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 3-4a+ | Cut and mortality [10 ³ kg ha ⁻¹ yr ⁻¹] NFI 3-4a+ |
|------------|--------------|---|--|---|--|---|--|
| 1 | <601 | 5.08 | 3.30 | 4.48 | 5.11 | 4.37 | 3.52 |
| | 601-1200 | 3.27 | 1.80 | 2.91 | 2.13 | 3.60 | 2.95 |
| | >1200 | 1.22 | 0.31 | 0.93 | 0.57 | 3.34 | 0.21 |
| 2 | <601 | 4.75 | 3.36 | 4.87 | 3.93 | 3.93 | 4.17 |
| | 601-1200 | 3.98 | 2.65 | 4.27 | 2.79 | 4.20 | 3.21 |
| | >1200 | 0.80 | 0.16 | 1.07 | 0.40 | 4.20 | 3.21 |
| 3 | <601 | 5.84 | 3.85 | 5.46 | 2.61 | 2.91 | 1.55 |
| | 601-1200 | 2.77 | 1.41 | 2.92 | 1.61 | 2.91 | 1.55 |
| | >1200 | 0.46 | 0.12 | 0.48 | 0.11 | 0.54 | 0.23 |
| 4 east | <601 | 4.66 | 6.41 | 5.09 | 2.18 | 2.93 | 1.96 |
| 4 west | <601 | 5.20 | 3.08 | 4.95 | 2.11 | 2.26 | 1.25 |
| 4 east | 601-1200 | 2.11 | 0.95 | 2.05 | 1.10 | 2.93 | 4.60 |
| 4 west | 601-1200 | 1.93 | 0.73 | 2.27 | 1.03 | 2.26 | 2.52 |
| 4 | >1200 | 0.25 | 0.14 | 0.34 | 0.13 | 0.50 | 0.09 |
| 5 | <601 | 5.39 | 2.29 | 3.96 | 2.72 | 4.98 | 2.58 |
| | 601-1200 | 3.97 | 1.40 | 3.79 | 1.11 | 4.70 | 2.00 |
| | >1200 | 0.83 | 0.26 | 1.12 | 0.16 | 0.48 | 0.28 |

Annual Gross Growth

Annual values of gross growth have been derived from the NFI 1 and NFI 2 datasets for the period 1985-1994, from the NFI 2 and NFI 3 datasets for the period 1995-2005 and from the NFI 3 and NFI 4a+ dataset for the period 2006-2012. Annual values of gross growth are constant in the intersurvey periods of NFI 1 to NFI 2, NFI 2 to NFI 3 and of NFI 3 to NFI 4a+, respectively. These annual values are averaged over 3 years (see Chapter 7.3.4.2), thereby affecting the values of gross growth of the years 1996, 1997 and 2006, 2007, respectively (see Table 7-19).

Annual Cut and Mortality

An average value for cut and mortality (CM) is derived from the NFI 1 and NFI 2 dataset for the period 1985-1994, from the NFI 2 and NFI 3 datasets for the period 1995-2005 and from the NFI 3 and NFI 4a+ dataset for the period 2006-2012. To calculate annual values of cut and mortality (CM_y) for the years 1985 to 1994, 1995 to 2005 and 2006 to 2012, respectively, the average amount of cut and mortality was weighted by the percentage of the relative harvesting amounts taken from the forest statistics (Table 7-18; SFSO 2013c; FOEN 2013k, and former editions 1985-2012). Relative harvesting amounts were calculated for each year per LFI-intersurvey period. As recommended in the Revised 1996 IPCC Guidelines (IPCC 1997a), moving three-year averages of the harvesting amounts from the

forest statistics were calculated in order to level out extreme events such as storm Vivian in 1990 and storm Lothar in 1999 (see Chapter 7.3.4.2).

Table 7-18 Annual harvesting amount in m³ merchantable timber specified for NFI production region as well as for coniferous and deciduous tree species for the period 1990-2012 as derived from forest statistics (SFSO 2013c; FOEN 2013k, and former editions 1985-2012). All values were averaged over three years as recommended in the revised 1996 IPCC guidelines (IPCC 1997a).

| Year | 1. Jura | | 2. Central plateau | | 3. Pre-Alps | | 4. Alps | | 5. Southern Alps | |
|------|-----------------------------|---------------------------|-----------------------------|---------------------------|-----------------------------|---------------------------|-----------------------------|---------------------------|-----------------------------|---------------------------|
| | Conif. [m ³] | Dec. [m ³] | Conif. [m ³] | Dec. [m ³] | Conif. [m ³] | Dec. [m ³] | Conif. [m ³] | Dec. [m ³] | Conif. [m ³] | Dec. [m ³] |
| 1990 | 669'756 | 364'296 | 1'400'390 | 582'340 | 963'683 | 138'833 | 851'765 | 65'707 | 38'790 | 24'026 |
| 1991 | 616'629 | 360'660 | 1'348'951 | 557'776 | 967'684 | 135'699 | 1'002'608 | 68'221 | 31'210 | 24'093 |
| 1992 | 573'269 | 361'633 | 1'328'880 | 556'023 | 966'390 | 133'405 | 1'034'064 | 71'000 | 31'106 | 25'943 |
| 1993 | 527'672 | 366'516 | 1'141'041 | 541'195 | 779'032 | 131'588 | 816'939 | 68'958 | 38'085 | 29'386 |
| 1994 | 575'928 | 379'505 | 1'225'395 | 554'916 | 752'565 | 132'571 | 701'336 | 67'181 | 43'628 | 31'723 |
| 1995 | 607'611 | 391'128 | 1'288'507 | 554'563 | 765'351 | 140'962 | 652'879 | 62'517 | 45'047 | 33'467 |
| 1996 | 597'544 | 393'817 | 1'241'999 | 556'409 | 742'348 | 147'125 | 604'935 | 61'095 | 46'972 | 35'501 |
| 1997 | 590'296 | 394'443 | 1'210'678 | 571'579 | 723'808 | 152'997 | 557'039 | 60'013 | 53'658 | 37'649 |
| 1998 | 575'006 | 399'476 | 1'191'359 | 590'606 | 744'730 | 156'410 | 579'223 | 77'391 | 53'319 | 40'188 |
| 1999 | 602'445 | 405'237 | 1'283'404 | 614'399 | 801'259 | 163'971 | 608'468 | 80'428 | 52'075 | 40'285 |
| 2000 | 733'872 | 402'682 | 2'196'853 | 733'718 | 1'300'811 | 184'017 | 562'665 | 78'246 | 38'806 | 38'572 |
| 2001 | 680'175 | 374'861 | 2'426'715 | 722'713 | 1'514'372 | 181'804 | 513'772 | 62'014 | 29'343 | 36'651 |
| 2002 | 626'798 | 351'805 | 2'448'000 | 674'298 | 1'603'283 | 168'724 | 491'872 | 60'187 | 24'903 | 35'522 |
| 2003 | 481'195 | 327'776 | 1'698'975 | 535'598 | 1'254'485 | 144'789 | 542'312 | 62'065 | 30'195 | 35'667 |
| 2004 | 551'910 | 316'752 | 1'617'068 | 509'352 | 1'135'069 | 147'134 | 534'976 | 65'377 | 32'781 | 35'617 |
| 2005 | 622'087 | 326'862 | 1'751'762 | 549'665 | 1'108'437 | 162'449 | 530'563 | 67'811 | 34'189 | 34'890 |
| 2006 | 681'354 | 357'113 | 1'788'551 | 606'050 | 1'082'363 | 191'691 | 524'433 | 75'116 | 36'300 | 39'261 |
| 2007 | 727'255 | 397'149 | 1'726'102 | 667'116 | 1'090'739 | 213'537 | 568'604 | 79'224 | 47'235 | 41'950 |
| 2008 | 744'843 | 430'545 | 1'549'750 | 704'695 | 1'093'245 | 228'233 | 618'331 | 83'231 | 53'102 | 45'453 |
| 2009 | 699'189 | 448'946 | 1'339'493 | 709'282 | 1'013'811 | 226'469 | 654'511 | 85'013 | 57'413 | 43'359 |
| 2010 | 650'428 | 471'929 | 1'173'993 | 717'138 | 963'166 | 232'425 | 687'652 | 90'799 | 56'610 | 46'159 |
| 2011 | 621'118 | 489'838 | 1'100'727 | 721'806 | 951'347 | 241'980 | 695'223 | 97'139 | 59'529 | 49'219 |
| 2012 | 566'782 | 488'626 | 970'748 | 719'003 | 825'019 | 225'988 | 665'506 | 94'480 | 51'475 | 50'757 |

Growing Stock: Calculation of Time Series

In order to develop a consistent time series, annual growing stocks (GS) are calculated backward or forward starting from the growing stock 2005, determined from NFI 3.

A backward calculation is used for the time period 1985-2004, meaning that the annual growing stock equals the growing stock 2005 minus the cumulated gains of the annual gross growths and plus the cumulated annual amounts of cut and mortality (CM_y).

Growing stocks for inventory years after 2005 are determined using a forward calculation, i.e. adding the cumulated annual gross growths to the growing stock 2005, and subtracting the cumulated annual amounts of cut and mortality (CM_y).

$$GS_{iy} = GS_{2005} - \sum_y [\text{annual gross growth}_y] + \sum_y [CM_y] \quad \text{for } iy < 2005$$

$$GS_{iy} = GS_{2005} \quad \text{for } iy = 2005$$

$$GS_{iy} = GS_{2005} + \sum_y [\text{annual gross growth}_y] - \sum_y [CM_y] \quad \text{for } iy > 2005$$

where the "iy" indicates the inventory year and "y" refers to the years between 2005 and the inventory year.

An overview of the values of gross growth, cut & mortality and calculated growing stock for the period 1990 to 2012 specified for all spatial strata are displayed in Table 7-19.

All work steps and data required to reproduce the calculation of emission factors for productive forests (CC12) in the period 1990-2012 are summarized in FOEN (2014b).

Table 7-19 Annual carbon data of living biomass for productive forest (CC12) disaggregated for NFI region (NFI) and altitude zone (Alt.), 1990-2012, three-year-averages. Highlighted data for 1990 as displayed in Table 7-4.

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 126.87 | 128.07 | 129.34 | 130.66 | 132.01 | 133.24 | 134.39 | 133.80 | 133.14 | 132.41 |
| 1 | 2 | 124.88 | 125.82 | 126.88 | 128.02 | 129.25 | 130.35 | 131.35 | 131.98 | 132.58 | 133.13 |
| 1 | 3 | 84.73 | 85.35 | 86.06 | 86.85 | 87.72 | 88.50 | 89.21 | 89.97 | 90.68 | 91.36 |
| 2 | 1 | 134.18 | 134.69 | 135.35 | 136.06 | 137.13 | 138.01 | 138.79 | 139.29 | 139.78 | 140.21 |
| 2 | 2 | 146.77 | 147.47 | 148.33 | 149.22 | 150.49 | 151.58 | 152.55 | 153.26 | 153.97 | 154.66 |
| 2 | 3 | 101.21 | 101.95 | 102.72 | 103.51 | 104.40 | 105.24 | 106.05 | 106.56 | 106.97 | 107.29 |
| 3 | 1 | 135.06 | 136.58 | 138.13 | 139.71 | 141.54 | 143.38 | 145.08 | 147.32 | 149.42 | 151.36 |
| 3 | 2 | 147.43 | 148.52 | 149.61 | 150.72 | 152.30 | 153.93 | 155.49 | 156.74 | 158.01 | 159.21 |
| 3 | 3 | 119.32 | 119.69 | 120.06 | 120.43 | 121.20 | 122.02 | 122.82 | 123.67 | 124.57 | 125.44 |
| 4 | 1 | 94.81 | 95.34 | 95.69 | 95.92 | 96.37 | 96.96 | 97.88 | 99.57 | 101.40 | 102.99 |
| 4 | 2 | 104.42 | 105.10 | 105.52 | 105.87 | 106.59 | 107.51 | 108.45 | 109.33 | 110.32 | 111.14 |
| 4 | 3 | 96.41 | 96.60 | 96.50 | 96.34 | 96.59 | 97.06 | 97.61 | 98.11 | 98.75 | 99.35 |
| 5 | 1 | 70.67 | 72.40 | 74.13 | 75.78 | 77.29 | 78.70 | 80.03 | 81.22 | 82.10 | 82.65 |
| 5 | 2 | 76.70 | 78.18 | 79.69 | 81.15 | 82.50 | 83.78 | 85.01 | 86.58 | 88.10 | 89.58 |
| 5 | 3 | 76.70 | 77.83 | 79.03 | 80.22 | 81.33 | 82.38 | 83.40 | 84.53 | 85.65 | 86.82 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 131.57 | 130.45 | 129.64 | 129.12 | 129.10 | 128.99 | 129.41 | 129.51 | 129.39 | 129.30 |
| 1 | 2 | 133.61 | 133.79 | 134.17 | 134.74 | 135.72 | 136.57 | 136.97 | 137.22 | 137.41 | 137.64 |
| 1 | 3 | 92.00 | 92.46 | 93.01 | 93.66 | 94.52 | 95.30 | 96.49 | 97.71 | 98.99 | 100.29 |
| 2 | 1 | 140.40 | 138.51 | 136.24 | 134.09 | 133.77 | 133.68 | 132.77 | 131.83 | 131.17 | 130.93 |
| 2 | 2 | 155.10 | 153.34 | 151.11 | 148.95 | 148.70 | 148.67 | 147.52 | 146.40 | 145.65 | 145.40 |
| 2 | 3 | 107.54 | 107.16 | 106.64 | 106.12 | 106.12 | 106.19 | 108.05 | 111.06 | 115.25 | 119.51 |
| 3 | 1 | 153.16 | 154.11 | 154.78 | 155.44 | 156.77 | 158.25 | 159.20 | 159.80 | 160.12 | 160.54 |
| 3 | 2 | 160.20 | 159.64 | 158.47 | 157.10 | 156.87 | 156.97 | 158.02 | 158.91 | 159.67 | 160.61 |
| 3 | 3 | 126.18 | 125.85 | 125.07 | 124.10 | 123.89 | 123.92 | 124.53 | 125.25 | 126.08 | 127.05 |
| 4 | 1 | 104.50 | 106.09 | 107.99 | 109.95 | 111.82 | 113.65 | 114.67 | 114.93 | 114.73 | 114.44 |
| 4 | 2 | 111.87 | 112.71 | 113.77 | 114.88 | 115.89 | 116.88 | 119.01 | 120.10 | 121.15 | 122.11 |
| 4 | 3 | 99.89 | 100.53 | 101.29 | 102.09 | 102.78 | 103.49 | 104.73 | 106.01 | 107.30 | 108.54 |
| 5 | 1 | 83.20 | 83.82 | 84.51 | 85.24 | 85.97 | 86.70 | 87.90 | 89.31 | 90.90 | 92.55 |
| 5 | 2 | 91.06 | 92.59 | 94.16 | 95.76 | 97.35 | 98.93 | 100.07 | 101.17 | 102.27 | 103.37 |
| 5 | 3 | 88.01 | 89.33 | 90.75 | 92.22 | 93.63 | 95.02 | 96.32 | 97.52 | 98.68 | 99.80 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|--|------|--------|--------|--------|--|--|--|--|--|--|--|
| CC12: carbon stock in living biomass (stockCl,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 129.22 | 129.14 | 129.04 | | | | | | | |
| 1 | 2 | 137.90 | 138.17 | 138.44 | | | | | | | |
| 1 | 3 | 101.64 | 103.01 | 104.38 | | | | | | | |
| 2 | 1 | 131.00 | 131.21 | 131.52 | | | | | | | |
| 2 | 2 | 145.53 | 145.82 | 146.25 | | | | | | | |
| 2 | 3 | 123.81 | 128.14 | 132.49 | | | | | | | |
| 3 | 1 | 160.94 | 161.24 | 161.55 | | | | | | | |
| 3 | 2 | 161.65 | 162.69 | 163.77 | | | | | | | |
| 3 | 3 | 128.10 | 129.18 | 130.28 | | | | | | | |
| 4 | 1 | 113.98 | 113.36 | 112.71 | | | | | | | |
| 4 | 2 | 122.98 | 123.79 | 124.59 | | | | | | | |
| 4 | 3 | 109.73 | 110.91 | 112.10 | | | | | | | |
| 5 | 1 | 94.11 | 95.59 | 96.99 | | | | | | | |
| 5 | 2 | 104.43 | 105.39 | 106.30 | | | | | | | |
| 5 | 3 | 100.92 | 101.99 | 103.08 | | | | | | | |

(Table 7-19 continued)

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|------|------|------|------|------|------|------|------|------|------|
| CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 3.60 | 3.60 | 3.60 | 3.60 | 3.60 | 3.60 | 3.53 | 3.45 | 3.37 | 3.37 |
| 1 | 2 | 3.21 | 3.21 | 3.21 | 3.21 | 3.21 | 3.21 | 3.15 | 3.09 | 3.04 | 3.04 |
| 1 | 3 | 1.95 | 1.95 | 1.95 | 1.95 | 1.95 | 1.95 | 1.90 | 1.85 | 1.80 | 1.80 |
| 2 | 1 | 4.63 | 4.63 | 4.63 | 4.63 | 4.63 | 4.63 | 4.60 | 4.57 | 4.54 | 4.54 |
| 2 | 2 | 4.63 | 4.63 | 4.63 | 4.63 | 4.63 | 4.63 | 4.61 | 4.59 | 4.56 | 4.56 |
| 2 | 3 | 1.60 | 1.60 | 1.60 | 1.60 | 1.60 | 1.60 | 1.49 | 1.39 | 1.28 | 1.28 |
| 3 | 1 | 4.56 | 4.56 | 4.56 | 4.56 | 4.56 | 4.56 | 4.45 | 4.34 | 4.23 | 4.23 |
| 3 | 2 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 |
| 3 | 3 | 2.48 | 2.48 | 2.48 | 2.48 | 2.48 | 2.48 | 2.49 | 2.49 | 2.50 | 2.50 |
| 4 | 1 | 3.24 | 3.24 | 3.24 | 3.24 | 3.24 | 3.24 | 3.31 | 3.37 | 3.44 | 3.44 |
| 4 | 2 | 2.49 | 2.49 | 2.49 | 2.49 | 2.49 | 2.49 | 2.50 | 2.50 | 2.50 | 2.50 |
| 4 | 3 | 1.81 | 1.81 | 1.81 | 1.81 | 1.81 | 1.81 | 1.84 | 1.87 | 1.90 | 1.90 |
| 5 | 1 | 2.74 | 2.74 | 2.74 | 2.74 | 2.74 | 2.74 | 2.51 | 2.27 | 2.04 | 2.04 |
| 5 | 2 | 2.20 | 2.20 | 2.20 | 2.20 | 2.20 | 2.20 | 2.19 | 2.18 | 2.18 | 2.18 |
| 5 | 3 | 1.61 | 1.61 | 1.61 | 1.61 | 1.61 | 1.61 | 1.67 | 1.73 | 1.79 | 1.79 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|------|------|------|------|------|------|------|------|------|------|
| CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 3.37 | 3.37 | 3.37 | 3.37 | 3.37 | 3.37 | 3.32 | 3.27 | 3.22 | 3.22 |
| 1 | 2 | 3.04 | 3.04 | 3.04 | 3.04 | 3.04 | 3.04 | 3.13 | 3.22 | 3.31 | 3.31 |
| 1 | 3 | 1.80 | 1.80 | 1.80 | 1.80 | 1.80 | 1.80 | 1.88 | 1.96 | 2.03 | 2.03 |
| 2 | 1 | 4.54 | 4.54 | 4.54 | 4.54 | 4.54 | 4.54 | 4.57 | 4.60 | 4.63 | 4.63 |
| 2 | 2 | 4.56 | 4.56 | 4.56 | 4.56 | 4.56 | 4.56 | 4.60 | 4.63 | 4.67 | 4.67 |
| 2 | 3 | 1.28 | 1.28 | 1.28 | 1.28 | 1.28 | 1.28 | 2.41 | 3.54 | 4.67 | 4.67 |
| 3 | 1 | 4.23 | 4.23 | 4.23 | 4.23 | 4.23 | 4.23 | 4.13 | 4.02 | 3.92 | 3.92 |
| 3 | 2 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.15 | 4.08 | 4.00 | 3.92 | 3.92 |
| 3 | 3 | 2.50 | 2.50 | 2.50 | 2.50 | 2.50 | 2.50 | 2.63 | 2.75 | 2.88 | 2.88 |
| 4 | 1 | 3.44 | 3.44 | 3.44 | 3.44 | 3.44 | 3.44 | 3.15 | 2.86 | 2.57 | 2.57 |
| 4 | 2 | 2.50 | 2.50 | 2.50 | 2.50 | 2.50 | 2.50 | 2.64 | 2.72 | 2.80 | 2.80 |
| 4 | 3 | 1.90 | 1.90 | 1.90 | 1.90 | 1.90 | 1.90 | 1.99 | 2.07 | 2.16 | 2.16 |
| 5 | 1 | 2.04 | 2.04 | 2.04 | 2.04 | 2.04 | 2.04 | 2.32 | 2.60 | 2.88 | 2.88 |
| 5 | 2 | 2.18 | 2.18 | 2.18 | 2.18 | 2.18 | 2.18 | 2.29 | 2.41 | 2.52 | 2.52 |
| 5 | 3 | 1.79 | 1.79 | 1.79 | 1.79 | 1.79 | 1.79 | 1.82 | 1.85 | 1.88 | 1.88 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|--|------|------|------|------|--|--|--|--|--|--|--|
| CC12: gain of living biomass (gainCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 3.22 | 3.22 | 3.22 | | | | | | | |
| 1 | 2 | 3.31 | 3.31 | 3.31 | | | | | | | |
| 1 | 3 | 2.03 | 2.03 | 2.03 | | | | | | | |
| 2 | 1 | 4.63 | 4.63 | 4.63 | | | | | | | |
| 2 | 2 | 4.67 | 4.67 | 4.67 | | | | | | | |
| 2 | 3 | 4.67 | 4.67 | 4.67 | | | | | | | |
| 3 | 1 | 3.92 | 3.92 | 3.92 | | | | | | | |
| 3 | 2 | 3.92 | 3.92 | 3.92 | | | | | | | |
| 3 | 3 | 2.88 | 2.88 | 2.88 | | | | | | | |
| 4 | 1 | 2.57 | 2.57 | 2.57 | | | | | | | |
| 4 | 2 | 2.80 | 2.80 | 2.80 | | | | | | | |
| 4 | 3 | 2.16 | 2.16 | 2.16 | | | | | | | |
| 5 | 1 | 2.88 | 2.88 | 2.88 | | | | | | | |
| 5 | 2 | 2.52 | 2.52 | 2.52 | | | | | | | |
| 5 | 3 | 1.88 | 1.88 | 1.88 | | | | | | | |

(Table 7-19 continued)

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | -2.41 | -2.33 | -2.29 | -2.25 | -2.37 | -2.46 | -4.12 | -4.10 | -4.10 | -4.21 |
| 1 | 2 | -2.27 | -2.15 | -2.07 | -1.98 | -2.11 | -2.21 | -2.52 | -2.50 | -2.48 | -2.56 |
| 1 | 3 | -1.34 | -1.24 | -1.16 | -1.09 | -1.18 | -1.24 | -1.15 | -1.14 | -1.12 | -1.16 |
| 2 | 1 | -4.13 | -3.97 | -3.93 | -3.56 | -3.75 | -3.85 | -4.09 | -4.08 | -4.11 | -4.36 |
| 2 | 2 | -3.93 | -3.78 | -3.74 | -3.36 | -3.55 | -3.66 | -3.90 | -3.87 | -3.88 | -4.12 |
| 2 | 3 | -0.86 | -0.82 | -0.81 | -0.71 | -0.76 | -0.79 | -0.98 | -0.97 | -0.96 | -1.03 |
| 3 | 1 | -3.04 | -3.00 | -2.97 | -2.73 | -2.72 | -2.85 | -2.22 | -2.24 | -2.29 | -2.43 |
| 3 | 2 | -3.06 | -3.05 | -3.04 | -2.57 | -2.51 | -2.59 | -2.90 | -2.88 | -2.96 | -3.16 |
| 3 | 3 | -2.11 | -2.11 | -2.11 | -1.71 | -1.66 | -1.69 | -1.63 | -1.59 | -1.64 | -1.76 |
| 4 | 1 | -2.71 | -2.89 | -3.01 | -2.79 | -2.66 | -2.47 | -1.62 | -1.55 | -1.85 | -1.93 |
| 4 | 2 | -1.81 | -2.07 | -2.14 | -1.77 | -1.57 | -1.47 | -1.61 | -1.51 | -1.69 | -1.77 |
| 4 | 3 | -1.62 | -1.91 | -1.97 | -1.56 | -1.34 | -1.25 | -1.34 | -1.23 | -1.29 | -1.36 |
| 5 | 1 | -1.01 | -1.01 | -1.08 | -1.23 | -1.33 | -1.40 | -1.32 | -1.40 | -1.49 | -1.49 |
| 5 | 2 | -0.71 | -0.69 | -0.74 | -0.84 | -0.92 | -0.97 | -0.62 | -0.66 | -0.70 | -0.70 |
| 5 | 3 | -0.48 | -0.41 | -0.41 | -0.50 | -0.56 | -0.58 | -0.54 | -0.61 | -0.61 | -0.60 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | -4.49 | -4.18 | -3.89 | -3.39 | -3.48 | -3.71 | -2.91 | -3.17 | -3.34 | -3.31 |
| 1 | 2 | -2.86 | -2.65 | -2.46 | -2.06 | -2.19 | -2.38 | -2.73 | -2.97 | -3.12 | -3.08 |
| 1 | 3 | -1.35 | -1.25 | -1.16 | -0.93 | -1.02 | -1.13 | -0.69 | -0.74 | -0.76 | -0.72 |
| 2 | 1 | -6.42 | -6.81 | -6.69 | -4.86 | -4.63 | -5.01 | -5.48 | -5.54 | -5.29 | -4.87 |
| 2 | 2 | -6.33 | -6.79 | -6.72 | -4.82 | -4.59 | -4.96 | -5.76 | -5.75 | -5.41 | -4.92 |
| 2 | 3 | -1.66 | -1.80 | -1.80 | -1.27 | -1.21 | -1.31 | -0.55 | -0.53 | -0.48 | -0.41 |
| 3 | 1 | -3.28 | -3.56 | -3.58 | -2.90 | -2.76 | -2.84 | -3.17 | -3.43 | -3.59 | -3.50 |
| 3 | 2 | -4.72 | -5.33 | -5.52 | -4.39 | -4.05 | -4.05 | -3.03 | -3.11 | -3.16 | -2.98 |
| 3 | 3 | -2.83 | -3.28 | -3.46 | -2.72 | -2.46 | -2.41 | -2.02 | -2.04 | -2.05 | -1.91 |
| 4 | 1 | -1.84 | -1.53 | -1.48 | -1.56 | -1.61 | -1.64 | -2.45 | -2.60 | -2.76 | -2.85 |
| 4 | 2 | -1.66 | -1.44 | -1.39 | -1.50 | -1.51 | -1.52 | -1.46 | -1.56 | -1.68 | -1.75 |
| 4 | 3 | -1.26 | -1.14 | -1.09 | -1.20 | -1.19 | -1.18 | -0.74 | -0.80 | -0.87 | -0.92 |
| 5 | 1 | -1.43 | -1.35 | -1.31 | -1.32 | -1.32 | -1.29 | -1.12 | -1.19 | -1.29 | -1.23 |
| 5 | 2 | -0.65 | -0.60 | -0.58 | -0.59 | -0.59 | -0.59 | -1.15 | -1.30 | -1.43 | -1.41 |
| 5 | 3 | -0.47 | -0.37 | -0.32 | -0.38 | -0.40 | -0.41 | -0.52 | -0.65 | -0.72 | -0.76 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|--|------|-------|-------|-------|--|--|--|--|--|--|--|
| CC12: loss of living biomass (lossCl,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | -3.30 | -3.31 | -3.32 | | | | | | | |
| 1 | 2 | -3.05 | -3.04 | -3.05 | | | | | | | |
| 1 | 3 | -0.69 | -0.67 | -0.66 | | | | | | | |
| 2 | 1 | -4.56 | -4.42 | -4.32 | | | | | | | |
| 2 | 2 | -4.54 | -4.37 | -4.24 | | | | | | | |
| 2 | 3 | -0.36 | -0.34 | -0.32 | | | | | | | |
| 3 | 1 | -3.52 | -3.62 | -3.61 | | | | | | | |
| 3 | 2 | -2.88 | -2.88 | -2.84 | | | | | | | |
| 3 | 3 | -1.83 | -1.81 | -1.78 | | | | | | | |
| 4 | 1 | -3.03 | -3.19 | -3.22 | | | | | | | |
| 4 | 2 | -1.85 | -1.91 | -1.92 | | | | | | | |
| 4 | 3 | -0.97 | -0.98 | -0.98 | | | | | | | |
| 5 | 1 | -1.31 | -1.40 | -1.48 | | | | | | | |
| 5 | 2 | -1.47 | -1.56 | -1.61 | | | | | | | |
| 5 | 3 | -0.76 | -0.80 | -0.79 | | | | | | | |

Separation of Above and Belowground Living Biomass

Carbon stock of total living biomass can be separated using the ratios listed in Table 7-20. Under the UNFCCC both pools are merged, under the Kyoto Protocol the pools are reported separately (see Chapter 11.3.1.1).

Table 7-20: Ratio the separate total living biomass into above and belowground living biomass. The ratios are retrieved from the NFI (Brändli 2010).

| NFI region | Altitude [m] | Ratio above-/belowground Living Biomass |
|-------------|--------------|---|
| 1 | <601 | 0.22 |
| | 601-1200 | 0.27 |
| | >1200 | 0.35 |
| 2 | <601 | 0.22 |
| | 601-1200 | 0.24 |
| | >1200 | 0.40 |
| 3 | <601 | 0.23 |
| | 601-1200 | 0.28 |
| | >1200 | 0.37 |
| 4 | <601 | 0.25 |
| | 601-1200 | 0.30 |
| | >1200 | 0.40 |
| 5 | <601 | 0.28 |
| | 601-1200 | 0.32 |
| | >1200 | 0.40 |
| Switzerland | <601 | 0.23 |
| | 601-1200 | 0.27 |
| | >1200 | 0.39 |

7.3.4.7 Productive Forests (CC12): Carbon Stocks in Dead Wood, Litter and in Soils

Dead Wood - Carbon Stock

The influence of wood decay on wood density and on carbon content of dead wood has been investigated by Dobbertin and Jüngling (2009) for two dominant tree species in Swiss forests: Norway spruce (*Picea abies*) and beech (*Fagus sylvatica*). They found a significant decrease in relative wood density with increasing decay stage for Norway spruce (30%) and beech (60%) compared to fresh wood. Only small differences in carbon content in dry matter were found between tree species and between fresh wood and dead wood (1.2 - 1.4%), but carbon content remained stable for dead wood across the four decay classes for each species.

The total amount of carbon in the total dead wood pool (TDW) in Switzerland consists of three components:

$$\text{TDW} = \text{CWD} + \text{LIS} + \text{DRoots}$$

where

- CWD (coarse woody debris) contains all wood of dead trees with a diameter of at least 12 cm,
- LIS contains lying small diameter dead wood with a diameter of at least 7 cm determined with the line intersect method and
- DRoots consist of dead coarse roots.

A time series of carbon stocks in dead wood is derived from the soil carbon model Yasso07 (see description in Chapter 7.3.4.8). The value for dead wood stock for 2012 is shown in Table 7-21. Values for dead wood stocks since 1990 are displayed in Table 7-22 under dead organic matter, encompassing dead wood and litter.

Soil and Litter (Organic Soil Horizons) in Mineral Soils - Carbon Stock

Nussbaum et al. (2012) provided updated data for carbon stocks of litter (organic soil horizons L - litter, F - fermentation and H - humus) and soil organic carbon in Swiss forests.

1033 sites of a database stored at WSL distributed among different forest types throughout Switzerland were chosen for this study. Further information on the C content of L horizons was taken from Moeri (2007). By using this dataset and robust geostatistical methods, the authors produced a map of organic carbon stocks of Swiss forest soils. The data for litter and soil carbon stocks are stratified for the five NFI production regions and three elevation levels (Table 7-21).

In the organic soil horizons (litter) of mineral soils in productive and unproductive forests an average carbon stock of 16.7 t C ha⁻¹ was estimated.

In the same study, an average carbon stock in mineral forest soils of 79.9 t C ha⁻¹ in 0-30 cm topsoil was estimated.

Table 7-21 Total dead wood (TDW) stock in Swiss productive forests (CC12) with diameter > 7 cm per spatial stratum in t C ha⁻¹ for 1990 (Didion et al. 2013) and carbon stock in organic soil horizons (litter; used for CC12, CC13) and in soil organic carbon (SOC) of forest soils (used for CC11, CC12, CC13) in mineral soil horizons (0-30 cm) stratified for five NFI production regions and three altitudinal levels (Nussbaum et al. 2012). The average values ± standard error are given.

| NFI region | Altitude [m] | Carbon in dead wood TDW 1990 [t C ha ⁻¹] | Carbon in organic soil horizon (litter) [t C ha ⁻¹] | SOC of mineral topsoil 0-30 cm [t C ha ⁻¹] |
|-------------|--------------|--|---|--|
| 1 | <601 | 5.44 ± 0.09 | 9.51 ± 1.57 | 82.65 ± 3.34 |
| 1 | 601-1200 | 6.79 ± 0.27 | 7.53 ± 0.70 | 102.03 ± 3.56 |
| 1 | >1200 | 5.90 ± 0.08 | 7.76 ± 1.74 | 121.34 ± 5.39 |
| 2 | <601 | 9.12 ± 0.11 | 8.70 ± 0.68 | 55.40 ± 1.55 |
| 2 | 601-1200 | 8.91 ± 0.13 | 11.42 ± 1.45 | 62.12 ± 1.68 |
| 2 | >1200 | 8.91 ± 0.13 | 11.42 ± 1.45 | 122.00 ± 7.07 |
| 3 | <601 | 10.78 ± 0.49 | 7.51 ± 1.25 | 66.10 ± 2.06 |
| 3 | 601-1200 | 7.94 ± 0.15 | 16.29 ± 1.55 | 57.91 ± 2.00 |
| 3 | >1200 | 9.14 ± 0.11 | 26.21 ± 4.77 | 95.78 ± 3.27 |
| 4 | <601 | 8.31 ± 0.36 | 3.15 ± 0.47 | 66.47 ± 2.44 |
| 4 | 601-1200 | 7.29 ± 0.09 | 19.99 ± 2.64 | 74.39 ± 2.42 |
| 4 | >1200 | 7.99 ± 0.12 | 33.37 ± 3.53 | 69.48 ± 1.85 |
| 5 | <601 | 2.83 ± 0.08 | 8.22 ± 1.62 | 102.37 ± 4.07 |
| 5 | 601-1200 | 2.92 ± 0.06 | 11.03 ± 2.11 | 108.99 ± 4.09 |
| 5 | >1200 | 2.82 ± 0.06 | 30.77 ± 5.43 | 107.08 ± 4.11 |
| Switzerland | | 7.30 ± 0.03 | 16.73 ± 0.83 | 79.93 ± 1.52 |

Total Dead Organic Matter DOM- Carbon Stock

According to the Good Practice Guidance LULUCF (IPCC 2003) annual values of carbon stock in dead organic matter are calculated as the sum of carbon in dead wood and of carbon in the organic soil horizons (litter) of mineral forest soils.

Table 7-22 shows annual data of DOM in productive forests (CC12) for 1990-2012. This dataset combines annual estimates of dead wood and litter in productive forests. A time series of litter is derived by adding the temporal changes derived from Yasso07 to the estimates of Nussbaum et al. (2012) as described in Chapter 7.3.4.8 (see also Table 7-21).

Table 7-22 Carbon stock in dead organic matter for CC12, 1990-2012. Highlighted data for 1990 as displayed in Table 7-4.

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: carbon stock in dead organic matter (stockCd,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 14.95 | 14.97 | 14.99 | 15.01 | 15.01 | 15.00 | 15.06 | 15.22 | 15.51 | 15.86 |
| 1 | 2 | 14.32 | 14.53 | 14.73 | 14.91 | 15.06 | 15.21 | 15.31 | 15.33 | 15.23 | 15.06 |
| 1 | 3 | 13.66 | 13.60 | 13.55 | 13.50 | 13.44 | 13.39 | 13.35 | 13.34 | 13.35 | 13.35 |
| 2 | 1 | 17.82 | 17.81 | 17.80 | 17.80 | 17.77 | 17.74 | 17.76 | 17.87 | 18.09 | 18.35 |
| 2 | 2 | 20.33 | 20.29 | 20.25 | 20.21 | 20.16 | 20.11 | 20.13 | 20.28 | 20.58 | 20.95 |
| 2 | 3 | 20.33 | 20.29 | 20.25 | 20.21 | 20.16 | 20.11 | 20.13 | 20.28 | 20.58 | 20.95 |
| 3 | 1 | 18.29 | 18.74 | 19.14 | 19.51 | 19.81 | 20.08 | 20.27 | 20.32 | 20.15 | 19.84 |
| 3 | 2 | 24.23 | 24.02 | 23.84 | 23.67 | 23.50 | 23.35 | 23.22 | 23.10 | 23.00 | 22.90 |
| 3 | 3 | 35.36 | 35.38 | 35.41 | 35.44 | 35.44 | 35.44 | 35.53 | 35.78 | 36.23 | 36.79 |
| 4 | 1 | 11.46 | 11.74 | 12.00 | 12.23 | 12.43 | 12.60 | 12.79 | 13.01 | 13.26 | 13.51 |
| 4 | 2 | 27.27 | 27.29 | 27.29 | 27.30 | 27.30 | 27.30 | 27.32 | 27.34 | 27.37 | 27.40 |
| 4 | 3 | 41.36 | 41.37 | 41.37 | 41.37 | 41.35 | 41.35 | 41.36 | 41.38 | 41.43 | 41.47 |
| 5 | 1 | 11.05 | 10.96 | 10.87 | 10.79 | 10.72 | 10.65 | 10.60 | 10.57 | 10.55 | 10.55 |
| 5 | 2 | 13.95 | 13.94 | 13.92 | 13.91 | 13.90 | 13.89 | 13.88 | 13.85 | 13.79 | 13.73 |
| 5 | 3 | 33.60 | 33.60 | 33.61 | 33.62 | 33.61 | 33.62 | 33.60 | 33.54 | 33.43 | 33.28 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: carbon stock in dead organic matter (stockCd,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 16.21 | 16.52 | 16.80 | 17.07 | 17.33 | 17.58 | 17.74 | 17.75 | 17.58 | 17.31 |
| 1 | 2 | 14.86 | 14.69 | 14.52 | 14.37 | 14.24 | 14.13 | 13.99 | 13.78 | 13.48 | 13.13 |
| 1 | 3 | 13.36 | 13.37 | 13.37 | 13.38 | 13.38 | 13.41 | 13.40 | 13.35 | 13.26 | 13.14 |
| 2 | 1 | 18.59 | 18.80 | 18.99 | 19.19 | 19.38 | 19.58 | 19.68 | 19.62 | 19.35 | 18.98 |
| 2 | 2 | 21.33 | 21.66 | 21.96 | 22.25 | 22.53 | 22.80 | 22.98 | 23.00 | 22.82 | 22.54 |
| 2 | 3 | 21.33 | 21.66 | 21.96 | 22.25 | 22.53 | 22.80 | 22.98 | 23.00 | 22.82 | 22.54 |
| 3 | 1 | 19.48 | 19.15 | 18.85 | 18.61 | 18.39 | 18.22 | 18.07 | 17.96 | 17.89 | 17.86 |
| 3 | 2 | 22.81 | 22.73 | 22.64 | 22.56 | 22.49 | 22.43 | 22.37 | 22.28 | 22.15 | 22.02 |
| 3 | 3 | 37.37 | 37.89 | 38.35 | 38.79 | 39.19 | 39.58 | 39.83 | 39.80 | 39.42 | 38.86 |
| 4 | 1 | 13.75 | 13.97 | 14.17 | 14.36 | 14.55 | 14.76 | 14.87 | 14.83 | 14.59 | 14.24 |
| 4 | 2 | 27.42 | 27.44 | 27.44 | 27.46 | 27.48 | 27.52 | 27.52 | 27.44 | 27.26 | 27.03 |
| 4 | 3 | 41.52 | 41.55 | 41.58 | 41.61 | 41.64 | 41.69 | 41.68 | 41.58 | 41.34 | 41.03 |
| 5 | 1 | 10.56 | 10.56 | 10.56 | 10.56 | 10.57 | 10.57 | 10.57 | 10.55 | 10.50 | 10.45 |
| 5 | 2 | 13.65 | 13.59 | 13.53 | 13.48 | 13.43 | 13.39 | 13.37 | 13.36 | 13.39 | 13.44 |
| 5 | 3 | 33.14 | 33.01 | 32.89 | 32.79 | 32.69 | 32.61 | 32.54 | 32.49 | 32.45 | 32.43 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|---|------|-------|-------|-------|--|--|--|--|--|--|--|
| CC12: carbon stock in dead organic matter (stockCd,i) [t C ha ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 17.04 | 16.81 | 16.59 | | | | | | | |
| 1 | 2 | 12.79 | 12.48 | 12.20 | | | | | | | |
| 1 | 3 | 13.03 | 12.93 | 12.84 | | | | | | | |
| 2 | 1 | 18.62 | 18.29 | 17.99 | | | | | | | |
| 2 | 2 | 22.26 | 22.01 | 21.78 | | | | | | | |
| 2 | 3 | 22.26 | 22.01 | 21.78 | | | | | | | |
| 3 | 1 | 17.86 | 17.85 | 17.84 | | | | | | | |
| 3 | 2 | 21.89 | 21.77 | 21.66 | | | | | | | |
| 3 | 3 | 38.26 | 37.71 | 37.22 | | | | | | | |
| 4 | 1 | 13.90 | 13.58 | 13.29 | | | | | | | |
| 4 | 2 | 26.80 | 26.58 | 26.37 | | | | | | | |
| 4 | 3 | 40.72 | 40.43 | 40.16 | | | | | | | |
| 5 | 1 | 10.39 | 10.34 | 10.30 | | | | | | | |
| 5 | 2 | 13.50 | 13.55 | 13.60 | | | | | | | |
| 5 | 3 | 32.43 | 32.42 | 32.42 | | | | | | | |

7.3.4.8 Productive Forests (CC12): Changes in Carbon stocks in Dead Wood, in Litter and in Soils

Switzerland used the soil carbon model Yasso07 to estimate temporal changes in carbon stocks in soil organic carbon, organic soil horizons (LFH; litter) and in dead wood (TDW) for productive forests (CC12). The implementation of Yasso07 (Tuomi et al. 2009, 2011) in the Swiss GHG inventory is described in detail in Didion et al. (2012, 2013). Didion et al. (2014) demonstrated the validity of the model for application in Swiss forests.

Yasso07 is a model of C cycling in mineral soil, litter and dead wood. For estimating stocks of organic C in mineral soil up to a depth of ca. 100 cm and the temporal dynamics of the C stocks, Yasso07 requires information on C inputs from dead organic matter (i.e. non-woody inputs, including foliage and fine roots, woody inputs, including standing and lying dead wood and dead roots) and climate (temperature, temperature amplitude and precipitation).

By default, Yasso07 does not provide separate estimates of carbon pool sizes for dead wood, litter and soil. In order to report estimates for each pool, the structure of Yasso07 was examined for deriving separate estimates (Didion et al. 2012). Dead wood, litter and soil pools could be correlated with modeled data based on the category of carbon input, i.e., non-woody and woody material, and the five carbon compartments in Yasso07, i.e. four chemical partitions (insoluble, soluble in ethanol, soluble in water or in acid and humus). The approach was validated using independent, measured data (see Didion et al. 2012).

Using annual data for climate and for C inputs obtained from the Swiss NFIs, Yasso07 was used for estimating the annual C stocks in soil, litter and dead wood. Annual C stock changes were calculated from C stocks that were averaged over three years following the recommendation in IPCC (2003; Chapter 7.3.4.2). For an overview, Figure 7-5 shows the mean stock change in Swiss forests for these three carbon pools (soil, litter, dead wood) and the aggregated Yasso total. Annual and stratified values for CC12 can be found in Table 7-23, where net change in dead organic matter encompasses changes in dead wood and in litter. Stocks and stock changes were validated as described in Didion et al. (2013).

Table 7-23 Net carbon stock change in dead organic matter (dead wood and litter) and in mineral soils for CC12, 1990-2012. Highlighted data for 1990 as displayed in Table 7-4.

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: net change in dead organic matter (changeCd,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | -0.01 | 0.08 | 0.03 | 0.06 | -0.05 | -0.05 | 0.05 | 0.22 | 0.35 | 0.33 |
| 1 | 2 | -0.15 | -0.07 | -0.08 | -0.04 | -0.12 | -0.10 | -0.02 | 0.07 | 0.13 | 0.09 |
| 1 | 3 | -0.10 | -0.03 | 0.00 | 0.04 | 0.00 | 0.03 | 0.02 | -0.05 | -0.18 | -0.29 |
| 2 | 1 | -0.16 | -0.03 | -0.05 | -0.01 | -0.13 | -0.12 | 0.00 | 0.19 | 0.32 | 0.29 |
| 2 | 2 | -0.04 | 0.03 | 0.00 | 0.03 | -0.09 | -0.08 | 0.04 | 0.22 | 0.39 | 0.39 |
| 2 | 3 | -0.04 | 0.03 | 0.00 | 0.03 | -0.09 | -0.08 | 0.04 | 0.22 | 0.39 | 0.39 |
| 3 | 1 | 0.83 | 0.78 | 0.66 | 0.58 | 0.38 | 0.33 | 0.26 | 0.12 | -0.17 | -0.40 |
| 3 | 2 | -0.09 | -0.02 | -0.01 | 0.02 | -0.08 | -0.05 | 0.10 | 0.32 | 0.56 | 0.64 |
| 3 | 3 | -0.38 | -0.31 | -0.26 | -0.21 | -0.27 | -0.21 | -0.12 | -0.03 | 0.05 | 0.06 |
| 4 | 1 | 0.59 | 0.56 | 0.45 | 0.40 | 0.25 | 0.22 | 0.27 | 0.36 | 0.44 | 0.41 |
| 4 | 2 | -0.07 | -0.03 | -0.05 | -0.03 | -0.10 | -0.05 | 0.00 | 0.06 | 0.10 | 0.07 |
| 4 | 3 | -0.05 | -0.02 | -0.06 | -0.02 | -0.10 | -0.04 | 0.02 | 0.11 | 0.17 | 0.14 |
| 5 | 1 | -0.39 | -0.29 | -0.24 | -0.20 | -0.19 | -0.15 | -0.08 | 0.02 | 0.12 | 0.17 |
| 5 | 2 | -0.33 | -0.24 | -0.19 | -0.15 | -0.15 | -0.10 | -0.06 | -0.02 | 0.01 | 0.01 |
| 5 | 3 | -0.03 | -0.01 | -0.02 | -0.01 | -0.06 | -0.01 | 0.06 | 0.13 | 0.20 | 0.19 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| CC12: net change in dead organic matter (changeCd,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 0.30 | 0.25 | 0.22 | 0.31 | 0.30 | 0.35 | 0.14 | -0.07 | -0.33 | -0.41 |
| 1 | 2 | 0.07 | 0.05 | 0.01 | 0.07 | 0.06 | 0.11 | 0.03 | -0.05 | -0.13 | -0.12 |
| 1 | 3 | -0.29 | -0.25 | -0.24 | -0.20 | -0.17 | -0.11 | -0.15 | -0.22 | -0.31 | -0.33 |
| 2 | 1 | 0.25 | 0.19 | 0.15 | 0.25 | 0.25 | 0.30 | 0.10 | -0.14 | -0.43 | -0.53 |
| 2 | 2 | 0.37 | 0.31 | 0.26 | 0.32 | 0.31 | 0.36 | 0.18 | -0.06 | -0.35 | -0.45 |
| 2 | 3 | 0.37 | 0.31 | 0.26 | 0.32 | 0.31 | 0.36 | 0.18 | -0.06 | -0.35 | -0.45 |
| 3 | 1 | -0.47 | -0.42 | -0.40 | -0.27 | -0.23 | -0.12 | -0.12 | -0.12 | -0.09 | -0.07 |
| 3 | 2 | 0.65 | 0.58 | 0.48 | 0.47 | 0.43 | 0.48 | 0.27 | -0.09 | -0.54 | -0.73 |
| 3 | 3 | 0.08 | 0.04 | -0.02 | -0.02 | -0.02 | 0.05 | 0.00 | -0.11 | -0.22 | -0.22 |
| 4 | 1 | 0.38 | 0.31 | 0.23 | 0.27 | 0.26 | 0.34 | 0.18 | -0.07 | -0.37 | -0.48 |
| 4 | 2 | 0.06 | 0.04 | 0.00 | 0.05 | 0.06 | 0.15 | 0.06 | -0.11 | -0.32 | -0.38 |
| 4 | 3 | 0.11 | 0.07 | 0.02 | 0.06 | 0.10 | 0.20 | 0.10 | -0.07 | -0.28 | -0.31 |
| 5 | 1 | 0.16 | 0.13 | 0.09 | 0.11 | 0.07 | 0.12 | 0.09 | 0.05 | -0.02 | -0.04 |
| 5 | 2 | 0.00 | -0.01 | -0.04 | -0.01 | -0.03 | 0.03 | 0.03 | 0.03 | 0.01 | 0.03 |
| 5 | 3 | 0.18 | 0.15 | 0.08 | 0.09 | 0.06 | 0.16 | 0.13 | 0.11 | 0.06 | 0.09 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|---|------|-------|-------|-------|--|--|--|--|--|--|--|
| CC12: net change in dead organic matter (changeCd,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | -0.34 | -0.29 | -0.28 | | | | | | | |
| 1 | 2 | -0.04 | -0.07 | -0.07 | | | | | | | |
| 1 | 3 | -0.28 | -0.29 | -0.26 | | | | | | | |
| 2 | 1 | -0.45 | -0.38 | -0.35 | | | | | | | |
| 2 | 2 | -0.36 | -0.32 | -0.29 | | | | | | | |
| 2 | 3 | -0.36 | -0.32 | -0.29 | | | | | | | |
| 3 | 1 | 0.05 | 0.01 | 0.00 | | | | | | | |
| 3 | 2 | -0.69 | -0.65 | -0.58 | | | | | | | |
| 3 | 3 | -0.13 | -0.19 | -0.17 | | | | | | | |
| 4 | 1 | -0.39 | -0.37 | -0.33 | | | | | | | |
| 4 | 2 | -0.30 | -0.31 | -0.29 | | | | | | | |
| 4 | 3 | -0.23 | -0.25 | -0.26 | | | | | | | |
| 5 | 1 | 0.00 | -0.01 | -0.01 | | | | | | | |
| 5 | 2 | 0.09 | 0.05 | 0.04 | | | | | | | |
| 5 | 3 | 0.19 | 0.12 | 0.10 | | | | | | | |

(Table 7-23 continued)

| NFI | Alt. | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| 1 | 2 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| 1 | 3 | 0.000 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.002 |
| 2 | 1 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| 2 | 2 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| 2 | 3 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| 3 | 1 | 0.003 | 0.004 | 0.004 | 0.005 | 0.005 | 0.005 | 0.006 | 0.006 | 0.006 | 0.005 |
| 3 | 2 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.001 | 0.001 |
| 3 | 3 | 0.000 | 0.000 | 0.000 | 0.000 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | 0.000 |
| 4 | 1 | 0.002 | 0.003 | 0.003 | 0.003 | 0.004 | 0.004 | 0.004 | 0.004 | 0.005 | 0.005 |
| 4 | 2 | 0.001 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| 4 | 3 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| 5 | 1 | -0.001 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.003 | -0.003 | -0.002 | -0.002 |
| 5 | 2 | -0.001 | -0.001 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 |
| 5 | 3 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.002 |

| NFI | Alt. | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 0.001 | 0.001 | 0.001 | 0.001 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.001 |
| 1 | 2 | 0.000 | 0.000 | 0.000 | 0.000 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| 1 | 3 | -0.002 | -0.002 | -0.002 | -0.002 | -0.003 | -0.003 | -0.003 | -0.003 | -0.003 | -0.003 |
| 2 | 1 | 0.000 | 0.000 | 0.000 | 0.000 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.000 |
| 2 | 2 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| 2 | 3 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| 3 | 1 | 0.005 | 0.005 | 0.004 | 0.004 | 0.004 | 0.004 | 0.004 | 0.004 | 0.004 | 0.004 |
| 3 | 2 | 0.001 | 0.002 | 0.002 | 0.002 | 0.002 | 0.003 | 0.003 | 0.003 | 0.002 | 0.002 |
| 3 | 3 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| 4 | 1 | 0.005 | 0.005 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| 4 | 2 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.000 |
| 4 | 3 | 0.001 | 0.001 | 0.001 | 0.001 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| 5 | 1 | -0.002 | -0.002 | -0.002 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 | -0.001 |
| 5 | 2 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.002 | -0.001 | -0.001 | -0.001 |
| 5 | 3 | 0.002 | 0.002 | 0.002 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 |

| NFI | Alt. | 2010 | 2011 | 2012 | | | | | | | |
|--|------|--------|--------|--------|--|--|--|--|--|--|--|
| CC12: net change in mineral soil (changeCs,i) [t C ha ⁻¹ yr ⁻¹] | | | | | | | | | | | |
| 1 | 1 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 1 | 2 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 1 | 3 | -0.003 | -0.003 | -0.003 | | | | | | | |
| 2 | 1 | 0.000 | -0.001 | -0.001 | | | | | | | |
| 2 | 2 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 2 | 3 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 3 | 1 | 0.004 | 0.004 | 0.004 | | | | | | | |
| 3 | 2 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 3 | 3 | 0.000 | 0.000 | -0.001 | | | | | | | |
| 4 | 1 | 0.005 | 0.005 | 0.005 | | | | | | | |
| 4 | 2 | 0.000 | 0.000 | 0.000 | | | | | | | |
| 4 | 3 | 0.001 | 0.001 | 0.001 | | | | | | | |
| 5 | 1 | -0.001 | -0.001 | -0.001 | | | | | | | |
| 5 | 2 | -0.001 | -0.001 | -0.001 | | | | | | | |
| 5 | 3 | 0.003 | 0.004 | 0.004 | | | | | | | |

A large source of uncertainties in simulated estimates of litter and soil carbon stocks is related to the calculation of the litter production of trees (de Wit et al. 2006), which determines the C inputs that drive the Yasso07 simulation. Data for soil and litter carbon are

retrieved from Nussbaum et al. 2012 (Table 7-21). Deadwood (TDW) stocks are reported based on estimates obtained with Yasso07 (annual values integrated in DOM in Table 7-23).

Carbon stock changes in the soil pool are small (Figure 7-5 SO_{csc} and Table 7-23), which corresponds to information from measurements (see Chapter 7.3.6). Carbon stock changes in litter are higher and more erratic than changes in the dead wood and soil pools (Figure 7-5 LFH_{csc}). This is expected since non-woody material decomposes faster than dead wood (Tuomi et al. 2011), and there is a higher interannual variability in the production of foliage (Etzold et al. 2011). The C stock change in the dead wood pool is to a large extent driven by the increase in the dead wood volume following the hurricane Lothar (1999). As Lothar occurred between the NFI2 (1993-1995) and NFI3 (2004-2007), it strongly affects the results of the change analysis for dead wood volume in the periode NFI2-3 and also between NFI3-4, where the mortality rate decreased again. Large-scale disturbance events like Lothar that occur between two consecutive NFIs strongly affect the estimates of annually accumulating mass of carbon in dead wood that drives the Yasso07 simulation. This bias is expected to disappear following the switch to a continuous sampling approach in the NFI4 (Braendli und Speich 2011).




Figure 7-5: Mean carbon stock change (CSC) based on averaged (avg) annual carbon stocks for soil organic carbon (SO), Litter (LFH), dead wood (CWD) and their sum (ALL). Note the different scale of the y-axis for SO_{csc,avg} in comparison to LFH_{csc,avg}, CWD_{csc,avg} and ALL_{csc,avg}. Negative values indicate a sink of carbon, positive values a carbon source.

7.3.4.9 Unproductive Forests (CC13): Stocks and Changes in Stocks of Living Biomass, Dead wood, Litter and Soil Carbon

Unproductive forests consist of brush forests and forest on unproductive areas. FOEN (2014f) shows some examples of unproductive forests in Switzerland.

For transparency reasons, both productive and unproductive forest areas are reported separately. However, there is only scarce information available about unproductive forests. In unproductive forests, wood is not harvested for economical reasons. Only in exceptional cases (e.g. wood log blocks a hiking trail) there can be an intervention where the log is moved, but not removed from the stand. Moreover, since yearly harvesting amounts from forest statistics (FOEN 2013k) are divided over the productive forests, total harvesting in Swiss forests is accounted for under CC12 remaining, thus all harvesting amounts are accounted for.

The national forest inventory does not incorporate unproductive stands in its regular inventory scheme because 1) the plots are difficult to access or it is not possible to carry out precise measurements (brush forests), (2) the plots are inaccessible or the NFI forest definition is not fulfilled (forest on unproductive areas).

- Brush forests: Since brush forests have no direct economical value in terms of wood harvest, an inventory of these stands has not been attributed high priority. During NFI3, some plots in brush forests have been visited for the first time, but only a limited number of parameters like tree species, stem diameter and crown cover have been determined. There are no NFI estimates available on carbon stocks of living biomass, since the allometric functions to calculate biomass from stem diameter in brush forests are not established yet.
- Forest on unproductive areas:
 - a) Inaccessible stands are forests which cannot be visited because of safety reasons (see description in Brändli 2010, p. 89). They are mainly located in the Alps and often grow on sites of low productivity: rocky sites, sites at high altitude near the tree line with a short vegetation period and low biological activity;
 - b) Unproductive forests not covered by NFI: after the review of its Initial Report (FOEN 2006h), Switzerland had to apply a forest definition for reporting activities under KP Art. 3.3 and Art. 3.4, which is different from the definition applied by the Swiss NFI and the Land Use Statistics AREA. The same definition is used for reporting under the UNFCCC and under the Kyoto Protocol. Because the country definition (NFI and AREA) was not in line with the specific requirements of the Kyoto Protocol forest definition, Switzerland had to develop an approach to classify certain AREA categories as forest. Those areas are not covered by the regular NFI and are situated in the threshold range between forests and alpine pastures with woody biomass of very low productivity. More specifically, it concerns the combination category "alpine pastures with a cluster of trees" in Table 7-6 (LU=242 and LC 47).

Unproductive Forests (CC13): Carbon Stocks in Living Biomass

Brush Forest

Brush forests in Switzerland mainly consist of *Alnus viridis* and horizontal *Pinus mugo* var. *prostrata*. No NFI data are available to derive their growing stock. Therefore, following assumptions were met to describe the stocks: 4'000 trees per ha, average height of 2.5 m and an average diameter at 1.3 m of 10 cm. Hence, an average growing stock (> 7 cm diameter) of 40 m³ ha⁻¹ was estimated. Multiplied by the mean BCEF for coniferous trees of 0.64 (see Table 7-28; Thürig and Herold 2013), an average biomass for brush forest of 25.7 t ha⁻¹, which translates to 12.9 t C ha⁻¹ (using the IPCC default carbon content of 50%) was estimated.

Forest on Unproductive Areas

Forest on unproductive areas in Switzerland is mainly located in the Alps and the Southern Alps. In those forests, no NFI data are available to derive growing stocks. As those forests are assumed to grow preferably on bad site conditions, we assume an average growing stock (> 7 cm diameter) of 150 m³ ha⁻¹. Multiplied by the mean BCEF for coniferous trees of 0.64 (see Table 7-28; Thürig and Herold 2013), an average biomass for forest on unproductive areas of 96.4 t ha⁻¹ was estimated, which translates to 48.2 t C ha⁻¹ (using the IPCC default carbon content of 50%).

Carbon Stocks of Living Biomass at CC13: Weighted Means

The carbon content of unproductive forest was calculated as a weighted average of brush forest and forest on unproductive areas per spatial stratum:

$$[\text{weighted C content}]_i = \text{RSi} * \text{CS} + (1 - \text{RSi}) * \text{CI}$$

where RSi is the rate of the brush forest per spatial stratum i,

CS is the carbon content of brush forest (12.9 t C ha⁻¹),

CI is the carbon content of forest on unproductive areas (48.2 t C ha⁻¹).

Table 7-24 shows the carbon content per spatial stratum in t C ha⁻¹.

Table 7-24 Rate of brush forest and forest on unproductive areas and the resulting weighted carbon content in t C ha⁻¹ of Swiss unproductive forests (CC13) specified for all spatial strata. The area of forest on unproductive sites is derived from NFI 2 (Brassel and Brändli 1999).

| NFI region | Altitude [m] | Brush forest [ha] | Forest on unproductive area [ha] | Total unproductive forest [ha] | Rate of brush forest | Weighted C content [t C ha ⁻¹] |
|------------|--------------|-------------------|----------------------------------|--------------------------------|----------------------|--|
| 1 | <601 | 25 | 356 | 381 | 0.07 | 45.90 |
| | 601-1200 | 1 | 1'780 | 1'781 | 0.00 | 48.20 |
| | >1200 | 1 | 178 | 179 | 0.01 | 48.03 |
| 2 | <601 | 25 | 534 | 559 | 0.05 | 46.64 |
| | 601-1200 | 25 | 356 | 381 | 0.07 | 45.90 |
| | >1200 | 1 | 0 | 1 | 1.00 | 12.86 |
| 3 | <601 | 25 | 356 | 381 | 0.07 | 45.90 |
| | 601-1200 | 50 | 3'204 | 3'254 | 0.02 | 47.68 |
| | >1200 | 2'100 | 1'780 | 3'880 | 0.54 | 29.08 |
| 4 | <601 | 100 | 356 | 456 | 0.22 | 40.47 |
| | 601-1200 | 1'925 | 4'984 | 6'909 | 0.28 | 38.37 |
| | >1200 | 36'925 | 7'120 | 44'045 | 0.84 | 18.58 |
| 5 | <601 | 200 | 534 | 734 | 0.27 | 38.59 |
| | 601-1200 | 2'550 | 3'560 | 6'110 | 0.42 | 33.46 |
| | >1200 | 16'875 | 5'162 | 22'037 | 0.77 | 21.14 |

Unproductive Forests (CC13): Carbon Stocks in Litter, Soil and Dead Wood

As stated on previous page CC13, consists of very different forests and data are hardly available. Carbon stocks in litter and mineral soil under unproductive forests reveal therefore a very high spatial heterogeneity. Brush forests are mainly situated on soils comparable to the ones under productive forests, others are situated at stony sites with very thin layers of organic material. Data on carbon stocks of litter and soil are not available. We therefore assumed carbon stocks of soil carbon and litter to be the same as for productive forests CC12 and are derived from Nussbaum et al. (2012; Table 7-21).

So far, there are no data available for dead wood stocks in unproductive forests (CC13). We conservatively assume no dead wood on CC13 sites. Therefore, the amount of dead organic matter (litter and dead wood) on CC13 sites equals the carbon stock in litter of mineral forest soils. Dead organic matter values for CC13 are listed in Table 7-4.

Unproductive Forests (CC13): Changes in Carbon Stocks of Living Biomass

There are a few case studies on carbon stocks, but just like in neighboring countries with forests in mountainous regions, there are no repeated forest inventory data available for these unproductive forests (also known as “mountain forest without harvest”) available.

As no harvesting is conducted in unproductive forests, gross growth and cut and mortality of unproductive forest are assumed to be in balance. This approach is confirmed by two studies in which basal area and crown cover is used as a proxy for the stock of living biomass. An increase in basal area or crown cover, respectively, is positively correlated with an increase in living biomass (e.g. Nowak and Crane 2002). Living biomass in brush forests is increasing during the stage of establishment: the stand develops from a stand with grasses, herbs and some shrubs towards a stand dominated by shrubs and with a denser crown cover. A decrease in crown cover in unproductive forests is observed when natural disturbances like avalanches or rock fall partially damage the stand.

- Huber and Thürig (2014) analyzed the available data on diameters of the terrestrial inventory NFI3 and NFI4a+. The authors found that the number of trees has increased over this 6 years period. Since no allometric functions are available for these stands, it is not possible to calculate stocks from these data. The authors calculated an increase in the mean basal area from 4.59 m² ha⁻¹ in 2006 to 5.47 m² ha⁻¹ in 2012.
- Ginzler (2014) analyzed the crown cover density of 135 aerial photographs between 2006 (NFI3) and 2011 (NFI4a) and found no statistical change in crown cover density of well-established, existing shrub forests. The terrestrial NFI data, however, showed a slight increase in the basal area of trees in brush forests.
- In addition, the study of Huber and Frehner (2013) shows that the expansion of *Green Alder* in eastern Switzerland has doubled in the past 75 years. Especially in the Alps or at unproductive sites, brush forests are expanding as summer pastures are abandoned. At these sites, an increase in crown cover is observed which correlates with an increment in carbon stocks. A literature review by Huber and Frehner (2012; for an overview see FOEN 2014f) shows that *Green Alder* has in general a strong annual gross growth, not only in very young stands, and that stands of *Green Alder* can be very vital at an age of over 100 years.

Considering the observed dynamics in brush forests, we conclude that living biomass in unproductive forests is not a net source of carbon and report living biomass to be in balance (conservative estimate, Tier 1 approach). In Table 7-3 and in the CRF-tables, this is transcribed into gains = losses = 0.

Unproductive Forests (CC13): Changes in Carbon Stocks of Litter, Mineral Soils, Dead Wood and organic soils

There are no repeated measurements of carbon stocks in litter, mineral soils and dead wood.

Above, transparent and verifiable information has been given that in Switzerland living biomass in brush forest is increasing. An increase in biomass leads to an increase in litter and dead wood production, which again can lead to an accumulation in soil carbon. Based on that, we can conclude that litter, mineral soils and dead wood are not a net source of carbon. Carbon changes in litter, mineral soils and dead wood are conservatively reported to be zero (Tier 1 approach). This assumption is further supported by the following arguments:

- The areas of CC13 occur on higher altitudes where microbiological processes in soils are slow (Hagedorn et al. 2010; Davidson and Janssens 2006).
- In addition, unproductive forests grow on poor or rocky sites with thin or no organic layer. Brush forest protect the soils; in particular, Alder brush is not even destroyed by

avalanches, rock fall when rock size is small or medium (Huber and Frehner 2014). By stabilizing soils, brush forests act as a perfect protection against soil erosion (Richard 1995; Stangle 2004).

- Further, by fixing nitrogen with its nitrogen-fixing root nodules, *Green Alder* has an ameliorative effect on the soil. Amelioration of soils enables an increase in biomass production which on the other hand increases the amount of litter and dead wood and finally leads to accumulation of soil carbon.
- Since there is no active logging on these unproductive stands, there is also no human impact on the soils, litter and dead wood.

By providing this transparent and verifiable information (survey of peer-reviewed literature and reasoning based on sound knowledge of likely system responses; as requested by the ERT, see Table 1-12), we conclude that the requirements of IPCC GPG Chapter 4.2.3.1 (IPCCC 2003) application of Tier 1 are fulfilled.

For conversions within forest land (CC13 to CC12 and CC12 to CC13), no changes in carbon stocks of litter and soil carbon of mineral soils are calculated because carbon stocks of litter and soil carbon are equal for CC12 and CC13. The reason why these stocks are assumed to be equal is (1) because data are not available (see above) and (2) because it leads to a conservative estimate of changes in these pools. With the exception of brush forests, it is very likely that carbon stocks of litter and soil carbon are smaller under unproductive forests CC13 than under productive forests CC12. As the area changing from CC13 to CC12 is larger than from CC12 to CC13 (see Table 7-9), applying the stock-change method (see Table 7-3) with equal stocks for litter and soil carbon under CC12 and CC13 is a conservative estimate.

Emissions from organic soils are accounted for using a Tier 1 approach. A conservative estimate (Tier 1; no changes) is applied for all pools of unproductive forests.

7.3.4.10 Afforestations (CC11)

Living Biomass: Growing Stock and Changes in Growing Stock

The average growing stock and growth of afforestations were empirically assessed from NFI 1 and NFI 2 data, specifically with those stands that were approximately 10 years old in the first NFI and 20 years old in the second NFI. The average growing stock of those 20 year old stands was derived from NFI 2. The NFI data were therefore stratified per altitudinal level. The growing stock of forest stands below 600 m was on average $90 \text{ m}^3 \text{ ha}^{-1}$. The growing stock on sites between 600 and 1200 m was assumed to be one-third smaller ($60 \text{ m}^3 \text{ ha}^{-1}$) than on sites below 600 m, and two-thirds smaller on sites above 1200 m ($30 \text{ m}^3 \text{ ha}^{-1}$). As trees below 12 cm diameter at breast height (DBH) were not measured in the NFI, the growing stock of 10 year old stands below 600 m was assumed to be $2 \text{ m}^3 \text{ ha}^{-1}$. Within the first few years of stand age, the growing stock was assumed to develop exponentially. The development of the growing stock between 10 and 20 years on sites below 600 m was simulated by calibrating a logistical growth function. To simulate the development of growing stock on intermediate and poor sites, growing stock was assumed to develop one-third slower on intermediate, and two-thirds slower on poor sites. The annual growth was calculated as the difference between growing stocks of two following years. These assumptions are not valid for single stands, but can be applied as a rough simplification. Table 7-25 shows the simulated growing stock and growth for the three altitudinal levels.

Table 7-25 Estimated average growing stock and annual growth of forest stands in stem-wood over bark including stock up to 20 years (CC11) specified per altitudinal zone. Bench marks derived from NFI 1 and NFI 2 (see text above) in bold letters.

| Stand age [yr] | < 601 m altitude | | 601 - 1200 m altitude | | > 1200 m altitude | |
|-------------------|---|---|---|---|---|---|
| | Growing stock [m ³ ha ⁻¹] | Growth [m ³ ha ⁻¹ year ⁻¹] | Growing stock [m ³ ha ⁻¹] | Growth [m ³ ha ⁻¹ year ⁻¹] | Growing stock [m ³ ha ⁻¹] | Growth [m ³ ha ⁻¹ year ⁻¹] |
| 0-9 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10 | 2 | 2 | 0 | 0 | 0 | 0 |
| 11 | 7 | 5 | 0 | 0 | 0 | 0 |
| 12 | 13 | 6 | 1 | 1 | 0 | 0 |
| 13 | 19 | 6 | 5 | 4 | 0 | 0 |
| 14 | 27 | 8 | 10 | 5 | 0 | 0 |
| 15 | 35 | 8 | 16 | 6 | 1 | 1 |
| 16 | 44 | 9 | 23 | 7 | 5 | 4 |
| 17 | 54 | 10 | 31 | 8 | 10 | 5 |
| 18 | 66 | 12 | 40 | 9 | 16 | 6 |
| 19 | 78 | 12 | 50 | 10 | 23 | 7 |
| 20 | 90 | 12 | 60 | 10 | 30 | 7 |

To convert the estimated growing stock (m³ ha⁻¹) and growth (m³ ha⁻¹ year⁻¹), both expressed in volume units, into tonnes of carbon, the following equations were applied:

Carbon stock in living biomass = Average growing stock * BCEF * C content

Growth of living biomass = Average growth * BCEF * C content

Where

- C content: Carbon to total biomass ratio. The IPCC default of 50% was applied (IPCC 2003; p 3.25)
- BCEF: Biomass conversion and expansion factor converting the volume of growing stock and the volume of net annual increment to total tree biomass and total tree biomass growth, respectively; an average value for coniferous and deciduous trees is taken from Burschel et al. (1993).

Table 7-26 Carbon stock in living biomass (stem-wood over bark including stock without branches) and growth of living biomass in afforestations (CC11) specified per altitudinal zone. BCEF taken from Burschel et al. (1993).

| Altitude [m] | Average growing stock [m ³ ha ⁻¹] | Average growth [m ³ ha ⁻¹ yr ⁻¹] | BCEF | Carbon content | Carbon stock in living biomass [t C ha ⁻¹] | Growth of living biomass [t C ha ⁻¹ yr ⁻¹] |
|-----------------|---|---|------|----------------|---|--|
| <601 | 21.7 | 4.5 | 0.7 | 0.5 | 7.84 | 1.63 |
| 601-1200 | 11.8 | 3 | 0.7 | 0.5 | 4.3 | 1.09 |
| >1200 | 4.25 | 1.5 | 0.7 | 0.5 | 1.61 | 0.57 |

Litter and Dead Wood (DOM): Carbon Stock and Carbon Stock Changes

In Switzerland, afforestations (CC11) occur mostly on grasslands and settlements (see Table 7-9 in Chapter 7.2.3) where there is no litter and no dead wood (IPCC 2003, p. 3.105).

Therefore, assuming no carbon stock in dead organic matter on afforestation sites, we follow the Tier 1 approach in terms of IPCC good practice (IPCC 2003, Sect. 3.1.5) and consistently

report no changes in the litter and dead wood pool after a land-use change to afforestation (see Chapter 11.3.1.2 for a deepened discussion)

Soil: Carbon Stock and Carbon Stock Changes

The estimates for soil carbon stocks from Nussbaum et al. (2012) are used for afforestations (CC11; see Table 7-4 and Table 7-21). Based on these carbon stocks, carbon stock changes are calculated with the stock-change method (see 7.1.3.2 and Table 7-3).

7.3.4.11 Organic Soils

Organic Soils - Carbon Stock

No specific information is available related to carbon stocks in organic soils under forest land. Therefore, the value calculated for cropland and permanent grassland based on Leifeld et al. (2003, 2005) is adopted for forest land, including CC11, CC12 and CC13. The approach uses measured carbon stocks in Swiss organic soils. The mean soil organic carbon stock (0-30 cm) for organic soils is $240 \pm 48 \text{ t C ha}^{-1}$.

Organic Soils - Changes in Carbon Stocks due to Drainage

Drainage of forests is not a permitted practice in Switzerland (Swiss Confederation 1991). There are no nation-wide survey data available. It is possible that small parts of the Swiss forest have been drained before 1990 or have been established on drained areas. We conservatively report all organic forest soils to be drained (which is definitely an overestimation).

In order to calculate CO₂ emissions due to drainage, we used equation 3.2.15 of the GPG for LULUCF (IPCC 2003) and applied the default emission factor of $0.68 \text{ t C ha}^{-1} \text{ yr}^{-1}$.

7.3.4.12 N₂O Emissions from N Fertilization and Drainage of Soils

Fertilization of forests is prohibited by the Swiss forest law and adherent ordinances (Swiss Confederation 1991, 1992). Additionally, the "Ordinance on Chemical Risk Reduction" (Swiss Confederation 2005) prohibits the application of fertilizers, including liming, in forests. Therefore, no emissions are reported in CRF-table 5(I A).

There are no data available on non-CO₂ emissions from organic soils in Switzerland. As in IPCC (2003) only a basis for future methodological development is included, Switzerland does not provide such estimates. No non-CO₂ emissions are estimated ("NE") in CRF-table 5(II A) (see also Chapter 7.6.4.4).

7.3.4.13 Emissions from Wildfires

Data on wildfires affecting Swiss forest land are obtained from cantonal authorities and are compiled by FOEN (FOEN 2013k). Table 7-27 shows the annual number of fires and the burnt area from 1990 to 2012.

As controlled burning is not allowed in Switzerland all fires are assigned to "wildfires". The number and area of the fires are assigned to productive forests. This is a conservative estimate, since the "available fuel" of productive forests is higher than the carbon stocks of afforestations and unproductive forests.

Using the default emission factor of $7.10 \text{ g (kg combusted biomass)}^{-1}$, an emission factor for CH₄ of $0.903 \text{ Mg CH}_4 \text{ ha}^{-1}$ is calculated (IPCC 2003, equation 3.2.20 and table 3A.1.16).

For N_2O , the default emission factor of $0.11 \text{ g (kg combusted biomass)}^{-1}$ is applied (IPCC 2003, Table 3A.1.16).

The mass of available fuel considered for calculating the emissions, depends on the greenhouse gas reported:

- (1) For reporting CH_4 and N_2O emissions from wildfires, the mass of available fuel encompasses carbon stock of living biomass, litter and dead wood.
- (2) For reporting CO_2 emissions from wildfires, the mass of available fuel only encompasses carbon stock of litter. Losses in living biomass and dead wood due to wildfires are already reflected in the NFI dataset and included in CRF-table 5A. Yearly values of these losses are included in the data shown in Table 7-19 under "loss of living biomass" and in Table 7-23 under "net change in dead organic matter", respectively.

On average, the amount of living biomass amounts to $119.40 \text{ t C ha}^{-1}$ or $238.81 \text{ t biomass ha}^{-1}$. This value has been derived from the mean growing stock in NFI 1, NFI 2, NFI 3 and NFI 4a (Brassel and Brändli 1999, Brändli 2010, Abegg et al. 2012).

On average in Swiss forests, the amount of litter amounts to 16.7 t C ha^{-1} or $33.40 \text{ t biomass ha}^{-1}$ (Nussbaum et al. 2012). The amount of dead wood amounts on average to 5.23 t C ha^{-1} or $10.46 \text{ t biomass ha}^{-1}$. These values are derived from Table 7-21 as weighted averages over all spatial strata.

The fraction of the biomass combusted is 0.45 (IPCC 2003, Table 3A.1.12). Inserting these values in equation 3.2.20 of IPCC (2003), the emissions shown in Table 7-27 are calculated.

CH_4 and N_2O emissions caused by wildfires are reported in CRF-table 5(V). CO_2 emissions caused by wildfires are included in CRF-table 5A (as described above) and 5(V). In Table 5(V), the emissions from all forest fires are reported under 5(V)A1, because it is not known which fires occur on forest land remaining forest land and which on land converted to forest land. Consequently, 5(V)A2 has the notation key "IE".

Table 7-27 Productive forest land affected by wildfires (FOEN 2013k) and resulting GHG emissions 1990-2012.

| Year | Number | Area burnt [ha] | CH ₄ [Mg] | N ₂ O [Mg] | CO ₂ [Mg] |
|------|--------|-----------------|----------------------|-----------------------|----------------------|
| 1990 | 216 | 1102 | 995.24 | 15.42 | 25'358.04 |
| 1991 | 157 | 148 | 133.66 | 2.07 | 3'405.62 |
| 1992 | 111 | 52 | 46.96 | 0.73 | 1'196.57 |
| 1993 | 99 | 42 | 37.93 | 0.59 | 966.46 |
| 1994 | 52 | 293 | 264.62 | 4.10 | 6'742.20 |
| 1995 | 56 | 438 | 395.57 | 6.13 | 10'078.79 |
| 1996 | 61 | 233 | 210.43 | 3.26 | 5'361.55 |
| 1997 | 77 | 1511 | 1364.62 | 21.14 | 34'769.52 |
| 1998 | 88 | 249 | 224.88 | 3.48 | 5'729.72 |
| 1999 | 31 | 9 | 8.13 | 0.13 | 207.10 |
| 2000 | 41 | 36 | 32.51 | 0.50 | 828.39 |
| 2001 | 39 | 37 | 33.42 | 0.52 | 851.40 |
| 2002 | 75 | 410 | 370.28 | 5.74 | 9'434.48 |
| 2003 | 189 | 564 | 509.36 | 7.89 | 12'978.16 |
| 2004 | 46 | 20 | 18.06 | 0.28 | 460.22 |
| 2005 | 97 | 47 | 42.45 | 0.66 | 1'081.51 |
| 2006 | 70 | 101 | 91.22 | 1.41 | 2'324.10 |
| 2007 | 64 | 234 | 211.33 | 3.27 | 5'384.56 |
| 2008 | 47 | 53 | 47.87 | 0.74 | 1'219.58 |
| 2009 | 52 | 42 | 37.93 | 0.59 | 966.46 |
| 2010 | 59 | 25 | 22.58 | 0.35 | 575.27 |
| 2011 | 77 | 167 | 150.82 | 2.34 | 3'842.83 |
| 2012 | 56 | 22 | 19.87 | 0.31 | 506.24 |

7.3.4.14 NMVOC Emissions

Estimates for annual biogenic emissions of NMVOC in Switzerland for forests (and natural grassland) are available in SAEFL (1996a): The values are 92.0 Gg for coniferous forests, 2.4 Gg for deciduous forests and 0.61 Gg for forest fires. These numbers are based on a study from Andreani-Aksoyoglu and Keller (1995). Approximately 97% of the total emissions are monoterpene and the rest consists of isoprene (Keller et al. 1995).

7.3.5 Uncertainties and Time-Series Consistency

Uncertainties

For living biomass, the following information on uncertainty related to 2012 was used:

- Stem wood of growth and cut & mortality (C&M) in NFI 4a+ and differences NFI 3-4a+:
mean growth $11.38 \text{ m}^3 \text{ ha}^{-1}$, mean C&M $-8.97 \text{ m}^3 \text{ ha}^{-1}$, resulting mean net change in stem volume $2.41 \text{ m}^3 \text{ ha}^{-1}$
relative uncertainty: 2% for growth and 5% for C&M (Thürig 2014)
resulting relative uncertainty of mean net change in volume: 20.9%
- Carbon content in solid wood: 5-10% (background: Lamtom and Savidge 2003, assessment of carbon content in wood; Monni et al. 2007, 2%)
- Wood density: guess 10-20% (background: Lamtom and Savidge 2003)
- Biomass expansion function (for the Swiss GHGI, allometric functions for individual trees are applied): The uncertainty is estimated to be 30% (Monni et al. 2007, Appendix 1, 2.7-21.3%; Vanninen and Mäkelä 1999; Cronan 2003; Helmisaari and Hallbäcken 1998).

Thus, the total uncertainty of net carbon stock change in living biomass ($U_{\text{liv.biom}}$) in terms of carbon per unit area can be calculated as:

- addition of relative uncertainties to derive uncertainty for gains and losses following equation 6.4 in chapter 'Quantifying Uncertainties in Practice' (IPCC 2000)
$$U_{\text{liv.biom}} = (20.9^2 + 10^2 + 15^2 + 30^2)^{0.5} = 40.7\%$$
- calculation of the absolute uncertainty, based on the mean gain of $0.72 \text{ t C ha}^{-1} \text{ yr}^{-1}$:
$$U_{\text{liv.biom}} = 0.29 \text{ t C ha}^{-1} \text{ yr}^{-1}.$$

The uncertainty in the estimates of annual stock changes derived with the Yasso07 model originates from the following sources:

- spatial climate data interpolation
- C input estimates obtained from the NFI (measurement errors, allometries, etc.)
- decomposition parameters used in the Yasso07 model

The uncertainty associated with the climate data could not be estimated.

The uncertainty associated with C inputs (dead wood production and litterfall) was estimated based on estimates of uncertainty in a) litter turnover rates (Wutzler and Mund 2007), b) wood densities of deadwood in different decay stages (Dobbertin and Jüngling 2009), and c) spatial uncertainty in the NFI data approximated based on the estimation error for tree volume reported for the NFI (see chapter 1.4 in Brändli 2010). Based on the mean C inputs and the estimated uncertainty, a distribution of possible values was obtained. Finally, the combined uncertainty from these sources was calculated:

- The uncertainty in the Yasso07 parameters was estimated based on a Markov Chain Monte Carlo approach (see also Tuomi et al. 2011). A distribution of possible parameter values was provided by A. Lehtonen, Finnish Forest Research Institute METLA .
- The uncertainty of Yasso07 estimates on C stocks and C stock changes in different pools, resulting from the uncertainty of C inputs and of model parameters, was obtained through Monte Carlo simulations: 30 values for C inputs and 30 parameter combinations were selected randomly and the combined uncertainty in Yasso07 estimates of C stocks and C stock changes in the soil, litter, and dead wood pools was calculated as (described in Didion et al. 2013).

- Based on this approach the absolute uncertainty of the estimates of C stock changes are (Didion et al. 2013):

$$U_{\text{Soil}} = 0.000081, U_{\text{Litter}} = 0.0065 \text{ and } U_{\text{Deadwood}} = 0.0047 \text{ t C ha}^{-1} \text{ yr}^{-1}.$$

The absolute uncertainty of the total C stock change is:

$$U_{\text{tot}} = (U_{\text{liv.biom}}^2 + U_{\text{Soil}}^2 + U_{\text{Litter}}^2 + U_{\text{Deadwood}}^2)^{0.5} = 0.29 \text{ t C ha}^{-1} \text{ yr}^{-1}.$$

The mean total C stock change in 2012 is the sum of the mean changes of living biomass, soil, litter and deadwood: $0.72 + 0.00 - 0.03 - 0.22 = 0.47 \text{ t C ha}^{-1} \text{ yr}^{-1}$. Thus, the resulting relative uncertainty of the C stock changes for forest land is 63%.

Combined uncertainties of the activities under the Kyoto Protocol are shown in Table 11-8.

The emission factor uncertainty for wildfires is 70%. This is the default value given for non-CO₂ emissions in the Good Practice Guidance (IPCC 2003, section 3.2.1.4.2.4). It is used here also for the CO₂ emissions as the fraction of the biomass combusted is quite uncertain for temperate forests (IPCC 2003, Table 3A.1.12): mean=0.45, SD=0.16.

Uncertainties of activity data of Forest land are described in Chapter 7.2.5. Table 7-5 lists the relative uncertainties in the LULUCF sector: an uncertainty of 63% was calculated for Afforestations, 50% for Deforestations and 63% for Forest Management.

Time-Series Consistency

Consistent time series of annual growing stocks were calculated backward or forward starting from the growing stock 2005, as derived from NFI 3 (see Chapter 7.3.4.6).

Consistent time series of dead wood, litter and soil carbon were calculated with the model Yasso07 (see Didion et al. 2013 and Chapter 7.3.4.8).

7.3.6 Category-Specific QA/QC and Verification

Estimation of Growing Stock, Gains and Losses of Living Biomass

Biomass Conversion and Expansion Factors

For transparency reasons and for comparison with the Biomass Conversion and Expansion Factors (BCEFs) used in former submissions (FOEN 2012 and before), updated BCEFs were calculated. Please note: These BCEFs (Table 7-28) were not used for the calculations but are only provided for verification purposes and transparency. BCEFs for gains and losses vary between the four inventories NFI 1-4a (Table 7-28). The values for gains range from 0.59 to 0.62 for conifers and from 0.82 to 0.86 for broadleaves. The values for losses show a similar range: 0.59 to 0.60 for conifers and 0.79 to 0.83 for broadleaves. The BCEFs and the differences between gains and losses, respectively, are in the same range as those reported in the Austrian NIR 2012 (Umweltbundesamt 2012).

Table 7-28 Comparison of BCEFs used in FOEN (2012 and before: 'BCEF 2012') and BCEFs 2013 (weighted averages, not used for calculation) for conifers and broadleaves, NFI 1-4a.

| | | BCEF 2012 Growing stock | BCEF 2013 Growing stock | BCEF 2013 Gains | BCEF 2013 Losses | Difference |
|-------------|-----------|----------------------------|----------------------------|--------------------|---------------------|------------|
| Conifers | NFI 1 | | 0.62 | | | |
| | NFI 1-2 | | | 0.59 | 0.60 | -0.01 |
| | NFI 2 | | 0.61 | | | |
| | NFI 2-3 | | | 0.61 | 0.59 | 0.02 |
| | NFI 3* | 0.64 | 0.62 | | | |
| | NFI 3-4 | | | 0.62 | 0.60 | 0.02 |
| | NFI 4a | | 0.63 | | | |
| | Austria** | | | 0.62 | 0.61 | 0.01 |
| Broadleaves | NFI 1 | | 0.81 | | | |
| | NFI 1-2 | | | 0.82 | 0.79 | 0.03 |
| | NFI 2 | | 0.82 | | | |
| | NFI 2-3 | | | 0.85 | 0.81 | 0.04 |
| | NFI 3 | 0.83 | 0.83 | | | |
| | NFI 3-4 | | | 0.86 | 0.83 | 0.03 |
| | NFI 4a | | 0.83 | | | |

* For conifers, BCEFs 2013 for growing stock is smaller than BCEF 2012 because of the new parameters for coniferous roots (cf. Thürig and Herold 2013).

** In Umweltbundesamt (2012), only BEFs are published. For comparison in this report, BEFs are converted to BCEFs by multiplication with wood densities (0.38 for conifers, 0.54 for broadleaves as given in Umweltbundesamt 2012: Table 215).

Brush Forests

Düggelin and Abegg (2011) calculated values of total growing stock (also < 7 cm) for brush forest. They measured an average growing stock of 166 m³ ha⁻¹ for *Pinus mugo* stands and 74 m³ ha⁻¹ for stands with *Alnus viridis*.

Litter of Afforestations CC11

In an experiment by Zimmermann and Hiltbrunner (2012; COST E639-project "Turnover and stabilization of soil organic matter: effect of land-use change in alpine regions") litter accumulation in a 40 year old afforestation with Norway Spruce was determined. The authors found accumulation rates of 0.17-0.20 t C ha⁻¹ yr⁻¹. An overview of other studies is given in Chapter 11.3.1.2.

Carbon Balance of two Mountain Forest Ecosystems in Switzerland: Net Ecosystem Exchange and Soil Respiration

Measurements of the net ecosystem exchange (NEE) and of soil respiration were conducted at a montane mixed forest over 5 years (Lägeren; 2005–2009; NFI production region 2), and at a subalpine coniferous forest over 12 years (Davos; 1997–2009; Swiss Plateau, NFI production region 4).

(1) Etzold et al. (2011) determined the net ecosystem exchange (NEE) by eddy covariance (EC) measurements. EC measurements, as well as biometric estimates indicate that both sites with two different mountain forest types were significant C sinks in the respective periods. During 2005 to 2009 NEE of the Lägeren forest ranged from -366 to -662 g C m⁻² yr⁻¹ (mean: -415 g C m⁻² yr⁻¹), and of the Davos forest from -47 to -274 g C m⁻² yr⁻¹ (mean: -154 g C m⁻² yr⁻¹).

(2) Rühr and Eugster (2009) measured soil respiration rates at these two Swiss forest sites. Modeled changes in soil C storage with the dynamic soil carbon model Yasso07 (see also Thürig et al. 2005) gave comparable results with measured soil respiration. The authors found that soils at the alpine site Davos acted as a significant C sink. Soils at the Lägeren site were neither a significant C sink nor a significant C source. This domestic study confirms the broadly spread knowledge that it is very difficult to detect short term changes in soil C stocks, since the uncertainty of the measurement is often higher than the actual change of the annual estimates (e.g. Falloon and Smith 2003).

Changes in Soil Carbon Stocks

SOC Dataset of the Swiss Soil Monitoring Network

The objective of the Swiss Soil Monitoring Network (<http://www.nabo.admin.ch/?lang=en>; NABO) is to assess soil quality in the long term and to validate appropriate soil protection measures. The network was established in 1985. Currently, it comprises 105 observation sites throughout Switzerland. For the statements below, the NABO sites had been classified according to the 18 LULUCF combination categories (CC).

Changes in SOC content of forest soils are being measured since 1985 at soil monitoring benchmark sites in the Swiss Soil Monitoring Network (SAEFL 1993). Repeated soil inventories at the soil monitoring sites are carried out every 5 years. Four replicate bulked soil samples from the upper soil layer 0-20 cm are taken at the monitoring sites (10m x 10m). For each bulked sample 25 single cores are taken at the site according to a stratified random sampling scheme. Further details can be found in SAEFL (2000a).

SOC of the archived soil samples was measured with a modified Walkley and Black method (ACW/ART 1998) in the same laboratory since 2006. Since 2012 we introduced in parallel the SOC analysis by a C/N- Analyser (ThruSpec, Leco company) and measured soil samples of almost all sites with both methods. Strong correlations between both methods were found, with slightly higher values measured by the C/N-Analyser as this analytical method provides total organic carbon (TOC). The resulting regression to transfer SOC-Walkley and Black measurements to TOC showed a high coefficient of determination of 97%. Therefore, we harmonized the dataset correcting the SOC values (Walkley and Black) to the measurements of the C/N-Analyser. For further SOC analysis of the 5th and 6th soil campaign we will only use the C/N-Analyser method. Thus, the SOC dataset (n = 1'884 measurements) presented here and in Chapters 7.4.6 and 7.5.6 is not subject to systematic methodological errors caused by different laboratories or methods. To assure the reliability and accuracy of the measurements, sampling quality, sampling preparation, chemical extraction, analysis and sample storage in the soil archive is evaluated. SOC measurements of a soil sample were repeated if a SOC value deviated more than a certain degree from the values of the other three bulked soil samples of the same sampling campaign.

The spatial variation of bulk density is included in calculating the carbon pools. Bulk density measurements and soil skeleton (> 2mm) were measured at the monitoring sites in the 4th (2000-04) and 5th (2005-09) re-sampling campaign (n=4 in each campaign per site), but not in the previous campaigns. As the mass of the fine earth (FE) is the relevant fraction for the element pools in the soil, the bulk density refers to the mass FE. The measured skeleton fraction of the volume sample is subtracted before. The temporal changes of the top soil bulk density between the 4th and 5th campaign were quite small and they differ between -0.2 and 0.1 g/cm³. We presumed that the bulk density of the first three sampling campaigns ranged within the values measured in the 4th and 5th re-sampling campaign, i.e. propagated the variability of the measurements through Latin Hypercube sampling (n = 1000 simulation runs) assuming a normal distribution of the bulk density and SOC measurements for each site.

The SOC pools for the forest top soils (0-20 cm) ranged between 35.4 t C ha⁻¹ (min) and 135.8 t C ha⁻¹ (max) and were in average 70.6 t C ha⁻¹. In these numbers we exclude one

coniferous forest site that revealed large SOC pools up to 191 t C ha^{-1} . Figure 7-6 shows that in average, SOC pools did not change monotonously during the measurement period between 1989 and 2009 in the sampled forest soils. At some of the forest monitoring sites higher values were found in the 3rd re-sampling campaign.




Figure 7-6 Time series of measured SOC pool changes in the top soil (0-20 cm) at the 28 NABO forest sites from the 1st to the 5th re-sampling campaigns. SOC pools were centred by the median SOC pool of all re-samplings of the monitoring site. Each pool value presents the median of four bulked soil samples per campaign with measured SOC and bulk density. The error bars indicate the 25% and 75% percentiles resulting from the spatial variation of the sites and the errors along the measurement chain. The altitude of the forest sites ranges between 380 and 1690 m a.s.l.

Detailed studies at monitoring sites showed that short-term temporal variation of soil properties can result from different site conditions at the sampling date, e.g. regarding soil moisture, soil temperature and bulk density (Keller et al. 2006). For instance, at two forest sites six re-samplings within three years revealed short-term variation of the SOC content between $\pm 1.8 \%$ and $\pm 0.6 \%$. Therefore, the majority of the measured temporal variation for all forest sites is interpreted as natural variation (noise) and not as real SOC changes (signal). This hypothesis is also supported by the fact that the soil samples in the 3rd resampling campaign were taken earlier in spring time as in the other sampling campaigns and hence, soil moisture content of the samples was higher in average. This might explain the large temporal variation, in particular at coniferous forest sites with a pronounced organic layer. Using a robust linear regression approach for the SOC pool data of the forest soils, the 95% confidence interval for the SOC pool was $\pm 1.5 \text{ t C ha}^{-1}$. In order to capture as good as possible the natural variation of these site-specific characteristics, standard operation procedures and quality assurance were implemented since the 4th soil campaign. Further work will focus on the correction of the measured C pools to equivalent mass of the fine earth $< 2 \text{ mm}$. In this way, we presume that the 95% confidence interval of the mean SOC pool can be reduced to some degree.

The 95% confidence interval obtained for the 28 forest monitoring sites is an order of magnitude larger than the modelled SOC pool changes for Swiss forest soils (Chapter 7.3.4.8). Didion et al. (2013), who applied the Yasso07 model to Swiss forest soils, estimated a soil sink effect of $-0.0008 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for 2009 to 2010. For the monitoring period of 20 years this rate of change would correspond to $-0.016 \text{ t C ha}^{-1}$. Modeled SOC pool changes are not inconsistent with the repeated soil inventories in the NABO network. As indicated by the 95% confidence interval for the 28 forest monitoring sites, the noise is about hundred times larger than the modeled signal.

Uncertainty Estimates

For comparison, the uncertainty for the stock of litter amounts to approximately 60% (Moeri 2007). The uncertainty of mortality data from the Finnish GHG inventory was assessed to be 30% (Monni et al. 2007). The uncertainty reported by Finland, where the Yasso07 model is also applied, was 46.8% for the 2010 emission factor in South Finland, 26.2% in North Finland, and 24.1% for the net change in the whole country (Statistics Finland 2012: Chapter 7.2.4.2).

7.3.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in category 5A.

For this submission, NFI4a+ data covering the period 2009-2012 were applied as described in Thürig (2014; see Chapter 7.3.4.1). These data are an update of Abegg et al. (2012) which had been used in the previous submission (FOEN 2013).

The calculation of carbon stock changes in the different pools has been adapted such that it is harmonized for reporting under UNFCCC and under the KP, that no double counting is possible and that the most conservative accounting method is applied. The calculation approach is explained in 7.1.3.2 and Chapter 11.3.1. and transparently shown in Table 7-3 and Table 11-7. The V- and W-factors are no longer used for describing the calculation methods. Instead, the terms 'gain-loss' approach and 'stock-change' approach were introduced in line with the GPG LULUCF in order to be more clear and comprehensible. The calculation procedure was modified in the following transitions and C pools:

- Land-use type conversions from non-forest to CC12 (productive forest), from non-forest to CC13 (unproductive forest) and from CC13 to CC12; living biomass:
Previous submission: stock-change.
This submission: gain-loss.
Rationale: The stock differences being dispersed over a conversion time of 20 years led to very high growth rates (Implied Emission Factor for gains). With this modification the C sink on areas converted from non-forest to forest land strongly decreased.
- Land-use type conversion from afforestation CC11 to CC12: living biomass, dead organic matter and soil:
Previous submission: stock-change.
This submission: gain-loss.
Rationale: Afforestation is a transient land-use type by definition. It is supposed that it converts smoothly to CC12 during the 20 years.

Updated estimates of yearly changes in carbon stocks of soil organic carbon, dead wood and litter, modeled with Yasso07, are described in Didion et al. (2013). The accuracy of the estimates of C stocks and CSC was improved. As a result the a posteriori calibration of deadwood stocks (Chapter 7.3.4.9 in FOEN 2013) became redundant and, hence, a source of uncertainty was eliminated.

For wildfires on forest land, the emission factor for CH₄ was revised (see description in Chapter 7.3.4.13). Now, the default emission factor is used.

7.3.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines (IPCC 2006) represents the largest improvement. To accomplish this transition, other projects or planned improvements may be postponed in order to give first priority to changes related to the new reporting guidelines.

In 2011, Agroscope started a three-year running research project that aims to identify (drained) fens and raised bogs under different land uses beyond the national inventories of bogs and fens in order to improve the AD estimates of organic soils ("Area and location of organic soils and peatlands in Switzerland"). An ongoing study at Agroscope measures soil carbon stocks in organic soils under forest of three different vegetation units. These data will become available during 2014 and will help to validate the results of the above-mentioned project (ground truthing).

Further research is underway to estimate yearly values for gross growth. First results of a study investigating the relationship between climate and gross growth (Thürig et al. 2009) are expected in the course of 2014.

Switzerland intends to make better use of the SOC data provided by the Swiss Soil Monitoring Network in forthcoming submissions. A contract was signed in 2011 that arranges a close collaboration for the period 2012-2014.

The application and the parametrization of the model Yasso07 to get better estimates of temporal changes in soil carbon, LFH layer and dead wood will be further improved. Contact with researchers from other countries also applying Yasso07 for LULUCF reporting has been established.

Projects in a new national research programme (Sustainable Use of Soil as a Resource: "SOM control", <http://www.nfp68.ch/E>) aims at identifying the drivers of soil organic matter storage in Swiss forest soils. The objectives are to assess how forest productivity and tree species composition affect SOM storage, to investigate if and how land-use history affects C pools in soils, to estimate the influence of climate, temperature and precipitation on SOM stocks, to link SOM stocks to physico-chemical parameters controlling SOM stabilization and to model SOM and evaluate the residuals to measured SOM stocks.

7.4 Category 5B – Cropland

7.4.1 Description

Tier 2 Key category 5B1

CO₂ from 5B1 Cropland remaining Cropland (2012: level and trend).

Tier 2 Key category 5B2

CO₂ from 5B2 Land converted to Cropland (2012: trend).

Swiss croplands belong to the cold temperate wet climatic zone. Carbon stocks in aboveground living biomass and carbon stocks in mineral and organic soils are considered. Croplands (CC21) include annual crops and leys in arable rotations (see Table 7-2 and Table 7-6). Because arable cropping mainly occurs in the temperate Swiss Central Plateau and no elevation-dependent soil carbon stocks are available for Swiss croplands (Leifeld et al. 2005), no stratification of carbon stocks has been applied.

In 2012, 5B1 Cropland remaining Cropland was a net source of 674.69 Gg CO₂ due to (I) a decrease in living biomass between 2011 and 2012 and due to (II) emissions from organic soils. Average living biomass was increasing slightly over the period 1990-2012. However, annual fluctuations in net carbon stock changes of biomass are considerable (see Table 7-30). Since carbon stocks on mineral soils are assumed to be in balance (i.e. no carbon stock changes occur on mineral soils) all soil emissions in 5B1 were generated by carbon mineralization in organic soils, mainly in the lowest altitudinal zone (z1: 98%). Overall, organic soils account for 2.7% of cropland area in Switzerland.

5B2 Land converted to Cropland was a small net source of 22.26 Gg CO₂ in 2012.

7.4.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.4.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.4.4 Methodological Issues

7.4.4.1 Carbon in Living Biomass

Annual biomass carbon stocks are shown in Table 7-29. They are calculated as area-weighted means of standing stocks at harvest for the seven most important annual crops (wheat, barley, maize, silage maize, sugar beet, fodder beet, potatoes) and as cumulated annual harvested biomass for leys.

The annual mean standing biomass carbon stock per hectare is calculated as:

$$\text{Biomass cropland} = \sum_i (A_i / A_t) * C_i$$

where A_f = Area of crop type f , A_t = total cropping area and C_f = yield (annual crops, leys) for the particular crop (t C ha^{-1}). Annual values for A_f , A_t and C_f were published by the Swiss Farmers Union (SBV 2013).

The resulting mean biomass stock for Swiss cropland over the inventory time period is 4.69 ± 0.34 (1 SD) t C ha^{-1} .

Table 7-29 Annual values for arable crop yields (i.e. carbon stocks) and area-weighted mean (t C ha^{-1}) (SBV 2013), assuming a carbon fraction of 0.5 (IPCC default).

| crop | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--------------------------------------|------|------|------|------|------|------|------|------|------|------|
| CC21: yield [t C ha^{-1}] | | | | | | | | | | |
| Barley | 2.36 | 2.47 | 2.48 | 2.54 | 2.19 | 2.29 | 2.69 | 2.69 | 2.86 | 2.18 |
| Wheat | 2.36 | 2.54 | 2.35 | 2.53 | 2.34 | 2.56 | 2.84 | 2.55 | 2.63 | 2.24 |
| Maize | 3.51 | 3.42 | 3.53 | 3.78 | 3.72 | 3.55 | 3.64 | 3.94 | 3.87 | 3.78 |
| Silage maize | 7.37 | 6.59 | 7.15 | 6.72 | 6.11 | 6.03 | 4.98 | 7.08 | 6.88 | 6.49 |
| Sugar beet | 7.41 | 6.91 | 7.04 | 7.63 | 6.72 | 6.78 | 7.83 | 7.76 | 7.42 | 7.48 |
| Fodder beet | 6.70 | 6.51 | 6.64 | 6.77 | 5.66 | 5.49 | 6.41 | 6.53 | 6.06 | 5.79 |
| Potatoes | 4.47 | 4.39 | 4.65 | 4.83 | 3.65 | 3.88 | 5.36 | 5.05 | 4.44 | 3.87 |
| Leys | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 |
| Mean | 4.34 | 4.30 | 4.39 | 4.44 | 4.12 | 4.28 | 4.51 | 4.72 | 4.67 | 4.43 |

| crop | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--------------------------------------|------|------|------|------|------|------|------|------|------|------|
| CC21: yield [t C ha^{-1}] | | | | | | | | | | |
| Barley | 2.55 | 2.38 | 2.68 | 2.35 | 2.92 | 2.61 | 2.64 | 2.57 | 2.58 | 2.73 |
| Wheat | 2.53 | 2.35 | 2.42 | 2.16 | 2.62 | 2.46 | 2.49 | 2.57 | 2.58 | 2.60 |
| Maize | 4.10 | 3.80 | 3.92 | 1.83 | 4.09 | 4.10 | 3.30 | 4.32 | 4.12 | 4.42 |
| Silage maize | 6.68 | 6.45 | 4.93 | 5.96 | 6.52 | 8.23 | 7.01 | 8.02 | 8.09 | 7.60 |
| Sugar beet | 8.74 | 6.51 | 8.52 | 7.89 | 8.60 | 8.50 | 7.30 | 8.37 | 8.73 | 9.37 |
| Fodder beet | 6.71 | 5.75 | 5.95 | 5.67 | 6.13 | 6.15 | 6.25 | 6.21 | 6.30 | 6.72 |
| Potatoes | 4.67 | 4.13 | 4.30 | 3.71 | 4.34 | 4.26 | 3.60 | 4.59 | 4.71 | 5.12 |
| Leys | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 | 6.11 |
| Mean | 4.71 | 4.51 | 4.54 | 4.41 | 4.89 | 4.97 | 4.70 | 5.07 | 5.13 | 5.19 |

| crop | 2010 | 2011 | 2012 | mean 1990-2012 | | | | | | |
|--------------------------------------|------|-------|------|----------------|--|--|--|--|--|--|
| CC21: yield [t C ha^{-1}] | | | | | | | | | | |
| Barley | 2.56 | 2.75 | 2.76 | 2.56 | | | | | | |
| Wheat | 2.48 | 2.72 | 2.50 | 2.50 | | | | | | |
| Maize | 3.61 | 4.13 | 3.86 | 3.76 | | | | | | |
| Silage maize | 7.47 | 8.42 | 8.06 | 6.91 | | | | | | |
| Sugar beet | 8.03 | 10.38 | 9.58 | 7.98 | | | | | | |
| Fodder beet | 6.49 | 6.65 | 5.58 | 6.22 | | | | | | |
| Potatoes | 4.26 | 5.04 | 4.56 | 4.43 | | | | | | |
| Leys | 6.11 | 6.11 | 6.11 | 6.11 | | | | | | |
| Mean | 4.99 | 5.44 | 5.23 | 4.69 | | | | | | |

7.4.4.2 Carbon in Soils

Soil carbon stocks in mineral soils under cropland are calculated based on Leifeld et al. (2003, 2005). The approach correlates measured soil organic carbon stocks (t ha^{-1}) for arable land and leys with soil texture after correction for soil depth and stone content. Area upscaling uses the Swiss digital soil map (SFSO 2000a), and average stocks are calculated as weighted means using the area of arable land and leys. The mean soil organic carbon stock (0-30 cm) for cropland is $53.40 \pm 5 \text{ t C ha}^{-1}$.

It should be noted that current carbon stocks are not only the result of the conditions for productivity and carbon turnover under different land-use types, but are also determined by farmers' decisions to use a site in a specific way due to the demands of a crop or the suitability of a site, e.g. regarding machine use (see Leifeld et al. 2003: 65).

Soil carbon stocks in organic soils under cropland are calculated based on Leifeld et al. (2003, 2005). The approach uses measured carbon stocks in Swiss organic soils. The mean soil organic carbon stock (0-30 cm) for cultivated organic soils is $240 \pm 48 \text{ t C ha}^{-1}$.

7.4.4.3 Changes in Carbon Stocks

Carbon stocks in living biomass intermittently increased from 4.34 t C ha^{-1} in 1990 to 5.23 t C ha^{-1} in 2012 (Table 7-29; SBV 2013). The difference in biomass stock between a specific year and the preceding year is reported as gain or loss of carbon (see Table 7-30: "change"). The resulting values are in the range between -0.32 and $0.49 \text{ t C ha}^{-1} \text{ yr}^{-1}$ with an average of $0.041 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for the inventory time period.

Changes of carbon stocks in mineral soils are assumed to be zero for cropland remaining cropland.

The annual net carbon stock change in organic soils was estimated to $-9.52 \text{ t C ha}^{-1}$ according to measurements in Europe including Switzerland as compiled by Leifeld et al. (2003, 2005) and rechecked by ART (2009b).

In the case of land-use change, the net changes in biomass and soil are calculated as described in Chapter 7.1.3.

Table 7-30 Annual carbon data for cropland (CC21), 1990-2012. Annual carbon stocks are broken down for arable crops in Table 7-29. Highlighted data for 1990 as displayed in Table 7-4.

| cropland | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|-------|--------|-------|-------|--------|-------|-------|-------|--------|--------|
| CC21: carbon stocks and changes in living biomass [t C ha^{-1}] | | | | | | | | | | |
| stock | 4.34 | 4.30 | 4.39 | 4.44 | 4.11 | 4.28 | 4.51 | 4.72 | 4.67 | 4.43 |
| change | 0.053 | -0.033 | 0.086 | 0.047 | -0.321 | 0.165 | 0.227 | 0.215 | -0.051 | -0.239 |

| cropland | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|-------|--------|-------|--------|-------|-------|--------|-------|-------|-------|
| CC21: carbon stocks and changes in living biomass [t C ha^{-1}] | | | | | | | | | | |
| stock | 4.71 | 4.51 | 4.54 | 4.41 | 4.89 | 4.97 | 4.70 | 5.07 | 5.13 | 5.19 |
| change | 0.282 | -0.206 | 0.033 | -0.131 | 0.486 | 0.075 | -0.269 | 0.368 | 0.061 | 0.056 |

| cropland | 2010 | 2011 | 2012 | mean 1990-2012 |
|--|--------|-------|--------|----------------|
| CC21: carbon stocks and changes in living biomass [t C ha^{-1}] | | | | |
| stock | 4.99 | 5.44 | 5.23 | 4.69 |
| change | -0.198 | 0.450 | -0.206 | 0.041 |

7.4.4.4 N₂O Emissions from Land-Use Conversion to Cropland

N₂O emissions as a result of the disturbance associated with land-use conversion to cropland are reported in CRF-table 5(III). The emissions are calculated with default values proposed by IPCC (2003, following Equations 3.3.14 and 3.3.15, and Chapter 3.3.2.3.1.2):

$$\text{Emission (N}_2\text{O)} = -\text{deltaC}_s \cdot 1 / (\text{C} : \text{N}) \cdot \text{EF1} \cdot 44 / 28 \quad [\text{Gg N}_2\text{O}]$$

where:

deltaC_s: soil carbon change induced by land-use conversion to cropland [Gg C]

C:N: C:N ratio = 9.8 in grassland soils (Leifeld et al. 2007)

EF1: IPCC default emission factor = $0.0125 \text{ kg N}_2\text{O-N (kg N)}^{-1}$

deltaC_s is calculated according to the methodology described in Chapter 7.1.3. If deltaC_s is zero or positive (carbon gain) there are no N₂O emissions provoked by a land-use conversion to cropland.

On mineral soils this is the case for wetlands converted to cropland and other land converted to cropland (see CRF-table 5B2); consequently, NO is reported in CRF-table 5(III) for these land-use conversions.

On organic soils the carbon stock differences for land-use conversions to cropland are zero or – in the case of other land) – positive (cf. Table 7-4). No N₂O emissions are generated.

The country specific C/N ratio of 9.8 for grassland proposed by Leifeld et al. (2007) is used because the largest part of the area converted to cropland consisted of grassland (cf. CRF-table 5B2).

7.4.4.5 Carbon Emissions from Agricultural Lime Application

The total annual amount of limestone input to agricultural soils (CRF-table 5(IV)) is between 51'300 Mg and 74'050 Mg. It was estimated by Agroscope in 2009 for the period 1990-2008 (see Table 7-31). For 2009–2012 the same value as for 2008 is used: An inquiry in 2013 including the most important producers of lime products suggests that the consumption of limestone remained constant in this period (Agroscope 2014a).

Dolomite is probably applied only in small quantities. The available data do not allow to differentiate Ca(CO₃) and CaMg(CO₃)₂.

The availability of a country specific emission factor for agricultural lime application has been investigated, but no domestic measurement data could be found. Consequently, the IPCC default carbon conversion factor for carbonate containing lime of 0.12 Mg C per Mg Ca(CO₃) or CaMg(CO₃)₂ (IPCC 2003) has been used. The resulting carbon emissions associated with liming range from 22.57 to 32.58 Gg CO₂ yr⁻¹.

In the CRF-table 5(IV) all emissions are reported under Cropland and for Grassland the notation key IE is used because it is not known on which areas the lime was applied.

Table 7-31 Amount of limestone applied on agricultural soils and resulting CO₂ emissions 1990-2012 (Agroscope 2014a).

| Year | Limestone Mg | Emission Gg CO ₂ |
|------|-----------------|--------------------------------|
| 1990 | 51'300 | 22.57 |
| 1991 | 51'342 | 22.59 |
| 1992 | 52'383 | 23.05 |
| 1993 | 53'425 | 23.51 |
| 1994 | 54'467 | 23.97 |
| 1995 | 55'508 | 24.42 |
| 1996 | 56'550 | 24.88 |
| 1997 | 57'592 | 25.34 |
| 1998 | 58'633 | 25.80 |
| 1999 | 59'675 | 26.26 |
| 2000 | 60'717 | 26.72 |
| 2001 | 61'758 | 27.17 |
| 2002 | 62'800 | 27.63 |
| 2003 | 63'842 | 28.09 |
| 2004 | 69'883 | 30.75 |
| 2005 | 70'925 | 31.21 |
| 2006 | 71'967 | 31.67 |
| 2007 | 73'008 | 32.12 |
| 2008 | 74'050 | 32.58 |
| 2009 | 74'050 | 32.58 |
| 2010 | 74'050 | 32.58 |
| 2011 | 74'050 | 32.58 |
| 2012 | 74'050 | 32.58 |

7.4.5 Uncertainties and Time-Series Consistency

A range of possible carbon stock changes in mineral soils has been determined by the Swiss Soil Monitoring Network (NABO). The upper and lower margin of the 95% confidence interval for carbon stock changes under cropland is $0 \pm 0.52 \text{ t C ha}^{-1}$ (Keller 2013). This absolute uncertainty is used to calculate relative uncertainties for 5B1 and 5B2 by dividing with the mean net emission per hectare of 5B1 and 5B2, respectively. In 2012, the mean net emissions were $0.474 \text{ t C ha}^{-1}$ for 5B1 and $0.363 \text{ t C ha}^{-1}$ for 5B2 (calculated from CRF-table 5B). The resulting relative uncertainties are 110% for 5B1 and 143% for 5B2, respectively (Table 7-5).

In the uncertainty analysis, these values were chosen for the overall emission factor uncertainties for CO₂ in sectors 5B1 and 5B2 as they dominate the other sources of uncertainty by far.

- Uncertainties for soil carbon stocks are given together with the mean value in the text above: 9% for mineral soils and 20% for cultivated organic soils. They take into account uncertainties in measured carbon contents and predicted soil bulk densities, i.e., they consider only uncertainties in emission factors.
- The relative uncertainty in yield determination has been estimated at 13% for biomass carbon from agricultural land (Leifeld and Fuhrer 2005). Data on biomass yields for different elevations and management intensities as published by FAL/RAC (2001) are based on many agricultural field experiments and have a high reliability.
- The uncertainty of the carbon stock change in organic soils is 23% as reported by Leifeld et al. (2003: 56).

For the uncertainty of the emission factor for N_2O on land converted to cropland a default value of 90% is used (Table 7-5).

The amount of total lime application in agriculture is mainly based on expert judgement; the resulting number is uncertain. A relative uncertainty of $\pm 40\%$ can be used as an approximation (Agroscope 2014a). For the emission factor of lime a lower uncertainty of $\pm 25\%$ was chosen, because it is a plain chemical process.

Uncertainties of activity data of Cropland are described in Chapter 7.2.5.

7.4.6 Category-Specific QA/QC and Verification

Changes in Living Biomass

In 2012 an assessment of the appropriateness of the estimated pools of carbon in living biomass was conducted (ART 2012a). It came to the conclusion that almost all carbon stocks and carbon stock changes are in the expected range of the IPCC Guidelines and Good Practice Guidance. Nevertheless there is room for improvements. However, given the relatively low significance of the respective emissions a major effort in this area is hardly justified. Consequently, the biomass carbon pools will eventually be recalculated only in the course of the new planned Tier 3 approach for quantification of carbon stocks and carbon stock changes in agricultural soils (see also Chapter 7.4.8).

Changes in SOC Pools

A SOC pool dataset provided by the Swiss Soil Monitoring Network (NABO; see Chapter 7.3.6) supports the Tier 1 assumption that changes of carbon stocks in mineral soils are zero for cropland remaining cropland (cf. UNFCCC 2009: §79; UNFCCC 2010: §72; UNFCCC 2011: §94). The SOC pool changes measured at 38 cropland monitoring sites in the Swiss Soil Monitoring Network show a slight trend towards decreasing SOC pools since 1985 (Figure 7-7). However, this trend is not statistically significant yet. The range of the calculated SOC pools is quite large ($27.7\text{--}598 \text{ t C ha}^{-1}$), as three cropland soil monitoring sites are on peat soils. These three sites are excluded in Figure 7-7. Average SOC pool in the topsoils (0–20 cm) of the remaining 35 cropland monitoring sites for all soil sampling campaigns was 45.4 t C ha^{-1} . At the three cropland sites on peat soils the temporal variation of SOC content during the last 20 years ranged between $\pm 0.4\%$, corresponding to SOC pool changes larger than $\pm 10 \text{ t C ha}^{-1}$. However, for the majority of cropland sites the temporal variation found was smaller ($\pm 0.2\%$), the confidence interval of the SOC pool changes calculated by robust regression was $\pm 0.4 \text{ t C ha}^{-1}$ over the last 20 years. This finding is in agreement with the detailed study mentioned in Chapter 7.3.6 (Keller et al. 2006), where six re-samplings of two cropland sites within three years revealed natural SOC content variation of $\pm 0.23\%$ in the topsoil. The temporal variation of the SOC content and SOC pools at the cropland sites are rather small and possible future changes can be detected by this soil monitoring design.

Yet, the results suggest that Swiss cropland mineral soils did not act as a net carbon source or sink during the last 20 years. The results of the 6th soil campaign (2010–2014), that will be available in 2015 for the majority of the monitoring sites, will provide more evidence if the slight trend of decreasing SOC in cropland soils will be confirmed. In addition, SOC pools of the whole soil profile have to be determined as top soil SOC pools might be changed from changing land management practices such as the ploughing depth.




Figure 7-7 Time series of measured SOC pool changes in the top soil (0-20 cm) at 35 NABO cropland sites from the 1st to the 5th re-sampling campaigns. SOC pools were centred by the median SOC pool of all re-samplings of the monitoring site. Each pool value presents the median of four bulked soil samples per campaign with measured SOC and bulk density. The error bars indicate the 25% and 75% percentiles resulting from the spatial variation of the sites and the errors along the measurement chain. The altitude of the cropland sites ranges between 209 and 945 m a.s.l.

Short-term Land-Use Changes in Arable Rotations

Short-term land-use changes between "grassland" and cropland are to be expected for leys in arable rotations. However, leys are allocated to cropland by the Swiss Land Use Statistics (AREA) and are thus not considered grasslands in the common sense (i.e. permanent grassland). Furthermore, only long-term changes between cropland and grassland are considered relevant for carbon stock changes in soils. Since only long-term land-use changes are registered by the Swiss Land Use Statistics (AREA), carbon stock changes in soils associated with land-use changes between cropland and grassland and vice versa are adequately reported in the GHG inventory.

7.4.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in category 5B.

7.4.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines represents the largest improvement. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented. Other projects and planned improvements may be postponed in order to give first priority to changes related to the new reporting guidelines.

In 2011, Agroscope started a three-year running research project that aims to identify (drained) fens and raised bogs under different land uses beyond the national inventories of bogs and fens in order to improve the AD estimates of organic soils ("Area and location of organic soils and peatlands in Switzerland").

Switzerland intends to make better use of the SOC data provided by the Swiss Soil Monitoring Network (NABO) in forthcoming submissions. A contract was signed in 2011 that arranges a close collaboration for the period 2012-2014.

Furthermore, information about carbon stock changes in soils for cropland remaining cropland will become available from Agroscope research activities. A pilot study to evaluate possible Tier 3 methodological approaches for quantification of carbon stocks and carbon stock changes in agricultural soils (cf. UNFCCC 2009: §79; UNFCCC 2010: §72; UNFCCC 2011: §94) has recently been terminated (Köck et al. 2013).

Ongoing efforts to combine SOC measurements on the level of soil fractions with modelled pools (Zimmermann et al. 2007; Leifeld et al. 2009) will be combined with the planned Tier 3 approach for an independent check of emission and removal rates for cropland and grassland conversions.

The Research Institute of Organic Agriculture FiBL in CH-Frick runs a project with focus on the determination of sources and sinks of greenhouse gases in Swiss arable soils (project duration 2012-2014; co-funded by FOEN).

A new study on GHG emissions from peatlands and organic soils under different land use (Agroscope in collaboration with the University of Basel, 2013-2016, financed by FOEN) will improve the robustness of domestic emission factor estimates for soils rich in organic matter in the medium term.

Projects in a new national research programme ("Sustainable Use of Soil as a Resource", <http://www.nfp68.ch/E>) focus on (1) sustainable management of organic soils, (2) agricultural management and below ground carbon inputs, and (3) an integrated modelling framework to monitor and predict trends of agricultural management and their impact on soil functions at multiple scales. In the long term, results from these studies are expected to improve the reporting of emissions by sources and removals by sinks in category 5B.

7.5 Category 5C – Grassland

7.5.1 Description

Tier 2 Key category 5C1

CO₂ from Grassland remaining Grassland
(2012: level and trend).

Tier 2 Key category 5C2

CO₂ from Land converted to Grassland
(2012: level and trend)

Swiss grasslands belong to the cold temperate wet climatic zone.

Grasslands are subdivided into permanent grassland (CC31), shrub vegetation (CC32), vineyards, low-stem orchards ('Niederstammobst') and tree nurseries (CC33), copse (CC34), orchards ('Hochstammobst'; CC35), stony grassland (CC36), and unproductive grassland (CC37), see Table 7-2 and Table 7-6. Carbon stocks in living biomass and carbon stocks in soils have been estimated for every subclass and have been considered accordingly in the calculation scheme.

In the CRF-table 5C2, the land-use types CC32, CC33, CC34 and CC35 are merged under the notation 'woody' and CC36 and CC37 are merged under 'unproductive' (see Table 7-2).

In 2012, 5C1 Grassland remaining Grassland was a net source of 134.74 Gg CO₂. Carbon stocks in mineral soils and carbon stocks in living biomass under constant land use are considered to be in balance (i.e. no carbon stock changes do occur). The highest contribution were thus generated by carbon mineralization in organic soils under permanent grasslands in the lowest altitudinal zone (z1: 173.2 Gg CO₂), although only 0.47% of all grassland soils in Switzerland are organic soils. Contributions of other grassland source categories remaining grassland were of minor importance.

5C2 Land converted to Grassland was a net source of 169.07 Gg CO₂ in 2012. The highest individual contribution came from 5C2.1 Forest Land converted to Grassland being responsible for a net source of 323.19 Gg CO₂. Most of this source (68%) is due to net changes in living biomass from deforestation. Land-use change source categories 5C2.2 to 5C2.5 were small net sinks due to sequestration of CO₂ in soils and biomass in the course of the conversion to grassland.

7.5.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.5.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.5.4 Methodological Issues

7.5.4.1 Carbon in Living Biomass

Permanent Grassland (CC31)

Permanent grasslands range in altitude from < 300 m to 3000 m above sea level. Because both biomass productivity and soil carbon rely on the prevailing climatic and pedogenic conditions, grassland stocks were calculated separately for three altitude zones (corresponding to those used in category 5A Forest Land).

Standing stocks for permanent grasslands (t C ha^{-1}) are calculated as the annual cumulative yield of differentially managed grasslands (meadows, pastures, alpine meadows and pastures) based on FAL/RAC (2001; Table 7-32), assuming a carbon fraction of 0.5 (IPCC default). Mean standing above-ground biomass stocks were taken for each of the altitudinal zones because the spatial distribution of grassland management types is not known.

Table 7-32 Annual yields of differentially managed permanent grassland (CC31). Each value represents the mean of two fertilization levels.

| Management | Altitude [m] | Annual yield [t C ha ⁻¹] |
|---------------------------|--------------|--------------------------------------|
| Meadow | <601 | 5.88 |
| | 601-1200 | 4.38 |
| | >1200 | 3.25 |
| Pasture | <601 | 4.63 |
| | 601-1200 | 3.75 |
| | >1200 | 2.75 |
| Alpine pasture and meadow | 601-1200 | 3.75 |
| | >1200 | 0.75 |

Data for root biomass C was compiled by ART (2011a) based on published data of Swiss grassland. Carbon stocks in roots are in the range of 1.82–5.70 t C ha^{-1} depending on altitude. Root biomass is added to above-ground biomass to derive the total living biomass for CC31. Table 7-33 shows the living biomass of permanent grassland for the three altitudinal zones presented as the cumulated annual yield including roots.

Table 7-33 Root biomass C_{root} and total living biomass C_l of permanent grassland (CC31).

| Altitude [m] | C_{root} [t C ha ⁻¹] | C_l [t C ha ⁻¹] |
|--------------|---|-------------------------------|
| <601 | 1.82 | 7.08 |
| 601-1200 | 2.04 | 6.00 |
| >1200 | 5.70 | 7.95 |

Shrub Vegetation (CC32) and Copse (CC34)

Due to the lack of more precise data, the living biomass of shrub vegetation and copse was assumed to be equal to the living biomass of brush forest as described in Chapter 7.3.4.9, where brush forest is assumed to contain 12.9 t C ha^{-1} .

Vineyards, Low-stem Orchards and Tree Nurseries (CC33)

Low-stem orchards are small fruit trees distinguished from CC35 ('orchards') by a maximum stem-height of 1 m and a much higher stand density. Only low-stem orchards and vineyards are considered in the following because no stand densities for tree nurseries are available. Data from SFSO (2002) indicate a very small contribution of tree nurseries (1'378 ha) as compared to the sum of vineyards (15'436 ha, SFSO 2005) and low-stem orchards (240 ha, based on Widmer 2006).

The standing carbon stock of living biomass per ha (CI) for CC33 is therefore calculated as:

$$CI = [(CI \text{ vineyards} * \text{area vineyards}) + (CI \text{ low-stem orchards} * \text{area low-stem orchards})] / (\text{area vineyards} + \text{area low-stem orchards})$$

CI of vineyards is 3.61 t C ha^{-1} , calculated based on the mean stand density (5'556 vines ha^{-1}) and woody biomass of a plant including roots (0.65 kg C; Ruffner 2005).

For small fruit trees on low-stem orchards, no literature value was found for biomass expansion factors. Therefore, following assumptions were met. Diameter at breast height (DBH) of such trees was assumed to be around 10 cm and the stem height was assumed to be around 1 m. The bole shape of low-stem apple trees can be approximated by a cylinder shape.

$$\text{Stem wood volume} = r^2 * \pi * \text{height} = (5 \text{ cm})^2 * 3.1 * 100 \text{ cm} = 7.75 \text{ dm}^3$$

Based on expert knowledge (Kaufmann 2005), the percentage of branches was estimated as 100%, and the percentage of roots was estimated as 30% of the stem wood volume. This results in a BEF of 2.3. A wood density of 0.55 kg dm^{-3} (Vorreiter 1949) and the default carbon content of 50% were assumed. With these assumptions the carbon content of a tree of the type low-stem ('Niederstamm') is calculated as follows:

$$\begin{aligned} \text{C low-stem} &= \text{stem wood volume} * \text{BEF} * \text{wood density} * \text{carbon content} \\ &= 7.75 \text{ dm}^3 * 2.3 * 0.55 \text{ kg/dm}^3 * 50\% \text{ C-content} = 4.9 \text{ kg C} \end{aligned}$$

The mean stand density of low-stem orchards is estimated at 2500 ha^{-1} (Widmer 2006), resulting in a CI of $12.25 \text{ t C ha}^{-1}$.

The resulting CI for CC33 is 3.74 t C ha^{-1} .

Orchards (CC35)

Orchards are loosely planted larger fruit trees ('Hochstammobst') with grass understory. CI of orchards trees is calculated as:

$$CI \text{ biomass} = (\text{carbon per fruit tree [t]} * \text{number of fruit trees} / \text{area orchards [ha]}) + \text{carbon in grass [t ha}^{-1}]$$

The carbon content of a large fruit tree with a DBH of 25 - 35 cm was calculated as follows:

$$C (\text{Hochstamm}) = \text{Stem wood volume} * \text{KE-Factor} = 225 \text{ kg C}$$

where:

Stem wood volume of an apple tree assuming a cylindrical stem with mean DBH of 30 cm and a stem height of 7 m amounts to 0.5 m^3 ;

$$\text{KE-Factor [t C m}^{-3}] = \text{BEF} * \text{Density} * \text{C-content} = 0.45, (\text{Wirth et al. 2004: 68, Table 16}).$$

From the total fruit-growing area of 41'480 ha (SFSO 2005), the area of small fruit trees (240 ha, see CC33) was subtracted, and the remaining area was divided by the number of large fruit trees. Large fruit trees were counted in 1991 (3'616'301) and 2001 (2'900'000; SFSO 2002), and the mean value was divided by 41'240 ha to obtain a mean stand density of 79

trees ha⁻¹. The resulting woody biomass of CC35 is thus 17.78 t C ha⁻¹. Because orchards typically have a grass understory, the biomass of CC31 was added to the woody biomass. Orchards are located below 1000 m a.s.l., so the mean of grass biomass of the classes <601 and 601-1200 m a.s.l. (i.e. 6.54 t C ha⁻¹; cf. Table 7-33) was taken to obtain a total biomass stock of 24.32 t C ha⁻¹ for CC35.

Stony Grassland (CC36)

Approximately 35% of the surface of CC36 (herbs and shrubs on stony surfaces) is covered by vegetation. No accurate data were available for this category. Therefore, the carbon content of brush forest (12.9 t C ha⁻¹) was multiplied by 0.35 to account for the 35% vegetation coverage. This results in a carbon content of 4.52 t C ha⁻¹.

Unproductive Grassland (CC37)

The category CC37 includes grass and herbaceous plants at watersides of lakes and rivers including dams and other flood protection structures, constructions to protect against avalanches and rock slides, and alpine infrastructure (e.g. for skiing). For none of these land-use types, biomass data are currently available. Therefore, the mean value of permanent grasslands in all altitude zones, 7.01 t C ha⁻¹ (cf. Table 7-33), is arbitrarily chosen as the preliminary biomass value for CC37.

7.5.4.2 Carbon in Soils

Permanent Grassland (CC31)

Carbon stocks in grassland soil refer to a depth of 0-30 cm.

Soil carbon stocks in mineral soils under permanent grassland CC31 are calculated based on Leifeld et al. (2003, 2005). The approach correlates measured soil organic carbon stocks (t ha⁻¹) for permanent grasslands with soil texture and elevation after correction for soil depth and stone content. Area upscaling makes use of the Swiss digital soil map (SFSO 2000a) and topography. Mean Cs values calculated for grasslands CC31 are given in Table 7-34. It should be noted that the current C stocks are not only the result of the conditions for productivity and C turnover under different land-use types, but are also determined by farmers' decisions to use a site in a specific way due to the demands of a crop or the suitability of a site, e.g. regarding machine use (see Leifeld et al. 2003: 65).

Table 7-34 Mean carbon stocks under permanent grassland on mineral soils (0-30 cm).

| Altitude [m] | Cs [t C ha ⁻¹] |
|--------------|----------------------------|
| <601 | 62.02 ± 13 |
| 601-1200 | 67.50 ± 12 |
| >1200 | 75.18 ± 9 |

Soil carbon stocks in organic soils under permanent grassland are calculated based on Leifeld et al. (2003, 2005). The approach uses measured carbon stocks in Swiss organic soils without differentiation among cropland and grassland. The mean soil organic carbon stock (0-30 cm) for organic soils is 240 ± 48 t C ha⁻¹.

Shrub Vegetation (CC32)

Due to the lack of data, the values of CC31 (Table 7-34) were used as the mineral soil carbon stocks for this category (0-30 cm).

The mean soil organic carbon stock (0-30 cm) for organic soils is 240 t C ha^{-1} .

Vineyards, Low-stem Orchards and Tree Nurseries (CC33)

The category includes carbon stocks in soils of vineyards, low-stem orchards and tree nurseries. In accordance to carbon stocks in biomass, only vineyards and low-stem orchards are considered. Both land-use types are assumed to have no grass undercover. Therefore, the soil carbon values of cropland, i.e. $53.40 \text{ t C ha}^{-1}$ (mineral soils, 0-30 cm) and 240 t ha^{-1} (organic soils, 0-30 cm) are taken for CC33 (see Chapter 7.4.4.2).

Copse (CC34)

Due to the lack of data, the values of CC31 (Table 7-34) were used as the mineral soil carbon stocks for this category (0-30 cm).

The mean soil organic carbon stock (0-30 cm) for organic soils is 240 t C ha^{-1} .

Orchards (CC35)

Cs of orchards was calculated in accordance to the biomass calculation. No specific Cs orchards values are available, and the mean value of grassland mineral soil carbon stocks from the two lower altitudinal zones (i.e. $64.76 \text{ t C ha}^{-1}$; cf. Table 7-34) was taken for mineral soils (0-30 cm), and the value of 240 t ha^{-1} for organic soils (0-30 cm).

Stony Grassland (CC36)

Soil organic carbon stocks under herbs and shrubs on stony surfaces were calculated according to the procedure described in Chapter 7.5.4.1, i.e. it is assumed that not more than 35% of the area of CC36 is covered with vegetation and thus only 35% of the area bears a mineral soil while the remainder is bare rock. Land use of this category mostly belongs to 'grassland' and 'unproductive land' and likely includes many of the former alpine grasslands (SFSO 2005). These grasslands are mainly located at altitudes $> 1200 \text{ m a.s.l.}$ Thus, using the respective value from Table 7-34, the carbon stock Cs of CC36 is calculated as:

$\text{Cs of CC36} = 0.35 * \text{Cs permanent grassland} > 1200 \text{ m} = 26.31 \text{ t C ha}^{-1} \text{ (0-30 cm)}$

The mean soil organic carbon stock (0-30 cm) for organic soils is 240 t C ha^{-1} . It is assumed that the small area covered by organic soils in CC36 (cf. CRF 5C1 'stony'), albeit entitled 'stony grassland', does not contain significant contributions from stones because bogs are free of stones as a matter of nature and fens usually contain, if any, only fine mineral sediments.

Unproductive Grassland (CC37)

The category CC37 'unproductive grasslands' includes grass and herbaceous plants at watersides of lakes and rivers including dams and other flood protection structures, constructions to protect against avalanches and rock slides, and alpine infrastructure (e.g. for skiing). For none of these land-use types, Cs data are currently available. Soil carbon stocks of CC37 'unproductive grassland' were arbitrarily set as the mean value of carbon stocks under permanent grassland on mineral soils (Table 7-34) in accordance to the procedure followed for biomass. Cs of CC37 is thus $68.23 \text{ t C ha}^{-1}$.

The mean soil organic carbon stock (0-30 cm) for organic soils is 240 t C ha^{-1} .

7.5.4.3 Changes in Carbon Stocks

Changes of carbon stock in biomass and in mineral soils are assumed to be zero for constant land use.

The annual net carbon stock change in organic soils on managed grassland (CC31, CC33 and CC35) was estimated to $-9.52 \text{ t C ha}^{-1}$ according to measurements in Europe including Switzerland as compiled by Leifeld et al. (2003, 2005) and rechecked by ART (2009b). For weakly managed grasslands (CC32, CC34, CC36 and CC37) the emission from organic soils was estimated to $5.30 \text{ t C ha}^{-1} \text{ yr}^{-1}$ according to available domestic data (ART 2011b).

In the case of land-use change, the net changes in biomass and soil of CC31, CC32, CC33, CC34, CC35, CC36, and CC37 are calculated as described in Chapter 7.1.3.

7.5.4.4 Carbon Emissions from Agricultural Lime Application

All CO_2 emissions caused by agricultural lime application are included under the category 5B Cropland (see Chapter 7.4.4.5, CRF-table 5(IV)).

7.5.4.5 NMVOC Emissions

Estimates for annual biogenic emissions of NMVOC (CRF-table 5) for forests and natural grassland in Switzerland are available in SAEFL (1996a): the value for natural grassland (unproductive vegetation) is 0.51 Gg .

7.5.4.6 Emissions from Wild Fires

Data on wildfires affecting Swiss grassland are obtained from cantonal authorities and are compiled by the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL 2013). Table 7-35 shows the annual burnt area from 1990 to 2012. The Swissfire database differentiates between 'grassland' and 'unmanaged land'. As a conservative assumption the sum of both categories is reported. Controlled burning is no common practice in Switzerland. Therefore, all fires are assigned to "wildfires".

The resulting CH_4 and N_2O emissions were calculated according to Equations 3.2.19/3.2.20 of IPCC (2003):

$$\begin{aligned} \text{Emission } \text{CH}_4 &= (\text{carbon released}) * (\text{emission ratio}) * 16/12 \\ \text{Emission } \text{N}_2\text{O} &= (\text{carbon released}) * (\text{emission ratio}) * 28/12 \\ \text{carbon released} &= (\text{burnt area}) * (\text{available fuel}) * (\text{combustion efficiency}) \end{aligned}$$

The "available fuel" was calculated as the area-weighted carbon stock in living biomass for all grassland categories (CC31 to CC37) in 2012: $7.68 \text{ Mg C ha}^{-1}$. Applying a default combustion efficiency of 0.74 (IPCC 2003, Table 3A.1.12, savanna grassland) the average amount of carbon that could be released by wildfires on grasslands is $5.68 \text{ Mg C ha}^{-1}$.

For CH_4 , the emission ratio of savannas was used: 0.004 (IPCC 1996, Table 4-15). The resulting mean emission is $0.030 \text{ Mg CH}_4 \text{ ha}^{-1}$.

For N_2O , a N/C ratio of 0.015 (IPCC 1996, pp. 4.83) and an emission ratio of 0.007 (IPCC 2003, Table 3A.1.15) result in a mean emission of $0.00094 \text{ Mg N}_2\text{O ha}^{-1}$.

The resulting annual emissions 1990-2012 on burnt areas in 5C grasslands are shown in Table 7-35 and are reported in CRF-table 5(V).

Table 7-35 Area of Grassland affected by wildfires (WSL 2013) and resulting CH₄ and N₂O emissions, 1990-2012.

| Year | Area burnt ha | CH ₄ Mg yr ⁻¹ | N ₂ O Mg yr ⁻¹ |
|------|------------------|--|---|
| 1990 | 637 | 19.315 | 0.598 |
| 1991 | 22 | 0.676 | 0.021 |
| 1992 | 6 | 0.183 | 0.006 |
| 1993 | 17 | 0.513 | 0.016 |
| 1994 | 175 | 5.292 | 0.164 |
| 1995 | 82 | 2.485 | 0.077 |
| 1996 | 43 | 1.297 | 0.040 |
| 1997 | 373 | 11.302 | 0.350 |
| 1998 | 73 | 2.203 | 0.068 |
| 1999 | 19 | 0.574 | 0.018 |
| 2000 | 22 | 0.662 | 0.020 |
| 2001 | 8 | 0.227 | 0.007 |
| 2002 | 257 | 7.799 | 0.241 |
| 2003 | 138 | 4.181 | 0.129 |
| 2004 | 4 | 0.130 | 0.004 |
| 2005 | 4 | 0.116 | 0.004 |
| 2006 | 6 | 0.171 | 0.005 |
| 2007 | 88 | 2.656 | 0.082 |
| 2008 | 29 | 0.879 | 0.027 |
| 2009 | 3 | 0.083 | 0.003 |
| 2010 | 1 | 0.040 | 0.001 |
| 2011 | 56 | 1.706 | 0.053 |
| 2012 | 4 | 0.133 | 0.004 |

7.5.5 Uncertainties and Time-Series Consistency

A range of possible carbon stock changes in mineral soils has been determined by the Swiss Soil Monitoring Network (NABO). The upper and lower margin of the 95% confidence interval for carbon stock changes under grassland is $0 \pm 0.57 \text{ t C ha}^{-1}$ (Keller 2013). This absolute uncertainty is used to calculate relative uncertainties for 5C1 and 5C2 by dividing with the mean net emission per hectare of 5C1 and 5C2, respectively. In 2012, the mean net emissions were $0.027 \text{ t C ha}^{-1}$ for 5C1 and $0.846 \text{ t C ha}^{-1}$ for 5C2 (calculated from CRF-table 5C). The resulting relative uncertainties are 2084% for 5C1 and 67% for 5C2, respectively (Table 7-5).

In the uncertainty analysis, these values were chosen for the overall emission factor uncertainties for CO₂ in sectors 5C1 and 5C2 as they dominate the other sources of uncertainty by far:

- Uncertainties for soil carbon stocks are given together with the mean value in the text above: 12-21% for mineral soils and 20% for organic soils. They take into account uncertainties in measured carbon contents and predicted soil bulk densities, i.e., they consider only uncertainties in emission factors.
- The uncertainty of the carbon stock change in organic soils of intensly managed grassland is 23% as reported by Leifeld et al. (2003: 56). For weakly managed grassland this uncertainty is 117% according to ART 2011b.
- The relative uncertainty in yield determination has been estimated at 13% for biomass carbon from both cropland and grassland (Leifeld and Fuhrer 2005). Data on biomass yields for different elevations and management intensities as published by FAL/RAC (2001) are based on many agricultural field experiments and have a high reliability.

Uncertainties of activity data of Grassland are described in Chapter 7.2.5. For wildfires, the emission factor uncertainties of CH₄ and N₂O were set to 70% (identical to forest land).

7.5.6 Category-Specific QA/QC and Verification

Changes in Living Biomass

The assumption of a constant carbon stock in living biomass has been reconsidered (UNFCCC 2007: §97). According to Schneider (2010) yields on meadows and pastures did not increase since 1990. Neither management nor the share of clover did significantly change over the past 20 years. Consequently, the current approach has been reconfirmed.

In 2012 an assessment of the appropriateness of the estimated pools of carbon in living biomass was conducted (ART 2012a). It came to the conclusion that almost all carbon stocks and carbon stock changes are in the expected range of the IPCC Guidelines and Good Practice Guidance. Nevertheless there is room for improvements. However, given the relatively low significance of the respective emissions a major effort in this area is hardly justified. Consequently, the biomass carbon pools will eventually be recalculated only in the course of the new planned Tier 3 approach for quantification of carbon stocks and carbon stock changes in agricultural soils (see also Chapter 7.5.8).

Changes in SOC Pools

A SOC pool dataset provided by the Swiss Soil Monitoring Network (NABO; see Chapter 7.3.6) supports the Tier 1 assumption that changes of carbon stocks in mineral soils are zero for grassland remaining grassland (cf. UNFCCC 2007: §97). The SOC pool measured at 33 grassland monitoring sites in the Swiss Soil Monitoring Network showed in average a slight increase during the period 1985 to 2000 and a slight decrease thereafter (Figure 7-8). SOC pools ranged between 20.9 and 183.2 t C ha⁻¹, the average SOC pool for the 33 grassland monitoring sites was 77.9 t C ha⁻¹ (0-20 cm). Two alpine grassland sites above 1200 m a.s.l. showed remarkable SOC pools of about 120 and 173 t C ha⁻¹ (0-20 cm). The confidence interval of the mean SOC pool versus time was ± 0.9 t C ha⁻¹. In total, the results of the soil monitoring data indicate that Swiss grassland mineral soils did not act as a net source or sink of carbon during the last 20 years.




Figure 7-8 Time series of measured SOC pool changes in the top soil (0-20 cm) at the 33 NABO grassland sites from the 1st to the 5th re-sampling campaigns. SOC pools were centred by the median SOC pool of all re-samplings of the monitoring site. Each pool value presents the median of four bulked soil samples per campaign with measured SOC and bulk density. The error bars indicate the 25% and 75% percentiles resulting from the spatial variation of the sites and the errors along the measurement chain. The altitude of the grassland sites ranges between 265 and 2340 m a.s.l.

The slight increase and decrease will be subject for further detailed analysis. Partly, it may be attributed to natural variation of soil sampling (see Chapter 7.3.6). Furthermore, we presume that at grassland sites with intensive management and large manure application the temporal change of the SOC content is partly related to the nitrogen input fluxes and nitrogen content in soil. Therefore, the total nitrogen content of the archived soil samples will be measured and the correlation to the SOC content analysed. Moreover, management data of the monitoring sites gathered directly from the farmers since 1985 permit the calculation of annual nutrient fluxes of the sites (Keller et al. 2005). In this way temporal changes in nutrient management of the grassland sites can be separated from other effects that may cause temporal changes of SOC contents in grassland soils.

Short-term Land-Use Changes between Grassland and Cropland

See comments in Chapter 7.4.6.

7.5.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in the category 5C.

Emissions from wildfires on grassland are reported the first time (see Chapter 7.5.4.6).

7.5.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines represents the largest improvement. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented. Other projects and planned improvements may be postponed in order to give first priority to changes related to the new reporting guidelines.

In 2011, Agroscope started a three-year running research project that aims to identify (drained) fens and raised bogs under different land uses beyond the national inventories of bogs and fens in order to improve the AD estimates of organic soils ("Area and location of organic soils and peatlands in Switzerland").

Switzerland intends to make better use of the SOC data provided by the Swiss Soil Monitoring Network (NABO) in forthcoming submissions. A contract was signed in 2011 that arranges a close collaboration for the period 2012-2014.

Furthermore, information about carbon stock changes in soils for grassland remaining grassland will become available from Agroscope research activities. A pilot study to evaluate possible Tier 3 methodological approaches for quantification of carbon stocks and carbon stock changes in agricultural soils (including meadows and pastures) (cf. UNFCCC 2007: §97) has recently been terminated (Köck et al. 2013).

Ongoing efforts to combine SOC measurements on the level of soil fractions with modelled pools (Zimmermann et al. 2007; Leifeld et al. 2009) will be combined with the planned Tier 3 approach for an independent check of emission and removal rates for cropland and grassland conversions.

A new study on GHG emissions from peatlands and organic soils under different land use (Agroscope in collaboration with the University of Basel, 2013-2016, financed by FOEN) will improve the robustness of domestic emission factor estimates for soils rich in organic matter in the medium term.

Projects in a new national research programme ("Sustainable Use of Soil as a Resource", <http://www.nfp68.ch/E>) focus on (1) sustainable management of organic soils, (2) agricultural management and below ground carbon inputs, and (3) an integrated modelling framework to monitor and predict trends of agricultural management and their impact on soil functions at multiple scales. In the long term, results from these studies are expected to improve the reporting of emissions by sources and removals by sinks in category 5C.

7.6 Category 5D – Wetlands

7.6.1 Description

The categories 5D1 Wetlands remaining Wetlands and 5D2 Land converted to Wetlands are not key categories.

Wetlands consist of surface waters (CC41) and unproductive wet areas such as shore vegetation, fens or (raised) bogs (CC42), see Table 7-2 and Table 7-6.

7.6.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.6.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.6.4 Methodological Issues

7.6.4.1 Carbon in Living Biomass

Surface Waters (CC41)

Surface waters have no carbon stocks by definition.

Unproductive Wetland (CC42)

CC42 consists of unmanaged or weakly managed grassland, bushes or tree groups. The pool of living biomass was estimated to 6.50 t C ha^{-1} (Mathys and Thürig 2010).

7.6.4.2 Carbon in Soils

In general, the soil carbon stock for surface waters (CC41) is zero. However, for CC41 situated in areas with organic soil (see Chapter 7.2.3.1 and Table 7-7), the soil carbon stock is set to 240 t C ha^{-1} (0-30 cm). These surface waters are assumed to be shallow ponds as integrated parts of fens or bogs.

Land cover in CC42 explicitly includes peatlands protected by Federal Legislation (Swiss Confederation 1991a, 1994) as well as reed. More than 10% of the unproductive wetland are located on organic soils (cf. Table 7-7) as defined in Chapter 7.2.3.1. In this case the carbon stock in soils is 240 t C ha^{-1} (0-30 cm). Currently, no specific soil data are available for CC42 on mineral soils. As a first guess, it is suggested that the soil carbon stock of unproductive wetlands is similar to permanent grassland on mineral soils (mean value: $68.23 \text{ t C ha}^{-1}$; 0-30 cm).

7.6.4.3 Changes in Carbon Stocks

Changes of carbon stock in biomass and in mineral soils are assumed to be zero for wetlands remaining wetlands.

The emission from organic soils under CC41 is assumed to be zero because the respective areas are not drained.

The emission from organic soils under CC42 was estimated to $5.30 \text{ t C ha}^{-1} \text{ yr}^{-1}$ according to domestic data (ART 2011b). This value is used for weakly managed ecosystems such as fens and unmanaged ecosystems such as raised bogs. Bogs and fens are protected to a large part by Federal Ordinances (Swiss Confederation 1991a, 1994) and drainage is therefore not allowed. However, the impact of old drainages constructed before 1990 probably leads to a certain emission.

In the case of land-use change, the net changes in biomass and soil of both CC41 and CC42 are calculated as described in Chapter 7.1.3.

For land converted to unproductive wetland (CC42) a conversion time of one year was chosen for the carbon stock change in living biomass and in dead organic matter (see Table 7-3). This was done because at the moment when the land-use change is detected in the sense of changes in the vegetation cover on the AREA aerial photographs the land-use change has already occurred (cf. UNFCCC 2009: §82). For carbon stock changes in soil the conversion time is 20 years.

7.6.4.4 Non CO₂ Emissions from Drainage of Soils and Wetlands

The reporting of non-CO₂ emissions from drainage of soils and wetlands is not mandatory. Due to the lack of data Switzerland decided not to prepare emission estimates for drainage of soils.

An estimate of $0.43 \text{ Gg CH}_4 \text{ yr}^{-1}$ emitted by reservoirs (flooded lands) is given by Hiller et al. (2014). The estimate encompasses 97 artificial lakes covering a total area of 10.6 kha. This emission is reported in CRF-table 5(II)D.

7.6.5 Uncertainties and Time-Series Consistency

As a best guess, a value of 50% was chosen for the overall emission factor uncertainty in sector 5D (Table 7-5).

The uncertainty of the emission factor for carbon stock losses in organic soils is 100% based on monitoring values compiled by ART (2011b).

Uncertainties of activity data of Wetlands are described in Chapter 7.2.5.

7.6.6 Category-Specific QA/QC and Verification

No category-specific QA/QC activities have been carried out.

7.6.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in the category 5D.

Emissions of CH₄ are reported for flooded lands (reservoirs) in CRF-table 5(II)D (Chapter 7.6.4.4). In former submissions, this emission source was not reported.

7.6.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines (IPCC 2006) represents the largest improvement. To accomplish this transition, other projects or planned improvements may be postponed in order to give first priority to changes related to the new reporting guidelines.

In 2011, Agroscope started a three-year running research project that aims to identify (drained) fens and raised bogs under different land uses beyond the national inventories of bogs and fens in order to improve the AD estimates of organic soils ("Area and location of organic soils and peatlands in Switzerland").

A new study on GHG emissions from peatlands and organic soils under different land use (Agroscope in collaboration with the University of Basel, 2013-2016, financed by FOEN) will improve the robustness of domestic emission factor estimates for soils rich in organic matter in the medium term.

Lake Wohlen, a hydroelectric reservoir on the Swiss Plateau, has been shown to be a large emitter of CH₄ (DeSontro et al. 2010). Sediments cores from the reservoir were sampled in March 2014 by the University of Bern. The project (2014-2015, co-financed by FOEN) focuses on the question if Lake Wohlen is a representative system within Switzerland.

7.7 Category 5E – Settlements

7.7.1 Category Description

Tier 2 Key category 5E2

CO₂ from Land converted to Settlements
(2012: level and trend)

The category 5E1 Settlements remaining Settlements is not a key category.

Settlements consist of buildings/constructions (CC51), herbaceous biomass in settlements (CC52), shrubs in settlements (CC53), and trees in settlements (CC54) as shown in Table 7-2 and Table 7-6.

7.7.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.7.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.7.4 Methodological Issues

7.7.4.1 Carbon in Living Biomass

Buildings and Constructions (CC51)

Buildings/constructions contain no carbon by default.

Herbaceous Biomass, Shrubs and Trees in Settlements (CC52, CC53, CC54)

Results of the research project “Baumbiomasse in der Landschaft” are used for carbon stocks in living biomass as follows: 9.54 t C ha⁻¹ for CC52, 15.43 t C ha⁻¹ for CC53 and 20.72 t C ha⁻¹ for CC54 (Mathys and Thürig 2010: Table 7).

7.7.4.2 Carbon in Soils

The carbon stock in soil for CC51 (buildings and construction) was set to zero.

In case of land-use changes to CC51 or from CC51, only 50% of the difference between carbon stocks before and after the change is reported as a source or sink, respectively. The reason for this is that the soil organic matter on construction sites is stored temporarily and is later used for replanting the surroundings or for vegetating dumps, for example. According to paragraph 7 of the "Ordinance against deterioration of soils" (Swiss Confederation 1998) the soil material excavated on a construction site must be treated in such a way that it can be used as a soil again. When the material is re-used (e.g. for re-cultivations) the fertility of the soil must not be affected. This regulation ensures that a large part of the soil organic matter is preserved on land converted to and from CC51. Switzerland has chosen the factor 0.5 (i.e.

50% stabilised SOC fraction which is not likely to be oxidised in the medium term) to reflect this domestic soil protection measure (see discussion in Leifeld et al. 2003: 67). Thus, the equation 7.6 presented in Chapter 7.1.3.2 is adjusted as follows if a=CC51 or b=CC51:

$$\Delta C_{s,i,ba} = [0.5 * (stockC_{s,i,a} - stockC_{s,i,b}) / CT] * A_{i,ba}$$

where:

| | |
|------------|--|
| a | land-use category after conversion (CC = a) |
| b | land-use category before conversion (CC = b) |
| ba | land use conversion from b to a |
| i | spatial stratum |
| $A_{i,ba}$ | area of land (ha) converted from b to a in the spatial stratum i |
| CT | conversion time (yr), see Table 7-3. |

The carbon stock in mineral soil for CC52, CC53 and CC54 is 53.40 t C ha⁻¹ (0-30 cm, same value as for cropland).

For organic soils the carbon stock is 240 t C ha⁻¹ (0-30 cm, same value as for the other land-use categories on organic soils).

7.7.4.3 Changes in Carbon Stocks

Changes of carbon stock in biomass and in mineral soils are assumed to be zero for settlements remaining settlements.

On organic soils, the following emission factors were applied:

- 9.52 t C ha⁻¹ yr⁻¹ for CC52. This corresponds to the value used for cropland because CC52 areas are managed (gardens, parks) (Leifeld et al. 2003, 2005 and rechecked by ART 2009b).
- 5.30 t C ha⁻¹ yr⁻¹ for CC53 and CC54. This corresponds to the value used for weakly managed grasslands (ART 2011b).

In the case of land-use change, the net changes in biomass and soil of CC51, CC52, CC53, and CC54 are calculated as described in Chapter 7.1.3.

7.7.5 Uncertainties and Time-Series Consistency

As a best guess, a value of 50% was chosen for the overall emission factor uncertainty in sector 5E (Table 7-5).

Uncertainties of activity data of Settlements are described in Chapter 7.2.5.

7.7.6 Category-Specific QA/QC and Verification

No.

7.7.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in the category 5E.

7.7.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines (IPCC 2006) represents the largest improvement. To accomplish this transition, other projects or planned improvements may be postponed in order to give first priority to changes related to the new reporting guidelines.

7.8 Category 5F – Other Land

7.8.1 Description

The categories 5F1 Other Land remaining Other Land and 5F2 Land converted to Other Land are not key categories.

As shown in Table 7-2 and Table 7-6 other land (CC61) covers non-vegetated areas such as glaciers, rocks and shores.

7.8.2 Information on Approaches Used for Representing Land Areas and on Land-use Databases Used for the Inventory Preparation

See Chapter 7.2.

7.8.3 Land-use Definitions and the Classification Systems Used and their Correspondence to the LULUCF Categories

See Chapter 7.2.

7.8.4 Methodological Issues

By definition, other land has no carbon stocks. Coherently, changes of carbon stock in biomass and in soil are assumed to be zero for other land remaining other land.

In the case of land-use change, the net C changes in biomass and soil are calculated as described in Chapter 7.1.3.

7.8.5 Uncertainties and Time-Series Consistency

As a first guess, a value of 50% was chosen for the overall emission factor uncertainty in sector 5F2 (Table 7-5).

Uncertainties of activity data of Other Land are described in Chapter 7.2.5.

7.8.6 Category-Specific QA/QC and Verification

No.

7.8.7 Category-Specific Recalculations

The completion of the AREA surveys in 2013 (see Chapter 7.2) led to a recalculation in the category 5F.

7.8.8 Category-Specific Planned Improvements

In the course of the next inventory preparation, the implementation of the new reporting guidelines (IPCC 2006) represents the largest improvement.

8 Waste

8.1 Overview

8.1.1 Greenhouse Gas Emissions

Within the waste sector, emissions from four source categories are considered:

- 6A Solid Waste Disposal on Land,
- 6B Wastewater Handling,
- 6C Waste Incineration,
- 6D Other Waste




Figure 8-1 Switzerland's greenhouse gas emissions in the waste sector 1990–2012.

Table 8-1 Trend of total GHG emissions from waste management in Switzerland 1990-2012.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 63 | 60 | 56 | 50 | 40 | 35 | 31 | 28 | 26 | 23 |
| CH ₄ | 734 | 731 | 706 | 654 | 600 | 600 | 578 | 569 | 550 | 528 |
| N ₂ O | 210 | 213 | 214 | 210 | 211 | 213 | 214 | 217 | 219 | 218 |
| Sum | 1007 | 1004 | 976 | 914 | 852 | 847 | 823 | 814 | 795 | 768 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 19 | 16 | 13 | 12 | 12 | 12 | 12 | 12 | 13 | 12 |
| CH ₄ | 497 | 477 | 468 | 422 | 421 | 404 | 396 | 375 | 349 | 326 |
| N ₂ O | 225 | 224 | 227 | 234 | 239 | 238 | 244 | 249 | 268 | 274 |
| Sum | 741 | 717 | 708 | 669 | 673 | 654 | 653 | 637 | 629 | 612 |

| Gas | 2010 | 2011 | 2012 |
|-------------------------|------|------|------|
| CO ₂ eq (Gg) | | | |
| CO ₂ | 13 | 12 | 12 |
| CH ₄ | 314 | 299 | 287 |
| N ₂ O | 281 | 287 | 292 |
| Sum | 608 | 598 | 591 |

As illustrated in Table 8-1, in the waste sector a total of 591 Gg CO₂ equivalents were emitted in the year 2012. 29.3% stem from category 6A Solid Waste Disposal on Land, 40.8% of the emissions stem from category 6B Wastewater Handling, 7.3% from 6C Waste Incineration and 21.2% from 6D Others.

The total greenhouse gas emissions in the waste sector show a decrease of 41.3% from 1990 until 2012. From 1990 – 2007, the greenhouse gas emissions had been clearly dominated by the ones from the source category 6A Solid Waste Disposal on Land. In 2008, emissions of 6A Solid Waste Disposal on Land and 6B Wastewater Handling became equivalent. Since 2009 6B Wastewater Handling has been the most important source category.

While CO₂ and CH₄ emissions in the waste sector are decreasing since 1990, N₂O emission continue to augment, mainly due to the increasing number of inhabitants and related emissions caused by waste water handling. CH₄ emissions have decreased from 1990 until 2012 by 60.9%. N₂O emissions increased by 39.2% from 1990 until 2012. In 2012, N₂O has replaced CH₄ as the most important greenhouse gas in the waste sector. CO₂ is of minor importance in the waste sector. CO₂ emissions have remained rather constant since 2003 at a low level of about 12Gg. Since 1990 they have been reduced by 80.5%.

The relative trends of the gases can be seen in Figure 8-2.

Please note that according to IPCC Good Practice Guide all emissions from waste-to-energy, where waste material is used directly as fuel or converted into a fuel, are reported under the sector 1 Energy (see also Figure 8-4). Therefore the largest share of waste-related emissions in Switzerland is not reported under sector 6 Waste, as the box below shows.




Figure 8-2 Trend of total GHG emissions from waste management in Switzerland 1990-2012.

Box: Waste related GHG emissions in Switzerland

The respective GHG emissions are reported in different chapters within the National Inventory. The following figure provides an overview on all waste related GHG emissions in Switzerland, not only the ones reported in the present Chapter 8 (see also Figure 8-4).




Figure 8-3 Waste related GHG emissions from 1990-2012.

8.1.2 Overview on Waste Management in Switzerland

The goals and principles regarding waste management in Switzerland are stated in the Guidelines on Swiss Waste Management (BUS 1986) and in the Waste Concept for Switzerland (SAEFL 1992). The four principles are:

- The generation of waste shall be avoided as far as possible.
- Pollutants from manufacturing processes and in products shall be reduced as far as possible.
- Waste shall be recycled wherever this is environmentally beneficial and economically feasible.
- Waste shall be treated in an environmentally sound way. In the long term only materials of final storage quality shall be disposed of in landfills.

Figure 8-4 gives a general overview of the type of treatment and amounts of waste fractions treated in the respective sectors in the NIR of Switzerland in 2012 including imports and exports. Only waste fractions that are emission-relevant are shown. It illustrates where the processes related to the waste management system are reported in the NIR.

1 Energy: In accordance with the IPCC provisions (IPCC 1997c) emissions from the combustion of waste-to-energy activities, if waste is used as an alternative fuel or used for energy production, emissions are reported in 1A Fuel Combustion Activities. This applies for municipal solid waste incineration plants (MSWIP), special waste incineration plants (SWIP) where energy is being recovered, as well as the cement industry where special waste and sewage sludge are used as an alternative fuel in the cement production. The fermentation of biomass is also reported in sector 1 Energy, as biogas is used for co-generation of heat and electricity. The energy production from renewable goods, such as the use of waste wood in wood-fired power stations is reported under 1A4ai and 1A4bi.

4 Agriculture: Since 2003 it is forbidden to use sewage sludge as a fertilizer. In 2012, within sector 4 Agriculture only compost used as fertilizer was emission-relevant (due to N₂O emissions as described in chapter 6.5.2 table 6-18).

6 Waste: Only if waste is not used for energy production purposes, it is reported under the sector 6 Waste. Solid waste disposal on land does not occur anymore in Switzerland as incineration is the mandatory disposal option for burnable waste since 2000. The emissions related to the waste water management are reported under 6B. Emissions related to the digestion process which stem from processes that are not related to the energy production, such as the storage of fermented biomass are reported under 6D. 6D also describes emissions related to composting. 6C is only a small fraction, consisting of illegal waste incineration, sewage sludge incineration and burning of residues in agriculture and forestry. Special waste incineration without energy recovery such as cable incineration or hospital waste incineration no longer takes place. For this reason it is crossed out in the figure. These waste fractions are also incinerated in MWSIP and therefore reported under the sector 1 Energy.




Figure 8-4 Overview on the type of treatment and amounts of waste fractions treated in the respective sectors in Switzerland in 2012

Abbreviations: MSWIP: Municipal Solid Waste Incineration Plant, SWIP: Special Waste Incineration Plant, SSIP: Sewage Sludge Incineration Plant

In the following the treatment and amounts of relevant waste fractions of Switzerland in 2012 are outlined (see also Figure 8-4). Recycled amounts are not shown in the figure because they are not emission-relevant.

- **Municipal Solid Waste:** Switzerland has a very high recycling rate, 50% of the municipal solid waste is collected separately and recycled (FOEN 2013h). While 2'790 Gg MSW were recycled, 2'787 Gg were incinerated in 2012. Additional 220 Gg MSW were imported into Switzerland for incineration (in the first place from neighbouring

countries such as Germany, France, Austria and Italy). The import of waste into Switzerland needs to be authorized by the Federal Office for the Environment. A part of the separately collected plastic fractions from households and industry which can not be recycled, is used as a alternative fuel in the cement production (45 Gg in 2012).

- Construction Waste:** It is assumed that about 1.5t construction waste is produced per inhabitant¹⁵. Thus, about 12'000 Gg construction waste was generated in Switzerland in 2012¹⁶. From this quantity 8'400 Gg (about 70%) was recycled. About half of the recycling took place at the construction site, e.g. by reusing material left after breaking up the road cover. The other half was separated at the construction site and recycled individually, e.g. used glass, used metals, used concrete etc, EMPA 2004a A minor amount e.g. 425 Gg reflecting the burnable part of construction waste was incinerated in MSWIP as shown in Figure 8-4 (internal numbers provided by the waste section of FOEN). The remaining, inert construction waste (about 3'125 Gg) was disposed of in landfills for inert waste.
- Special Waste:** Special waste refers to a highly divers waste fraction containing hospital wastes, batteries, electronic waste, hazardous industrial sludges, contaminated soils, etc.). According to the yearly reported special waste statistics (FOEN 2013i) about 1'749 Gg special waste was generated in Switzerland in 2012. About 326 Gg special waste was recycled, 294 Gg were biologically treated, 522 Gg landfilled and approximately 426 Gg exported for landfilling (FOEN 2013i). Only the amount of incinerated special waste is emission-relevant and therefore shown in figure 8-4. In 2012, 91 Gg were incinerated in MWSIP (EMIS 2014/1 A 1 a_Kehrichtverbrennungsanlagen), 263 Gg in SWIP (EMIS 2014/1 A 1 a_Sondermüllverbrennungsanlagen 20140123). In 2012 126 Gg special waste was used as an alternative fuel in the cement production (EMIS 2014/ 1 A 2 f i_Zementwerke Feuerung).
- Sewage Sludge:** Since 2006 it is forbidden to use sewage sludge as a fertilizer in agriculture due to the content of organic contaminants, heavy metals and other substances. About 210 Gg (dry matter) sewage sludge was generated in 2012 (FOEN 2013h). 56 Gg were incinerated in MSWIP, 104 Gg incinerated in SWIP (without energy recovery) (internal numbers provided by the waste section of FOEN) and 52 Gg used as a fuel in the cement industry (EMIS 2014/ 1 A 2 f i_Zementwerke Feuerung) .
- Biomass:** the term biomass refers to a broad range of materials such as garden waste, grass, waste wood, liquid manure and production from the food industry or further fractions, depending on the process concerned. In 2012 23 Gg residues from agriculture, forestry and private gardens were burned without energy recovery (EMIS 2014/6C1 Abfallverbrennung in der Land- und Forstwirtschaft). 1'737 Gg biomass was fermented (EMIS 2014/ 1 A 1 a und 6 D_Vergärung IG und LW). The amount of biomass composted in large-scale composting facilities (industrial composting) was 759 Gg wet matter in 2012. It is assumed that composted households waste adds up to 10 percent of industrial composting. (EMIS 2014/ 6 D_Kompostierung Industrie and EMIS 2014/ 6 D_Kompostierung Haushalte). Quantities of biomass refer to the wet matter. 84 Gg of biomass such as used wood or animal fat, was used as an alternative fuel in the cement industry (EMIS 2014/ 1 A 2 f i_Zementwerke Feuerung). Compost used as a fertilizer amounted to 3.424t N (see chapter). 3'032 TJ wood was used for energy production purposes, for example in wood-fired power stations, chimneys, or pellets heatings systems (SFOE 2013b).

¹⁵ It is assumed that this estimation in FOEN 2010j still applies for the year 2012.

¹⁶ Inhabitants in Switzerland in 2012: 7.997 million (SFSO 2013a)

8.2 Source Category 6A – Solid Waste Disposal on Land

8.2.1 Source Category Description

Tier 1 key category 6A

CH₄ emissions from managed waste disposal on land (trend)

Tier 2 key category 6A CH₄

CH₄ emissions from managed waste disposal on land (level and trend)

The source category 6A1 Managed Waste Disposal on Land comprises all emissions from managed solid waste landfill sites.

As incineration is the mandatory disposal option for burnable waste since 2000, input into managed solid waste landfill sites have dropped down to zero. Emissions thus stem from landfilling before 2000. Emissions from the source category 6A2 Unmanaged Waste Disposal Sites are included in source category 6A1 Managed Waste Disposal on Land. This is motivated by the fact that in Switzerland officially no unmanaged waste disposal sites exist. The effective quantity of waste not properly treated in landfills is estimated to be very small. However, no reliable data is available.

In Switzerland, in 2012 six managed biogenic active landfills are equipped to recover landfill gas (SFOE 2013a). The landfill gas is generally used in co-generation plants in order to produce electricity and heat. Some landfill gas is used to generate heat only. A very small amount of the landfill gas is flared.

Table 8-2 Specification of source category 6A Solid Waste Disposal on Land. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 6A | Source | Specification | Data Source |
|-----|--------------------------------|--|--|
| 6A1 | Managed Waste Disposal on Land | Emissions from managed solid waste landfill sites. | EMIS 2014/1A1a & 6A1 Kehrlichtdeponien |
| 6A2 | Unmanaged Waste Disposal Sites | Emissions from all other waste disposal sites not included in source category 6A1. (included in 6A1) | |
| 6A3 | Others | Not occurring in Switzerland | |

8.2.2 Methodological Issues, Managed Waste Disposal on Land (6A1)

Methodology

The emissions are calculated in four steps:

- The rate of CH₄ generation over time is based on the First Order Decay model (FOD) according to IPCC (IPCC 1997a-c). The following equation is applied to calculate the CH₄ generation in the year t:

$$\text{CH}_4 \text{ generated in the year } t [\text{Gg/year}] = \sum_x [A \cdot k \cdot M(x) \cdot L_0(x) \cdot e^{-k(t-x)}] \cdot (1-\text{OX})$$

where

t = current year

x = the year of waste input, $x \leq t$

A = $(1-k)/k$, norm factor (fraction)

k = methane generation rate [1/yr]

M(x) = the amount of waste disposed in year x

| | |
|------------|--|
| $L_0(x)$ = | methane generation potential ($MCF(x) \cdot DOC(x) \cdot DOC_F \cdot F \cdot 16/12$) [Gg CH ₄ / Gg waste] |
| $MCF(x)$ = | methane correction factor (fraction) |
| $DOC(x)$ = | degradable organic carbon [Gg C/ Gg waste] |
| DOC_F = | portion of DOC, that is converted to landfill gas (fraction) |
| F = | portion of CH ₄ in landfill gas (fraction) |
| $16/12$ = | factor to convert C to CH ₄ . |
| OX = | oxidation factor (fraction) |

The following general assumptions are made:

$MCF(x)$ = constant = 1 (default value according to IPCC for managed solid waste disposal sites)

OX = 0.1 (default value according to IPCC 1997a-c)

DOC_F = 0.6 (default value according to IPCC 1997a-c)

F = 0.5 (default value according to IPCC 1997a-c)

The degradable organic carbon $DOC(x)$ is calculated based on country specific waste composition for municipal solid waste, construction waste and sewage sludge and default values for $DOC(x)$ adopted from IPCC Guidelines, table 6-3. Assumptions for the composition are summarized in Table 8-3.

Table 8-3 Composition of landfilled Municipal Solid Waste (MSW), Construction Waste (CW) and Sewage Sludge (SS) used to derive DOC (Source 1A1 a / 6A1 EMIS Kehrichtdeponien)

| Fraction IPCC | MSW 1993 | MSW 2003 | CW 1993 | CW2003 | SS1993 | SS2003 |
|--|----------|----------|---------|--------|--------|--------|
| Paper, textiles and cardboard % | 28 | 20 | 0 | 0 | 0 | 0 |
| Garden waste and non-food organic putrescible % | 5 | 2 | 0 | 0 | 100 | 100 |
| Food waste % | 22 | 27 | 0 | 0 | 0 | 0 |
| Wood and straw % | 0 | 0 | 67 | 67 | 0 | 0 |
| Other (glass, metals, plastics, minerals, etc.) do not contribute to methane generation)% | 45 | 49 | 33 | 33 | 0 | 0 |

The methane generation rate k is based on expert judgement taking into account the country specific conditions as well as the type of waste disposed of in landfills (EMIS 2014/1A1a & 6A1 Kehrichtdeponien).

For the calculation of CH₄ generation three different categories of waste are distinguished. The three categories are i) municipal solid waste, ii) construction waste, and iii) sewage sludge.

The following parameters are applied for the calculation of CH₄ generated.

Table 8-4 Parameters used for FOD model

| | k [1/yr] | L_0 [Gg CH ₄ / Gg waste] | resulting DOC [-] |
|-----------------------|------------|---------------------------------------|---|
| municipal solid waste | 0.139 | 0.050 | 1990-1992: 0.15 1993-2002: linear interpolation 2003-2012: 0.12 |
| construction waste | 0.046 | 0.080 | 0.20 |
| sewage sludge | 0.069 | 0.068 | 0.17 |

- ii) In a second step, the amount of CH₄ that is recovered and used as fuel for co-generation units as well as for flaring is subtracted from the CH₄ generated in landfills (resulting from step 1).

$$\text{CH}_4 \text{ emissions}_{\text{step ii)}} = \text{CH}_4 \text{ emissions}_{\text{step i)}} - (\text{CH}_4 \text{ emissions}_{\text{step i)}} * \text{FI(t)} - \text{Qco-gen(t)}$$

where

FI(t) = portion of generated methane that is flared in the present year (fraction)

Qco-gen(t) = CH₄ which is recovered in co-generation units in the present (Gg)

- iii) In the third step CH₄ emissions from on-site open burning are added. This results in the overall CH₄ emissions from landfill sites.

$$\text{CH}_4 \text{ emissions}_{\text{step iii)}} = \text{CH}_4 \text{ emissions}_{\text{step ii)}} + \text{Qopen(t)}$$

where

Qopen(t) = CH₄ which is emitted from open burning in the present year (Gg)

- iv) In the fourth and last step the emissions of the other gases are calculated. The respective emissions are considered as proportional to the CH₄ burnt (co-generation and flaring), or to the waste quantity burnt (open burning), respectively.

Emission Factors

Emission factors for CO₂, CH₄, CO, NMVOC and SO₂ are country specific based on measurements and expert estimates, documented in EMIS 2014/1A1a & 6A1 Kehrichtdeponien. CO₂ emissions from non-biogenic waste are included, while the CO₂ emissions from biogenic waste are excluded from total emissions.

The following table presents the emission factors used in 6A1:

Table 8-5 Emission Factors for 6A1 Managed Waste Disposal Sites on Land in 2012.

| Source | CO ₂ biogen | CO ₂ fossil | CH ₄ | NO _x | CO | NMVOC | SO ₂ |
|---|--------------------------------|---------------------------|-----------------|-----------------|----|-------|-----------------|
| 6A1 Managed Waste Disposal on Land | t / t CH ₄ produced | | | | | | |
| Direct emissions from landfill | 3.00 | 0 | 1 | | | | |
| | kg / t CH ₄ burned | | | | | | |
| Flaring | 2750 | 0 | | 1 | 17 | | |
| | kg / t waste burned | | | | | | |
| Open burning | 663 | 608 | 6 | 2.5 | 50 | 16 | 0.8 |

Activity data

One set of activity data for Managed Waste Disposal on Land (6A1) are the waste quantities disposed on landfills and the municipal solid waste burnt on-site.

Activity data for Managed Waste Disposal on Land (6A1) are taken from EMIS 2014/1A1a & 6A1 Kehrichtdeponien.

Table 8-6 Activity data in 6A1: Waste disposed of on Managed Landfill Sites from 1990 to 2012 (source EMIS 2014/1A1a & 6A1 Kehrichtdeponien).

| Source / Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 6A1 Managed Waste Disposal on Land | | | | | | | | | | | |
| Municipal solid waste (MSW) | Gg | 637.0 | 637.0 | 637.0 | 637.0 | 581.2 | 531.9 | 482.7 | 472.8 | 463.0 | 465.3 |
| Construction waste | Gg | 147.0 | 170.5 | 170.5 | 123.5 | 59.1 | 47.3 | 35.5 | 35.5 | 41.4 | 41.6 |
| Sewage sludge | Gg (dry) | 58.8 | 58.8 | 58.8 | 48.8 | 39.0 | 27.7 | 16.3 | 12.2 | 8.1 | 6.1 |
| Open burned waste | Gg | 17.2 | 20.0 | 20.0 | 18.2 | 10.9 | 9.7 | 8.5 | 8.3 | 8.2 | 5.5 |
| Total waste quantity | Gg | 860.0 | 886.3 | 886.3 | 827.5 | 690.2 | 616.6 | 542.9 | 528.7 | 520.6 | 518.5 |

| Source / Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|----------|-------|-------|------|------|------|------|------|------|------|------|
| 6A1 Managed Waste Disposal on Land | | | | | | | | | | | |
| Municipal solid waste (MSW) | Gg | 288.8 | 184.8 | 80.8 | 52.2 | 23.7 | 13.6 | 3.5 | 1.5 | 1.2 | 0.0 |
| Construction waste | Gg | 30.7 | 17.7 | 4.8 | 3.4 | 2.0 | 1.4 | 0.8 | 0.0 | 0.0 | 0.0 |
| Sewage sludge | Gg (dry) | 4.1 | 3.9 | 3.6 | 2.6 | 1.6 | 1.0 | 0.3 | 0.0 | 0.0 | 0.0 |
| Open burned waste | Gg | 3.5 | 2.2 | 0.9 | 0.6 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total waste quantity | Gg | 327.0 | 208.6 | 90.1 | 58.8 | 27.5 | 16.1 | 4.7 | 1.5 | 1.2 | 0.0 |

| Source / Parameter | Unit | 2010 | 2011 | 2012 |
|---|----------|------|------|------|
| 6A1 Managed Waste Disposal on Land | | | | |
| Municipal solid waste (MSW) | Gg | 0.0 | 0.0 | 0.0 |
| Construction waste | Gg | 0.0 | 0.0 | 0.0 |
| Sewage sludge | Gg (dry) | 0.0 | 0.0 | 0.0 |
| Open burned waste | Gg | 0.0 | 0.0 | 0.0 |
| Total waste quantity | Gg | 0.0 | 0.0 | 0.0 |

Table 8-6 documents the amounts of municipal solid waste, construction waste, sewage sludge and open burnt waste disposed of on managed waste disposal sites over the time period 1990 – 2012.

The continuous reduction happened due to changes in the legislative framework, making incineration the mandatory disposal option for burnable waste and banning its disposal on landfills from 1 January 2000. The amounts of burnable waste disposed of on managed waste disposal sites dropped down to zero in 2009.

The other activity data for Managed Waste Disposal on Land (6A1) are CH₄ direct emissions and CH₄ flared (Table 8-7). The landfill gas recovered and used as fuel for co-generation units is reported under 1A1 Energy in accordance to the IPCC Good Practice Guide. The sum of landfill gas flared and landfill gas used in co-generation units is reported as being recovered in CRF-table 6.A,C.

Table 8-7 Activity data in 6A1: CH₄ direct emissions and CH₄ flared from 1990 to 2012 (source EMIS 2014/1A1a & 6A1 Kehrichtdeponien).

| Source / Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| 6A1 Managed Waste Disposal on Land | | | | | | | | | | | |
| CH ₄ direct emissions | Gg | 32.7 | 32.3 | 30.8 | 28.2 | 25.4 | 25.0 | 23.8 | 23.2 | 22.2 | 21.0 |
| CH ₄ flared | Gg | 4.2 | 4.2 | 4.3 | 4.3 | 4.2 | 4.1 | 4.0 | 3.9 | 3.7 | 3.6 |
| CH ₄ used in co-generation units (reported under 1A1a) | Gg | 4.9 | 5.7 | 7.6 | 10.4 | 12.6 | 12.1 | 12.1 | 11.5 | 11.3 | 11.4 |

| Source / Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| 6A1 Managed Waste Disposal on Land | | | | | | | | | | | |
| CH ₄ direct emissions | Gg | 19.0 | 18.0 | 17.2 | 15.0 | 15.0 | 14.1 | 13.5 | 12.4 | 11.1 | 10.1 |
| CH ₄ flared | Gg | 3.4 | 3.1 | 2.8 | 2.5 | 2.3 | 2.0 | 1.8 | 1.6 | 1.4 | 1.3 |
| CH ₄ used in co-generation units (reported under 1A1a) | Gg | 11.3 | 10.0 | 8.1 | 7.7 | 5.3 | 4.1 | 2.7 | 2.1 | 1.8 | 1.5 |

| Source / Parameter | Unit | 2010 | 2011 | 2012 |
|---|------|------|------|------|
| 6A1 Managed Waste Disposal on Land | | | | |
| CH ₄ direct emissions | Gg | 9.4 | 8.5 | 7.5 |
| CH ₄ flared | Gg | 1.2 | 1.0 | 0.9 |
| CH ₄ used in co-generation units (reported under 1A1a) | Gg | 1.0 | 0.9 | 0.9 |

The CH₄ generated in landfills decreased since 1990 due to the fact that waste quantities disposed of in landfills have been decreasing. Together with the relative increase of CH₄ recovery from 1990 until 2012 this is the reason for CH₄ emissions from the source category 6A being a key source regarding trend.

8.2.3 Uncertainties and Time-Series Consistency

Uncertainty in CH₄ emissions from Solid Waste disposal on land in 6A

Uncertainty of direct CH₄ emissions from sanitary landfills is estimated to be 58.3%.

For the amount of waste disposed of on a landfill an uncertainty of 30% is assumed. This is because most of the emissions in the nineties stem from waste disposed of in the eighties, when waste statistics were reported less accurately. An uncertainty of 50% is assumed for the emission factor (EMIS 2014/1A1a & 6A1 Kehrlichtdeponien).

Combined uncertainty of CO₂ emissions is estimated to be 40% in emissions estimates according to an uncertainty assessment based on expert judgments.

Qualitative estimate of uncertainties of non-key source emissions in 6A

A preliminary uncertainty assessment based on expert judgment results in medium confidence in emission estimates.

Consistency: The time series is consistent.

8.2.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

8.2.5 Source-Specific Recalculations

Value for 2011 of sewage gas production has changed in SFOE 2013a. The AD of 2011 (CH₄ direct emissions and CH₄ flared) have been recalculated.

8.2.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

8.3 Source Category 6B – Wastewater Handling

8.3.1 Source Category Description

Tier 1 Key category 6B

N₂O emissions from waste water handling (level).

Tier 2 Key category 6B

N₂O emissions from waste water handling (level and trend).

The source category 6B1 Industrial Waste Water comprises all emissions from liquid waste handling and sludge from industrial processes such as food processing, textiles, or pulp and

paper production. Food processing may result in effluents with a high load of organics. In order to reduce the load of polluted waste water (and to meet the regulatory standards as well as to reduce discharge fee) the effluent is pre-treated on site. The treatment is generally anaerobic, in order to use the biogas as source for heat and power generation. Currently, there are 22 industrial waste water pre-treatment plants. For emission calculations, industrial waste water is not identified as separate waste water stream, but joined to the domestic waste water treatment. Industrial waste water comprises, for example, metal-containing waste waters from electroplating plants, waste water from the food processing industry or waste water from car-washing places. Depending on the contaminants, an on-site pre-treatment is necessary so that the legal threshold values are met and that disruptions on the public sewage treatment plant can be precluded. After the pre-treatment process, the industrial wastewater is discharged in the sewage system and cleaned in a public wastewater treatment plant.

The source category 6B2 Domestic and Commercial Waste Water comprises all emissions from liquid waste handling and sludge from housing and commercial sources (including gray water and night soil). In Switzerland, MWWTP treat waste water from single cities or several cities/municipalities together. Waste water in general is treated in three steps: 1. Mechanical treatment, 2. Biological treatment, and 3. Chemical treatment. The treated waste water flows into a receiving system (lake, river or stream). As mentioned above, the pre-treated effluents from industries are also handled for final treatment in these waste water treatment plants. Switzerland's wastewater management infrastructure - comprising 849 sewage plants and 40'000-50'000 km of public sewers - is now practically complete (FOEN 2012g). In 2012, 285 waste water treatment plants use sewage gas for energy production.

Table 8-8 Specification of source category 6B Wastewater Handling. "EMIS 2014/..." refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 6B | Source | Specification | Data Source |
|-----|-------------------------------------|---|--|
| 6B1 | Industrial Waste Water | Emissions from handling of liquid wastes and sludge from industrial processes. | AD: EMIS 2014/6B1 Kläranlagen Industriell and SFOE 2013 EF: EMIS 2014/6B1 Kläranlagen Industriell |
| 6B2 | Domestic and Commercial Waste Water | Emissions from liquid waste handling and sludge from housing and commercial sources | AD: EMIS 2014/6B2 Kläranlagen Kommunal and SFOE 2013 EF: EMIS 2014/6B2 Kläranlagen Kommunal |
| 6B3 | Others | Not occurring in Switzerland | |

The emissions related to waste water treatment are included in various categories as illustrated in Figure 8-5 below. The system boundaries of category 6B contain all emissions from direct waste water handling including direct emissions of sewage gas (leakage), torching, combined heat and power production (CHP), furnaces (only heat production), emissions from the combustion of heating oil in boilers and engines, and natural upgrading into gas quality (intended for introduction into the natural gas network and/or use as fuel).




Figure 8-5 System boundaries of emissions related to wastewater treatment. CHP= Combined Heat and Power Production

Sewage sludge is usually dried in sewage plants in Switzerland before being disposed of. Emissions associated with sewage sludge drying are assumed to be negligible. The discharge of sewage sludge in agriculture has been phased out since 2003 and is generally forbidden since 2008, therefore this process is crossed out in the figure. The same applies to the disposal on land (6A). All sewage sludge is incinerated either in MSW incineration plants (1A1a), SS incineration plants (6C) or used as alternative fuel in the cement industry (1A2f).

8.3.2 Methodological Issues, Wastewater Handling (6B)

Methodology

For industrial waste water treatment (6B1) a country specific method is used. For domestic and commercial waste water treatment (6B2), a country specific method is used, with the exception of N_2O . The N_2O emissions are calculated based on the national protein consumption according to the IPCC default method. The GHG emissions are calculated by multiplying the number of inhabitants connected to waste water treatment plants by emission factors.

Emission Factors

Emission factors for CO_2 , CH_4 , N_2O , CO , NMVOC and SO_2 are country specific based on measurements and expert estimates, documented in the EMIS 2014/6B1 Industrielle Abwässer and EMIS 2014/6B2 Kommunale Kläranlagen. To calculate emission factors, the total sewage gas production for Switzerland is taken into account. It is assumed that 2% of the total amount of sewage gas is flared and 0.75% of the total amount is leaking. It is further assumed that 5% of the upgraded gas leaks as well. With these assumptions EF in kg/TJ are calculated. All emission factors have been converted and standardized into g/inhabitants.

In case of emissions related to industrial waste water treatment, total mass flows of emissions are divided by the population “artificially”, because activity data in our database correspond to the connected population.

Emission factors are adapted on a yearly basis due to respective changes in the amounts of total production of sewage gas produced, population growth and eventual changes in the fraction of inhabitants connected to waste water treatment plants.

EF for N₂O is derived based on the IPCC-default method (EMIS 2014 6B2 Kläranlage kommunal). The IPCC standard values were used for the percentage of nitrogen in proteins and the emission factor of N₂O in waste water. The annual protein consumption in Switzerland is taken from the statistics of the Secretariat of the Swiss Farmers Association (SBV 2013). As these numbers take into account swiss conditions, results seem to be more accurate for national conditions than by applying the FAOSTAT database.

The following formula taken from the revised 1996 IPCC GL is used (adjusted for per capita emissions):

$$\text{N}_2\text{O(s)} = \text{Protein} * \text{FracNPR} * \text{EF6}$$

According to the Swiss Farmer's Union, total protein consumption in Switzerland raised from 236'000 t in 1990 to 305'000 t in 2011 (SBV 2012). The value for 2012 has not been published yet and is therefore extrapolated by using the value of 2011 and the population in 2012. Protein consumption factors thus range around 34 kg/ inhabitant and year. Using an N fraction of 0.16 kg N per kg protein (FracNPR; IPCCdefault) an emission factor of around 97 g N₂O per inhabitant results for the year 2012. N₂O-emissions are thus only an approximation. Expert interviews reveal that currently there is no better data or more accurate values available for Switzerland¹⁷. Measurements of emissions have only been carried out for certain processes in specific waste water treatment plants and results are not transferable to other plants. There might be more specific values available in the future.

CH₄ emissions reported are due to flaring, leakage from sewage gas upgrading as well as leakage from piping systems. As mentioned before it is assumed that 5% of the upgraded gas is lost. A specific EF for the amount of upgraded biogas of 19'945 kg/TJ is assumed. This results in an EF for the amount of upgraded biogas of 19'945 kg/TJ x 0.05 = 997.25 kg/TJ. The total amount of biogas lost is allocated to the (connected) inhabitants, which results in an EF of 51.4 [g/connected inhabitant] or 49.8 [g/inhabitant] in 2012.

Although the emission factor for CH₄ in kg/ TJ has not changed since 2007, rather abrupt changes in EF [g/ inhabitant] can be observed since 2007. This is related to the changes in input mass flows, i.e. sewage gas produced in each year. Additionally, the amount of sewage gas fed into the natural gas pipes has increased significantly from 2007 to 2012 from 4.6 GWh in 2007 to 26.0 GWh in 2012 (SFOE 2013a). Thus, higher emissions are mainly related to the growing amounts of upgraded sewage gas.

The emission factors used in 6B1 and 6B2 are summarized in the following table.

Table 8-9 Emission Factors for 6B1 Industrial Waste Water and 6B2 Domestic and Commercial Waste Water in 2012.

| Source | CO ₂ biog. | N ₂ O | CH ₄ | NO _x | CO | NMVOC | SO ₂ |
|----------------------------|-----------------------|------------------|-----------------|-----------------|-----|-------|-----------------|
| | kg/inhabitant | | | g/inhabitant | | | |
| 6B1 Industrial Waste Water | 2.2 | NO | 5.5 | 2.5 | 3.6 | 0.07 | 0.23 |
| 6B2 Municipal Waste Water | 14.8 | 97 | 50 | 24 | 42 | 0.5 | 3 |

The activity data for N₂O emissions of municipal waste water is the total number of inhabitants (not connected inhabitants), in line with IPCC. For industrial waste water it is assumed that N₂O emissions are not occurring (EMIS 2014/6B1 Industrielle Abwässer).

¹⁷ Telephone interview Denise Fussen with Pascal Wunderlin, PhD EAWAG, 11.09.2012

In 2012, 96.8 % of inhabitants are connected to public waste water treatment. Emissions from waste water of the inhabitants not connected to public waste water treatment are not considered, as their contribution is of minor importance.

Several waste water treatment plants also accept co-substrates (organic wastes from the food processing industry) and add them to the digestion process. As they are rich in energy content a considerable part of the sewage gas stems therefrom. A part of the emissions are thus related to the addition of co-substrates.

Activity data

Activity data in 6B correspond to the number of connected inhabitants (except for N₂O as mentioned above). The number of inhabitants connected to the system (ICS) is calculated as a product of number of inhabitants and the service level. The latter is assumed to rise from 96.7% in 2007 to 97% in 2020 with interpolated values in between. Activity data for Domestic and Commercial Waste Water (6B2) are extracted from EMIS 2014/6B2 Kläranlage kommunal from SFOE 2013a. For 6B1 the same activity data were adopted.

Table 8-10 Activity data 6B2 Domestic and Commercial Waste Water; Population and fraction connected to waste water treatment plants.

| Source/Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 6B2 Domestic and Commercial Waste Water | | | | | | | | | | | |
| Population | inhabitants in 1000 | 6'796 | 6'880 | 6'943 | 6'989 | 7'037 | 7'081 | 7'105 | 7'113 | 7'132 | 7'167 |
| Fraction connected to waste water treatment plants | % | 90.0 | 91.0 | 91.5 | 92.0 | 93.0 | 93.5 | 94.0 | 94.5 | 95.0 | 95.3 |
| connected inhabitants | inhabitants in 1000 | 6'116 | 6'261 | 6'353 | 6'430 | 6'544 | 6'621 | 6'679 | 6'722 | 6'775 | 6'830 |

| Source/Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 6B2 Domestic and Commercial Waste Water | | | | | | | | | | | |
| Population | inhabitants in 1000 | 7'209 | 7'285 | 7'343 | 7'405 | 7'454 | 7'501 | 7'558 | 7'619 | 7'711 | 7'801 |
| Fraction connected to waste water treatment plants | % | 95.4 | 95.7 | 96.0 | 96.3 | 96.6 | 96.7 | 96.7 | 96.7 | 96.7 | 96.7 |
| connected inhabitants | inhabitants in 1000 | 6'877 | 6'972 | 7'049 | 7'131 | 7'201 | 7'253 | 7'309 | 7'368 | 7'458 | 7'547 |

| Source/Parameter | Unit | 2010 | 2011 | 2012 |
|--|---------------------|-------|-------|-------|
| 6B2 Domestic and Commercial Waste Water | | | | |
| Population | inhabitants in 1000 | 7'878 | 7'912 | 7'997 |
| Fraction connected to waste water treatment plants | % | 96.8 | 96.8 | 96.8 |
| connected inhabitants | inhabitants in 1000 | 7'623 | 7'658 | 7'742 |

8.3.3 Uncertainties and Time-Series Consistency

Uncertainty in N₂O emissions from 6B2

Activity data is highly reliable (estimated uncertainty 1.3%). The uncertainty for the emission factor is estimated to be 50%, according to EMIS 2014/6B1 Kläranlage Industriell.

Qualitative estimate of uncertainties of non-key category emissions in 6B

A preliminary uncertainty assessment based on expert judgment results in medium uncertainty of emissions estimates for CH₄ and low uncertainty for N₂O.

The time series is consistent.

8.3.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

8.3.5 Source-Specific Recalculations

Values of connected inhabitants (AD) have not been correctly interpolated from 2008-2011. Recalculations for those years were carried out.

The amount of sewage gas produced has been updated for the years 2007-2011 (SFOE 2013a). Updated values have been used in the model to calculate EF for all pollutants.

The Swiss Farmers Association (SBV) has changed its method of calculation for protein consumption for the years 2008 onwards. The EF of N₂O been recalculated for the years 2008-2011.

8.3.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

8.4 Source Category 6C – Waste Incineration

8.4.1 Source Category Description

Source category 6C Waste Incineration is **not a key category**.

There is a long tradition in Switzerland to incinerate waste. The waste heat generated during the incineration has to be recovered if technically and economically feasible. In accordance with the IPCC provisions (IPCC 1997c) emissions from the combustion of waste-to-energy activities are dealt with in 1A Fuel Combustion Activities.

Emissions from open burning of branches in agriculture and forestry have formerly (before Submission 2013) been reported in category 4F. However field burning of agricultural residues does not occur in Switzerland. Concerned waste fractions related to agriculture and forestry are burned only after they have been translocated from their place of origin. Therefore they are now reported under sector 6 Waste. Accordingly, the source category has been moved to 6C1 and is now reported under waste incineration as Burning of Residues in Agriculture and Forestry.

The following sources are included in source category 6C:

Table 8-11 Overview on waste incineration sources reported under 6C. "EMIS 2014/..." refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| Waste incineration | Specification | Data Source |
|---|--|--|
| Hospital waste incineration | Emissions from incinerating hospital waste in hospital incinerators | AD, EF: EMIS 2014/6C2 Spitalabfall-Verbrennung |
| Illegal waste incineration | Emissions from illegal incineration of municipal solid wastes at home Emissions from waste incineration at construction sites (open burning) | AD, EF: EMIS 2014/6C2 Abfallverbrennung illegal |
| Insulation material from cables | Emissions from incinerating cable insulation materials | AD, EF: EMIS 2014/6C2 Kabelabbrand |
| Sewage sludge | Emissions from sewage sludge incineration plants | AD, EF: EMIS 2014/6C2 Klärschlamm-Verbrennung |
| Crematoria | Emissions from the burning of bodies in crematoria | AD, EF: EMIS 2014/6C Krematorien |
| Burning of residues in agriculture and forestry | Emissions from burning of agricultural and silvicultural waste | AD, EF: EMIS 2014/6C1 Abfallverbrennung in der Land- und Forstwirtschaft |

8.4.2 Methodological Issues

Methodology

For the calculation of the greenhouse gas emissions a country specific Tier 2 method is used. In general, the GHG emissions are calculated by multiplying the waste quantity incinerated by emission factors. For crematoria, the GHG emissions are calculated by multiplying the number of cremations by emission factors.

For sewage sludge incineration plants the respective waste quantities are based on reliable statistical data (updated every two years until 2006). The emission factors are based on emission declarations from an incineration plant in 2002 that covered approx. one third of the Swiss capacities. Due to the lack of better or newer data these EF are kept constant since then and no increase in flue gas cleaning standards is assumed.

For hospital waste incineration, illegal waste incineration and incineration of insulation material, the waste quantities used are based on rough expert estimates.

The emissions of burning of residues in agriculture and forestry are calculated by multiplying the annual estimate of branches burned (in Gg of wood equivalent) by emission factors (IPCC default method).

Emission Factors

Emission factors for CO₂, CH₄, N₂O, CO, NMVOC and SO₂ are country specific based on measurements and expert estimates, documented in the EMIS 2014/6C database. The emission factors of burning of residues in agriculture and forestry are calculated based on EMEP/EEA 2013 except for CH₄ und N₂O which are based on EMEP/CORINAIR (EEA 2002), EMIS 2014/6C1 Abfallverbrennung in der Land- und Forstwirtschaft.

In the years with no specific data for activity data or emission factors the respective data are interpolated.

The following table presents the emission factors used in 6C:

Table 8-12 Emission Factors for 6C Waste Incineration in 2012 (source EMIS 2014/6C, EEA 2007 several EMIS comments see Table 8-11).

| 6C Waste Incineration | | | | | | | |
|--|-----------------------------------|------------------------------------|-----------------------------------|------------------------------------|------------------------|----------------------------|------------------------------------|
| Source | CO₂ t/t | CH₄ kg/t | N₂O g/t | NO_x kg/t | CO kg/t | NM VOC kg/t | SO₂ kg/t |
| Hospital waste incineration | 0.9 | 0 | 60 | 1.5 | 1.4 | 0.3 | 1.3 |
| Illegal waste incineration | 0.61 | 6 | 150 | 2.5 | 50.0 | 16 | 0.75 |
| Insulation material cables | 1.3 | 0 | 0 | 1.3 | 2.5 | 0.5 | 6 |
| Sewage sludge plants | 0 | 0.10 | 800 | 0.7 | 0.19 | 0.0050 | 0.47 |
| Burning of natural residues in agriculture | NA | 6.8 | 180 | 1.38 | 48.8 | 1.5 | 0.03 |
| Burning of natural residues in forestry | NA | 6.8 | 180 | 1.38 | 48.8 | 1.5 | 0.03 |
| Burning of natural residues in private households | NA | 6.8 | 180 | 1.38 | 48.8 | 1.5 | 0.03 |
| | CO₂ t/crem. | CH₄ kg/crem. | N₂O g/crem. | NO_x kg/crem. | CO kg/crem. | NM VOC kg/crem. | SO₂ kg/crem. |
| Crematoria | NA | 0 | 0 | 0.21 | 0.16 | 0.013 | 0 |

Additional information on the emission factor CO₂:

For all waste incineration options, the CO₂ emissions only from non-biodegradable waste are taken into account.

- Hospital waste incineration plants: Mainly waste of fossil origin. Default value for the CO₂ emission factor taken from CORINAIR (1992). Since 2002, no emissions from hospital waste incineration occur, as all special hospital waste incinerator plants have been closed and hospital waste is incinerated in municipal solid waste incineration plants (accounted for in 1A1).
- Illegal waste incineration: The main source of non-biodegradable CO₂ emissions is plastic. Until 2002 it is assumed that 40% of the waste mix is of fossil origin. In the FOCAWIN study by EMPA (Mohn 2011), the share of fossil matter in municipal waste in 2011 has been measured to be 47.8% on average. It is assumed that the same share applies for illegal waste burning. The value of 47.8% is implemented for the year 2011. Until 2002 the value is assumed to be 40%. In between 2002 and 2011 the values are linearly interpolated. The assumptions for the share of fossil/biogenic and C-content of waste are the same as for MSW incineration plants (EMIS 2014/6 C 2 (6 C c UNECE)_Abfallverbrennung illegal). According to EMIS 2014/ 1A1 a_Kehrichtverbrennungsanlagen, p. 5, the emission factors for CO₂ were estimated for 1990 and 2004 by the Waste Division of the FOEN.
- Insulation materials: The CO₂ emission factor is based on measurements of the flue gas treatment of a cable disassembling site where O₂ was measured in the flue gas. Assuming that the ratio of CO₂/O₂ is the same as in municipal solid waste incineration plants, a fraction of 7% of CO₂ results. Based on these assumptions, an EF of 1.3kg/cable can be derived. Since 1995, no emissions from incinerating cable insulation materials occur.
- Sewage sludge plants: Sewage sludge is biodegradable waste. Emission factor for CO₂ is 0. It is assumed that the share of fossil fuel used during the start-ups is negligible.

Additional information on other emission factors:

- Sewage sludge plants: Gradual technical improvements lead to reductions in NMVOC, CO, SO₂ and CH₄ emissions. Since 2002, emission factors are being kept constant because there is no newer data available (EMIS 2014 6C 2 Klärschlammverbrennung).
- Crematoria: NMVOC and CO emissions were reduced by technical improvements. Emission factors therefore depend on the number of technically improved crematoria.

The emission factors of 2011 were calculated by linearly extrapolating estimations for 2008 by taking into account an increase in the number of technically improved crematoria (EMIS 2014 6C Krematorien).

Activity Data

The activity data for Waste Incineration (6C) are the quantities of waste incinerated.

Table 8-13 Activity data for the different emission sources within source category 6C Waste Incineration.

| Source/Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Hospital Waste Incineration | Gg | 30.0 | 27.5 | 25.0 | 22.5 | 20.0 | 17.5 | 15.0 | 12.5 | 10.0 | 7.5 |
| Illegal waste | Gg | 32.3 | 31.7 | 31.0 | 29.7 | 27.3 | 26.2 | 25.0 | 24.6 | 24.2 | 25.1 |
| Insulation material cables | Gg | 7.5 | 6.0 | 4.5 | 3.0 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sewage sludge | Gg dry | 57.0 | 53.9 | 50.7 | 47.6 | 44.4 | 50.2 | 56.0 | 59.6 | 63.2 | 63.8 |
| Burning of natural residues in agriculture | Gg | 16.5 | 16.2 | 16.0 | 15.7 | 15.5 | 15.2 | 15.0 | 14.7 | 14.5 | 14.2 |
| Burning of natural residues in forestry | Gg | 28.8 | 28.0 | 27.1 | 26.2 | 25.4 | 24.5 | 23.6 | 22.8 | 21.9 | 21.0 |
| Burning of natural residues in private households | Gg | 6.1 | 5.8 | 5.6 | 5.3 | 5.1 | 4.9 | 4.6 | 4.4 | 4.1 | 3.9 |
| Total | Gg | 178.2 | 169.1 | 159.9 | 150.1 | 139.1 | 138.5 | 139.2 | 138.6 | 137.9 | 135.5 |
| Cremations | Numb. | 37'513 | 37'407 | 37'939 | 38'884 | 39'620 | 40'968 | 41'932 | 43'468 | 43'456 | 44'180 |

| Source/Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Hospital Waste Incineration | Gg | 5.0 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Illegal waste | Gg | 24.9 | 24.1 | 23.8 | 22.9 | 22.3 | 21.7 | 22.6 | 22.1 | 22.4 | 20.7 |
| Insulation material cables | Gg | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sewage sludge | Gg dry | 64.3 | 70.2 | 76.0 | 86.5 | 97.0 | 94.9 | 92.7 | 93.1 | 93.5 | 93.9 |
| Burning of natural residues in agriculture | Gg | 14.0 | 13.8 | 13.5 | 13.3 | 13.0 | 12.8 | 12.5 | 12.3 | 12.0 | 11.8 |
| Burning of natural residues in forestry | Gg | 20.2 | 19.3 | 18.4 | 17.6 | 16.7 | 15.9 | 15.0 | 14.1 | 13.3 | 12.4 |
| Burning of natural residues in private households | Gg | 3.6 | 3.4 | 3.2 | 2.9 | 2.7 | 2.4 | 2.2 | 1.9 | 1.7 | 1.5 |
| Total | Gg | 132.0 | 133.2 | 134.9 | 143.2 | 151.7 | 147.6 | 145.0 | 143.5 | 142.9 | 140.2 |
| Cremations | Numb. | 45'104 | 45'681 | 45'979 | 47'488 | 46'128 | 48'169 | 48'083 | 49'413 | 50'885 | 52'402 |

| Source/Parameter | Unit | 2010 | 2011 | 2012 |
|---|--------|--------|--------|--------|
| Hospital Waste Incineration | Gg | 0.0 | 0.0 | 0.0 |
| Illegal waste | Gg | 21.0 | 20.3 | 20.3 |
| Insulation material cables | Gg | 0.0 | 0.0 | 0.0 |
| Sewage sludge | Gg dry | 94.3 | 94.7 | 95.1 |
| Burning of natural residues in agriculture | Gg | 11.5 | 11.4 | 11.3 |
| Burning of natural residues in forestry | Gg | 11.5 | 11.4 | 11.3 |
| Burning of natural residues in private households | Gg | 1.2 | 1.2 | 1.2 |
| Total | Gg | 139.6 | 139.1 | 139.2 |
| Cremations | Numb. | 52'813 | 52'530 | 54'167 |

8.4.3 Uncertainties and Time-Series Consistency

Qualitative estimate of uncertainties of (non-key source) emissions in 6C

A preliminary uncertainty assessment based on expert judgment results in high uncertainty for CO₂ und CH₄ and low uncertainty for N₂O of emissions estimates.

The time series is consistent.

8.4.4 Source-Specific QA/QC and Verification

In addition to the general QA/QC measures described in Section 1.6 the activity data and emission factors were verified.

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

8.4.5 Source-Specific Recalculations

Following recalculation were carried out in 6C:

Crematoria: The Swiss Cremations Association (SVFB) has corrected AD of the year 2010. Recalculations have been made.

Sewage Sludge Incineration: Emission factors are now being kept constant since 2002 because there are no new data available. Before, a linear decrease until 2020 has been assumed. This led to recalculation from 2003 - 2011.

Sewage Sludge Incineration: AD between 2007 and 2011 have changed due to revised projections. The projections of emissions of air pollutants in Switzerland have been fully revised over the past year in order to provide consistency with the recently established energy scenarios of Energieperspektiven 2050 (Prognos 2012a).

Forestry and Agriculture: The process „Field Burning of Agricultural and silvicultural Residues“ has been remodeled. Previously only the open burning of branches was considered. The former process has been split up into three processes „Incineration of biomass, Agriculture“, „Incineration of biomass, Silviculture“ and „Incineration of biomass; Private gardens“. This led to recalculations of AD for the whole time series.

8.4.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

8.5 Source Category 6D – Other Waste

8.5.1 Source Category Description

Tier 1 Key category 6D

CH₄ from composting and digesting organic waste (trend)

Tier 2 Key Category 6D

CH₄ from composting and digesting organic waste (level and trend)

The source category 6D Other Waste comprises the GHG emissions from car shredding plants, and the process related GHG emissions from composting and from digesting organic waste.

Within the composting activity four types of composting means are distinguished, i.e. i) hall composting, ii) field edge composting, iii) box composting and iv) windrow composting. Composting covers the GHG emissions from centralized composting plants with a capacity of more than 100 tons of organic matter/year. Backyard composting is also common practice in Switzerland. It is assumed that the quantities treated in small composting facilities such as gardens, backyards etc., add up to 10% of those treated in industrial composting plants (EMIS 2014 6D Kompostierung Industrie and EMIS 2014 6D Kompostierung Haushalte).

The digestion of organic waste takes place under anaerobic conditions. The digested matter (solid left-overs after completion of a process of anaerobic microbial degradation of organic matter) is being composted. The biogas is used for combined heat and power generation or upgraded and used as fuel. In 6D Other Waste the emissions from the composting of the digested matter as well as the methane losses due to biogas up-grading are included. The emissions related to the use of biogas in co-generation plants and emissions from biogas-upgrading are reported under the sector 1 Energy.

Table 8-14 Specification of source category 6D Other Waste. "EMIS 2014/..." refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 6B | Source | Specification | Data Source |
|----|--------------------------|---|---|
| 6D | Car shredding plants | Emissions from car shredding plants | AD, EF: EMIS 2014/6D Shredder Anlagen |
| 6D | Composting and digesting | Process related emissions from composting and digesting organic waste | AD, EF: EMIS 2014/6D Kompostierung Industrie, EMIS 2014/6D Kompostierung Haushalte, EMIS 2014/1A1a und 6D Vergärung IG und EMIS 2014/1A1a und 6D Vergärung LW |

8.5.2 Methodological Issues

Methodology

For the emissions from car shredding a country specific method is used. The GHG emissions are calculated by multiplying the quantity of scrap by the emission factors. For all years the same constant emission factors have been applied.

For the emissions from composting a country specific method is used. The GHG emissions are calculated by multiplying the quantity of waste by the emission factors. For all years, the same constant emission factors have been applied.

For the emissions from digesting a country specific method is used. Digestion plants lead to GHG emissions from the storage of fermentable waste, losses due to leakages and diffusion, power water (storage of liquid fermented waste), rotting (storage of solid fermented waste) and flaring. The GHG emissions are calculated by multiplying the quantity of fermented waste by the emission factors. For all years, constant emission factors have been applied, except for CH₄. The methane emissions from biogas upgrading are calculated by multiplying the biogas quantity upgraded by the percentage of methane losses.

Because of the increase in composting and digesting organic waste the source category 6D Other Waste is a key source regarding trend.

Emission Factors

Emission factors for car shredding, composting and digestion are country specific based on measurements and expert estimates, documented in the EMIS 2014/6D database.

The following table presents the emission factors used in 6D:

Table 8-15 Emission Factors for 6D Others in 2012.

| Source | Unit | CH ₄ | N ₂ O | NO _x | CO | NM VOC | SO ₂ |
|--|------------------------------|-----------------|------------------|-----------------|-------|--------|-----------------|
| Shredding | g/t scrap | | | | 5 | 200 | |
| Composting | g/t composted waste | 5'000 | 70 | | | 1'700 | |
| Fermentation (industrial, storage) | g/t fermentable waste | 95 | 12 | | | 70 | |
| Fermentation (industrial, losses) | g/t biogas losses | 426'903 | | | | | |
| Fermentation (industrial, power water) | g/t fermented waste (liquid) | 1'896 | 2 | | | 400 | |
| Fermentation (industrial, rotting) | g/t fermented waste (solid) | 1'043 | 98 | | | 230 | |
| Fermentation (industrial, flaring) | g/t CH ₄ | 42 | | 4'066 | 2'054 | 82 | 616 |
| Fermentation (agricultural, losses) | g/t biogas losses | 426'903 | | | | | |
| Fermentation (agricultural, rotting) | g/t fermented waste (solid) | 1'100 | 98 | | | 230 | |
| Fermentation (agricultural, flaring) | g/t CH ₄ | 246 | | 4'066 | 2'054 | 82 | 616 |
| Biogas up-grade | g/GJ | 1075 | | | | | |

The fermentation process is split into different activities for industrial and agricultural biogas plants, respectively. These are following activities: storage of fermentable waste, losses due to leakages and diffusion, power water (storage of liquid fermented waste), rotting (storage of solid fermented waste) and flaring. For agricultural plants, reliable data on the emission factors exist for the storage of fermentable waste and the storage of the liquid fermented waste. For the purposes of a very rough initial estimate, the emission factors given here are taken from the corresponding subprocesses of the industrial and trades biogas installations. For each activity, new AD and EF have been reported, based on the newest data available. However, uncertainties are relatively high (EMIS 2014/1A1a und 6D Vergärung Industrie und Gewerbe und EMIS 2014/1A1a und 6D Vergärung Landwirtschaft).

The NMVOC emissions from car shredding stem from residues of fuels in the tanks of shredded cars (EMIS 2014/6D Shredder). For emission factors of NMVOC constant values are used since 2005 (EMIS 2014/6D Shredder Anlagen).

CH₄ emissions from biogas-upgrading occur due to leakage and are assumed to be 5 % of the total biogas production (value for 1990 – 2007). Since no data or better estimates are available the emission factor is being kept constant for the following years (until 2035).

Emissions from composting encompass CO₂, CH₄, NH₃, N₂O and NMVOC and are based on measured or estimated values reported in literature.

Activity data

Activity data for shredding and composting are extracted from EMIS 2014/6D Shredder Anlagen and Kompostierung Industrie. Activity data of fermentation are extracted from EMIS 2014/1A1a und 6D Vergärung IG und EMIS 2014/1A1a und 6D Vergärung LW.

Activity data for composting are based on periodically collected reliable statistical data, but latest available data is from the year 2007. Since then values have been interpolated. The quantities for backyard composting are estimated values, i.e. 10% of the amount of waste from composting plants.

Activity data of fermentation are based on information on the number of biogas plants (EMIS 2014/1A1a und 6D Vergärung IG und EMIS 2014/1A1a und 6D Vergärung LW) and encompass activity data of all relevant processes, e.g. storage of fermentable waste, losses due to leakages and diffusion, power water (storage of liquid fermented waste), rotting (storage of solid fermented waste) and flaring.

The biogas used as fuel for co-generation units is reported under 1A1 Energy in accordance to the IPCC Good Practice Guide.

Table 8-16 Activity data in 6D Other Waste.

| Source/Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|--|------|------|------|------|------|------|------|------|------|-------|-------|
| Shredding | Gg | 280 | 284 | 288 | 292 | 296 | 300 | 300 | 300 | 300 | 300 |
| Composting | Gg | 260 | 300 | 320 | 350 | 370 | 400 | 450 | 480 | 500 | 510 |
| Fermentation (ind., storage, fermentable waste) | Gg | 0 | 0 | 9 | 9 | 17 | 27 | 26 | 33 | 40 | 56 |
| Fermentation (ind., losses, biogas) | Gg | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.02 | 0.03 | 0.04 | 0.05 | 0.07 |
| Fermentation (ind., power water, fermented waste liquid) | Gg | 0 | 0 | 5 | 5 | 10 | 15 | 15 | 18 | 22 | 31 |
| Fermentation (ind., rotting, fermented waste solid) | Gg | 0 | 0 | 3 | 3 | 6 | 9 | 9 | 11 | 14 | 19 |
| Fermentation (ind., flaring, CH ₄) | Gg | 0.00 | 0.00 | 0.01 | 0.01 | 0.02 | 0.03 | 0.05 | 0.06 | 0.08 | 0.10 |
| Fermentation (agr., losses, biogas) | Gg | 0.28 | 0.28 | 0.27 | 0.26 | 0.25 | 0.23 | 0.22 | 0.21 | 0.22 | 0.25 |
| Fermentation (agr., rotting, fermented waste solid) | Gg | 6 | 6 | 6 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| Fermentation (agr., flaring, CH ₄) | Gg | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Biogas up-grade | GJ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3053 | 5'684 | 8'526 |

| Source/Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|--|------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Shredding | Gg | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 |
| Composting | Gg | 640 | 650 | 730 | 732 | 733 | 735 | 737 | 739 | 742 | 746 |
| Fermentation (ind., storage, fermentable waste) | Gg | 60 | 74 | 83 | 82 | 90 | 107 | 137 | 163 | 176 | 224 |
| Fermentation (ind., losses, biogas) | Gg | 0.07 | 0.09 | 0.10 | 0.10 | 0.10 | 0.13 | 0.16 | 0.19 | 0.20 | 0.25 |
| Fermentation (ind., power water, fermented waste liquid) | Gg | 33 | 41 | 46 | 46 | 50 | 60 | 76 | 91 | 98 | 125 |
| Fermentation (ind., rotting, fermented waste solid) | Gg | 20 | 25 | 28 | 28 | 31 | 37 | 47 | 56 | 60 | 77 |
| Fermentation (ind., flaring, CH ₄) | Gg | 0.10 | 0.12 | 0.15 | 0.14 | 0.15 | 0.17 | 0.23 | 0.27 | 0.30 | 0.39 |
| Fermentation (agr., losses, biogas) | Gg | 0.30 | 0.33 | 0.36 | 0.40 | 0.47 | 0.62 | 0.80 | 1.01 | 0.92 | 0.74 |
| Fermentation (agr., rotting, fermented waste solid) | Gg | 7 | 7 | 7 | 8 | 9 | 12 | 18 | 28 | 33 | 37 |
| Fermentation (agr., flaring, CH ₄) | Gg | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Biogas up-grade | GJ | 20'084 | 25'768 | 20'842 | 23'116 | 33'347 | 41'305 | 42'442 | 51'916 | 73'137 | 86'779 |

| Source/Parameter | Unit | 2010 | 2011 | 2012 |
|--|------|---------|---------|---------|
| Shredding | Gg | 300 | 300 | 300 |
| Composting | Gg | 750 | 753 | 757 |
| Fermentation (ind., storage, fermentable waste) | Gg | 307 | 371 | 507 |
| Fermentation (ind., losses, biogas) | Gg | 0.34 | 0.41 | 0.55 |
| Fermentation (ind., power water, fermented waste liquid) | Gg | 171 | 207 | 283 |
| Fermentation (ind., rotting, fermented waste solid) | Gg | 105 | 127 | 173 |
| Fermentation (ind., flaring, CH ₄) | Gg | 0.52 | 0.60 | 0.80 |
| Fermentation (agr., losses, biogas) | Gg | 0.57 | 0.62 | 0.72 |
| Fermentation (agr., rotting, fermented waste solid) | Gg | 45 | 49 | 59 |
| Fermentation (agr., flaring, CH ₄) | Gg | 0.12 | 0.13 | 0.16 |
| Biogas up-grade | GJ | 124'295 | 172'042 | 241'768 |

8.5.3 Uncertainties and Time-Series Consistency

Uncertainty in CH₄ emissions from composting and digestion 6D

The uncertainty of the CH₄ emissions in Category 6D from composting and digesting of organic waste is estimated to be 100% (expert estimate). The uncertainty of the related activity data is estimated to be 10% (expert estimate), because waste statistics are rather reliable.

Qualitative estimate of uncertainties of non-key source emissions in 6D

A preliminary uncertainty assessment based on expert judgment results in medium confidence in emissions estimates.

The time series is consistent.

8.5.4 Source-Specific QA/QC and Verification

The time series have been compared between the current and the previous submission as well as available energy data. All activity data, implied emission factors and emissions undergo the triple check:

- the results for 2012 are compared with the results 2011 within the current CRF
- the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
- the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013

8.5.5 Source-Specific Recalculations

Following recalculations were made in 6D:

Biogas upgrading: Emission factor of CH₄ had been wrong by a factor of 1000 and has been corrected. Recalculations were made from 1997, when biogas-upgrading activities started, until 2011.

Fermentation: Until submission 2013, AD for the years 2009-2011 have been extrapolated based on the values of the years 2007/08. Now actual data from SFOE 2013a have been used for estimating AD. Furthermore, activity data for the years before 2009 have been recalculated due to updated data in SFOE 2013a. This led to recalculation of AD for the whole time series 1990-2011 (agricultural biogas production) and the years 1999-2011 (industrial biogas production).

8.5.6 Source-Specific Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. In order to accomplish this transition, other projects are postponed in order to give first priority to changes related to the new reporting guidelines.

9 Other

9.1 Overview

9.1.1 Greenhouse Gas Emissions

Within the sector 7 Other emissions from various sources are considered:

- Fire damage estates,
- Fire damage motor vehicles




Figure 9-1 Switzerland's greenhouse gas emissions in the sector 7 Other 1990–2012.

Table 9-1 Trend of total GHG emissions from 7 Other in Switzerland 1990-2012.

| Gas | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 11.0 | 11.2 | 11.3 | 11.5 | 11.7 | 11.9 | 12.1 | 12.3 | 12.5 | 12.7 |
| CH ₄ | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| N ₂ O | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| Sum | 12.1 | 12.3 | 12.5 | 12.7 | 12.9 | 13.1 | 13.3 | 13.5 | 13.7 | 13.9 |

| Gas | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|---------------------------------|------|------|------|------|------|------|------|------|------|------|
| CO ₂ equivalent (Gg) | | | | | | | | | | |
| CO ₂ | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 | 12.9 |
| CH ₄ | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| N ₂ O | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| Sum | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 |

| Gas | 2010 | 2011 | 2012 |
|--------------------------|------|------|------|
| CO ₂ eq. (Gg) | | | |
| CO ₂ | 12.9 | 12.9 | 12.9 |
| CH ₄ | 0.6 | 0.6 | 0.6 |
| N ₂ O | 0.6 | 0.6 | 0.6 |
| Sum | 14.1 | 14.1 | 14.1 |

In the sector Other a total of 14.1 Gg CO₂ equivalents was emitted in the year 2012. 93% of the emissions stem from “fire damage estates”, the rest from “fire damage motor vehicles”. The total greenhouse gas emissions of this sector show an increase of 16% from 1990 until 2012.

9.2 Source Category – Other non-specified

9.2.1 Source Category Description

Tier 1 and 2 key category 7

Source category 7 Other is **not a key category**.

The sources reported in 7 Other are depicted in Table 9-2.

Table 9-2 Specification of source category 7 Other N-specified. EMIS 2014/... refers to the internal documentation of the emission database and contains further references regarding specific AD and/or EF.

| 7 | Source | Specification | Data Source |
|----|----------------------------|---|---|
| 7D | Fire damage estates | Emissions from fires in buildings. | EMIS 2014/7D “Brand- und Feuerschäden Immobilien” |
| 7D | Fire damage motor vehicles | Emissions from fires in motor vehicles. | EMIS 2014/7D “Brand- und Feuerschäden Motorfahrzeuge” |

9.2.2 Methodological Issues: Fire damage estates and motor vehicles

Methodology

a) Fire damage estate

Based on average damage sums from insurances between 1992 and 2001, the average damage sum per fire case is estimated to be 15'600 CHF representing 780 kg of flammable material per fire case. Further assuming that not the whole amount burns down due to the intervention of the fire brigade, an amount of 400kg of burnt material per fire case is estimated. On average between 1992 and 2001, 20'650 cases of fire incidents happened each year. For emission calculation, a constant number of yearly 20'000 fire cases is assumed (EMIS 2014/7D “Brand- und Feuerschäden Immobilien”).

b) Fire damage motor vehicles

Based on data from a Swiss insurance company with 25% market share in 2002, the number of reported vehicle fires was extrapolated to 100%. Based on this estimate and the total vehicle number of Switzerland it was estimated that one fire case per 790 vehicles occurs per year and this was assumed to remain constant from 1990-2012. Multiplied with the actual vehicle number, the number of burnt vehicles in Switzerland per year is obtained (EMIS 2014/7D “Brand- und Feuerschäden Motorfahrzeuge”).

Emission Factors (Fire damages)

Fire damages in estates: Emission factors for CO₂, CO, NO_x and SO₂ are country specific based on measurements and expert estimates originally completed for illegal waste

incineration. It is assumed that emissions are similar to emissions of fire damage in estates (EMIS 2014/7D "Brand- und Feuerschäden Immobilien").

The fraction between fossil and biogenic CO₂ emissions is assumed to remain constant since 2000 with 80% being fossil and 20% biogenic CO₂ emissions. Before 2000, it is assumed that the fraction of fossil CO₂ emissions from burnt goods has been increasing linearly from 20% in 1950 to 80% in 2000.

Fire damages in motor vehicles: Emission factors for CO₂, CO, NO_x and SO₂ are country specific values based on measurements and expert estimates originally gained from the combustion of cables, documented in EMIS 2014/7D "Brand- und Feuerschäden Motorfahrzeuge".

Emissions for CH₄ from fire damage in motor vehicles are reported as well, while N₂O emissions have not been estimated for this source.

Table 9-3 Emission Factors for fire damages in 2012 (EMIS 2014/7D).

| Source 7 Other | CO ₂ biogenic | CO ₂ fossil | CH ₄ | N ₂ O | NO _x | CO | NM VOC | SO ₂ |
|----------------------------|-----------------------------|---------------------------|-----------------|------------------|-----------------|-------|--------|-----------------|
| t / t burned good | | | | | | | | |
| Fire damage estates | 0.40 | 1.5 | 0.003 | 0.00025 | 0.0020 | 0.100 | 0.016 | 0.001 |
| Fire damage motor vehicles | NO | 1.5 | 0.005 | NE | 0.0013 | 0.002 | 0.002 | 0.005 |

Activity data (Fire damages)

Activity data is the weight of burnt goods, calculated according to the following rule of proportion: 400 kg of burnt goods per incidence of fire cases in estates (EMIS 2014/7D "Brand- und Feuerschäden Immobilien") and 100 kg of burnt goods per incidence of burnt vehicles (EMIS 2014/7D "Brand- und Feuerschäden Motorfahrzeuge"). Activity data for estates is not estimated on a year to year basis but is assumed to be at a constant 8 Gg for the whole time period since 1990.

Table 9-4 Activity data: Burnt goods from 1990 to 2012 (source EMIS 2014/7D).

| Source / Parameter | Unit | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----------------------------|------|------|------|------|------|------|------|------|------|------|------|
| 7 Burnt goods | | | | | | | | | | | |
| Fire damage estates | Gg | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 |
| Fire damage motor vehicles | Gg | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.6 |

| Source / Parameter | Unit | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 |
|----------------------------|------|------|------|------|------|------|------|------|------|------|------|
| 7 Burnt goods | | | | | | | | | | | |
| Fire damage estates | Gg | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 | 8.0 |
| Fire damage motor vehicles | Gg | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |

| Source / Parameter | Unit | 2010 | 2011 | 2012 |
|----------------------------|------|------|------|------|
| 7 Burnt goods | | | | |
| Fire damage estates | Gg | 8.0 | 8.0 | 8.0 |
| Fire damage motor vehicles | Gg | 0.6 | 0.6 | 0.6 |

9.2.3 Uncertainties and Time Series consistency

Uncertainty of CO₂, CH₄ and N₂O emissions is estimated to be high (according to Table 1-14).

9.2.4 Source-Specific QA/QC and Verification

In addition to the general QA/QC measures described in Section 1.6 subsequent source-specific activities have been carried out:

- Cross check of activity data in EMIS comments.
- Verification of NMVOC emission factor for car shredding. The time series have been compared between the current and the previous submission. All activity data, implied emission factors and emissions undergo the triple check.
 - the results for 2012 are compared with the results 2011 within the current CRF.
 - the CRF-tables for the year 2011 are compared between the current CRF-tables and the CRF-tables of submission 2013.
 - the CRF-tables for the base year 1990 are compared between the current CRF-tables and the CRF-tables of submission 2013.

9.2.5 Source-Specific Recalculations

7D Fire Damages Motor Vehicles: AD decreased in 2003-2015 due to revised projections in 2015 (linear interpolation between last set value in the year 2002 and projections for 2015).

9.2.6 Source-Specific Planned Improvements

There are no source-specific planned improvements in this category.

10 Recalculations

10.1 Explanations and Justifications for Recalculation

10.1.1 GHG Inventory

The Inventory Development Plan (IDP, see in FOEN 2014a) is regularly updated, mainly based on the various “Reports of the individual review of the greenhouse gas inventory of Switzerland” (e.g. UNFCCC 2010, 2011, 2012, 2013) and the outcome of domestic reviews. The IDP represents the main instrument for continuous improvement of the Swiss GHG inventory in subsequent inventory cycles. It includes suggestions and recommendations for recalculations that have an impact on emission levels in the corresponding sectors.

The processing of the expert review team’s recommendations in the course of inventory preparation and compilation led to several recalculations (see also Table 1-12). Further recalculations had to be carried out due to improvements in some sectors. The details are explained below. An extensive list with all detailed recalculations and specifics of the recalculations is compiled by the EMIS experts and available to the reviewers (in German/French only).

1 Energy

- 1A: Activity data of all fuels of the overall time series have been recalculated based on the new data from SFOE 2013. This includes data on 2011 for fossil fuels and waste, data for biomass on 1997, 2009 – 2011 and data of other fuels 1990 – 2011.
- 1A: Activity data of wood consumption have been recalculated for the overall time series based on the new data from SFOE 2013b.
- 1A: SO_x emission factor value of gas oil has been updated for 2010 based on sulphur analyses of the gas oil for the year 2010 (Directorate General of Customs) resulting in a revised value for 2011 as well.
- 1A: CO₂ emission factor for bituminous coal has been corrected from 94t CO₂/TJ to 92.7 t CO₂/TJ for the entire time series.
- 1A1a: Activity data of biogas production from waste dumpsites have been recalculated for 2011 based on new values in the Swiss statistics of renewable energies SFOE 2013a.
- 1A1a: Emission factor of N₂O for waste incineration has been updated for the entire time series based on a new study realized by EMPA (2011).
- 1A1a: Emission factor of CH₄ for biomass fermentation has been corrected for the entire time series.
- 1A1a / 6D: Activity data for fermentation of biogenic waste in industrial and trades biogas installations and in agricultural biogas installations has been updated for the years 1999 to 2011 based on new data from the Swiss statistics of renewable energies SFOE 2013a.
- 1A1a: N₂O default emission factor from the IPCC guidelines has been introduced for the whole time series for fermentation of biogenic waste in industrial and trades biogas installations and in agricultural biogas installations.
- 1A1a: Activity data regarding bituminous coal has been corrected over the whole time series as it was not consistent with other years.

- 1A1a: Activity data regarding gas oil has been corrected over the whole time series based on data from the Swiss overall energy statistics (SFOE 2013).
- 1A1a: Activity data regarding other fuels has been corrected for 2007 and 2008 based on correction of the energy content of waste.
- 1A1b: Activity data for 2011 has been corrected for boilers refinery, residual fuel oil as there was a mistake in the database in Submission 2013.
- 1A1b: CO₂ Emission factor for 1990 to 2011 has been updated based on measurements that led to an average emission factor of 59'800 g/GJ compared to the previous emission factor of 59'300 g/GJ that was based on expert judgement.
- 1A1c: Reporting of the biogenic CO₂ and precursor emissions from the charcoal production has been shifted from source category 2D3 to 1A1c. (Please note that the reporting of the CH₄ emissions from the charcoal production has already been shifted from 2D3 to 1A1c within the resubmission of Switzerland's Greenhouse Gas Inventory 1990–2011(FOEN2013g).)
- 1A1c: Activity data from 2004 to 2012 has been corrected based on corrected data from small producers of charcoal.
- 1A1c: Emission factors for charcoal production have been corrected over the whole timeseries. The emission factors are newly based on Revised 1996 IPCC Guidelines for CO, NMVOC, CH₄, NO_x and USEPA (1995) for CO₂.
- 1A2: N₂O emission factor of liquefied petroleum gas has been changed for the whole time series from 0.6 g/GJ to the IPCC 2006 emission factor of 0.1 g/GJ. In previous submissions, the emission factor for gas oil was used for liquefied petroleum gas because they have been reported jointly. This has been corrected in the present submission.
- 1A2: Since SO_x emission factor values for 2010 and 2011 of gas oil have been updated based on sulphur analyses of gas oil for the year 2010 (Directorate General of Customs) also the SO_x emission factor values of liquefied petroleum gas for 2010 and 2011 have been revised as for all air pollutants the same EF are assumed as for gas oil.
- 1A2a: Activity data for bituminous coal in iron foundries has been revised due to corrected production shares of cupola and electric furnaces in iron foundries for 2010 and 2011. This change resulted in new activity data of bituminous coal in heat furnances for the same years.
- 1A2c: Activity data of steam production from the cracker-by-products has been updated for 1990 - 1999 based on new net calorific values. For calculation of the light virgin naphtha consumed as cracker feedstock the so far used net calorific value of gasoline has been replaced by the value for naphtha according to the 2006 IPCC guidelines resulting in revised activity data for 1990-1999.
- 1A2c: Activity data for liquefied petroleum gas has been updated for 1990-1999 and 2011 based on new activity data from the cracker process and resulting steam production.
- 1A2fi: Activity data of one of the two glass wool production plants have been revised for 1991-2004 based on effective production data for 1996-2004 resulting in revised gas consumption.
- 1A2fi: Activity data of all fuels for 1991-2011 have been updated based on new production data from brick and tile and glasswool production.

- 1A2fi: Activity data from gas oil and natural gas have been revised for 1990-2011 by the subtraction of nonroad from "Heizkessel GLD, HEL" and "Heizkessel GLD, Gas" instead of "Industrie Heizkessel weitere, HEL" and "Industrie Heizkessel weitere, Gas" within the Energy model.
- 1A2fi: Newly also CH₄ emissions of the cement production are reported with a CH₄ emission factor of 5 g/t cement from 1990 to 1995 and 4 g/t cement from 2002 onwards.
- 1A2fi: Activity data of brick and tile production has been updated for 2001-2006 using effective production data instead of interpolated data. In addition, also so far interpolated fuel consumptions for these three plants have been replaced by effective values for 2000 and 2007-2011 resulting in overall revised fossil and biogenic fuel consumptions for 1991-2011 and 2000-2011.
- 1A2fi: Activity data of biogas from wastewater treatment plants has been updated for 2008, 2009 and 2011 based on recalculations in SFOE 2013a.
- 1A2fii: Diesel and gasoline consumption is based on INFRAS (2008) and on an update carried out in 2013 based on the latest numbers on growth of population and economy (Prognos 2012a, Keller/INFRAS 2013). The consumption has been recalculated accordingly. Numbers from 2005 onwards are affected.
- 1A3ai: The kerosene consumption (AD) for international flights (bunker fuels) has been recalculated in the Swiss energy statistics (SFOE 2012) for the year 2011. Therefore, all gases and pollutants have been recalculated for 2011 as well.
- 1A3b: The entire time series has been recalculated based on the latest numbers on growth of population and economy (Prognos 2012a, ARE 2012). Vehicle kilometres from 1993 are slightly lower in total; fleet compositions have changed, with slight impacts on implied emission factors; fuel consumption in tank tourism has been recalculated; the modelled share of biofuels has been reduced to be consistent with real-world developments. The overall impacts of these recalculations on emissions are low.
- 1A3c, 1A3d, 1A4aii, 1A4bii, 1A4cii, 1A5b (mobile off-road machinery): The activity data have so far been taken from INFRAS (2008). For this submission, the latest numbers on growth of population and economy (Prognos 2012a, Keller/INFRAS 2013) have been integrated in the off-road model. This leads to an increase of the fuel consumption from 2005 onwards.
- 1A3dii: The activity data of the international marine bunkers has been updated for all pollutants in 2011.
- 1A3dii: The emission factors have been recalculated for the year 2011 due to interpolation of the emission factors between 2010 and 2015 for the gases CH₄ and N₂O. The EF of 2015 has not been changed but the interpolation was by mistake not applied in former submissions.
- 1A3di: Recalculations due to new activity data in 2004, 2008 and 2011 have been made which also affects the "tank tourism" of diesel fuel in 1A3b.
- 1A4a/1A4b: Activity data for engines and gasturbines in 1A4 households and services have been updated for 2011 based on updated statistical data.
- 1A4a: Activity data for natural gas in the commercial and institutional sector has been updated for 1990, 1995, 1997, 1998 and 2011 based on recalculations in SFOE 2013.
- 1A4a: Activity data for natural gas has been updated based on a recalculation of SFOE 2013 for the year 2011.

- 1A4a: Activity data for gas oil and natural gas consumption has been updated for the overall time series based on changes in the energy model resulting from changes in the non-road transport model for boats and natural gas consumption in industry.
- 1A4b: Activity data for natural gas in the residential sector has been updated for 1990, 1995, 1997, 1998 and 2011 based on recalculations in SFOE 2013.
- 1A4b: Activity data for gas oil has been updated for the whole time series based on recalculations in the energy consumption.
- 1A4b: CO and SO₂ Emission factors for gas oil and natural gas of 2011 have been corrected in the energy model.
- 1A4b: Reporting of charcoal production has been shifted to sector 1A4b. Before, it was reported under 2D3.
- 1A4b: A new process for bonfires has been introduced in this submission.
- 1B2a v: Emission factor for CO₂ and NMVOC were corrected to exclude emissions from Liechtenstein that overestimated emissions by 0.5%.
- 1B2b: As recommended in ARR 2012 and 2013 Switzerland has included in its inventory the emissions from the only small plant for natural gas production (Finsterwald) from 1985-1994. The default emission factors from IPCC Guidelines 2006 is used.
- 1B2b: Activity data from 1990-1997 has been updated based on data from SFOE 2013.
- 1B2b: A new model for emissions in swiss gas transport system based on a new study realized by Quantis has been developed.

2 Industrial Processes

- 2A1: A calculation error has been corrected resulting in new EFs for NO_x, CO, NMVOC and SO₂.
- 2A3 Brick and tile production: Interpolated activity data have been replaced by effective production data resulting in updated AD for 2001-2006. Revision of CO₂ EF for 1990-2011.
- 2A5: Activity data has been revised for 2008-2011 due to updated projection values.
- 2A7 Glass wool production: Revision of activity data and cullet ratios resulting in revised EF values for CO₂ for 1991-2004.
- 2B2: Revision of NO_x EF values for the entire time series.
- 2B5 Acetic acid production: Revision of activity data from 1991-2011. In addition also revision of the EF values for CH₄, CO and NMVOC and introduction of an EF for CO₂ for the entire time series.
- 2B5 Sulphuric acid production: Updating of activity data for 2009-2011. Revision of EF values for SO₂ for the entire time series.
- 2C1 Iron foundries: Revision of activity data in iron foundries for 2010 and 2011. The EF value for NMVOC for 2012 has been updated resulting in revised interpolated values 1991-2011 as well.
- 2C1 Steel production: Revision of EF values for CO₂ for the entire time series.
- 2C5e Battery recycling: Activity data for 2011 has been revised due to corrected data from the plant operator.

- 2D2 Food and Drink: Activity data of meat smokehouses has been updated for the years 2007-2011.
- 2F1: Improvement of model calculations of stock for all gases from the Refrigeration and Air-conditioning source categories for the entire time series (recharge of equipment considering minimal technical charge and related frequency of service).
- 2F1: Bus air-conditioning added to the calculation of mobile air-conditioning and causes recalculations in the entire time series for HFC134a.
- 2F1: In Transport Refrigeration export of retiring equipment is now included in model calculations of trucks and the lifetime of railway is elevated. This affects data in the years 2000-2011.
- 2F4: The use of spray products consider now earlier aerosol loss of production to avoid 1% in double counting of aerosol emissions (HFC134a and HFC152a) in the years 1998-2011.
- 2G Claus units in refineries: Activity data of the years 1990–1997 has been revised due to recalculations in the Swiss overall energy statistics.

3 Solvent and Other Product Use

- 3A1: Activity data for paint applications in households has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.
- 3B2: Activity data and EF value for dry cleaning have been updated for 2012 resulting as well in revised interpolated values for 2007-2011 and 1991-2011, respectively.
- 3C: Activity data for paint and ink manufacturing has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.
- 3C: Activity data for manufacturing of rubber has been updated for 2011 and EF value has been updated for 2012 resulting as well in revised interpolated values for 2008-2010 and 1998-2011, respectively.
- 3C: Activity data for manufacturing of polyester has been updated for 2010 and 2011 and EF values have been updated for 2010 and 2012 resulting as well in revised interpolated values for 2008-2009 and 2008, 2009 and 2011, respectively.
- 3C: Activity data for manufacturing of polystyrene has been updated for 2011 resulting in revised interpolated values for 2008-2010 as well.
- 3C: Activity data for manufacturing of PVC has been updated for 2010 and 2011 and EF values have been updated for 2004 and 2012 resulting as well in revised interpolated values for 2009 and 1991-2003 and 2005- 2011, respectively.
- 3D5: Activity data for print industry has been revised due to updated projection for 2015. This has led to a revision of the AD for 2011.
- 3D5: Activity data for impregnating of glass wool have been revised for 1991-2004 based on effective production data for 1996-2004.
- 3D5: Activity data for production and use of tobacco products has been updated for the years 2005-2011.
- 3D5: Activity data for use of concrete additives has been revised and updated for 1990, 1998, 2001 and 2008-2011, respectively. This has resulted in revised interpolated values in between. EF values have been corrected to interpolated values and updated for 1999-2006 and 2008-2011, respectively.
- 3D5: Activity data for car underbody sealant has been revised and updated for 1990, 1998 and 2012, respectively, resulting in revised interpolated values in between. EF

values have been corrected to interpolated values and updated for 1999-2003 and 2012, respectively, resulting in revised interpolated values for 2005-2011 as well.

- 3D5: Activity data and EF value for de-icing of airplanes and other de-icing have been updated for 2012 resulting in revised interpolated values for 2008-2011 as well.
- 3D5: Activity data and EF values for use of cooling lubricants have been updated for 2008-2011 and 2012, respectively, resulting as well in revised interpolated values from 2005 onwards.
- 3D5: Activity data for use lubricants has been updated for 2008-2011 resulting in revised interpolated values for 2005-2007 as well. EF values have been corrected to interpolated values and updated for 1999-2003 and 2012, respectively resulting in revised interpolated values for 2008-2011 as well.

4 Agriculture

- 4A,B: New more precise livestock population data have been used for the years 1994 and 2006. Previously only rounded values have been available. The effects on overall emissions is negligible.
- 4A,B: Preliminary estimates for energy requirements for non cattle populations for the years 2010 and 2011 have been revised. The effect of the recalculation on overall greenhouse gas emissions is considered negligible.
- 4A,B: Milk yield of mature dairy cattle in the year 2011 has been slightly revised due to an update of the provisional number from the Swiss Farmers Union.
- 4B,D: The nitrogen excretion rate of mature dairy cattle of the year 2011 has been revised in order to be consistent with the AGRAMMON model (Kupper et al. 2013). The reduced nitrogen excretion rate resulted in an overall emission reduction (including source category 4B and 4D) of less than 10 Gg CO₂ equivalent.
- 4D: The area of cultivated organic soils has been revised due to new projections in the LULUCF-sector. The mean change of overall greenhouse gas emissions is below 0.02 Gg CO₂ equivalent.
- 4D: Activity data for compost for the years 2008-2011 has been revised due to an error correction. New emission estimates are lower by approximately 7-8 Gg CO₂ equivalent.
- 4D: The ammonia emission factor for recycling fertilizers (liquid digestate) has been adjusted for the years 2008 - 2011 due to the increasing use of trailing hoses during land application as fertilizer. Impact on FracGASF and on overall emissions is negligible.
- 4D: A general recalculation for the year 2011 has been carried out due to some updates of crop yield data from the Swiss Farmers Union (SBV 2012). The respective changes are only of minor importance for total emission estimates.

5 Land Use, Land-Use change and Forestry and KP

- 5: The completion of the AREA surveys in 2013 (see Chapter 7.2.7, SFSO 2013) led to a recalculation in category 5 LULUCF.
- 5A1 and 5A2: The application of the gain-loss approach and the stock-change approach with regard to the calculation of carbon stock changes following land-use changes has been modified (see Chapter 7.1.3.2 and Chapter 7.3.7). This led to a large reduction of carbon sinks on land converted to forest land.

- 5A: New values of stocks, gains and losses of living biomass were derived from most recent NFI4a+ data (Thürig 2014). Implied emission factors for productive forests in the period 2006-2012 were affected (see Chapter 7.3.7).
- 5A: The Yasso07 values 1990-2012 for annual net changes in the dead wood pool, in the LFH litter pool and in mineral soils were updated (Didion et al. 2013). See Chapter 7.3.7.
- 5(II)D: Emissions of CH₄ are reported for flooded lands/reservoirs (was NO in former submissions). See Chapter 7.6.7.
- 5(V)A: For wildfires on forest land, the emission factor for CH₄ was revised. See Chapter 7.3.7.
- 5(V)C: Emissions of CH₄ and N₂O are reported due to wildfires on grassland (was NO in former submissions). See Chapter 7.5.7.

6 Waste

- 6A: Value for 2011 of sewage gas production has changed in SFOE 2013a. The AD of 2011 (CH₄ direct emissions and CH₄ flared) have been recalculated.
- 6B2 domestic waste water: Values of connected inhabitants (AD) have not been correctly interpolated from 2008-2011. Recalculations for those years were carried out.
- 6B2 domestic waste water: The amount of sewage gas produced has been updated for the years 2007-2011 (SFOE 2013a). Updated values have been used in the model to calculate EF for all pollutants.
- 6B2 domestic waste water: The Swiss Farmers Association (SBV) has changed its method of calculation for protein consumption for the years 2008 onwards. The EF of N₂O been recalculated for the years 2008-2011.
- 6C Crematoria: The Swiss Cremations Association (SVFB) has corrected AD of the year 2010. Recalculations have been made.
- 6C Sewage Sludge Incineration: Emission factors are now being kept constant since 2002 because there are no new data available, instead of a linear decrease until 2020. This led to recalculation of emissions from 2003-2011.
- 6C Sewage Sludge Incineration: AD between 2007 and 2011 have changed due to revised projections.
- 6C1 Forestry and Agriculture: The process „Field Burning of Agricultural and silvicultural Residues“ has been remodeled. Before, only the open burning of branches was considered. The former has been split up in three processes „Incineration of biomass, Agriculture“, „Incineration of biomass, Silviculture“ and „Incineration of biomass; Private gardens“. This led to recalculations of AD for the whole time series.
- 6D biogas upgrading: Emission factor of CH₄ had been wrong by a factor of 1000 and has been corrected. Recalculations were made from 1997, when biogas-upgrading activities started, until 2011.
- 6D Fermentation: Until submission 2013, AD for the years 2009-2011 have been extrapolated based on the values of the years 2007/08. Now actual data from SFOE 2013a have been used for estimating AD. Furthermore, activity data for the years before 2009 have been recalculated due to updated data in SFOE 2013a. This led to recalculation of AD for the whole time series 1990-2011 (agricultural biogas production) and the years 1999-2011 (industrial biogas production).

7 Other

- 7D: AD of Fire Damages Motor Vehicles decreased slightly in 2003-2015 due to revised projections in 2015 (linear interpolation between last set value in 2002 and 2015).

10.1.2 KP- LULUCF Inventory

A recalculation of the years 2008, 2009, 2010 and 2011 was carried out. The methodological improvements affect the activities reported under KP Art. 3.3 as well as under KP Art. 3.4. The improvements are described in detail in Chapter 7.3.7 (Recalculations 5A LULUCF Forest Land) and in Chapter 11.3.1.4 (Kyoto specific recalculations).

10.2 Implications for Emission Levels 1990 and 2011

10.2.1 GHG Inventory

Table 10-1 shows the recalculation results for the base year 1990. It results in a decrease of the total emissions in CO₂ equivalents (without emissions/removals from CO₂ from LULUCF) of 159.55 Gg CO₂ eq. This corresponds to a decrease of the latest submission compared to the previous submission (2012,v2.1) by 0.30% of the national total. If the LULUCF sector is included, there is a increase of 1074.77 Gg CO₂ eq (2.15%) due to substantial recalculations in the LULUCF sector.

Table 10-1 Overview of implications of recalculations on 1990 data. Emissions are shown before the recalculation according to the previous submission in 2013 "Prev." (FOEN 2013g) and after the recalculation according to the present submission "Latest". The differences "Differ." are defined as latest minus previous submission.

| Recalculation | CO ₂ | | | CH ₄ | | | N ₂ O | | | Sum (CO ₂ , CH ₄ and N ₂ O) | | |
|------------------------------------|---------------------------------|--------|---------|-----------------|---------|---------|------------------|---------|---------|--|--------|----------|
| | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. |
| Emissions for 1990 | | | | | | | | | | | | |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | | | | | | | | CO ₂ equivalent (Gg) | | |
| 1 Energy | 41'178 | 41'199 | 20.54 | 621.6 | 502.1 | -119.49 | 283.2 | 287.6 | 4.38 | 42'083 | 41'989 | -94.58 |
| 2 Ind. Processes (without F-gases) | 3'059 | 3'006 | -53.26 | 9.6 | 1.5 | -8.09 | 68.1 | 68.1 | 0.00 | 3'137 | 3'076 | -61.35 |
| 3 Solvent and Other Product Use | 360 | 360 | 0.06 | | | | 110.1 | 110.1 | 0.00 | 470 | 470 | 0.06 |
| 4 Agriculture | | | | 3'307.1 | 3'307.1 | 0.00 | 2'785.0 | 2'785.0 | 0.01 | 6'092 | 6'092 | 0.01 |
| 5 LULUCF | -3'174 | -1'962 | 1'212.0 | 8.2 | 30.3 | 22.14 | 10.4 | 10.5 | 0.19 | -3'156 | -1'921 | 1'234.32 |
| 6 Waste | 63 | 63 | 0.00 | 737.0 | 734.3 | -2.66 | 210.6 | 209.6 | -1.04 | 1'011 | 1'007 | -3.70 |
| 7 Other | 11 | 11 | 0.00 | 0.6 | 0.6 | 0.00 | 0.6 | 0.6 | 0.00 | 12 | 12 | 0.00 |
| Sum (without F-gases) | 41'498 | 42'677 | 1'179.3 | 4'684 | 4'576 | -108.10 | 3'468 | 3'472 | 3.53 | 49'650 | 50'725 | 1'074.77 |

| Recalculation | HFC | | | PFC | | | SF ₆ | | | Sum (synthetic gases) | | |
|------------------------------------|---------------------------------|--------|---------|-------|--------|---------|-----------------|--------|---------|---------------------------------|--------|---------|
| | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. |
| Emissions for 1990 | | | | | | | | | | | | |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | | | | | | | | CO ₂ equivalent (Gg) | | |
| 2 Ind. Processes (only syn. gases) | 0.02 | 0.02 | 0.00 | 100.2 | 100.2 | 0.00 | 143.6 | 143.6 | 0.00 | 243.85 | 243.85 | 0.00 |

| Recalculation | Sum (all gases) | | |
|---|---------------------------------|---------|----------|
| | Prev. | Latest | Differ. |
| Emissions for 1990 | | | |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | |
| Total CO ₂ eq Em. with LULUCF | 49'894 | 50'969 | 1'074.77 |
| | 100% | 102.15% | 2.15% |
| Total CO ₂ eq Em. without LULUCF | 53'049 | 52'890 | -159.55 |
| | 100% | 99.70% | -0.30% |

For 2011, the recalculation results in a decrease of the total emissions in CO₂ equivalents (without emissions/removals from LULUCF) of 190.21 Gg CO₂ eq. This corresponds to an decrease of 0.38% of the national total in the latest submission compared to the previous submission. If the LULUCF sector is included, a decrease of 1323.31 Gg CO₂ eq. (2.83%) results due to major recalculations in the LULUCF sector.

Table 10-2 Overview of implications of recalculations on 2011 data. Emissions are shown before the recalculation according to the previous submission in 2013 "Prev." (FOEN 2013) and after the recalculation according to the present submission "Latest". The differences "Differ." are defined as latest minus previous submission.

| Recalculation Emissions for 2011 | CO ₂ | | | CH ₄ | | | N ₂ O | | | Sum (CO ₂ , CH ₄ and N ₂ O) | | |
|-------------------------------------|---------------------------------|--------|----------|-----------------|--------|---------|------------------|--------|---------|--|--------|----------|
| | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | | | | | | | | CO ₂ equivalent (Gg) | | |
| 1 Energy | 39'466 | 39'479 | 12.46 | 257.0 | 253.9 | -3.16 | 266.3 | 212.4 | -53.91 | 39'990 | 39'945 | -44.62 |
| 2 Ind. Processes (without F-gases) | 2'332 | 2'186 | -145.81 | 8.6 | 2.3 | -6.28 | 54.1 | 54.1 | 0.00 | 2'394 | 2'242 | -152.08 |
| 3 Solvent and Other Product Use | 155 | 158 | 2.45 | | | | 44.1 | 44.1 | 0.00 | 199 | 202 | 2.45 |
| 4 Agriculture | | | | 3'159 | 3'155 | -4.16 | 2'445 | 2'417 | -27.77 | 5'604 | 5'572 | -31.93 |
| 5 LULUCF | -3'417 | -1'914 | 1'502.48 | 1.2 | 12.2 | 10.99 | 4.4 | 4.4 | 0.05 | -3'411 | -1'897 | 1'513.52 |
| 6 Waste | 12 | 12 | 0.00 | 309.2 | 299.4 | -9.83 | 265.4 | 286.7 | 21.26 | 587 | 598 | 11.43 |
| 7 Other | 13 | 13 | -0.11 | 0.6 | 0.6 | -0.01 | 0.6 | 0.6 | 0.00 | 14 | 14 | -0.12 |
| Sum (without F-gases) | 38'562 | 39'934 | 1'371.46 | 3'736 | 3'723 | -12.44 | 3'080 | 3'019 | -60.38 | 45'377 | 46'676 | 1'298.64 |

| Recalculation Emissions for 2011 | HFC | | | PFC | | | SF ₆ | | | Sum (synthetic gases) | | |
|-------------------------------------|---------------------------------|----------|---------|-------|--------|---------|-----------------|--------|---------|---------------------------------|----------|---------|
| | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. | Prev. | Latest | Differ. |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | | | | | | | | CO ₂ equivalent (Gg) | | |
| 2 Ind. Processes (only syn. gases) | 1'171.45 | 1'195.50 | 24.05 | 39.4 | 40.0 | 0.62 | 164.4 | 164.4 | 0.00 | 1'375.18 | 1'399.85 | 24.67 |

| Recalculation Emissions for 2011 | Sum (all gases) | | |
|---|---------------------------------|---------|----------|
| | Prev. | Latest | Differ. |
| Source and Sink Categories | CO ₂ equivalent (Gg) | | |
| Total CO ₂ eq Em. with LULUCF | 46'752 | 48'076 | 1'323.31 |
| | 100% | 102.83% | 2.83% |
| Total CO ₂ eq Em. without LULUCF | 50'163 | 49'973 | -190.21 |
| | 100% | 99.62% | -0.38% |

10.2.2 KP- LULUCF Inventory

The methodological improvements are described in Chapter 11.3.1.4. (Kyoto specific recalculations) and Chapter 7.3.7 (Recalculations LULUCF Forest Land).

10.3 Implications for Emissions Trends, including Time Series Consistency

10.3.1 GHG Inventory

Due to recalculations, the emission trend 1990–2011 reported in the present 2014 submission (FOEN 2014) has slightly changed. Compared to 1990, 2011 emissions (national total without emissions/removals from LULUCF) showed a decrease of 5.44% before recalculation (previous submission). After recalculation, the decrease turns out to be higher with a change 1990–2011 of 5.51% (latest submission).

Table 10-3 Change of the emission trend 1990–2011 due to recalculation. “Previous” refers to data reported in FOEN (2013g), whereas “latest” refers to the present submission.

| Recalculation | 1990 | | 2011 | | change 2011/1990 | |
|--------------------|-------------------------|--------|----------|--------|------------------|--------|
| Submission | previous | latest | previous | latest | previous | latest |
| Unit | CO ₂ eq (Gg) | | | | % | |
| Total excl. LULUCF | 53'049 | 52'890 | 50'163 | 49'973 | -5.44% | -5.51% |

All-time series in the present submission are consistent.

10.3.2 KP- LULUCF Inventory

As for KP-LULUCF 2008-2012 data is submitted, recalculations were done for 2008, 2009, 2010 and 2011 data and there are no implications for emission trends.

10.4 Recalculations, Including in Response to the Review Process, and Planned Improvements to the Inventory

10.4.1 Recalculations GHG Inventory

Many recalculations have been carried out in response to recommendations proposed in review reports. All relevant recalculations are listed in Chpt. 10.1.1. Further information on improvements due to the ERT recommendations are found in Table 1-12.

10.4.2 Recalculations KP-LULUCF Inventory

For recalculations see Chapter 10.1.2. Further information on improvements due to the ERT recommendations are found in Table 1-12. The methodological improvements are described in detail in Chapter 11.3.1.4. (Kyoto specific improvements) and Chapter 7.3.7 (Improvements LULUCF Forest Land).

10.4.3 Planned Improvements

In the coming year, the transition to the new reporting guidelines represents the largest improvement for the entire inventory. All methods will be adapted to the 2006 IPCC Guidelines (IPCC 2006). Within this general recalculation a number of optional country specific methods will be explored and eventually implemented. Other projects are eventually postponed in order to give first priority to changes related to the new reporting guidelines.

PART 2

11 KP-LULUCF

Switzerland has chosen to account annually for emissions and removals from the LULUCF sector (FOEN 2006h, Sect. G). In addition to the mandatory submission of the inventory years 2008, 2009, 2010, 2011 and 2012, data for the years 1999-2007 are available and shown in Switzerland's NIR on a voluntary basis. Switzerland has elected to account for Forest Management under the voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol (FOEN 2006h, Sect. F). Switzerland applies the condition of "direct human-induced" in relation to Afforestation and Deforestation very strictly for both activities (see Chapter 11.1.3, FOEN 2010d, FOEN 2010h). Table 11-1 shows the activity coverage and the carbon pools reported for the mandatory activities under Article 3, paragraph 3 and for Forest Management under paragraph 4 of the Kyoto Protocol. The areas and change in areas between the previous and the current inventory year are shown in Table 11-2. Table 11-3 summarizes the results of the KCA for LULUCF activities under the Kyoto Protocol.

Table 11-1: NIR1 – Summary-Table.

TABLE NIR 1. SUMMARY TABLE
Activity coverage and other information relating to activities under Article 3.3 and elected activities under Article 3.4

| Activity | Change in carbon pool reported ⁽¹⁾ | | | | Greenhouse gas sources reported ⁽²⁾ | | | | | | |
|------------------------|--|----------------------|--------|-----------|--|------------------------------|---|--|-----------------|--------------------------------|-----------------|
| | Above-ground biomass | Below-ground biomass | Litter | Dead wood | Soil | Fertilization ⁽³⁾ | Drainage of soils under forest management | Disturbance associated with land-use conversion to croplands | Liming | Biomass burning ⁽⁴⁾ | |
| | | | | | | N ₂ O | N ₂ O | N ₂ O | CO ₂ | | CO ₂ |
| Article 3.3 activities | Afforestation and Reforestation Deforestation | R | R | NR | R | NO | | | NO | NO | NO |
| | | R | R | R | R | | | R | NO | NO | NO |
| | | R | R | R | R | NO | NR | | NO | R,IE | R |
| Article 3.4 activities | Cropland Management Grazing Land Management Revegetation | NA | NA | NA | NA | | | NA | NA | NA | NA |
| | | NA | NA | NA | NA | | | | NA | NA | NA |
| | | NA | NA | NA | NA | | | | NA | NA | NA |
| | | NA | NA | NA | NA | | | | NA | NA | NA |

⁽¹⁾ Indicate R (reported), NR (not reported), IE (included elsewhere) or NO (not occurring), for each relevant activity under Article 3.3 or elected activity under Article 3.4. If changes in a carbon pool are not reported, it must be demonstrated in the NIR that this pool is not a net source of greenhouse gases. Indicate NA (not applicable) for each activity that is not elected under Article 3.4. Explanation about the use of notation keys should be provided in the text.

⁽²⁾ Indicate R (reported), NE (not estimated), IE (included elsewhere) or NO (not occurring) for greenhouse gas sources reported, for each relevant activity under Article 3.3 or elected activity under Article 3.4. Indicate NA (not applicable) for each activity that is not elected under Article 3.4. Explanation about the use of notation keys should be provided in the text.

⁽³⁾ N₂O emissions from fertilization for Cropland Management, Grazing Land Management and Revegetation should be reported in the Agriculture sector. If a Party is not able to separate fertilizer applied to Forest Land from Agriculture, it may report all N₂O emissions from fertilization in the Agriculture sector.

⁽⁴⁾ If CO₂ emissions from biomass burning are not already included under changes in carbon stocks, they should be reported under biomass burning; this also includes the carbon component of CH₄. Parties that include CO₂ emissions from biomass burning in their carbon stock change estimates should report IE (included elsewhere).

Table NIR 1.1 Additional information
Selection of parameters for defining "Forest" under the Kyoto Protocol

| Parameter | Range | Selected value |
|---------------------|-------------|----------------|
| Minimum land area | 0.05 - 1 ha | 0.06 |
| Minimum crown cover | 10 - 30 % | 20.00 |
| Minimum height | 2 - 5 m | 3.00 |

Table 11-2: NIR 2 – Land Transition Matrix Inventory Year 2012: the total Swiss area amounts too 4'128.42 kha.
A time series of the total values for Article 3.3 and Article 3.4 activities is displayed in Table 11-6.

Table NIR 2. LAND TRANSITION MATRIX
Areas and changes in areas between the previous and the current inventory year ^{(1), (2), (3)}

| From previous inventory year / To current inventory year | | Article 3.3 activities | | Article 3.4 activities | | | Other ⁽⁵⁾ | Total area at the beginning of the current inventory year ⁽⁶⁾ |
|--|---|---------------------------------|---------------|--------------------------------|----------------------------------|--------------------------------------|---------------------------|--|
| | | Afforestation and Reforestation | Deforestation | Forest Management (if elected) | Cropland Management (if elected) | Grazing Land Management (if elected) | Revegetation (if elected) | |
| Article 3.3 activities | Afforestation and Reforestation | 2.44 | NO | | | | | 2.44 |
| | Deforestation | | 6.81 | | | | | 6.81 |
| | Forest Management ⁽⁴⁾ (if elected) | | 0.31 | 1'231.22 | | | | 1'231.53 |
| | Cropland Management ⁽⁴⁾ (if elected) | NA | NA | | NA | NA | NA | NA |
| | Grazing Land Management ⁽⁴⁾ (if elected) | NA | NA | | NA | NA | NA | NA |
| Article 3.4 activities | Revegetation ⁽⁴⁾ (if elected) | NA | | | NA | NA | NA | NA |
| | Other ⁽⁵⁾ | 0.05 | NA | 1.44 | NA | NA | NA | 2'887.64 |
| Total area at the end of the current inventory year | | 2.49 | 7.13 | 1'232.66 | NA | NA | NA | 4'128.42 |

⁽¹⁾ This table should be used to report land area and changes in land area subject to the various activities in the inventory year. For each activity it should be used to report area change between the previous year and the current inventory year. For example, the total area of land subject to Forest Management in the year preceding the inventory year, and which was deforested in the inventory year, should be reported in the cell in column of Deforestation and in the row of Forest Management.

⁽²⁾ Some of the transitions in the matrix are not possible and the cells concerned have been shaded.

⁽³⁾ In accordance with section 4.2.3.2 of the IPCC good practice guidance for LULUCF, the value of the reported area subject to the various activities under Article 3.3 and 3.4 for the inventory year should be that on 31 December of that year.

⁽⁴⁾ Lands subject to Cropland Management, Grazing Land Management or Revegetation which, after 2008, are subject to activities other than those under Article 3.3 and 3.4, should still be tracked and reported under Cropland Management, Grazing Land Management or Revegetation, respectively.

⁽⁵⁾ "Other" includes the total area of the country that has not been reported under an Article 3.3 or an elected Article 3.4 activity.

⁽⁶⁾ The value in the cell of row "Total area at the end of the current inventory year" corresponds to the total land area of a country and is constant for all years.

Table 11-3 NIR 3 – Summary Overview for Key Categories for LULUCF Activities under the KP in 2012. A detailed description of the KCA for Article 3.3 and 3.4 activities is given in Chapter 11.6.1.

| KEY CATEGORIES OF EMISSIONS AND REMOVALS | GAS | CRITERIA USED FOR KEY CATEGORY IDENTIFICATION | | | COMMENTS ⁽³⁾ |
|--|-----|---|---|------------------------------|--|
| | | Associated category in UNFCCC inventory ⁽¹⁾ is key (indicate which category) | Category contribution is greater than the smallest category considered key in the UNFCCC inventory ^{(1), (4)} (including LULUCF) | Other ⁽²⁾ | |
| Specify key categories according to the national level of disaggregation used ⁽¹⁾ | | | | | |
| Forest Management | CO2 | Forest land remaining forest land | Yes | Since the total Swiss forest | Associated category in UNFCCC inventory is KC level and KC trend (Tier 2; 2012). |
| Afforestation and Reforestation | CO2 | Conversion to forest land | No | Natural forest regeneration | |
| Deforestation | CO2 | Conversion to settlements | Yes | see NIR Chapter 11.6.1 | Associated category in UNFCCC inventory is KC level and KC trend (Tier 2; 2012). |

⁽¹⁾ See section 5.4 of the IPCC good practice guidance for LULUCF.

⁽²⁾ This should include qualitative consideration as per section 5.4.3 of the IPCC good practice guidance for LULUCF or any other criteria.

⁽³⁾ Describe the criteria identifying the category as key.

⁽⁴⁾ If the emissions or removals of the category exceed the emissions of the smallest category identified as key in the UNFCCC inventory (including LULUCF), Parties should indicate YES. If not, Parties should indicate NO.

An overview of net CO₂ eq emissions and removals of activities under Article 3, paragraph 3 and Forest Management under Article 3, paragraph 4 of the Kyoto Protocol is shown in Figure 11-1 and Table 11-4. Differences in the annual emissions from Deforestation can mainly be attributed to the changes in the area of Deforestations (see Table 11-6; Figure 11-2). Year-to-year differences in removals from Afforestations are mainly due to changes in the afforested area (see Table 11-6). Another relevant factor is the application of a logistical growth curve for Afforestations younger than 20 years (see Chapter 11.3.1.1; Figure 11-2; see Table 7-25). Fluctuations in the contribution of Forest Management can mainly be explained by changes in the losses of living biomass, dead wood and litter, whereas fluctuations in the area of managed forest are relatively small (see Table 11-6). In 2001 and 2002, Forest Management was a small source of CO₂ eq due to the damage caused by the storm Lothar.




Figure 11-1 CO₂ eq emissions (positive sign) and removals (negative sign) from Afforestation and Deforestation under Article 3, paragraph 3 and Forest Management under Article 3, paragraph 4, 1999-2012.

Table 11-4 Overview on net CO₂ equivalent emissions (positive sign) and removals (negative sign) for activities under Article 3, paragraph 3 and Forest Management under Article 3, paragraph 4 of the Kyoto Protocol, 1999-2012.

| Greenhouse gas source and sink activities | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|---|---|----------------|---------------|---------------|-----------------|-----------------|-----------------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| A. Article 3.3 activities | 216.97 | 216.83 | 215.84 | 214.33 | 212.51 | 210.95 | 174.16 |
| A.1. Afforestation and Reforestation | -6.09 | -6.82 | -7.80 | -9.17 | -10.83 | -12.60 | -14.86 |
| A.1.1. Units of land not harvested since the beginning of the commitment period | -6.09 | -6.82 | -7.80 | -9.17 | -10.83 | -12.60 | -14.86 |
| A.1.2. Units of land harvested since the beginning of the commitment period | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| A.2. Deforestation | 223.06 | 223.66 | 223.63 | 223.50 | 223.34 | 223.55 | 189.02 |
| B. Article 3.4 activities | -3330.01 | -395.31 | 166.35 | 219.39 | -2434.18 | -2810.38 | -2663.60 |
| B.1. Forest Management incl. biomass burning | -3330.01 | -395.31 | 166.35 | 219.39 | -2434.18 | -2810.38 | -2663.60 |
| gains above ground living biomass | -9617.06 | -9624.02 | -9630.98 | -9637.94 | -9644.90 | -9651.86 | -9662.29 |
| gains below ground living biomass | -2839.47 | -2841.80 | -2844.13 | -2846.46 | -2848.79 | -2851.12 | -2856.62 |
| losses above ground living biomass | 7758.43 | 10051.40 | 10371.01 | 10243.29 | 8273.08 | 8004.19 | 8365.83 |
| losses below ground living biomass | 2252.11 | 2847.55 | 2925.45 | 2894.29 | 2392.50 | 2319.62 | 2410.48 |
| changes litter | -335.15 | -271.92 | -166.56 | -34.20 | -201.14 | -218.05 | -467.97 |
| changes dead wood | -555.68 | -563.91 | -495.37 | -423.47 | -435.68 | -418.28 | -458.88 |
| changes soil C min. soils | -2.28 | -2.95 | -3.46 | -3.79 | -4.07 | -4.50 | -5.04 |
| changes soil C org. soils | 8.68 | 8.68 | 8.68 | 8.69 | 8.69 | 8.69 | 8.70 |
| sum forest management excl. biomass burning | -3330.43 | -396.98 | 164.63 | 200.40 | -2460.30 | -2811.31 | -2665.78 |
| biomass burning | 0.42 | 1.67 | 1.71 | 18.99 | 26.12 | 0.93 | 2.18 |
| B.2. Cropland Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.3. Grazing Land Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.4. Revegetation (if elected) | NA | NA | NA | NA | NA | NA | NA |

| Greenhouse gas source and sink activities | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 |
|---|---|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| | Net CO ₂ equivalent emissions/removals (Gg CO ₂ eq) | | | | | | |
| A. Article 3.3 activities | 147.05 | 117.04 | 81.74 | 162.22 | 197.10 | 201.52 | 204.74 |
| A.1. Afforestation and Reforestation | -17.18 | -20.14 | -22.17 | -24.33 | -23.34 | -19.62 | -17.13 |
| A.1.1. Units of land not harvested since the beginning of the commitment period | -17.18 | -20.14 | -22.17 | -24.33 | -23.00 | -18.89 | -15.89 |
| A.1.2. Units of land harvested since the beginning of the commitment period | 0.00 | 0.00 | 0.00 | 0.00 | -0.34 | -0.73 | -1.25 |
| A.2. Deforestation | 164.23 | 137.19 | 103.91 | 186.56 | 220.45 | 221.14 | 221.87 |
| B. Article 3.4 activities | -2804.38 | -1901.42 | -1202.77 | -1419.28 | -2020.23 | -2063.62 | -2236.38 |
| B.1. Forest Management incl. biomass burning | -2804.38 | -1901.42 | -1202.77 | -1419.28 | -2020.23 | -2063.62 | -2236.38 |
| gains above ground living biomass | -9844.35 | -10013.44 | -10184.68 | -10196.75 | -10201.54 | -10206.14 | -10210.74 |
| gains below ground living biomass | -2925.08 | -2989.60 | -3055.14 | -3060.06 | -3062.02 | -3063.86 | -3065.70 |
| losses above ground living biomass | 8106.69 | 8426.45 | 8467.86 | 8105.81 | 7882.81 | 7839.72 | 7757.11 |
| losses below ground living biomass | 2288.18 | 2387.54 | 2417.30 | 2332.60 | 2285.48 | 2283.19 | 2264.70 |
| changes litter | -222.74 | 71.47 | 383.78 | 354.85 | 29.07 | 115.00 | 134.28 |
| changes dead wood | -214.87 | 202.23 | 761.98 | 1037.66 | 1039.35 | 954.69 | 876.33 |
| changes soil C min. soils | -5.60 | -5.63 | -5.05 | -4.07 | -3.27 | -2.69 | -2.12 |
| changes soil C org. soils | 8.71 | 8.72 | 8.73 | 8.74 | 8.74 | 8.74 | 8.74 |
| sum forest management excl. Biomass burning | -2809.05 | -1912.26 | -1205.22 | -1421.23 | -2021.39 | -2071.35 | -2237.40 |
| biomass burning | 4.68 | 10.84 | 2.45 | 1.95 | 1.16 | 7.73 | 1.02 |
| B.2. Cropland Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.3. Grazing Land Management (if elected) | NA | NA | NA | NA | NA | NA | NA |
| B.4. Revegetation (if elected) | NA | NA | NA | NA | NA | NA | NA |

The KP-CRF-table "Information table on accounting for activities under Article 3, paragraph 3 and 4 of the Kyoto Protocol" gives an overview of the CO₂ eq emissions and removals from Afforestation and Deforestation under Article 3, paragraph 3 and Forest Management under Article 3, paragraph 4 and also provides information on the extent to which GHG removals by sinks offsets the debit incurred under Article 3.3.

- In 2008 Forest Management in Switzerland caused removals of -1202.77 Gg CO₂ eq. The debit incurred from activities under Article 3.3 is 81.74 Gg CO₂ eq.
- In 2009 Forest Management in Switzerland caused removals of -1419.28 Gg CO₂ eq. The debit incurred from activities under Article 3.3 is 162.22 Gg CO₂ eq.
- In 2010 Forest Management in Switzerland caused removals of -2020.23 Gg CO₂ eq. The debit incurred from activities under Article 3.3 is 197.10 Gg CO₂ eq.
- In 2011 Forest Management in Switzerland caused removals of -2063.62 Gg CO₂ eq. The debit incurred from activities under Article 3.3 is 201.52 Gg CO₂ eq.
- In 2012 Forest Management in Switzerland caused removals of -2236.38 Gg CO₂ eq. The debit incurred from activities under Article 3.3 is 204.74 Gg CO₂ eq.

11.1 General Information

The inventory datasets on which the calculations are based (Swiss Land Use Statistics AREA and National Forest Inventory NFI) are described in Chapters 7.2.2 and 7.3.4.1, respectively.

Methodological issues and assumptions concerning the calculation of activity data and emission factors used for the reporting under Article 3, paragraphs 3 and 4 of the Kyoto Protocol, follow the IPCC good practice guidance and are described in Chapter 7.3.4 and in FOEN (2014c).

11.1.1 Definition of Forest and any other Criteria

The forest definition used under the Kyoto Protocol is defined in Switzerland's Initial Report (FOEN 2006h, Sect. E and Chapter 7.3.1 in this submission). Forest is defined as a minimum area of land of 0.0625 ha with crown cover of at least 20% and a minimum width of 25 m. The minimum height of the dominant trees must be 3 m or have the potential to reach 3 m at maturity in situ. The selected values are also listed in KP LULUCF Table NIR1 (see Table 11-1).

Some source categories were explicitly excluded from the category "Forest Land", although they may partly fulfil the requirements of the Swiss forest definition used under the Kyoto Protocol (see Chapter 7.2.2.2, Table 7-6). Those are:

- Vineyards, Low-Stem Orchards, Tree nurseries, Copses and Orchards in the land-use category "Grassland";
- Cemeteries and public parks in the land-use category "Settlements".

11.1.2 Elected Activities under Article 3, Paragraph 4, of the Kyoto Protocol

Switzerland has decided to account for Forest Management under the elective voluntary activities of Article 3, paragraph 4 of the Kyoto Protocol (FOEN 2006h, Sect. F). In accordance with the Annex to Decision 16/CMP.1, credits from Forest Management are capped in the first commitment period. For Switzerland the cap amounts to 1.83 Mt CO₂ yr⁻¹ (0.5 Mt C yr⁻¹), or 9.15 Mt CO₂ for the whole commitment period 2008-2012.

11.1.3 Description of how the Definitions of each Activity under Article 3.3 and each elected Activity under Article 3.4 have been implemented and applied consistently over Time

The Swiss definitions of Afforestation, Deforestation and Forest Management are published in Switzerland's Initial Report (see FOEN 2006h, Sect. E and F). Switzerland applies the condition of "direct human-induced" in relation to Afforestation and Deforestation very strictly for both activities (see FOEN 2010d, FOEN 2010h).

Afforestation

Afforestation is the conversion to forest of an area not fulfilling the definition of forest for a period of at least 50 years if the definition of forest in terms of minimum area (625 m²) is fulfilled, and the conversion is a direct human-induced activity.

Natural forest regeneration due to abandonment of land, mainly occurring in the Alpine area, is not considered to be a direct human-induced activity. Only Afforestations which can clearly be attributed as direct human-induced from aerial photographs (SFSO 2013; see also Chapter 7.2) are considered as Afforestation. Examples of direct human-induced Afforestations are shown in FOEN (2010h).

Deforestation

Deforestation is the permanent conversion of areas fulfilling the definition of forest in terms of minimum forest area (625 m²) to areas not fulfilling the definition of forest as a consequence of direct human influence.

Temporary removals of tree stand (e.g. for the construction of high-tension lines and pipelines) are not reported as Deforestation under the Kyoto Protocol because in those cases the forest stand has to be re-established. In the NFI methodology (Brändli 2010: 91) "forest aisles" under high-tension are explicitly classified as forests. These forest aisles underlie however a specific management, i.e. maximum tree height is limited to a certain height. The NFI dataset thus covers such areas with a specific Forest Management practice.

After approximately 12 years (see Chapter 7.2.2.1) it is possible to check if deforestations or other land-use changes have been correctly classified. Sigmaplan (2012a) screened the classification of all land-use changes classified as Deforestation under the Kyoto Protocol. They found that 86% of all these Kyoto Deforestations are still deforested after 20 years, whereas 14% of these Kyoto Deforestations were in fact removals of crown coverage limited in time and should be classified as "management interventions" rather than as real land-use changes. As no reclassification was done the area of Deforestations reported under KP Art. 3.3 is in fact a slight overestimation. Accordingly, emissions are overestimated since implied emission factors for Deforestations are higher than for Forest Management (see Table 5(KP-I)A.2 for Deforestations and Table 5(KP-I)B.1 for Forest Management).

Reforestation

Reforestation does not occur in Switzerland (FOEN 2006h, Sect. E; see also Chapter 11.4.1).

Forest management

Forest Management includes all activities serving the purpose of fulfilling the Federal Law on Forests (Swiss Confederation 1991, Art. 1c), i.e. the obligation to conserve forests and to ensure forest functions – such as wood production, protection against natural hazards,

preservation of biodiversity, purification of drinking water and maintenance of recreational value – in a sustainable manner.

11.1.4 Description of Precedence Conditions and/or Hierarchy among 3.4. Activities and how they have been consistently applied in determining how Land was classified.

Since Switzerland only elected Forest Management from the elective activities of Article 3, paragraph 4 of the Kyoto Protocol, the hierarchy among 3.4 activities does not affect Swiss reporting.

11.2 Land-related Information

11.2.1 Spatial Assessment Unit used for determining the Area of the units of Land

The spatial assessment unit for the submission of the KP LULUCF tables covers the entire territory of Switzerland, i.e. 4'128.42 kha (see).

All activity data for reporting the activities under the Kyoto Protocol are retrieved from the Swiss Land Use Statistics (SFSO 2013; see also Chapter 7.2). The Swiss Land Use Statistics AREA (SFSO 2006a) uses a regular sample grid with a grid size of 100 m to frame her fixed sample points with known coordinates. To each grid point a specific combination category (see Table 7-2) is assigned.

11.2.2 Methodology used to develop the Land Transition Matrix

The methodology used to develop the land transition matrix is described in detail in Chapter 7.2.3.2.

11.2.3 Maps / Database to identify the geographical Locations and the system of Identification Codes for the geographical Locations

All Afforestations and Deforestations are accounted for under Article 3, paragraph 3 and are not reported under Forest Management under Article 3, paragraph 4. Afforestations older than the general conversion period of 20 years, are still reported under Afforestations: CRF-table 5(KP-I)A.1.2. The calculation of changes in carbon stocks is described in Chapter 11.3.1.1. The changes in areas between the activities under Article 3, paragraph 3 and Article 3, paragraph 4 are listed in KP-Table NIR2 (see).

Forest areas under Forest Management are subdivided into productive forests (CC12) and unproductive forests (CC13; for a description see Chapter 7.3.4.9). Productive forests in Switzerland reveal a high heterogeneity in terms of elevation; growth conditions and tree species composition (see Chapter 7.2.3.1 and Figure 7-4). We therefore stratified Switzerland into five National Forestry Inventory production regions (L1: Jura, L2: Central Plateau, L3: Pre-Alps, L4: Alps, L5: Southern Alps), three altitudinal zones (Z1: <601 m, Z2: 601-1200 m, Z3: >1200 m) and two soil types (mineral soils and organic soils). In the KP CRF-tables, the stratification of the activity data into production region (L) and altitudinal level (Z) is indicated in the column "Subdivision".

Afforestation

Activity data for Afforestations are derived from the Swiss Land Use Statistics (AREA) (SFSO 2006a, 2013; see also Chapter 7.2.2.1). A detailed description of the identification of Afforestations fulfilling the Kyoto definition is provided in FOEN (2010h).

Deforestation

Data for Deforestations are derived from the Swiss Land Use Statistics (AREA). A detailed description of the identification of Deforestations under the Kyoto Protocol from the AREA dataset is given in FOEN (2010d) and Sigmaplan (2010a).

Not all changes from a forest combination category (Afforestation CC11, productive forest CC12 and unproductive forest CC13) to a non-forest combination category do correspond to the definition of Deforestation according to the Kyoto Protocol Art. 3.3. The following criteria identify conversions from a forest combination category to a non-forest combination category, which are not classified as Deforestations under the Kyoto Protocol Art. 3.3 (FOEN 2010d):

1. Non-permanent conversions are due to Forest Management practices, natural dynamics or hazards:
 - Tree loss is temporally limited: areas with loss of tree biomass, but where a change in land use cannot be identified. Natural regeneration, which is a common practice in Swiss Forest Management, is expected, but could not yet be recognized on the aerial photograph at the time the AREA survey (see Chapter 7.2.2.1) was conducted. Also, in the NFI methodology (Brändli 2010: 91) "forest aisles" under high-tension are explicitly classified as forests (see also Chapter 11.1.3). Further, a study by Sigmaplan (2012a) showed that, although the aspect of "temporal limitation" was considered when classifying Deforestations, at the end still 14% of these Kyoto Deforestations were in fact "short-term reduction of crown coverage" and should be classified as "management interventions" rather than as real land-use changes (see 11.1.3).
 - Tree loss is spatially limited: conversion is caused by an alteration of the surrounding stand, but the change does not affect the tree cover at the sample point.
2. Conversions of combination categories (see Table 7-2 and Table 7-6) not meeting the definition of Deforestation as defined under the Kyoto Protocol and in Switzerland's Initial Report (FOEN 2006h).
 - Areas smaller than the minimum area of 625 m².
 - Areas with a reduction in forest cover on the grid point but still fulfilling the Kyoto definition of Forest, i.e. having the potential to reach 3 m at maturity in situ.
3. No change in land use took place: reduction of tree cover without land-use change; former land use was mainly pasture
4. Tree loss is not human-induced: Conversion due to natural hazards and dynamics.

Forest Management

Since all forests in Switzerland are subject to Forest Management, the area of managed forest corresponds to the forest area (see FOEN 2006h, Sect. E) as derived from the Swiss Land Use Statistics (AREA; SFSO 2006a, 2013; see also Chapter 7.2). We report changes in pools for the following geographical locations:

- productive forest remaining productive forests (CC12 remaining)
- productive forest converted to unproductive forests (CC12 to CC13)
- unproductive forest remaining unproductive forests (CC13 remaining)

- unproductive forest converted to productive forests (CC13 to CC12).

Difference in area reported under KP Art. 3.4 Forest Management and area “Forest Land remaining Forest Land” reported under the UNFCCC

A direct comparison of the areas reported in the CRF-tables under the Convention "Forest Land remaining Forest Land" (Table 5A) and under "Forest Management" under the Kyoto Protocol (Table 5(KP-I)B.1) is not possible due to the different structure of these CRF-tables and due to different reporting requirements:

- Conversions to Forest Land which are not human-induced (natural regeneration) are not accounted for as “Afforestations under the Kyoto Protocol”. These areas are reported under KP Art. 3.4 Forest Management in table 5(KP-I)B.1 as soon as the definition of Forest is fulfilled. Under the Convention, these Afforestations are reported under land-use category 5A2 with a conversion time of 20 years.
- Afforestations under the Kyoto Protocol which are older than 20 years are always reported under Art. 3.3 (KP-CRF-table 5(KP-I)A.1.2: units of land harvested since the beginning of the commitment period). Thus, there is no reclassification of the units of lands reported under Art. 3.3. In contrast, under the Convention, the Afforestations older than 20 years are moved to the land-use category 5A1 “Forest Land remaining forest land”.
- Not all changes from a forest combination category (CC11, 12, 13) to a non-forest combination category correspond to the definition of Deforestation according to the Kyoto Protocol Art. 3.3. (see above). These areas remain under the KP Art. 3.4 activity Forest Management and are included in the areas as reported in table 5(KP-I)B.1.
- Reporting of land-use changes LUC: Since only the KP activity “Forest Management” is chosen under KP Art. 3.4, changes from other KP activities to forest land are not reported as LUC but are reported as CC12 or CC13 as soon as the KP definition of forest is fulfilled. Only conversions within the activity “Forest Management” are reported under the Kyoto Protocol, i.e. CC12 to CC13 and CC13 to 12. Under the Convention, LUC to forest land are reported in land-use category 5A2.

In a study by Meteotest (2013a) the reported activity data have been checked and compared. It could be shown that the differences in the CRF-tables can be explained and that the resulting budget of areas reported under the Convention and the Kyoto Protocol are identical.

The cross-check was updated for the present submission:

Table 11-5 Area budget (in kha) of KP-LULUCF and LULUCF under the Convention (UNFCCC) in the year 2012 for forest land and afforestations.

| activity | Table, Cells | area UNFCCC kha | area KP kha | Check Difference kha | remarks |
|--------------------------------|------------------------------|-----------------------|----------------|----------------------------|---------|
| All Forest Land | | | | | |
| Forest Management | 5(KP-I)B.1, C9 | | 1'232.660 | | a) |
| Afforestations <= 20 years | 5(KP-I)A.1.1, C10 | | 1.697 | | b) |
| Afforestations > 20 years | 5(KP-I)A.1.2, C10 | | 0.793 | | c) |
| Total area KP | | | 1'235.150 | | |
| Non-Kyoto loss of forest cover | | | -0.670 | | d) |
| Forest Land UNFCCC | 5.A, C10 | 1'234.480 | | | e) |
| Total | | 1'234.480 | 1'234.480 | 0.000 | |
| Afforestation, CC11 | | | | | |
| UNFCCC | 5.A, C31+C35+C39 +C43+C47 | 1.697 | | | f) |
| KP | 5(KP-I)A.1.1, C10 | | 1.697 | 0.000 | g) |

Remarks:

- a) KP forest management consists of CC12 and CC13 areas fulfilling the criteria of the KP.
- b) KP afforestations are afforested areas since 1990 cumulated over 20 years at most.
- c) KP afforestations "older than 20 years" (>20 years) is the area that has been afforested since more than 20 years. In the UNFCCC tables these areas belong to 5A1 (CC12 or CC13).
- d) The non-Kyoto loss of forest cover is the part of the total area of forest loss (reported under UNFCCC) not fulfilling the definition of deforestations according to the Kyoto Protocol (see NIR Chapter 11.2.3). For the comparison this area must be subtracted from the KP forest area.
- e) The total Forest Land in CRF 5A covers productive forests (CC12), unproductive forests (CC13) and afforestations (CC11). It is congruent with the forest area derived from the aerial photos of the AREA survey (NIR Chapter 7.2.2).
- f) The CC11 area in UNFCCC can be taken from CRF 5A2 by summing up the afforestation source categories.
- g) The cumulated (20 years) CC11 area of KP and UNFCCC are congruent.

Area reported under Afforestation, Deforestation and Forest Management

AREA data allow to clearly separate between the land areas subject to a specific activity. Absolute and cumulated activity data of Afforestations, Deforestations and forests under Forest Management are listed in Table 11-6. The total Swiss area remains constant and amounts to 4'128.42 kha.

Table 11-6 Activity data for activities under Article 3, paragraphs 3 and 4, 1990-2012. Afforestation, Deforestation data and values depicting the area of Forest Management are derived from the Swiss Land Use Statistics (AREA) (derived from SFSO 2006a, 2013). See also KP-CRF-Table NIR2 (Table 11-2).

| Year | Deforested area [kha] | Cumulated deforested area since 1990 [kha] | Afforested area [kha] | Cumulated afforested area since 1990 [kha] | Area Forest Management [kha] |
|------|-----------------------|--|-----------------------|--|------------------------------|
| 1990 | 0.31 | 0.31 | 0.27 | 0.27 | 1178.82 |
| 1991 | 0.31 | 0.62 | 0.26 | 0.53 | 1181.97 |
| 1992 | 0.31 | 0.93 | 0.26 | 0.79 | 1185.14 |
| 1993 | 0.31 | 1.24 | 0.23 | 1.02 | 1188.18 |
| 1994 | 0.33 | 1.58 | 0.18 | 1.20 | 1191.08 |
| 1995 | 0.34 | 1.92 | 0.13 | 1.33 | 1193.86 |
| 1996 | 0.34 | 2.26 | 0.12 | 1.44 | 1196.36 |
| 1997 | 0.35 | 2.61 | 0.09 | 1.53 | 1198.84 |
| 1998 | 0.38 | 2.98 | 0.06 | 1.59 | 1201.10 |
| 1999 | 0.37 | 3.36 | 0.06 | 1.65 | 1203.32 |
| 2000 | 0.37 | 3.73 | 0.06 | 1.71 | 1205.53 |
| 2001 | 0.36 | 4.09 | 0.06 | 1.77 | 1207.75 |
| 2002 | 0.36 | 4.44 | 0.06 | 1.82 | 1209.96 |
| 2003 | 0.35 | 4.79 | 0.06 | 1.88 | 1212.18 |
| 2004 | 0.34 | 5.14 | 0.06 | 1.94 | 1214.39 |
| 2005 | 0.28 | 5.41 | 0.10 | 2.04 | 1217.30 |
| 2006 | 0.23 | 5.64 | 0.08 | 2.12 | 1220.14 |
| 2007 | 0.18 | 5.82 | 0.08 | 2.20 | 1223.01 |
| 2008 | 0.12 | 5.93 | 0.07 | 2.27 | 1226.24 |
| 2009 | 0.25 | 6.19 | 0.06 | 2.33 | 1229.15 |
| 2010 | 0.31 | 6.50 | 0.05 | 2.38 | 1230.41 |
| 2011 | 0.31 | 6.81 | 0.05 | 2.44 | 1231.53 |
| 2012 | 0.31 | 7.13 | 0.05 | 2.49 | 1232.66 |

11.3 Activity-specific Information

11.3.1 Methods for Carbon Stock Change and GHG Emission and Removal estimates

11.3.1.1 Description of the Methodologies and the underlying Assumptions used

Emission factors for Afforestations, Deforestations and Forest Management were accounted for following the methodology described in chapter 7.1.3.2, Table 7-3 and using equations 7.1-7.6.

Annual values for carbon stocks and changes in the pools of living biomass (for a description of separation in above and belowground living biomass see Chapter 7.3.4.6), dead wood, litter and soil carbon of afforestations (CC11), productive forests (CC 12) and unproductive forests (CC13) are displayed in Table 7-4, Table 7-19 and Table 7-23. The methodological approach is based on Table 7-3 and elaborated in detail for each pool in Table 11-7. Under the Kyoto Protocol, the pool "dead organic matter" is separated into dead wood and litter. Supplementary methodological information can be found in FOEN (2014c).

Table 11-7 Application of the methodology described in equations 7.1-7.6 in Chapter 7.1.3.2 and in Table 7-3 for calculating changes in carbon pools for the Kyoto activities Afforestations (CC11) younger than 20 years (≤ 20 yr) and older than 20 years (>20 yr), Deforestations (DEF) and Forest Management (FM) with the 4 geographical locations: CC12 remaining, CC13 remaining, conversions from CC12 to CC13 (FM CC1213) and conversions from CC13 to CC12 (FM CC1312). In the case of Deforestation (LUC to CC51), losses in soil carbon are accounted for by reducing the soil carbon pool by 50%. A conversion time CT of 20 years is applied for all pools except for the loss of living biomass, litter and dead wood after Deforestation (CT=1 year). Suffixes used: l for living biomass, dw for dead wood, s for soil, li for litter, i for spatial stratum. CC11 (afforestation), CC12 (productive forests) and CC13 (unproductive forests) refer to the specific combination category (see Table 7-2).

| | Living biomass | Litter | Dead Wood | Soil-C |
|---|--|--|---|---|
| Afforestation CC11 ≤ 20 yr | gain-loss $\text{gainC}_{li,CC11} - \text{lossC}_{li,CC11}$ $= \text{gainC}_{li,CC11} - 0$ $= \text{gainC}_{li,CC11}$ | stock-change, CT=20 $(\text{stockC}_{li,CC11} - \text{stockC}_{li,CC31/51})/CT$ $= (0 - 0)/20 = 0$ | stock-change, CT=20 $(\text{stockC}_{dw,i,CC11} - \text{stockC}_{dw,i,CC31/51})/CT$ $= (0 - 0)/20 = 0$ | stock-change, CT=20 $(\text{stockC}_{s,i,CC11} - \text{stockC}_{s,i,CC31/51})/CT$ |
| Afforestation CC11 > 20 yr | gain-loss $\text{gainC}_{li,CC12} - \text{lossC}_{li,CC12}$ | gain-loss $\text{changeC}_{li,CC12}$ | gain-loss $\text{changeC}_{dw,i,CC12}$ | gain-loss $\text{changeC}_{s,i,CC12}$ |
| Deforestation DEF | stock change, CT=1 $(0 - \text{stockC}_{li,CC12})/CT$ $= -\text{stockC}_{li,CC12}$ | stock change, CT=1 $(0 - \text{stockC}_{li,CC12})/CT$ $= -\text{stockC}_{li,CC12}$ | stock change, CT=1 $(0 - \text{stockC}_{dw,i,CC12})/CT$ $= -\text{stockC}_{dw,i,CC12}$ | stock change, CT=20 $(0.5 \cdot \text{stockC}_{s,i,CC12} - \text{stockC}_{s,i,CC12})/CT$ $= -(0.5 \cdot \text{stockC}_{s,i,CC12})/20$ |
| FM CC12 remaining | gain-loss $\text{gainC}_{li,CC12} - \text{lossC}_{li,CC12}$ | gain-loss $\text{changeC}_{li,CC12}$ | gain-loss $\text{changeC}_{dw,i,CC12}$ | gain-loss $\text{changeC}_{s,i,CC12}$ |
| FM CC13 remaining | gain-loss $\text{gainC}_{li,CC13} - \text{lossC}_{li,CC13}$ $= 0$ | gain-loss $\text{changeC}_{li,CC13} = 0$ | gain-loss $\text{changeC}_{dw,i,CC13} = 0$ | gain-loss $\text{changeC}_{s,i,CC13} = 0$ |
| FM CC1213 | stock change, CT=20 $(\text{stockC}_{li,CC13} - \text{stockC}_{li,CC12})/CT$ | stock change, CT=20 $(\text{stockC}_{li,CC13} - \text{stockC}_{li,CC12})/CT = 0$ | stock change, CT=20 $(\text{stockC}_{dw,i,CC13} - \text{stockC}_{dw,i,CC12})/CT$ $= (0 - \text{stockC}_{dw,i,CC12})/20$ | stock change, CT=20 $(\text{stockC}_{s,i,CC13} - \text{stockC}_{s,i,CC12})/CT = 0$ |
| FM CC1312 | gain-loss $\text{gainC}_{li,CC12} - \text{lossC}_{li,CC12}$ | stock change, CT=20 $(\text{stockC}_{li,CC12} - \text{stockC}_{li,CC13})/CT = 0$ | stock change, CT=20 $(\text{stockC}_{dw,i,CC12} - \text{stockC}_{dw,i,CC13})/CT$ $= \text{stockC}_{dw,i,CC12}/20$ | stock change, CT=20 $(\text{stockC}_{s,i,CC12} - \text{stockC}_{s,i,CC13})/CT = 0$ |

Reforestation

Reforestation does not occur in Switzerland (FOEN 2006h, Sect. E).

Afforestation ≤ 20 years: Units of Land not harvested since the beginning of the Commitment Period

Most of the afforestations occur on permanent grasslands CC31 and on settlements CC51 (see Table 7-9).

Living Biomass

- Gross growth of living biomass of Afforestations follows a logistical growth function. Values are available for three altitudinal levels (Table 7-25). The total gross growth of the cumulative afforested area was determined by multiplying the afforested area of a specific year with the corresponding age-specific growth values. For estimates under the UNFCCC, a simplified approach was chosen and mean growth values were applied.

Soil Carbon

- Organic soils: In the case of organic soils, emissions due to drainage are calculated as described in Chapter 11.3.1.2.
- Mineral soils: In the case of land-use conversions to Afforestations, the difference in soil carbon stocks between land use before the conversion (CC31 and CC51) and Afforestations CC11 is considered.

The fact that soils are acting as small sinks under afforestation is supported by Jandl et al. (2007) who reviewed several studies on the effect of different forest management systems (including afforestations) on soil carbon sequestration and concluded that a long-term consequence of afforestation is the gradual incorporation of C in the mineral associated soil C pool.

Dead Wood and Litter

- On grasslands and areas with settlements there are no dead wood and no litter available and a zero stock is attributed. Assuming no dead wood nor litter on Afforestations, the difference in the carbon stocks of these pools are zero. This is a conservative estimate (in terms of IPCC good practice: IPCC 2003, Sect. 3.1.5), since there actually is a small pool of dead wood and litter under Afforestations. This conservative approach is confirmed in a study by Zimmermann and Hiltbrunner (2012) and by other literature (see Chapter 11.3.1.2).

Afforestation > 20 years: Units of Land harvested since the beginning of the Commitment Period

In KP-CRF-Table A.1.2, changes in carbon stocks of Afforestations older than 20 years are reported. After 20 years, Afforestations are subject to normal Forest Management and the first thinnings and treatments are conducted. There is however no reclassification of these afforested areas to Forest Management: all Afforestations after 1990 are reported under Article 3.3 (see Chapter 11.2.3). Emissions and removals for the carbon pools of Afforestations older than 20 years are calculated using the emission factors of productive (CC12) forests, since nearly all of the afforestations (99.9%) develop to productive forests (see methodological description under “Forest Management”).

Deforestation

- Change in carbon stock due to conversion: Total carbon stock of living biomass, dead wood and litter are immediately removed after Deforestation. Losses in soil carbon due to disturbance caused by Deforestation (conversion to buildings and constructions) are accounted for by reducing the soil carbon pool by 50% (Covington 1981; Rusch et al. 2009; see also Chapter 7.1.3.2) over a conversion period of 20 years (see Table 7-3).

Forest Management

Living biomass

- Total living biomass, calculated with single-tree allometric function, is separated into above- and belowground living biomass using the factors in Table 7-20 as described in Chapter 7.3.4.6.
- Gross growth of productive forests is used for “CC12 remaining”. Gross growth of unproductive forests is used for “CC13 remaining” and amounts to zero (see Chapter 7.3.4.9).
- Cut and mortality reflect yearly losses of living biomass in productive forests “CC12 remaining”. Unproductive forests are not systematically harvested (description Chapter 7.3.4.9). Thus losses of unproductive forests “CC13 remaining” are zero. Moreover, since yearly harvesting amounts from forest statistics (FOEN 2013k) are divided over the productive forests, total harvesting in Swiss forests is accounted for under CC12 remaining.

- For the conversions between different forest source categories (“CC13 to CC12” and “CC12 to CC13”) the method is chosen in such a way that no potential carbon losses are underestimated. For areas which changed from “CC12 to CC13” the difference in carbon stocks of living biomass was considered and a net loss in carbon stock of living biomass is reported; in the case of a conversion from “CC13 to CC12” a gain-loss approach has been applied, since applying a stock-change approach would lead to a considerable sink in living biomass in this category.

Dead wood, litter and soil

- A literature overview of the influence of forest management on soil carbon and litter is provided in Didion (2014) and summarized in Chapter 11.3.2.2.
- For productive forests “CC12 remaining”, values for yearly changes in carbon stock of dead wood, litter and soil are used (Table 7-23). Estimates of those yearly changes were derived from Yasso07 (see Chapter 7.3.4.8). For unproductive forests “CC13 remaining”, yearly changes in dead wood, litter and soil carbon stock are assumed to be zero (Chapter 7.3.4.9).
- For the conversions between different forest categories (“CC13 to CC12” and “CC12 to CC13”) the difference in carbon stock of dead wood, litter and soil carbon is taken into account (Table 7-23). For the conversion “CC12 to CC13”, we report a net loss in carbon stock for these pools, in the case of a conversion “CC13 to CC12”, we report a net gain (Table 7-4).
- In the case of organic soils, emissions due to drainage are calculated as described in Chapter 11.3.1.2.

Differences in accounting for “Forest Sector 5A1 and 5A2” under the UNFCCC and „Forest Management“ under KP Art. 3.4

For reporting changes in living biomass of Afforestations, under KP the age-specific growth value was multiplied with the afforested area of a specific year. For estimates under the UNFCCC, a simplified approach was chosen and mean growth values instead of age-specific growth values were applied.

Under KP Art. 3.4, natural forest regeneration is reported under “Forest Management” as CC12 or CC13 as soon as the KP definition of Forest is fulfilled and management activities have taken place. Changes within the activity “Forest Management” are reported under the Kyoto Protocol in the source categories CC12 to CC13 and CC13 to 12.

Under the UNFCCC, all changes in land use from non-forest land to forest land are reported in the land-use category 5A2 for a conversion time of 20 years.

11.3.1.2 Justification when omitting any Carbon Pool or GHG Emissions/Removals from Activities under Article 3.3 and elected Activities under Article 3.4

KP LULUCF Table NIR1 (Table 11-1) summarizes the activity coverage and the carbon pools reported. When using the conservative Tier 1 approach (IPCC 2003, Sect. 3.1.5) assuming a specific carbon pool to be in balance, the carbon pool is indicated as not reported (NR). This is the case for litter and dead wood under Afforestation. Also for all pools of unproductive forests CC13, we report no changes.

Change in Carbon Pool not Reported

Afforestation: litter and dead wood

Figure 5.4.2 in the GPG LULUCF (IPCC 2003) illustrates the issue that, when data are not available, a “good practice method” should be used and that financial resources for key categories should not be jeopardized and an appropriate tier level should be used. Because Afforestation is not a key category (see Chapter 11.6.1), a conservative estimate (Tier 1) for the pools litter and dead wood is compliant to IPCC guidelines (IPCC 2003) for the first twenty years when no management activity takes place. We provide verifiable information to justify this approach:

for the pools litter and dead wood is compliant to IPCC guidelines (IPCC 2003). We provide verifiable information to justify this approach:

- **Changes in litter after Afforestation:** Under the Kyoto Protocol, we conservatively report no changes in the litter pool after Afforestations. In an experiment by Zimmermann and Hiltbrunner (2012) litter accumulation of an Afforestation with Norway Spruce was determined 40 years after Afforestation. The authors found accumulation rates of $0.17\text{--}0.20 \text{ t C ha}^{-1} \text{ yr}^{-1}$. Other studies show even higher accumulation rates of $0.24\text{--}0.34 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for Afforestations with Norway spruce in Southern Alps (Thuille and Schulze 2006), $0.24 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for Afforestation with ash and maple (Alberti et al. 2008) and $0.36 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for Scotch pine (Vesterdal et al. 2002). Karhu et al. (2011) determined for Finisch forest stands mean annual rate of carbon accumulation in the litter over 18 years was 0.28 and 0.15 Mg ha^{-1} for Scots pine and birch, respectively.

In a literature overview, Jandl et al. (2007) showed, that the accumulation of a forest floor layer in, e.g., a conifer forest, is a C sink. The authors concluded that after afforestation, forest floors accumulate C quickly. A long-term consequence of afforestation is the gradual incorporation of C in the mineral-associated soil C pool.

- **Changes in dead wood after Afforestation:** Under the Kyoto Protocol, we conservatively report no changes in the litter pool after Afforestations. Zimmermann and Hiltbrunner (2012) showed that 40 years after Afforestation with Norway Spruce, dead wood volume amounted to 10.4 t C ha^{-1} . Thus, an annual increase in dead wood of $0.26 \text{ t C ha}^{-1} \text{ yr}^{-1}$ for afforestations with Norway Spruce can be derived from this case study, considering the fact that on grassland, the starting point of most afforestations, there is no dead wood available.
- Besides the results of the case studies listed above, we provide a reasoning based on sound knowledge of likely system responses (Grassi and Blujdea 2011): At stand level, the pools dead wood and litter of Afforestation on cropland and grassland cannot be a source, especially if previous land use did not have perennial woody biomass. In Afforestations the stands development follow exponential patterns, which can also be theoretically attributed to all other C pools.

Note that for afforestations older than 20 years, we report estimates of CSC in dead wood, litter and SOC.

Unproductive Forests CC13

These unproductive forests and the reasoning why these pools are not a source is described in detail in Chapter 7.3.4.9.

Based on the fact that unproductive forest land only covers 7.2% of the area under Forest Management (Table 7-10 LUC matrix and CRF-KP-Table 5(KP.I)B.2.) and based on the description of these stands given before, emissions or removals of any of the pools of unproductive forests cannot account for more than 25% of the activity Forest Management. According to IPCC 2003 Fig 3.1.1, note 4, 25% is the threshold that would require a higher

Tier. Since our resources are limited and we do not want to jeopardize financial resources for the key categories (IPCC 2003 Fig. 5.4.2.), Switzerland decided to use the Tier 1 approach and reports no changes in living biomass, litter, soil and dead wood of unproductive areas arguing that this is a conservative estimate. Emissions from organic soils are accounted for using default factors from IPCC 2013 (Tier 1).

Greenhouse Gas Sources Reported

- Fertilization of forests is prohibited by the Swiss forest law and adherent ordinances (Swiss Confederation 1991, 1992). Additionally, the “Ordinance on Chemical Risk Reduction” (Swiss Confederation 2005) prohibits the application of fertilizers, including liming, in forests. Thus, emissions from fertilization are reported as “not occurring”.
- Drainage of forests is not a permitted practice in Switzerland and since 1991 not a permitted practice in Switzerland (Swiss Confederation 1991). There are no nation-wide survey data available. It is possible that a small part of the Swiss forest has been drained before 1990 or has been established on drained areas. We conservatively report all organic forest soils to be drained which is definitely an overestimation. In order to calculate CO₂ emissions due to drainage, we used equation 3.2.15 of the GPG for LULUCF (IPCC 2003) and applied the default emission factor of 0.68 Mg ha⁻¹ yr⁻¹. N₂O-emissions due to drainage are not estimated as no data are available. Moreover, reporting is not mandatory according to the Annex of chapter 3 of the IPCC GPG for LULUCF (reported as “NE” in KP-Table NIR1; Table 11-1).
- Biomass burning: emissions of CO₂, CH₄ and N₂O are reported. The calculation of these emissions is described in Chapter 7.3.4.13 according to the methodology of the GPG LULUCF (IPCC 2003).

11.3.1.3 Information on whether or not indirect and natural GHG Emissions and Removals have been factored out

No anthropogenic GHG emissions and removals from elevated carbon dioxide concentrations, indirect nitrogen deposition or the dynamic effects of the age structure resulting from LULUCF activities under Article 3, paragraphs 3 and 4 prior to 01 January 1990 have been factored out.

The IPCC does not give specific methods for factoring out these effects. Besides this, there are no reliable country specific data available. Investigations on elevated CO₂ concentrations on growth showed complex relationships in the middle term. Some species showed an increase others a decrease and some no change in growth (Bader et al. 2013). Until recently, researchers were convinced that there is a distinct positive effect of nitrogen deposition and gross growth (Spiecker 1999; Jarvis and Linder 2000). Recent scientific publications, however, question this relationship and even show that nitrogen deposition, while leading to soil acidification, even can cause a reduction in growth (Hyvönen et al. 2008; Högberg et al. 2006; Braun et al. 2010; Gschwantner 2006; Meining et al. 2008). Such acidification processes are widely detected in Swiss forest soils.

11.3.1.4 Changes in Data and Methods since the previous Submission (Recalculations)

Table 1-12 lists the improvements made since last year based on the questions or recommendations of the UNFCCC Expert Review Team.

A recalculation of the years 2008, 2009, 2010 and 2011 was carried out.

The completion of the AREA surveys (see Chapter 7.2.2.1, SFSO 2013) led to a recalculation of all areas reported under Art. 3.3 and Art 3.4 Forest Management.

In detail, the changes in the calculation of the emission factors calculated for all areas reported under Art. 3.3 and Art 3.4 Forest Management are described in Chapter 7.3.7 and in Chapter 10.1.2.

The following Kyoto-specific methodological modification was made for this submission:

- The calculation of carbon stock changes in the different pools has been adapted such that it is harmonized between reporting under the UNFCCC and under the KP, that no double counting is possible and that the most conservative accounting method is applied. The calculation approach is explained in Chapter 7.1.3.2 and in Chapter 11.3.1. and transparently shown in Table 7-3 and Table 11-7.
- New available NFI 4a+ data covering the period 2009-2012 have been used in the calculations for changes in living biomass (Thürig 2014). Total living biomass has been separated into above- and belowground biomass (see Chapter 7.3.4.6)
- Updated estimates of yearly changes in carbon stocks of soil organic carbon, dead wood and litter, modeled with Yasso07, are described in Didion et al. (2013).

Additional information, references and documentation justifying not to report changes in litter and dead wood under afforestation is provided in Chapter 11.3.1.2.

The description of unproductive forests has been extended and also the reasoning why the carbon pools of the unproductive forest are not a source is supported with references (see section 7.3.4.9).

11.3.1.5 Uncertainty Estimates

An overview of the uncertainty estimates of activity data is discussed in detail in Chapter 7.2.5 and is shown in Table 7-10. Uncertainty estimates of emission factors for the reported activities under the Kyoto Protocol are shown in Table 7-5, overall uncertainties in Table 11-8.

A detailed description of the determination of the emission factor uncertainty of Forest Management can be found in Chapter 7.3.5. Table 7-5 lists the relative uncertainties in the LULUCF sector: an uncertainty of 63% was calculated for Afforestations, 50% for Deforestations and 63% for Forest Management.

Lands fulfilling the definition Forest (see Chapter 11.1.1) are accounted for under "Forest Management". This means, that lands under Forest Management which are due to natural regeneration are attributed the uncertainty of Forest Management.

Table 11-8 Uncertainty estimates of activity data and emission factors and the overall uncertainty of activities reported under the Kyoto Protocol Article 3.3 and Article 3.4

| Activity under KP | Associated category in UNFCCC inventory (Chapter 7.3) | Activity data uncertainty [%] | Emission factor uncertainty [%] | Combined uncertainty [%] |
|-------------------|---|-------------------------------|---------------------------------|--------------------------|
| Afforestation | 5A2 Land converted to Forest Land | 2 | 63 | 63 |
| Deforestation | mainly 5E2 Land converted to Settlements | 5 | 50 | 50 |
| Forest Management | 5A1 Forest Land remaining Forest Land | 2 | 63 | 63 |

11.3.1.6 Other methodological Issues

Methodology used for reporting under the Kyoto Protocol is described in detail in previous sections.

N₂O emissions as a result of the disturbance associated with land-use conversion (Deforestation) to Cropland are reported in KP-CRF-table 5(KP-II)3. The emissions are calculated according to the methodology described in Chapter 7.4.4.4.

11.3.1.7 The Year of the onset of an Activity, if after 2008

All activities reported started in 1990, i.e. before the beginning of the first commitment period.

11.3.2 Category-Specific QA/QC and Verification

In Chapter 7.3.6 category-specific QA/QC and verification items for forest land are described in detail. Differences between the forest areas reported in “Forest Sector 5A1 and 5A2” under UNFCCC and „Forest Management“ under KP Art. 3.4 are explained in Chapter 11.2.3.

11.3.2.1 Comparison of the Forest Areas reported in the CRF-tables and KP-CRF-tables

The relationship between forest land reported under LULUCF table 5.A and those reported for activities under KP-LULUCF forests has been documented in Meteotest (2013a) for the year 2010. An updated version of this study has been compiled in Table 11-5 in Chapter 11.2.3.

11.3.2.2 Impact of Forest Management on Changes in Carbon Stocks in Soil and in Litter

Accounting for forest management impacts on carbon storage in litter and soil in Swiss productive forests with Yasso07

To estimate C stocks and C stock changes in the reported litter and soil pools, Switzerland uses the C cycling model Yasso07 (cf. Didion et al. 2012, 2013). Inputs to the model include C deriving from the annually amount of leaves, fine-roots and fruits that are produced by living trees. The C inputs are obtained for each plot in the National Forest Inventory (NFI) that is simulated with Yasso07. The NFI plots have been repeatedly measured since the first inventory in 1985 and, hence, observed changes in the volume of living and dead biomass reflect, among other, the site-specific impact of forest management. Based on harvesting statistics and allometric relationships, the production of dead wood (incl. dead roots, stems, stumps and branches) and litter from living trees (i.e. controlled by forest management) and as harvest residues are estimated.

Thus, the Yasso07-model reflects the impact of forest management: forest management effects on C stocks in litter (including non-woody and woody material) and soil are fully accounted for in the Swiss GHGI (Didion 2014).

Literature Review

A detailed screening of the available scientific literature on the impact of forest management on carbon stock changes in litter and soils is provided in Didion (2014). The majority of studies indicated no significant effect of forest management on soil C stocks with the exception of clearcutting (e.g. Jandl et al. 2007). Since silvicultural practices in Switzerland are regulated by law and exclude intensive management options such as clearcuts, fertilization or liming (Swiss Confederation 1991, 1992), no or only minor forest management impacts on soil C stocks can be expected. The production of litter is directly affected by silvicultural practices since the removal of trees results in harvest residues and in a decrease in the amount of remaining foliage (e.g. Van Miegroet and Olsson 2011). Generally, the impact of forest management on litter production is temporary and losses of litter C can be rapidly replaced (Nave et al. 2010).

11.4 Article 3.3.

Figure 11-2 shows removals of CO₂ eq from Afforestations and emissions of CO₂ eq from Deforestations for the years 1999-2012. The corresponding values are listed in Table 11-4.




Figure 11-2 Removals (negative sign) and emissions (positive sign) of CO₂ eq from Afforestations (upper panel) and from Deforestations (lower panel) shown per carbon pool, 1999-2012. Belowground (BG) and aboveground (AG) living biomass is reported separately.

The order of magnitude of total removals or emissions of CO₂ eq from Afforestations and Deforestations is considerably different (Figure 11-2, Figure 11-3). Since carbon from living biomass is immediately removed after clear-cutting, Deforestation can be seen as a “quick carbon-losing process”. In contrast, due to the slow increase of living biomass, Afforestation is a “slow process with increasing importance” in terms of carbon accumulation. CO₂ emissions on organic soils under Afforestations are due to drainage (see Chapter 11.3.1.2).




Figure 11-3 Removals (negative sign) and emissions (positive sign) of CO₂ eq of Afforestations and Deforestations, 1999-2012.

11.4.1 Information that demonstrates that Activities under Article 3.3. began on or after 1 January 1990 and before December 2012 and are direct Human-induced.

The Swiss definitions of Afforestation and Deforestation only consider directly human-induced activities (see FOEN 2006h, Sect. E and FOEN 2010d).

Reforestation

For more than 100 years, the area of forest in Switzerland has been increasing (see Chapter 11.5.3), and a decrease in forest area as a result of Deforestation is prohibited by the Federal Law on Forests (Swiss Confederation 1991). Therefore, reforestation of areas not forested for a period of at least 50 years does not occur in Switzerland (FOEN 2006h, Sect. E). Switzerland only considers Afforestation and Deforestation under Article 3, paragraph 3.

Afforestation

Switzerland is very restrictive in reporting Afforestations under the Kyoto Protocol and only reports planted Afforestations (see Chapter 11.1.3; FOEN 2010h).

The annual rate of Afforestation since 1990 is assessed by AREA (Chapter 7.2.2). For reporting under the Kyoto Protocol, afforested areas since 1990 always remain in the "Afforestation" category. Therefore, the area of Afforestations is increasing since 1990 (see Table 11-6).

Afforestations older than 20 years are subject to normal Forest Management practices including harvesting (see Chapter 11.3.1.1). These areas are reported in CRF-tables 5(KP-I)A.1.2 and 5(KP-I)A.1.3.

Deforestation

In Switzerland, direct human-induced Deforestation is subject to authorization (Swiss Confederation 1991, Art. 5). For details concerning the classification of Deforestations under the Kyoto Protocol see Chapter 11.2.3). Only Deforestations carried out after 01 January 1990 are considered. For reporting under the Kyoto Protocol, deforested areas since 1990 remain in the Deforestation category. Therefore, the area of Deforestations is increasing since 1990 (see Table 11-6). Since Switzerland decided to only account for KP Art. 3.4 activity "Forest Management", these deforested areas are not subject to another KP Art. 3.4 activity.

11.4.2 Information on how Harvesting or Forest Disturbance that is followed by the Re-Establishment of Forest is distinguished from Deforestation

The Swiss definition of Deforestation only covers permanent conversions from forest land into non-forest land and is assessed by AREA applying the criteria discussed in chapter 11.2.3. This approach is confirmed by Sitmaplan (2012a).

They implicitly distinguish between permanent conversions and transient situations like harvesting or forest disturbance. Construction of e.g. pipelines and power supply lines within a forest area are transient situations (see Chapter 11.1.3 and 11.2.3; Brändli 2010). As described in FOEN (2010d), these non-permanent conversions are not classified as Deforestation under the Kyoto Protocol.

11.4.3 Information on the Size and Geographical Location of Forest Areas that have lost Forest Cover but which are not yet classified as Deforested

The AREA survey provides a detailed overview of land-use changes with regard to land cover and land use (see Chapter 7.2). Temporal changes of land cover can lead to a reclassification in AREA from a forest category to a non-forest category. In FOEN (2010d) and in Chapter 11.2.3 the criteria are listed which conversions from a forest combination category to a non-forest combination category are not identified as Kyoto Deforestation under the Kyoto Protocol.

11.5 Article 3.4

CO₂ eq emissions and removals from the reported pools and total CO₂ eq emissions and removals of the Kyoto Protocol activity Forest Management for the years 1999 until 2012 are shown in Figure 11-4. The corresponding values are listed in Table 11-4.




Figure 11-4 CO₂ eq emissions (positive sign) and removals (negative sign) from the reported carbon pools under Forest Management (upper panel) and the total CO₂ eq emissions and removals from Forest Management (lower panel), 1999-2012. Belowground (BG) and aboveground (AG) living biomass is reported separately.

The yearly fluctuations in the GHG emissions and removals from Forest Management can mainly be explained by changes in the losses of living biomass, dead wood and litter (see Table 11-4). Changes in the area of managed forest are relatively small (Table 11-6). In 2001 and 2002 Forest Management was a small source of CO₂ eq. and in 2000 a small sink compared to previous and following years. This was due to an elevated amount of losses in living biomass after storm Lothar, which ravaged Swiss forests in December 1999.

11.5.1 Information that demonstrates that Activities under Article 3.4. have occurred since 1 January 1990 and are Human-induced

According to the Swiss Federal Law on Forests, the extent and the spatial distribution of the total forest area in Switzerland has to be preserved (Swiss Confederation 1991, Art. 1) and thus, any change of the forested area has to be authorized. All Swiss forests are under continuous observation of the Swiss Forest Service and monitored by the NFI. Therefore, all forests in Switzerland are subject to Forest Management (FOEN 2006h, Sect. F).

11.5.2 Information relating to Cropland Management, Grazing Land Management and Revegetation, if elected, for the Base Year

Not applicable.

11.5.3 Information Relating to Forest Management

There is a long tradition of forest protection in Switzerland. The first federal Forest Act came into force in 1876, but it only covered the Alpine region. Its aim was to put a halt to the depletion of forests, to manage the remaining forest areas in a sustainable way, and to promote Afforestation. The Forest Act of 1902 covered the whole country. The Forest Act as well as an enabling overall economic development resulted in an increase of the forested area in Switzerland by nearly 50% compared to the mid-19th century (Figure 11-5). Also growing stock increased significantly due to changes in Forest Management practices. The Forest Act (Swiss Confederation 1991) that came into force in 1993 reaffirms the long-standing Swiss tradition of preserving both forest area and forest as a natural ecosystem. It prescribes sustainable Forest Management, prohibits clearing, and bans Deforestation unless it is replaced by an equal area of afforested land or an equivalent measure to improve biodiversity.




Figure 11-5: Historical mean growing stock and forest area in Switzerland since 1880.

In 2004, the Swiss national forest programme was published, outlining an action plan for the period 2004-2015 (SAEFL 2004b). It specifies five priority objectives: (1) the forest's

protective function is guaranteed, (2) the economic viability of the forestry sector is improved, (3) the value-added chain for wood is strengthened, (4) biodiversity is conserved and (5) forest soils, trees and drinking water are not threatened. These objectives encompass that CO₂ removals by sinks and emissions by sources in the forests shall be recognized in terms of compliance with the Kyoto Protocol while making better use of the potential of forests for timber production and fuel wood through economic incentives and implementing new technologies.

In November 2006, the Swiss government communicated in its initial report to the UNFCCC that Switzerland will be accounting for Forest Management under Article 3.4 of the Kyoto Protocol (FOEN 2006h).

To implement the objectives of the national forest programme (SAEFL 2004b), FOEN has formulated its wood resource policy (FOEN 2008h) which is coordinated with the other relevant sectoral policies (e.g. energy policy, regional development policy). This wood resource policy defines, among other things, the direction to be taken by federal policy in relation to wood promotion on completion of the "Wood 21" wood promotion programme which was terminated at the end of 2008. Under this programme, a wood action plan was started in 2009. The main focus in the implementation of the action plan lies on the ecologically and economically effective use of wood. With a view to the efficient use of wood, cascade use is prioritized, i.e. wood is used as material prior to its use for energy. In the case of energy use, greater overall efficiency of the conversion technology should be targeted.

11.5.4 Information that demonstrates that Emissions and Removals resulting from elected Article 3, Paragraph 4, Activities are not accounted for under Activities under Article 3, Paragraph 3

This information is requested in the Annex to 15/CMP.1 paragraph 9.c.

The reporting of Forest Management under article 3, paragraph 4 is clearly separated from the reporting of the activities under article 3, paragraph 3.

Units of lands with ARD (Afforestation, Reforestation and Deforestation) activities, are reported under Article 3, paragraph 3. These areas always remain under Article 3, paragraph 3. Afforestations older than 20 years are attributed to emissions factors of mature forests under forest management. These units of lands are reported in Table 5(KP-I)A.1.2 and not under forest management. Thus, there is no double counting of units of lands under article 3, paragraph 3 to Article 3, paragraph 4.

Table 11-5 Area budget (in kha) of KP-LULUCF and LULUCF under the Convention (UNFCCC) in the year 2012 for forest land and afforestations. Table 11-5 shows the clear distinction between units of land subject to activities under Article 3, paragraph 3 and Land subject to Forest Management under Article 3, paragraph 4.

11.5.5 Information that indicates to what extent Removals from Forest Management offsets the Debit incurred under Article 3, Paragraph 3

This information is requested in the Annex to 15/CMP.1 paragraph 9.d.

This information is shown in the summary KP-CRF-Table "Information table on accounting for activities under Articles 3.3 and 3.4 of the Kyoto Protocol.

11.6 Other Information

11.6.1 Key Category Analysis for Article 3.3. and 3.4. Activities

The results of the Tier 2 key category analysis including LULUCF are shown and explained in Chapter 1.5 and are displayed in Table 1-9 for the year 2012. The smallest UNFCCC category, and therefore also the smallest LULUCF category, considered key based on a Tier 2 level assessment is "6D Waste, Other, CH₄" with a contribution of 113.76 Gg CO₂ eq.

The following LULUCF activities under the Kyoto Protocol are listed in Kyoto Table NIR 3 (Table 11-3) because their associated LULUCF categories in the UNFCCC inventory are key categories under the level or trend assessment:

- **Forest Management** (-2'236.38 Gg CO₂ eq, encompasses all greenhouse gas emissions) is a key category under the Kyoto Protocol because its absolute contribution is higher than the smallest category considered key (113.76 Gg CO₂ eq for Tier 2) in the UNFCCC inventory. This activity is associated with the UNFCCC category „Forest Land remaining Forest Land“ (-2'134.56 Gg CO₂ eq, encompasses only CO₂ emissions). Since the total Swiss forest is considered as managed, there is a good agreement between the category under the Kyoto Protocol and the UNFCCC inventory category. According to Table 1-9, the UNFCCC category "Forest Land remaining Forest Land" is both level and trend key category under a Tier 2 assessment in 2012.
- **Afforestation and Reforestation** (-17.13 Gg CO₂ eq) is not a key category under the Kyoto Protocol because its absolute contribution is substantially lower than the smallest category considered key (113.76 Gg CO₂ eq for Tier 2) in the UNFCCC inventory. Natural forest regeneration due to abandonment of land is not reported as Afforestation under the Kyoto Protocol. The contribution of the associated UNFCCC category "Land converted to Forest Land" is 518.61 Gg CO₂ eq. It includes converted areas after natural regenerations, which are not reported as Afforestation under the Kyoto Protocol. The UNFCCC category "Land converted to Forest Land" is both level and trend key category under a Tier 2 assessment in 2012 (Table 1-9).
- **Deforestation** (221.87 Gg CO₂ eq) is a key category under the Kyoto Protocol because its contribution is higher than the smallest UNFCCC category considered key (113.76 Gg CO₂ eq for Tier 2). The associated UNFCCC category is „Land converted to Settlements“ (302.93 Gg CO₂ eq), but only a part of this UNFCCC category represents the activity Deforestation under the Kyoto Protocol. The UNFCCC category "Land converted to Settlements" is both level and trend key category under a Tier 2 assessment in 2012 (Table 1-9).

11.7 Information Relating to Article 6

Switzerland does not host Joint Implementation projects.

12.3 Discrepancies and Notifications

Switzerland's reports on discrepancies (R-2), CDM notifications (R-3), non-replacements (R-4) including reversal of storage and failure of certification and invalid units (R-5) have been uploaded on the UNFCCC Submission Portal.

During the reported year 2013, the Swiss Emissions Trading Registry had no discrepancies, no CDM notifications, no non-replacements including reversal of storage and failure of certification and no invalid units. Therefore the SIAR tables R-2, R-3, R-4 and R-5 are empty and no actions and changes have been taken to address discrepancies.

12.4 Publicly Accessible Information

In accordance to section E of the annex to decision 13/CMP.1 the Swiss Emissions Trading Registry makes non-confidential information available to the public via webpage or user-interface.

Non-confidential information is publicly available on the Swiss Emissions Trading Registry website www.national-registry.ch. The national allocation plan is accessible under 'National Allocation Plan'. All other information can be downloaded by selecting the menu item 'Reports'. The reports 'List of legal entities holding an account in the national registry', 'List of installations created in the Swiss national registry', 'List of accounts opened in the national registry', 'Annual summary of quantity per type of operation made in the national registry', 'Summary statement on the quantity of surrendered Allowances', 'Verified emissions table', 'Surrendered allowances table', and 'Compliance statement status table' are publicly accessible.

Data of transfers and holdings of individual accounts are considered as business secrets and the disclosure may prejudice their competitiveness. Information on acquiring and transferring units of companies (as legal persons) is therefore regarded as personal data. Article 19 of the Federal Act on Data Protection (FADP, SR 235.1 Bundesgesetz vom 19. Juni 1992 über den Datenschutz (DSG) 2) enacts that Federal bodies may disclose personal data if there is a legal basis for doing so or if there is an overriding public interest. In the present case these conditions are not fulfilled. Therefore, the registry of Switzerland cannot make the information on acquiring and transferring accounts publicly available and considers them as confidential. A statement on which information is considered as confidential can be found on the public website www.national-registry.ch.

All other information referred to in paragraphs 44 to 48 to the annex to decision 13/CMP.1 are made publicly available by the Swiss Emissions Trading Registry, if they are not covered by the above mentioned articles.

Information related to Article 6 projects is publicly accessible on the website <http://www.bafu.admin.ch/ji-e>. Switzerland does not host JI-projects and therefore no issuance of ERUs has taken place.

12.5 Calculation of the Commitment Period Reserve (CPR)

The commitment period reserve remains unchanged and is the same as defined in the update of the Initial Report (submitted on 20 December 2007; FOEN 2006h). The calculation of the commitment period reserve is based on the assigned amount (Method 1 in Table 12-2).

Table 12-2 Calculation of the commitment period reserve

| Method 1 (based on assigned amount) | Method 2 (based on latest reviewed submission) |
|---|--|
| 90 % of the assigned amount [t CO ₂ equivalent] | Total of 2011 emissions of sectors 1,2,3,4,6 times 5 [t CO ₂ equivalent] |
| 242 838 402 x 0.9 = 218 554 562 | 50 149 216 x 5 = 250 746 080 |

Method 1 results in the lower value. Accordingly the commitment period reserve of Switzerland is calculated as 218 554 562 tonnes CO₂ equivalent.

12.6 KP-LULUCF Accounting

According to the 'Report of the individual review of the annual submission of Switzerland submitted in 2012' (<http://unfccc.int/resource/docs/2013/arr/che.pdf>), Switzerland cancelled 144,158 Removal Units (RMUs) for its Deforestation activities into its Net Source Cancellation account (ITL notification ID: 1000340049), and subsequently issued 24,586 RMUs for its Afforestation and Reforestation activities, and 810,496 RMUs for its Forest Management activities in its Emissions Trading Registry. The transactions took place on 30 July 2013 (cancellation), and 8 August 2013 (issuance) respectively..

13 Information on Changes in National System

The initial Swiss national inventory system is described in detail in FOEN (2006h). The detailed description of the national inventory system is updated annually in the description of the quality management system (FOEN 2014a). Changes to the national system in accordance with 15/CMP.1, annex II, 30a-g are listed below.

Change of name or contact information (15/CMP.1 annex II.D 30a):

No changes.

Change of roles and responsibilities as well as change of the institutional, legal and procedural arrangements (15/CMP.1 annex II.D 30b):

The current arrangements for cooperation within the national inventory system are shown in Table 13-1. No changes.

Changes in the process of inventory compilation (15/CMP.1 annex II.D 30c):

No changes.

Change of key source identification and archiving (15/CMP.1 annex II.D 30d):

No changes.

Change of process for recalculations (15/CMP.1 annex II.D 30e):

No changes.

Changes to QA/QC plan, activities and procedures (15/CMP.1 annex II.D 30f):

No changes.

Change to official consideration and approval procedures (15/CMP.1 annex II.D 30g):

No changes.

Table 13-1: Formal arrangements for cooperation within the national inventory system. Items marked in bold have changed in the past year.

| Partner | Subject/Sector | Type of arrangement | Duration |
|--|---|---|------------------|
| <i>Institutions of the federal administration</i> | | | |
| Swiss Federal Office of Energy (SFOE) | Energy statistics | Agreement | open-ended |
| Federal Office of Civil Aviation (FOCA) | Aviation emissions | Agreement | open-ended |
| Swiss Federal Statistical Office (SFSO) | LULUCF (area surveys) | Agreement | open-ended |
| Agroscope research station | Agriculture emissions and removals | Agreement with the FOAG | open-ended |
| FOEN air pollution control and chemicals division (new name of the division) | - EMIS inventory data base & archive | Documentation of roles and responsibilities | open-ended |
| | - Energy emissions | | |
| | - Industrial process emissions (without F-gases) | | |
| | - Solvent and Other Product Use emissions | | |
| | - Waste emissions | | |
| | - Key category analysis | | |
| FOEN forest division | Forestry emissions and removals | Documentation of roles and responsibilities | 2014 |
| <i>Private companies</i> | | | |
| Carbotech | F-gas emissions | Contract | renewed annually |
| Sigmaplan / Meteotest | LULUCF data compilation | Contract | renewed annually |
| EBP / INFRAS | - NIR - Uncertainty analysis | Contract | 2009-2014 |
| Prognos | Allocation of energy data to specific industrial and commercial sectors | Contract | renewed annually |

14 Information on Changes in National Registry

Table 14-1: Changes in the national registry in accordance with §32 decision 15/CMP.1

| Annual Submission Item | Reporting |
|--|---|
| 15/CMP.1 annex II.E paragraph 32.(a): Change of name or contact | No change in the name or contact information of the registry administrator occurred during the reported period. |
| 15/CMP.1 annex II.E paragraph 32.(b): Change of cooperation arrangement | No change of cooperation arrangement occurred during the reported period. |
| 15/CMP.1 annex II.E paragraph 32.(c): Change of the database or the capacity of National Registry | No change to the database or to the capacity of the national registry occurred during the reported period |
| 15/CMP.1 annex II.E paragraph 32.(d): Change of conformance to technical standards | No change in the registry's conformance to technical standards occurred for the reported period |
| 15/CMP.1 annex II.E paragraph 32.(e): Change of discrepancies procedures | No change of discrepancies procedures occurred during the reported period. |
| 15/CMP.1 annex II.E paragraph 32.(f): Change of Security | No change of security measures occurred during the reporting period. |
| 15/CMP.1 annex II.E paragraph 32.(g): Change of list of publicly available information | No change to the list of publicly available information occurred during the reporting period |
| 15/CMP.1 annex II.E paragraph 32.(h): Change of Internet address | No change of the registry Internet address occurred during the reporting period |
| 15/CMP.1 annex II.E paragraph 32.(i): Change of data integrity measures | No change of data integrity measures occurred during the reporting period |
| 15/CMP.1 annex II.E paragraph 32.(j): | No change of test results occurred during the reporting period |

15 Information on Minimization of Adverse Impacts in Accordance with Article 3, Paragraph 14

The Convention (Art. 4 §8 and §10) and its Kyoto Protocol (Art. 2 §3 and Art. 3 §14) commit Parties to strive to implement climate policies and measures in such a way as to minimize adverse economic, social and environmental impacts on developing countries when responding to climate change.

Context

Switzerland strives to design climate change policies and measures in a way as to ensure a balanced distribution of mitigation efforts by implementing climate change response measures in all sectors and for different gases. Indirectly, this approach is deemed to minimize also the scope of potential adverse impacts on concerned actors (including developing countries). Though, due to Switzerland's size and share related to international trade – mainly concentrated on the EU – and greenhouse gas emissions, it is not assumed that Swiss climate change policies have any significant adverse economic, social and environmental impacts in developing countries. Additionally, the policies and measures are very much compatible and consistent with those of the European Union in order to avoid trade distortion, non-tariff barriers to trade and to set similar incentives. All major projects of law in Switzerland are accompanied by impact assessments, inter alia including evaluation of trade-related issues. In accordance with international law, this approach strives at ensuring that Switzerland is implementing those climate change response measures, which are least trade distortive and do not create unnecessary barriers to trade. Consistently, Switzerland notifies all proposed non-tariff measures having a potential impact on trade to the WTO, where specific concerns can be raised by other parties. Moreover, Switzerland belongs to the most important donors in the area of Aid for Trade.

The impact assessment is accompanied by a broad internal and external consultation process, inter alia inviting competent actors to provide advice on international economic, social and environmental aspects of proposed policies and measures. The open public consultation process, together with regular policy dialogues with other countries guarantee that all domestic and foreign stakeholders can raise concerns and issues about new policy initiatives, i.e. including those concerns about possible adverse impacts on other countries.

Progressive reduction or phasing out of market imperfections, fiscal incentives, tax and duty exemptions and subsidies in all greenhouse-gas-emitting sectors, taking into account the need for energy price reforms to reflect market prices and externalities

Environmental policy in Switzerland, including climate change policies, are guided by the "polluter pays" principles, as enshrined in the Federal Law on the Protection of the Environment. Accordingly, the internalization of external costs and adequate price signals are key aspects of Switzerland's climate change policy. Regarding greenhouse gas emissions, market-based instruments, such as the Swiss Emissions Trading Scheme, the supplemental use of Certified Emission Reductions from the Clean Development Mechanism or the levy for heating and process fuels are important measures to put a price on emissions of greenhouse gases (see Sixth National Communication for more details), that are then reflected in market prices and thus internalizing externalities.

Fiscal incentives, tax and duty exemptions and subsidies

Price-based measures are recognized as essential instruments for promoting the efficient use of resources and to reduce market imperfections. In 2001 Switzerland introduced a heavy vehicle fee (HVF). It is applied to passenger and freight transport vehicles of more than 3.5 tonnes gross weight. The impact of the HVF introduction was most clearly reflected by changes in traffic volume (truck-kilometres) but also in reduced air pollution, a renewal of the heavy vehicle fleet and an increase of load per vehicle, fewer trucks having transported more goods. Two thirds of the revenues are used to finance major railway infrastructure projects (such as the two base tunnels through the Alps), and one third is transferred to the cantons.

In 2008 Switzerland introduced a CO₂ levy on heating and process fuel to set an incentive for a more efficient use of fossil fuels, promote investment in energy-efficient technologies and the use of low-carbon or carbon-free energy sources. The 2013 amendment to the CO₂ act (Swiss Confederation 2012) still encompasses the imposition of a CO₂ levy on heating and process fuel. Companies, especially those industries with substantial CO₂ emissions from use of heating fuels, may apply for exemption from the CO₂ levy, provided the company commits to emission reductions. The company has to elaborate an emission reduction target, based on the technological potential and economic viability of various measures within the company. While the proceeds from the CO₂ levy were initially to be fully and equally refunded to the Swiss population and to the business community in proportion of wages paid, a parliamentary decision of June 2009 earmarked a third of the revenues from the CO₂ levy to CO₂ relevant measures in the building sector (Building refurbishment programme). The partial earmarking of revenues from the CO₂ tax is limited in the revised CO₂ act to a maximum of 300 million Swiss francs per year.

The economic impact of the Swiss climate policy was analysed in two studies¹⁸. The impact is considered to be very small.

Switzerland does not subsidize fossil fuels in general. There are some minor schemes in place though that may be regarded as fossil fuel subsidies. In international comparison, however, these schemes are limited: At the federal level, a few tax exemptions and reductions provide some form of support to users of fossil fuels. Farmers, foresters, fishermen and the fuel use of snow cats are exempt from the mineral oil tax that is normally levied on sales of mineral oils, while public transport companies benefit from a reduced rate. Some vehicles are also exempt from the performance-related Heavy Vehicle Fee (HFV), e.g. agricultural vehicles, vehicles used for the concessionary transport of persons or vehicles for police, fire brigade, oil and chemical emergency unit, civil protection and ambulances.

The need for energy prices reforms

World-wide subsidies for fossil fuels are estimated at 300-500 billion USD per annum, depending on the level of energy prices. This huge market distortion does not only produce severe fiscal problems for the countries concerned, it is also a major obstacle for enhanced investments in energy efficiency measures and renewable energies.

Switzerland as a member of the Friends of Fossil Fuels Subsidies Reform group supports the gradual and sustained reduction of unnecessary market-distortions. Switzerland under its Economic Development Cooperation supports partner countries in the design and implementation of energy tariff reforms, as an element of infrastructure financing programs. Switzerland has been an initiator of specialized international programs, including the World Bank's

¹⁸ Ecoplan (2009): Volkswirtschaftliche Auswirkungen der Schweizer Post-Kyoto-Politik, im Auftrag des BAFU. BAFU (2010): Synthesebericht zur Volkswirtschaftlichen Beurteilung der Schweizer Klimapolitik nach 2012.

Energy Sector Management Program ESMAP. The Energy Efficiency Governance Handbook has been produced with Swiss financing (IEA/EBRD 2010).

Removing subsidies associated with the use of environmentally unsound and unsafe technologies

Switzerland doesn't subsidize the use of environmentally unsound and unsafe technologies.

Strengthening the capacity of developing country Parties for improving efficiency in upstream and downstream activities relating to fossil fuels, taking into consideration the need to improve the environmental efficiency of these activities

Switzerland supports through different projects the enhancement of efficiency in industrial production, i.e. "cleaner production". These cleaner production projects promote eco-efficient means of production and better working conditions attained through technical improvements and behavioural changes in both management and staff in industrial companies and services. The resulting rise of economic and environmental efficiency and improved competitiveness is gained through the systematic optimisation of energy use, processing of raw material, more efficient use of resources and thus better protection of the environment.

Furthermore, there is a rising awareness and demand by consumers for environmentally sound products. In order to alleviate potential adverse economic impacts of corresponding national measures Switzerland promotes and supports the development of international standards, especially with regard to the sustainable use of natural resources (including agricultural commodities), e.g. through the creation of sustainability standards, financial incentives and favourable framework conditions in developing countries by consultancy services and technology transfer. Further information is contained in Chapter 7 of Switzerland's Sixth National Communication (FOEN 2014d).

Assisting developing country Parties which are highly dependent on the export and consumption of fossil fuels in diversifying their economies

Most developing and transition countries have, in recent years, taken important steps towards trade liberalisation, in order to align their trade policies with multilateral trade agreements. The Swiss State Secretariat for Economic Affairs (SECO) supports these efforts, because a multilaterally acknowledged and respected set of regulations for international transactions not only strengthens trade as such, but also creates more potent and legally secure markets to the benefit of all players.

The measures taken by SECO are aimed at creating the necessary conditions for earning additional income in the beneficiary countries and thereby contribute directly to the alleviation of poverty. SECO is focusing on three areas of intervention along the value chain: (i) International competitiveness (ii) Enabling framework conditions for trade (iii) Improving market access.

For example market access: Trade between developing and industrial countries is still insufficiently developed respectively not diversified enough. On one hand, the developing countries lack the necessary production capacities, transport infrastructure and know-how; on the other hand, tariff and non-tariff barriers to trade make direct access to markets more difficult.

Switzerland promotes access to Swiss markets by granting preferential tariffs on products from developing and emerging countries. In addition, SECO runs programmes for promoting imports to Switzerland and the rest of Europe. The easing of market entry for products from disadvantaged countries is an important contribution to the promotion and diversification of trade, the increase of export revenues and thus to the economic development of the partner countries. Switzerland supports developing and transition countries in the following areas:

- Generalized system of preferences (GSP)
- Swiss Import Promotion Program (www.sippo.ch)
- Development of new private voluntary social and environmental standards based on international multi-stakeholder approaches: private sustainability standards Better Cotton, 4C (Common Code for the Coffee Community), Roundtable for Sustainable Biofuels, etc.

Finally, Switzerland is a strong supporter of the EITI (Extractive Industries Transparency Initiative). We share a belief that the rational use of natural resource wealth is an important driving force for sustainable economic growth that contributes to sustainable development and poverty reduction. The sustainable management of natural resource wealth – as supported by EITI principle and criteria incl. regular publication and audit of revenues – is key to mobilize the funds for diversification strategies.

Changes compared to the latest submission

The reference regarding capacity-building and technology transfer has been updated and refers now to the 6th National Communication. Some minor editorial changes and clarifications have been done.

16 Other Information

This Chapter contains Switzerland's response to the Saturday Paper (UNFCCC 2013a). Together with this response, Switzerland has also submitted a resubmission of the CRFs to the UNFCCC in November 2013 (FOEN 2013g):

Bonn 07 September 2013

Potential Problems and Further Questions from the ERT formulated in the course of the 2013 review of the greenhouse gas inventories of Switzerland submitted in 2013

For the ERT,

Mr. Ole-Kenneth Nielsen, Lead Reviewer Ms. Medea Inashvili, Lead Reviewer

Inventory related potential problems

With reference to the Guidelines for review under Article 8 of the Kyoto Protocol, the ERT requests that additional information and/or revised estimates for the 2011 greenhouse gas (GHG) inventory corresponding to the potential problems identified in this paper (see attached tables) be forwarded to the ERT, through the UNFCCC secretariat, not later than by 21 October 2013.

Should Switzerland decide to submit by 21 October 2013, in response to some or all potential problems, revised estimates of its GHG emissions, the ERT requests that the revised estimates contain the following:

- Relevant background information and a descriptive summary of the revisions made by Switzerland in its 2013 inventory submission with respect to CO₂ emissions from fuel combustion, N₂O emissions from public electricity and heat production, CH₄ emissions from other energy industries, CH₄ and N₂O emissions from residential plants, fugitive CO₂ and CH₄ emissions from oil transport, all from the energy sector and HFC emissions from refrigeration and air conditioning equipment from the industrial processes sector;
- A complete resubmission of the 2013 CRF-tables, reflecting the revised estimates;
- Switzerland's revision of the calculation of the commitment period reserve, based on the recalculated emissions reported for 2011, if the calculation of the commitment period reserve is based on the inventory and not the assigned amount.

ATTACHMENT A

Overview of inventory potential problems identified for 2008-2011

Annex A sources

2013 GHG inventory review

Switzerland

Abbreviations:

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|---|-----------------|-----------------|---|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 1. Energy, 1.A Fuel combustion | CO ₂ | Level and trend | | X | |
| Description of problem identified: During the review the ERT identified that the CO ₂ EF used for natural gas in Switzerland (55 t CO ₂ /TJ) was lower than that of all neighbouring countries as well as lower than the IPCC default value (56.1 t CO ₂ /TJ). The ERT required further information regarding the source of the EF that in the NIR was presented as a country-specific EF. Switzerland informed the ERT that the EF originally came from the 1992 version of the CORINAIR Guidebook. Considering that the EF was lower than the IPCC default and lower than all neighbouring countries and that the EF was not country-specific and backed up by documentation the ERT concluded that the estimate of CO ₂ emissions from natural gas combustion in Switzerland is not in line with the IPCC GPG. | | | | | |
| Recommendation by ERT: The ERT recommends that Switzerland: <ul style="list-style-type: none"> • Use a country-specific EF. • If a country-specific EF is not available, the default value from the Revised 1996 IPCC Guidelines can be used. • Recalculate emissions of CO₂ from natural gas combustion for 2008-2011 • Provide the ERT with the reference of the EF used in the recalculation | | | | | |

Response / Information by Party:

Emissions from natural gas combustion have been recalculated for the entire time period 1990-2011, using the CO₂-EF of the 2006 IPCC guidelines of 56.1 t/TJ. The CO₂-EF is the same as in the revised 1996 IPCC guidelines (15.3 tC/TJ).

| CO ₂ emissions Sector 1A [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|-----------|-----------|-----------|-----------|-----------|
| Submission April 2013 | 3'731'116 | 6'448'315 | 6'186'794 | 6'909'500 | 6'129'790 |
| Submission September 2013 Saturdaypaper | 3'805'738 | 6'577'281 | 6'310'530 | 7'047'690 | 6'252'386 |

Potential problem unsolved? Rationale:

The ERT considered the response of Switzerland with regard to the potential underestimation of CO₂ emissions from natural gas combustion. Switzerland has revised the EF for natural gas combustion using the default EF from the IPCC guidelines. Estimations are performed in accordance to IPCC methodologies (Revised 1996 IPCC guidelines). The ERT concluded that this potential problem has been resolved in the course of the review.

Overview of inventory potential problems identified for 2009-2011**Annex A sources****2013 GHG inventory review****Switzerland****Abbreviations:**

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|---|------------------|-------------|---|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 1.Energy, 1.A.1 Energy industries, 1.A.1.a Public electricity and heat production | N ₂ O | non-KC | | X | |
| Description of problem identified: During the review the ERT raised a question regarding the N ₂ O IEF from waste incineration. In response, Switzerland informed the ERT that an error had been detected for the years 2009-2011. By mistake the N ₂ O EFs for fossil and biogenic municipal waste incineration are not identical for the year 2009 and onward. The N ₂ O EF used for biogenic waste is lower than the real EF. | | | | | |
| Recommendation by ERT: The ERT recommends that Switzerland: <ul style="list-style-type: none"> • Correct the N₂O EF used for biogenic municipal waste incineration • Recalculate emissions of N₂O from municipal waste incineration for 2009-2011 | | | | | |

Response / Information by Party:

The N₂O EF has been corrected so that for both the biogenic and the fossil fraction the same EF is used. This correction is reflected in the revised estimates of the N₂O emissions of municipal waste incineration in 1A1a Public electricity and heat production Biomass.

| N ₂ O emissions 1A1a Biomass [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|---|------|------|------|------|------|
| Submission April 2013 | 89 | 179 | 158 | 150 | 137 |
| Submission September 2013 Saturdaypaper | 89 | 179 | 165 | 155 | 141 |

Potential problem unsolved? Rationale:

The ERT considered the response of Switzerland with regard to the potential underestimation of N₂O emissions from biogenic waste combustion. Switzerland has revised the EF for biogenic waste combustion correcting the error identified during the review. The recalculation is performed in accordance with IPCC good practice guidance. The ERT concluded that this potential problem has been resolved in the course of the review.

Overview of inventory potential problems identified for 2011**Annex A sources****2013 GHG inventory review****Switzerland****Abbreviations:**

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|--|-----------------|---------------|---|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 1.Energy, 1.A.1 Energy industries, 1.A.1.c Other energy industries | CH ₄ | non-KC | X | | |
| Description of problem identified: In the NIR, Switzerland states that emissions from charcoal production are not included in the inventory. The Revised 1996 IPCC Guidelines contain a default CH ₄ EF for charcoal production. Also, the ERT noted that the charcoal production as reported in the NIR for 2011 (0.11 Gg) is significantly lower than the data available through the FAO statistical database (5.0 Gg). | | | | | |
| Recommendation by ERT: The ERT recommends that Switzerland: <ul style="list-style-type: none"> • Provide explanations of any discrepancies between the activity data reported in the NIR and the FAO data. • Estimate emissions of CH₄ from charcoal production. • Report the activity data and EFs used and the reference for these. | | | | | |

Response / Information by Party:

The activity data on charcoal production (table 4-25 in the NIR) used for estimating emissions from charcoal production are based on enquiries with the charcoal producers directly. They are considered complete.

The FAO data seem to be a very inhomogeneous time series: Constant production of 5Gg for 1999-2011, 0 Gg for 1986-1993, and values between 4Gg and 16Gg in between. Enquiries with the Swiss Federal Statistical Office, the Federal Office for Agriculture, the Swiss Federal Office of Energy and the Forest Division of the Federal Office for the Environment all resulted in the same answer, that no charcoal production in Switzerland is reported. An enquiry with the FAO as to how their data on charcoal production for Switzerland were derived is pending. However, according to the "Joint Wood Energy Enquiry JWEE", a database hosted by UNECE and FAO, charcoal production was reported 0 Gg. It seems as if FAO databases contain contradicting information.

Activity data was reported before in 2D3 (see also chapter 4.5.2.3 "Charcoal production (2D3)" of the NIR 2013). It is now transferred to 1A1c. CH₄ emissions from charcoal production were estimated based on the activity data provided in the NIR table 4-25 and the default CH₄-EF of the revised 1996 guidelines (1tCH₄/TJ).

| CH ₄ emissions charcoal production 1A1c Biomass [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|------|------|------|------|------|
| Submission April 2013 | - | - | - | - | - |
| Submission September 2013 Saturdaypaper | 1 | 3 | 3 | 3 | 3 |

Potential problem unsolved? Rationale:

The ERT considered the response of Switzerland with regard to the potential underestimation of CH₄ emissions from charcoal production. Switzerland has estimated CH₄ emissions using the activity data reported in the NIR combined with the default EF from the revised 1996 IPCC guidelines. The ERT considers that the activity data used by Switzerland are appropriate. However, the ERT recommends that Switzerland continues the communication with FAO to revise the production data in the FAO database. The recalculation is performed in accordance with IPCC good practice guidance. The ERT concluded that this potential problem has been resolved in the course of the review.

Overview of inventory potential problems identified for 2008-2011**Annex A sources****2013 GHG inventory review****Switzerland****Abbreviations:**

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|--|--------------------------------------|-------------|---|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 1.Energy, 1.A.4 Other sectors, 1.A.4.b Residential | CH ₄ and N ₂ O | non-KC | X | | |
| Description of problem identified: In connection with the issue related to charcoal production, the ERT enquired whether emissions from charcoal use were included in the emission inventory. Switzerland informed the ERT that emissions from charcoal use were not included. In the FAO statistical database the consumption of charcoal is available in case national statistics do not provide this information. | | | | | |
| Recommendation by ERT: The ERT recommends that Switzerland: <ul style="list-style-type: none"> • Estimate emissions of CH₄ and N₂O from charcoal use. • Provide explanations to any discrepancies between the activity data chosen for the emission estimation and the FAO data. • Report the activity data and EFs used and the reference for these. | | | | | |

Response / Information by Party:

Activity data for estimating emissions from charcoal combustion are based on production, import and export data. Production data are listed in table 4-25 of the NIR (see also above). Import and export data is based on the Swiss energy balance. Comparing the time series of the Swiss energy balance with the data of FAO provides similar numbers in terms of import for several years (of the order of 300-350 TJ), however, FAO data seems rather inhomogeneous, with production varying between 300 TJ and 1000 TJ. As discussed above, the data source for FAO data is currently unclear, while the energy balance is based on import/export statistics of the Federal Customs administration, which is considered a more reliable source.

Emission factors for CH₄ and N₂O are based on the 2006 IPCC guidelines (online version), being the same as those of the revised 1996 IPCC guidelines. CH₄-EF: 200 kg/TJ N₂O-EF: 1 kg/TJ.

| CH ₄ emissions charcoal use 1A4b [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|---|------|------|------|------|------|
| Submission April 2013 | - | - | - | - | - |
| Submission September 2013 Saturdaypaper | 62 | 71 | 69 | 69 | 69 |

| N ₂ O emissions charcoal use 1A4b [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|------|------|------|------|------|
| Submission April 2013 | - | - | - | - | - |
| Submission September 2013 Saturdaypaper | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |

Potential problem unsolved? Rationale:

The ERT considered the response of Switzerland with regard to the potential underestimation of CH₄ and N₂O emissions from charcoal combustion. Switzerland has estimated the emissions using charcoal production data and data for import and export combined with the default EFs from the IPCC guidelines. The recalculation is performed in accordance with IPCC good practice guidance. The ERT concluded that this potential problem has been resolved in the course of the review.

Overview of inventory potential problems identified for 2011**Annex A sources****2013 GHG inventory review****Switzerland****Abbreviations:**

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|--|-------------------------------------|-------------|---|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 1.Energy, 1.B.2 Fugitive emissions from oil and natural gas, 1.B.2.a iii Oil transport | CO ₂ and CH ₄ | non-KC | X | | |
| Description of problem identified: Switzerland is reporting activity data and emissions from oil transport as not occurring (NO). However, due to oil refining activities in Switzerland the transport of crude oil does occur. The previous ERT recommended that Switzerland provide verifiable information that emissions from oil transport do not occur in Switzerland. The current ERT raised a question on this matter during the review. In response, Switzerland acknowledged that the emissions occur and indicated that they would be estimated and reported for the 2014 annual submission. | | | | | |
| Recommendation by ERT: The ERT recommends that Switzerland: <ul style="list-style-type: none"> • Estimate emissions of CO₂ and CH₄ from oil transport using the imported amount of crude oil as activity data and either country-specific EFs or IPCC default values. • Report the activity data and EFs used and the reference for these. | | | | | |

Response / Information by Party:

The emissions are estimated based on the default EF provided in Table 4.2.4 of the 2006 IPCC reporting guidelines (CH₄ EF 5.4E-6 Gg/1000m³, CO₂ EF 4.9E-7Gg/1000m³). The density of the crude oil is taken as 0.82 t/m³, based on the annual statistics of the Swiss Petroleum Association (EV, 2012). Activity data are crude oil use according to CRF-table 1.B.2.a.

| CH ₄ emissions from oil transport [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|------|------|------|------|------|
| Submission April 2013 | NO | NO | NO | NO | NO |
| Submission September 2013 Saturdaypaper | 21 | 33 | 31 | 30 | 29 |

| CO ₂ emissions from oil transport [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|------|------|------|------|------|
| Submission April 2013 | NO | NO | NO | NO | NO |
| Submission September 2013 Saturdaypaper | 2 | 3 | 3 | 3 | 3 |

Potential problem unsolved? Rationale:

The ERT considered the response of Switzerland with regard to the potential underestimation of CO₂ and CH₄ emissions from oil transport. Switzerland has estimated the emissions using the amount of crude oil transported combined with CO₂ and CH₄ EFs from the 2006 IPCC guidelines. The recalculation is performed in accordance with IPCC good practice guidance. The ERT concluded that this potential problem has been resolved in the course of the review.

Overview of inventory potential problems identified for 2008-2011**Annex A sources****2013 GHG inventory review****Switzerland****Abbreviations:**

GPG: IPCC good practice guidance

AD: activity data, EF: emission factor, IEF: implied emission factor

KC: key category, ERT: Expert Review Team

| Sector, category, sub-category (with code) | Gas | KC / non-KC | Identified inventory problem in terms of: | | |
|--|------|-----------------|--|--|--|
| | | | Missing estimate | Estimate provided but not in line with GPG | Estimate provided but lack of transparency |
| 2.Industrial processes, 2.F Consumption of halocarbons and SF ₆ , 2.F.1 refrigeration and air-conditioning equipment | HFCs | Level and Trend | X (estimate existing, but not included in CRF) | | |
| Description of problem identified: Switzerland uses a tier 2 approach in line with the IPCC GPG by modelling emissions from refrigeration and air conditioning using national statistics and industry data. Activity data and country-specific emission factors used are provided by industry or through expert estimates. In the 2013 NIR, Switzerland states that in the data files used for calculating the emissions from mobile air-conditioning / buses in category 2.F.1 an error has occurred which results in emissions related to the equipment type mobile air-conditioning / buses not being taken into account in the total emission figure of the category 2.F.1. The Party estimates emissions from mobile air conditioning in buses to be between 25-28 Gg CO ₂ -eq annually. This leads to an underestimation of approximately 2 per cent for this category for the years 2008-2011. As this error was discovered very late in the inventory compilation process, the Party was not able to correct the data files in time for the 2013 annual submission. The NIR indicates that the correction of the calculation model is planned for the 2014 annual submission. | | | | | |
| Recommendation by ERT: The ERT recommends Switzerland to submit revised emission data for category 2.F.1 by including emission estimates from air conditioning in buses. | | | | | |

Response / Information by Party:

The emissions from mobile air-conditioning have been recalculated, including emissions from buses. The recalculation results in additional emissions of HFC 134a of the order of 25-28 Gg CO₂eq per year for the years 2008-2011.

| HFC 134a Emission Sector 2 [t] | 1990 | 2008 | 2009 | 2010 | 2011 |
|--|------|--------|--------|--------|--------|
| Submission April 2013 | 0.02 | 354.61 | 353.99 | 384.10 | 451.16 |
| Submission September 2013 Saturdaypaper | 0.02 | 375.34 | 374.20 | 403.26 | 472.44 |

Potential problem unsolved? Rationale:

The ERT considered the data resubmitted by Switzerland and concluded that this potential problem has been resolved in the course of the review.

References

Abegg, M., Speich, S., Brändli, U.-B., Lanz, A., Meile, R., Rösler, E., 2012: Fourth national forest inventory - Results of the NFI4 2009-2011 (LFI4a). Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.

<http://www.lfi.ch/resultate/aktuell.php> [23.03.2014]

ACW/ART 1998: Bestimmung des organisch gebundenen Kohlenstoffs (Corg). [Determination of soil organic carbon]. Referenzmethoden der Forschungsanstalten Agroscope. Band 1: Bodenuntersuchung zur Düngeberatung. Agroscope Changins-Wädenswil Research Station, Nyon & Wädenswil and Agroscope Reckenholz-Tänikon Research Station, Zürich. Version 01.02.1998.

www.bafu.admin.ch/ghginv-ref

Agrammon 2010: The Swiss ammonia model, Federal Office for the Environment FOEN. Internet version in German, French and English:

<http://www.agrammon.ch/documents-to-download/> [28.02.2014]

Technical description for download (updated periodically):

<http://www.agrammon.ch/technical-description> [28.02.2014]

Agricura 2012: Geschäftsbericht 2011/2012. Agricura, Bern.

Agroscope 2014: Treibhausgasinventar Sektor Landwirtschaft, Tabellen zur Berechnung der Treibhausgasemissionen der schweizerischen Landwirtschaft. Internes Dokument. [GHG inventory, sector Agriculture. Spreadsheets to calculate the GHG emissions of Switzerland's agriculture. Internal document]. Agroscope Reckenholz-Tänikon Research Station, Zürich.

www.bafu.admin.ch/ghginv-ref

Agroscope 2014a: Lime application in Swiss Agriculture. Internal documentation by Bretscher, D., Agroscope Reckenholz-Tänikon Research Station, Zürich.

www.bafu.admin.ch/ghginv-ref

Alberti, G., Peressotti, A., Piussi, P., Zerbi, G. 2008: Forest ecosystem carbon accumulation during a secondary succession in the Eastern Prealps of Italy. *Forestry* 81: 1-11.

<http://dx.doi.org/10.1093/forestry/cpm026>

Alcan 2002: Written communication from Kurt Buxmann (ALCAN) to Carbotech (confidential), 30.1.2002.

Alcan 2003: Written communication from François Veuthey (ALCAN) to Carbotech (confidential).

Amon, B., Amon, T., Boxberger, J., Alt, C. 2001: Emissions of NH₃, N₂O and CH₄ from dairy cows housed in a farmyard manure tying stall (housing, manure storage, manure spreading). *Nutrient Cycling in Agroecosystems* 60 (1-3): 103-113.

Andreani-Aksoyoglu, S., Keller, J. 1995: Estimates of monoterpene and isoprene emissions from the forests in Switzerland. *Journal of Atmospheric Chemistry* 20: 71-87.

<http://dx.doi.org/10.1007/BF01099919>

ARE 2002: Fahrleistungen der Schweizer Fahrzeuge. Ergebnisse der periodischen Erhebung Fahrleistungen (PEFA) 2000. Federal Office for Spatial Development, Bern.

www.bafu.admin.ch/ghginv-ref

ARE 2010: Nationales Personenverkehrsmodell – Basismodell 2005, Bern.

http://www.are.admin.ch/dienstleistungen/00906/index.html?lang=de&download=NHZLpZeg7t_lnp6l0NTU042l2Z6ln1acy4Zn4Z2qZpnO2YUq2Z6gpJCEdYN5gmym162epYbq2c_JiKbNoKSn6A-- [28.02.2014]

ARE 2012: Ergänzungen zu den schweizerischen Verkehrsperspektiven bis 2030. Bundesamt für Raumentwicklung (ARE). Bern
<http://www.are.admin.ch/dokumentation/publikationen/00015/00471/index.html?lang=de>
[19.02.2014]

ARE/SAEFL 2001: Le paysage sous pression, suite 2. [Landschaft unter Druck, 2. Fortschreibung]. Federal Office for Spatial Development, Bern and Swiss Agency for the Environment, Forests and Landscape, Bern.
<http://www.are.admin.ch/themen/raumplanung/00246/03637/index.html?lang=fr> [28.02.2014]

ARE/SFSO 2005: Mobilität in der Schweiz. Ergebnisse des Mikrozensus 2005 zum Verkehrsverhalten. [La mobilité en Suisse. Résultats du microrecensement 2005 sur le comportement de la population en matière de transports]. Federal Office for Spatial Development, Bern and Swiss Federal Statistical Office, Neuchâtel.
<http://www.bfs.admin.ch/bfs/portal/de/index.Document.91826.pdf> [German] [28.02.2014]
<http://www.bfs.admin.ch/bfs/portal/fr/index.Document.91826.pdf> [French] [28.02.2014]

ART 2008a: Uncertainty in agricultural CH₄ and N₂O emissions of Switzerland. Internal documentation by Bretscher, D. and Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich.
www.bafu.admin.ch/ghginv-ref

ART 2009b: Emission factor drained peatlands Switzerland – A brief analysis of recent studies and comparison to EF used in the Swiss GHG Inventory. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich.
www.bafu.admin.ch/ghginv-ref

ART 2011a: Summary of the available published data on root biomass and root carbon in Swiss grasslands. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich.
www.bafu.admin.ch/ghginv-ref

ART 2011b: First estimate on CO₂ emission factor of organic soils under unproductive wetland. Internal documentation by Leifeld, J., Agroscope Reckenholz-Tänikon Research Station, Zürich.
www.bafu.admin.ch/ghginv-ref

ART 2013a: Agricultural CH₄ and N₂O emissions in Switzerland: QA/QC. Internal documentation (with continual update) by Bretscher, D., Agroscope Reckenholz-Tänikon Research Station, Zürich.
www.bafu.admin.ch/ghginv-ref

ART 2012a: Carbon in Living Biomass and Dead Organic Matter in Croplands and Grasslands: Current state of the Swiss GHG-Inventory. Internal documentation by Bretscher, D., Agroscope Reckenholz-Tänikon Research Station, Zürich.

ART/SHL 2012: Categorization of livestock animals in Switzerland. D. Bretscher and T. Kupper. Agroscope Research Station Zürich (ART), Schweizerische Hochschule für Landwirtschaft Zollikofen (SHL). March 2012.

Bader, M.K., Leuzinger, S., Keel, S.G., Siegwolf, T.W., Hagedorn, F., Schleppi, P., Körner, C. 2013: Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project. *Journal of Ecology* 101 (6): 1509–1519.
<http://onlinelibrary.wiley.com/doi/10.1111/1365-2745.12149/abstract> [04.04.2014]

Battelle 1994: Methanfreisetzung bei der Erdgasnutzung in der Schweiz und Vergleich mit anderen Emittenten. Studie im Auftrag des Schweizerischen Vereins des Gas- und Wasserfaches SVGW. Battelle Ingenieurtechnik, GmbH, Eschborn.
www.bafu.admin.ch/ghginv-ref

Berthoud, F. 2004: Dokumentation der Methan- & Lachgastabelle. Eine Hilfeleistung zum Verstehen der Berechnungen und Berechnungsgrundlagen der landwirtschaftlichen Treibhausgasemissionen hin zu den Resultatwerten des Common Reporting Format des IPCC. Internal documentation. Agroscope FAL, Swiss Federal Research Station for Agroecology and Agriculture.

www.bafu.admin.ch/ghginv-ref

Brändli, U.-B. (Red.) 2010: Schweizerisches Landesforstinventar. Ergebnisse der dritten Aufnahme 2004-2006. [Results of the third Swiss national forest inventory 2004-2006]. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft, Birmensdorf. Bundesamt für Umwelt, Bern, 312 S.

<http://www.lfi.ch/publikationen/publ/lfi3-fr.php> [18.03.2014]

Braendli, U.-B., Speich, S. 2011: Das LFI4– eine kontinuierliche Erhebung. LFI-News 13.

<http://www.lfi.ch/publikationen/publ/lfi-info/LFIinfo13.pdf> [04.04.2014]

Brassel, P., Brändli, U.-B. 1999: Schweizerisches Landesforstinventar. Ergebnisse der Zweitaufnahme 1993-1995. [Results of the second Swiss national forest inventory 1993-1995]. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft, Birmensdorf. Bundesamt für Umwelt, Wald und Landschaft, Bern. Haupt, Bern, Stuttgart, Wien. [available in German, French and Italian]

<http://www.lfi.ch/publikationen/publ/lfi2-en.php> [18.03.2014]

Braun, M., Hurni, P., Spiess, E. 1994: Phosphor- und Stickstoffüberschüsse in der Landwirtschaft und Para-Landwirtschaft : Abschätzung für die Schweiz und das Rheineinzugsgebiet der Schweiz unterhalb der Seen. [Surplus de phosphore et d'azote dans l'agriculture et la para-agriculture: estimation pour la Suisse et pour le bassin versant hydrographique suisse du Rhin en aval des lacs]. Schriftenreihe der FAC Liebefeld 18. [in German, with English and French summary]

Braun, S., Thomas, V.F., Quiring, R., Flückiger, W. 2010: Does nitrogen deposition increase forest production? The role of phosphorus. Environmental Pollution 158(6): 2043-2052.

<http://dx.doi.org/10.1016/j.envpol.2009.11.030> **Burschel, P., Kürsten, E., Larson, B.C.**

1993: Die Rolle von Wald und Forstwirtschaft im Kohlenstoffhaushalt. Eine Betrachtung für Deutschland. Forstliche Forschungsberichte München 126.

BUS 1986: Leitbild für die Schweizerische Abfallwirtschaft. [Lignes directrices pour la gestion des déchets en Suisse]. Schriftenreihe Umweltschutz Nr. 51. Bundesamt für Umweltschutz BUS. [Les cahiers de l'environnement No 51. Office fédéral de la protection de l'environnement], Bern.

www.bafu.admin.ch/ghginv-ref

Carbotech 2014: Swiss Greenhouse Gas Inventory 2012: PFCs, HFCs and SF₆ Emissions. Confidential report no. 251.14. For internal use on behalf of the Federal Office for the Environment, Bern. Basel.

CARBURA 2010: Bunkers of diesel oil 1990-2009. Written communication from Matthias Rufer (CARBURA) to Jan Landert (INFRAS), 17.12.2010.

Cemsuisse 2010a: Bestimmung der Emissionsfaktoren an Mischproben, Interner Bericht von Wessling im Auftrag von cemsuisse, 2010

Cemsuisse 2013: Jahresbericht 2012. [Rapport annuel 2012]. Association of the Swiss Cement Industry, Bern.

<http://www.cemsuisse.ch/cemsuisse/ueberuns/publikationen/jahresberichte/index.html?lang=de> [German and French] [25.02.2014]

CH2011 2011: Swiss Climate Change Scenarios CH2011. Published by C2SM, MeteoSwiss, ETH, NCCR Climate, and OcCC, Zurich, Switzerland: 1-88.

<http://www.ch2011.ch/> [27.01.2014]

- Covington, W.W. 1981:** Changes in forest floor organic matter and nutrient content following clear cutting in northern hardwoods. *Ecology* 62(1): 41-48.
<http://www.jstor.org/pss/1936666> [28.02.2013]
- Cronan, C.S. 2003:** Belowground biomass, production, and carbon cycling in mature Norway spruce, Maine, U.S.A. *Canadian Journal of Forest Research* 33: 339-350.
<http://dx.doi.org/10.1139/x02-189>
- CSD 2013:** Swiss Greenhouse Gas Inventory Submission 2012. National Review of Sector Industrial Processes. On behalf of the Federal Office for the Environment, Bern. Basel
www.bafu.admin.ch/ghginv-ref
- Davidson, E.A., Janssens, I.A. 2006:** Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440: 165-173.
<http://dx.doi.org/10.1038/nature04514>
- de Keizer, C., Ramirez, A., van der Sluijs, J. 2007:** Uncertainty Ranges and Correlations Assumed in Tier 2 Studies of Several European Countries. In: *Proceedings of the 2nd International Workshop on Uncertainties in Greenhouse Gas Inventories*, 27-28 September 2007, Laxenburg, Austria: 35-39.
<http://www.ibspan.waw.pl/ghg2007/GHG-total.pdf> [28.02.2014]
- de Wit, H.A., Palosuo, T., Hysten, G., Liski, J. 2006:** A carbon budget of forest biomass and soils in southeast Norway calculated using a widely applicable method. *Forest Ecology and Management* 225 (1-3): 15-26.
<http://dx.doi.org/10.1016/j.foreco.2005.12.023>
- DelSontro, T., McGinnis, D. F., Sobek, S., Ostrovsky, I., Wehrli, B.:** Extreme methane emissions from a Swiss hydropower reservoir: contribution from bubbling sediments, *Environmental Science & Technology* 44: 2419-2425.
<http://dx.doi.org/10.1021/es9031369>
- Didion, M., Kaufmann, E., Thürig, E. 2012:** Estimation of carbon stocks and stock changes in soil, LFH layer and deadwood in Swiss forests with Yasso07. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
www.bafu.admin.ch/ghginv-ref
- Didion, M., Kaufmann, E., Thürig, E. 2013:** Data on soil carbon stock change, carbon stock and stock change in surface litter and in coarse deadwood prepared for the Swiss GHGI 1990-2012. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
www.bafu.admin.ch/ghginv-ref
- Didion, M. 2014:** Impacts of forest management on carbon stock changes in litter and soil in Swiss forests. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
www.bafu.admin.ch/ghginv-ref
- Didion, M., Rogiers, N., Frey, B., Thürig, E. 2014:** Validating the Yasso07 model for estimating deadwood and litter decomposition in Swiss forests. Submitted to *European Journal of Forest Research*.
- Dobbertin, M., Jüngling, E. 2009:** Totholzverwitterung und C-Gehalt. Zwischenergebnisse. [Wood density and carbon content with changing degree of dead wood decay: First results]. Swiss Federal Research Institute for Forest, Snow and Landscape Research, Birmensdorf.
www.bafu.admin.ch/ghginv-ref
- Düggelin, C., Abegg, M. 2011:** Modelle zur Biomasse- und Holzvolumenschätzung im Schweizer Gebüschwald. *Schweizerische Zeitschrift für Forstwesen* 162 (2): 32-40.
<http://dx.doi.org/10.3188/szf.2011.0032>

EAFV (Eidg. Anstalt für das forstliche Versuchswesen) / BFL (Bundesamt für Forstwesen und Landschaftsschutz) (eds.) 1988: Schweizerisches Landesforstinventar. Ergebnisse der Erstaufnahme 1982-1986. [Results of the first Swiss national forest inventory 1982-1986]. Ber. Eidgenöss. Forsch.anst. Wald Schnee Landsch. 305.
<http://www.lfi.ch/publikationen/publ/lfi1-en/php> [13.03.2013]

EEA 2002: EMEP/CORINAIR Emission Inventory Guidebook. European Environment Agency. 3rd edition October 2002 update.
<http://www.eea.europa.eu/publications/EMEP-CORINAIR3/page002.html> [28.01.2014]

EEA 2007: EMEP/CORINAIR 2007 Emission Inventory Guidebook (December 2007). European Environment Agency. Technical Report No. 16/2007.
<http://www.eea.europa.eu/publications/EMEP-CORINAIR5/page002.html> [28.02.2013]

EEA 2010: EMEP/EEA Emission Inventory Guidebook 2009, updated June 2010, Methodology for the calculation of exhaust emissions – SNAPs 070100-070500, NFRs 1A3bi-iv Passenger cars, light-duty trucks, heavy-duty vehicles including buses and motor cycles.
<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> [28.02.2014]

EMEP/EEA 2013: Air Pollutant Emission Inventory Guidebook 2013, European Environment Agency.
<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> [29.01.2014]

EC 2004: Commission decision 2004/156/EC of 29 January 2004, Establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council (notified under document number C(2004) 130). Official Journal of the European Union L59, 26.2.2004.
<http://rod.eionet.europa.eu/instruments/593> [28.02.2014]

EMIS 2014/(NFR-Code): Comments to EMIS database. Internal documents. Federal Office for the Environment, Bern.
 To find the EMIS comment that belongs to the specified NFR Code, see Table A - 1 (below References) [available in German]

EMPA 1999: Written communication from Dr. H.W. Jäckle (EMPA, Dübendorf) to Andreas Liechti (FOEN, Bern), 09.03.1999.
www.bafu.admin.ch/ghqinv-ref

EMPA 2004: EMPA Research Report 202114b. Swiss Laboratories for Material Science and Technology, Dübendorf.

EMPA 2004a: Baustoffmanagement 21. Stand des Wissens und Forschungsbedarf.
http://www.novatlantia.ch/fileadmin/downloads/projekte/raumundressourcen/Baustoffmanag_21.pdf

Etzold, S., Ruehr, N.K., Zweifel, R., Dobbertin, M., Zingg, A., Pluess, P., Häslar, R., Eugster, W. Buchmann, N. 2011: The Carbon Balance of Two Contrasting Mountain Forest Ecosystems in Switzerland: Similar Annual Trends, but Seasonal Differences. Ecosystems 14: 1289–1309. <http://dx.doi.org/10.1007/s10021-011-9481-3>
http://www.natkon.ch/pdf_files/publikationsseite/Etzold_etal_2011_Ecosystems.pdf [28.02.2013]

EV 2013: Jahresbericht 2012. Erdöl-Vereinigung [Rapports annuel 2012. L'Union Pétrolière]. Zürich.
http://www.erdoel-vereinigung.ch/UserContent/Shop/EV_JB12_DE_20130605.pdf, in German [18.02.2014]

Falloon, P., Smith, P. 2003: Accounting for changes in soil carbon under the Kyoto protocol: need for improved long-term data sets to reduce uncertainty in model projections. *Soil Use and Management* 19(3): 265-269.

<http://dx.doi.org/10.1079/SUM2003201>

FAL/RAC 2001: Grundlagen für die Düngung im Acker- und Futterbau 2001. [Principles of fertilization in crop and feed production]. Eidgenössische Forschungsanstalt für Agrarökologie und Landbau / Eidgenössische Forschungsanstalt für Pflanzenbau, Agrarforschung, June 2001, Zürich-Reckenholz, Nyon. [available in German and French]
www.bafu.admin.ch/ghginv-ref

FEDRO 2010: Swiss Automatic Road Traffic Counts (SARTC). Federal Roads Office, Bern.
<http://www.astra.admin.ch/verkehrsdaten/00299/00301/index.html?lang=en> [28.02.2014]

Fial 2013: Die Schweizer Nahrungsmittel-Industrie im Jahr 2012.
http://fial.ch/de/statistics/fial_Statistik%202012_de.pdf [25.02.2014]

Flisch, R., Sinaj, S., Charles, R., Richner, W. 2009: Grundlagen für die Düngung im Acker- und Futterbau 2009. Forschungsanstalt Agroscope Changins-Wädenswil ACW und Agroscope Reckenholz-Tänikon ART, Agrarforschung 16 (2).

FOCA 1991: Crossair confidential data 1991. Federal Office of Civil Aviation, Bern.

FOCA 1991a: L'aviation civile Suisse en 1990. Federal Office of Civil Aviation, Bern.
www.bafu.admin.ch/ghginv-ref

FOCA 2004: Unternehmensstatistik der Schweizerischen Helikopterunternehmen. Federal Office of Civil Aviation, Bern.
www.bafu.admin.ch/ghginv-ref

FOCA 2006: GHG emissions of Swiss civil aircraft in 1990 and 2004: data, proceeding and description of the calculations. Written communication from Theo Rindlisbacher and Paul Stulz (FOCA, Bern) to Andreas Liechti (FOEN, Bern), 20./22.02.2006.
www.bafu.admin.ch/ghginv-ref

FOCA 2006a: GHG emissions of Swiss civil aircraft in 1990, 1995, 2000, 2002, 2004 and 2005: data, proceeding and description of the calculations. Written communication from Theo Rindlisbacher (FOCA, Bern) to Paul Filliger (FOEN, Bern), 17.11.2006.
www.bafu.admin.ch/ghginv-ref

FOCA 2007: GHG emissions of Swiss civil aircraft in 2006. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 03.12.2007.
www.bafu.admin.ch/ghginv-ref

FOCA 2007a: Aircraft Piston Engine Emissions. Summary Report. Federal Office of Civil Aviation. Report Ref. 0 / 3/33/33-05-003 ECERT. Bern.
www.bafu.admin.ch/ghginv-ref

FOCA 2007b: Validation of ADAECAM (Advanced Aircraft Emission Calculation Method). Report on fuel calculation. Federal Office of Civil Aviation. Report Ref. 0/3/33/33-05-007.021. Bern, 10.08.2007.
www.bafu.admin.ch/ghginv-ref

FOCA 2008: GHG emissions of Swiss civil aircraft in 2007. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 01.12.2008.
www.bafu.admin.ch/ghginv-ref

FOCA 2009: GHG emissions of Swiss civil aircraft in 2008. Written communication from Theo Rindlisbacher (FOCA, Bern) to Beat Müller (FOEN, Bern), 04.11.2009.
www.bafu.admin.ch/ghginv-ref

FOCA 2009a: Guidance on the determination of helicopter emissions. Federal Office of Civil Aviation. Report Ref. 0/3/33/ 33-05-020. Edition 1. Bern. March 2009
www.bafu.admin.ch/ghginv-ref

FOCA 2010: GHG emissions of Swiss civil aircraft in 2009. Written communication from Theo Rindlisbacher (FOCA, Bern) to Sophie Hoehn (FOEN, Bern), 18.11.2010.

www.bafu.admin.ch/ghginv-ref

FOCA 2011: GHG emissions of Swiss civil aircraft in 2010. Written communication from Theo Rindlisbacher (FOCA, Bern) to Sophie Hoehn (FOEN, Bern), 09.11.2011.

www.bafu.admin.ch/ghginv-ref

FOCA 2012: GHG emissions of Swiss civil aircraft in 2011. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 07.09.2012.

www.bafu.admin.ch/ghginv-ref

FOCA 2013: GHG emissions of Swiss civil aircraft in 2012. Written communication from Theo Rindlisbacher (FOCA, Bern) to Anouk-Aimée Bass (FOEN, Bern), 07.11.2013.

www.bafu.admin.ch/ghginv-ref

FOEN 2006b: Switzerland's Greenhouse Gas Inventory 1990–2004, National Inventory Report and CRF-tables 2006. Submission of 31 May 2006 to the United Nations Framework Convention on Climate Change. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/00546/index.html?lang=en>

FOEN 2006c: Prozess EMIS (Luftschadstoff-Emissions-Inventar der Schweiz). Beschrieb des Prozesses (= Handbuch zur EMIS-Datenbank (Entwurf)). Internes Dokument. [Manual to EMIS database (draft). Internal document]. Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

FOEN 2006h: Switzerland's Initial Report under Article 7, paragraph 4 of the Kyoto Protocol. Federal Office for the Environment, Bern. Including the Update following the UNFCCC review (FCCC/IRR/2007/CHE).

<http://www.bafu.admin.ch/climatereporting/03211/index.html?lang=en>

FOEN 2007: Switzerland's Greenhouse Gas Inventory 1990–2005, National Inventory Report and CRF-tables 2007. Submission of 13 April 2007 to the United Nations Framework Convention on Climate Change. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/04333/index.html?lang=en>

FOEN 2007f: Definition der Kombinationskategorien (CC) für die LULUCF-Berichterstattung auf der Basis der AREA-Landnutzungs- und Landbedeckungskategorien. Internes Dokument, Version 2 vom 30.05.2007. [Definition of combination categories (CC) for LULUCF reporting based on AREA land-use/land-cover categories. Internal document, version 2 as of 30.05.2007]. Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

FOEN 2008: Switzerland's Greenhouse Gas Inventory 1990–2006, National Inventory Report and CRF-tables 2008. Submission of 15 April 2008 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/index.html?lang=en> [27.01.2014]

FOEN 2008h: Wood Resource Policy: Strategy, Objectives and Action Plan for the Resource Wood. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/publikationen/publikation/01002/index.html?lang=en>

FOEN 2010: Switzerland's Greenhouse Gas Inventory 1990–2008: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 1999–2008, SEF and SIAR tables from the National Registry. Submission of 15 April 2010 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/10195/index.html?lang=en>

FOEN 2010d: Deforestations in Switzerland as reported under the Kyoto Protocol Art. 3.3. Forest Division, Federal Office for the Environment, Bern.
www.bafu.admin.ch/ghginv-ref

FOEN 2010h: Afforestations in Switzerland as reported under the Kyoto Protocol Art. 3.3. Forest Division, Federal Office for the Environment, Bern.
www.bafu.admin.ch/ghginv-ref

FOEN 2010i: Pollutant Emissions from Road Transport, 1990 to 2035. Updated in 2010. Environmental studies no. 1021. Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/publikationen/publikation/01565/index.html?lang=en> [10.02.2014]

FOEN 2010j: Abfallmengen und Recycling 2009 im Überblick. [Overview on waste quantities and recycling in 2009.] Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/abfall/01517/01519/10457/index.html?lang=de>

FOEN 2011: Switzerland's Greenhouse Gas Inventory 1990–2009: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008-2009, SEF and SIAR tables from the National Registry. Submission of 15 April 2011 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/climatereporting/00545/11269/index.html?lang=en>

FOEN 2011k: CO₂-Emissionsfaktoren des schweizerischen Treibhausgasinventars. Zusammenstellung der CO₂-Emissionsfaktoren und Energieinhalte verschiedener Energieträger, die im Treibhausgasinventar verwendet werden. Federal Office for the Environment, Bern.
www.bafu.admin.ch/ghginv-ref

FOEN 2012: Switzerland's Greenhouse Gas Inventory 1990–2010: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008-2010, SEF and SIAR tables from the National Registry. Submission of 13 April 2012 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/climatereporting/00545/11894/index.html?lang=en>

FOEN 2012b: Anpassung an den Klimawandel in der Schweiz – Ziele, Herausforderungen und Handlungsfelder. Erster Teil der Strategie des Bundesrates. Federal Office for the Environment, Bern. <http://www.bafu.admin.ch/klimaanpassung/11529> [27.01.2014]

FOEN 2012g: Adressliste der Schweizer Kläranlagen mit Angaben zur Ausbaugrösse. Swiss Federal Office of Energy, Bern.
<http://www.bafu.admin.ch/gewaesserschutz/01295/01296/01298/index.html?lang=de>

FOEN 2013: Switzerland's Greenhouse Gas Inventory 1990–2011: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008-2011, SEF and SIAR tables from the National Registry. Submission of 15 April 2013 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/climatereporting/00545/12558/index.html?lang=en>

FOEN 2013g: Switzerland's Greenhouse Gas Inventory 1990–2011. Resubmission in response to the centralized review: CRF-tables, Kyoto Protocol LULUCF table 2008-2011. Submission of 20 September 2013 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/climatereporting/00545/11894/index.html?lang=en>

FOEN 2013h: Abfallmengen und Recycling 2012 im Überblick. [Overview on waste quantities and recycling in 2010.] Federal Office for the Environment, Bern.
<http://www.bafu.admin.ch/abfall/01517/01519/12949/index.html?lang=de>

FOEN 2013i: Hazardous Waste Statistics 2012. Various reports compiled by FOEN on its website.

<http://www.bafu.admin.ch/abfall/01517/01519/12949/index.html?lang=de> [20.02.2014]

FOEN 2013j: Waste Statistics for 2012. Urban and Construction Waste Section, Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref (selected data).

FOEN 2013k: Jahrbuch Wald und Holz 2013. Umwelt-Zustand Nr. 1332. [Annuaire La forêt et le bois 2013. Etat de l'environnement no 1332]. Federal Office for the Environment, Bern.

http://www.bafu.admin.ch/publikationen/publikation/01743/index.html?lang=de&show_kat=/publikationen [German]

http://www.bafu.admin.ch/publikationen/publikation/01743/index.html?lang=fr&show_kat=/publikationen [French]

FOEN 2014: Switzerland's Greenhouse Gas Inventory 1990–2012: National Inventory Report, CRF-tables, Kyoto Protocol LULUCF tables 2008-2012, SEF and SIAR tables from the National Registry. Submission of 15 April 2014 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/13193/index.html?lang=en>

FOEN 2014a: Description of the Quality Management System. Supplement to Switzerland's Greenhouse Gas Inventory 1990-2012. Submission of 15 April 2014 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00545/13193/index.html?lang=en>

FOEN 2014b: Handbuch Berechnung der Wald-Emissionsfaktoren. [Manual: Calculation of EFs for Forest Land]. Including 1 Excel data file. Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

FOEN 2014c: Handbuch Anleitung zum Ausfüllen der Kyoto Tabellen Wald. [Instruction manual for completing data of forest related activities in Kyoto Tables]. Federal Office for the Environment, Bern. www.bafu.admin.ch/ghginv-ref

FOEN 2014d: Switzerland's Sixth National Communication under the UNFCCC; Third National Communication under the Kyoto Protocol to the UNFCCC. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/climatereporting/00551/13139/index.html?lang=en>

FOEN 2014e: Switzerland's Informative Inventory Report 2014 (IIR), Submission under the UNECE Convention on Long-range Transboundary Air Pollution, Submission of March 2014 to the United Nations ECE Secretariat. Federal Office for the Environment, Bern.

<http://www.bafu.admin.ch/luft/11640/11641/11643/index.html?lang=de>

FOEN 2014f: Illustration of unproductive forests in Switzerland. Internal report by the Swiss Federal Office for the Environment.

www.bafu.admin.ch/ghginv-ref

FOEN 2014g: Climate Change in Switzerland: Indicators of driving forces, impact and response. Federal Office for the Environment FOEN, Federal Office of Meteorology and Climatology, Meteoswiss.

<http://www.bafu.admin.ch/publikationen/publikation/01709/index.html?lang=en> [27.01.2014]

Frey, B. 2011: Langfristige Streuabbauexperimente auf LWF-Flächen – Daten zur Validierung von Bodenkohlenstoffmodellierungen im Schweizer Wald. [Testing the Yasso07 model with long term litterbag data from five LTFER sites and two elevation gradients in the Swiss Prealps]. Final report. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.

www.bafu.admin.ch/ghginv-ref

Ginzler, C. 2014: Crown cover and basal area in unproductive forest: special analysis of plots covered by aerial photographs and terrestrial NFI. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.

www.bafu.admin.ch/ghginv-ref

Giuliani, S. 2013: Energiebedarfwerte der Nutztiere für die Periode 1990-2011. Written communication from Silvano Giuliani (SBV, Swiss Farmers Union) to Daniel Bretscher (ART, Reckenholz), 23.8.2013.

www.bafu.admin.ch/ghginv-ref

Gkatzoflias, D., Kouridis, C., Ntziachristos, L. 2012: COPERT 4. Computer programme to calculate emissions from road transport. User manual (version 9.0). European Environment Agency. http://www.emisia.com/docs/COPERT4v9_manual.pdf [27.02.2014]

Grassi, G., Blujdea, V. 2011: Pools to be reported under KP-LULUCF. Harmonized guidance and decision tree on the application of "not a source" principle. Joint Research Centre, Ispra. Presentation at "JRC technical workshop on LULUCF issues under the Kyoto Protocol", 21 November 2011.

Gschwantner, T. 2006: Zuwachsänderungen nach den Daten der Österreichischen Waldinventur und ihre klimatischen Ursachen. [Growth changes according to the data of the Austrian National Forest Inventory and their climatic causes]. BFW-Berichte 133, Wien, Bundesamt für Wald.

Hackl, A., Mauschwitz, G. 2003: Emissionen aus Anlagen der Österreichischen Zementindustrie IV. Jahresreihe 2000-2002. Weitra/Wien.

www.bafu.admin.ch/ghginv-ref

Hadorn, R., Wenk, C. 1996: Effect of different sources of dietary fibre on nutrient and energy utilization in broilers. 2. Energy and N-balance as well as whole body composition. Archiv für Geflügelkunde 60: 22-29.

Hagedorn, F., Martin, M., Rixen, C., Rusch, S., Bebi, P., Zürcher, A., Siegwolf, R.T.W., Wipf, W., Escape, C., Roy, J., Hättenschwiler, S. 2010: Short-term responses of ecosystem carbon fluxes to experimental soil warming at the Swiss alpine treeline. Biogeochemistry: 97: 7–19.

<http://dx.doi.org/10.1007/s10533-009-9297-9>

Hausberger S., Rexeis M., Zallinger M., Luz R. 2009: Emission Factors from the Model PHEM for the HBEFA Version 3.1., Graz; Report Nr. I-20/2009 Haus-Em 33/08/679, 2009.

Helmisaari, H.-S., Hallbäcken, L. 1998: Tree biomass below-ground. In: Andersson, F., Braekke, F.H., Hallbäcken, L. (eds.): Nutrition and growth of Norway spruce forests in a Nordic climatic and deposition gradient, TemaNord 1998: 566, Nordic Council of Ministers, Copenhagen: 80–90.

www.bafu.admin.ch/ghginv-ref

Hiller, R.V., Bretscher, B., DelSontro, T., Diem, T., Eugster, W., Henneberger, R., Hobi, S., Hodson, E., Imer, D., Kreuzer, M., Künzle, T., Merbold, L., Niklaus, P.A., Rihm, B., Schellenberger, A., Schroth, M.H., Schubert, C.J., Sigrist, H., Stieger, J., Buchmann, N., Brunner, D. 2014: Anthropogenic and natural methane fluxes in Switzerland synthesized within a spatially explicit inventory. Biogeosciences 11: 1941-1959.

<http://dx.doi.org/10.5194/bg-11-1941-2014>

Hiller, R.V., Neininger, B., Brunner, D., Gerbig, Ch., Bretscher, D., Künzle, T. 2014a (subm.): Airborne CH₄ flux measurements for validation of emissions from an agriculturally dominated area in Switzerland. Journal of Geophysical Research.

- Hindrichsen, I.K., Wettstein, H.-R., Machmüller, A., Bach Knudsen, K.E., Madsen, J., Kreuzer, M. 2006:** Methane emission, nutrient degradation and nitrogen turnover in dairy cows and their slurry at different milk production scenarios with and without concentrate supplementation. *Agricultural Ecosystems and Environment* 113: 150–161.
- Högbert, P., Fan, H., Quist, M., Binkley, D., Tamm, C.O. 2006:** Tree growth and soil acidification in response to 30 years of experimental nitrogen loading on boreal forest. *Global Change Biology* 12: 489-499.
<http://dx.doi.org/10.1111/j.1365-2486.2006.01102.x>
- Huber, B., Frehner M. 2012:** Forschungsprojekt Grünerle. Bericht erstellt im Auftrag des Bundesamtes für Umwelt (BAFU), Bern. Abenis AG Chur und Forstingenieurbüro Monika Frehner, Sargans.
- Huber, B., Frehner M. 2013:** Die Verbreitung und Entwicklung der Grünerlenbestände in der Ostschweiz. *Schweizerische Zeitschrift für Forstwesen* 164: 87-94.
<http://www.szf-jfs.org/toc/swif/164/4> [04.04.2014]
- Huber, T., Thürig, E. 2014:** Special analysis of stocks of living biomass in non-productive forest (CC13) on terrestrial NFI plots. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf. Commissioned by the Federal Office for the Environment FOEN, Bern.
www.bafu.admin.ch/ghginv-ref
- Hyvönen, R., Persson, T., Andersson, S., Olsson, B. Ågren, G.I., Linder, S. 2008:** Impact of long-term nitrogen addition on carbon stocks in trees and soils in northern Europe. *Biogeochemistry* 89: 121-137.
<http://dx.doi.org/10.1007/s10533-007-9121-3>
- IAI 2005:** Aluminium for future Generations. Sustainability Update 2004. International Aluminium Institute, London.
www.bafu.admin.ch/ghginv-ref
- IEA/EBRD 2010:** Energy Efficiency Governance – Handbook, second edition, IEA/OECD Paris and EBRD, London.
http://www.iea.org/publications/freepublications/publication/gov_handbook-1.pdf [28.02.2014]
- IEA 2005:** Energy statistics manual, IEA/OECD Paris.
http://www.iea.org/publications/freepublications/publication/statistics_manual-1.pdf [25.01.2014]
- IEA 2012:** Energy statistics of OECD countries, 2012 edition, IEA/OECD Paris.
<http://www.iea.org/stats/index.asp> [14.02.2014]
- IFEU/INFRAS 2009:** Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland, Endbericht. IFEU und INFRAS im Auftrag des Umweltbundesamts Dessau/Deutschland, FKZ 360 16 023. Heidelberg/Zürich/Bern. 31. Oktober 2009.
<http://www.umweltdaten.de/publikationen/fpdf-l/3937.pdf> [28.02.2014]
- INFRAS 2004:** Emission Factors for Passenger Cars and Light-Duty Vehicles. Handbook Emission Factors for Road Transport (HBEFA), Version 2.1. Swiss Agency for the Environment, Forests and Landscape, Bern, Umweltbundesamt, Berlin, Umweltbundesamt, Wien. <http://www.hbefa.net/e/index.html> [28.02.2014]

INFRAS 2008: Treibstoffverbrauch und Schadstoffemissionen des Offroad-Sektors. Studie für die Jahre 1980–2020. [Offroad fuel consumption and pollutant emissions in Switzerland. Study for the period from 1980 to 2020]. Im Auftrag des Bundesamts für Umwelt. [On behalf of the Federal Office for the Environment]. Umwelt-Wissen Nr. 0828. Bern.

<http://www.bafu.admin.ch/publikationen/publikation/01003/index.html?lang=de> [28.02.2014]

[in German with English, French and Italian abstract]

Offroad database [in English, French, German]:

<http://www.bafu.admin.ch/luft/00596/06906/offroad-daten/index.html?lang=en> [28.02.2014]

INFRAS 2010: The Handbook Emission Factors for Road Transport (HBEFA), version 3.1 (MS AccessXP runtime application and report). INFRAS in cooperation with further editors: FOEN/Switzerland; Umweltbundesamt Dessau/Germany; Umweltbundesamt Wien/Austria; Swedish Road Administration, ADEME/France; SFT/Norway. Bern. 30.01.2010.

<http://www.hbefa.net/e/index.html> [28.02.2014]

INFRAS 2010a: Reference approach and feedstocks-improvements, internal documentation for BAFU, November 2010. www.bafu.admin.ch/ghginv-ref

INFRAS 2011 (forthcoming): Handbuch Emissionsfaktoren des Strassenverkehrs 3.1. Dokumentation. Bern.

<http://www.hbefa.net/e/index.html> [28.02.2014]

INFRAS 2011a: Bunker fuels Bodensee/Genfersee, Datenverfügbarkeit und Berechnung der Bunker Fuels, September 2011.

www.bafu.admin.ch/ghginv-ref

INFRAS 2012: Verification of Swiss implied emission factors. INFRAS, Zürich.

www.bafu.admin.ch/ghginv-ref

Intertek 2008: Bestimmung der Heizwerte und CO₂-Emissionsfaktoren von Brenn- und Treibstoffen. Untersuchungsbericht Nr. 170375. Intertek Caleb Brett im Auftrag des Bundesamts für Umwelt BAFU. FOEN internal document. Schlieren/Bern.

www.bafu.admin.ch/ghginv-ref

Intertek 2012: Bestimmung der Heizwerte und CO₂-Emissionsfaktoren von Brenn- und Treibstoffen. Untersuchungsbericht Nr. 111377. Intertek im Auftrag des Bundesamts für Umwelt BAFU. FOEN internal document. Schlieren.

www.bafu.admin.ch/ghginv-ref

IPCC 1997a: Greenhouse Gas Inventory Reference Manual, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Reporting Instructions (Volume 1).

Intergovernmental Panel on Climate Change.

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs4.htm> [27.01.2014]

IPCC 1997b: Greenhouse Gas Inventory Reference Manual, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Workbook (Volume 2). Intergovernmental Panel on Climate Change.

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs5.htm> [27.01.2014]

IPCC 1997c: Greenhouse Gas Inventory Reference Manual, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Reference Manual (Volume 3). Intergovernmental Panel on Climate Change.

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> [27.01.2014]

IPCC 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC C GPG). Intergovernmental Panel on Climate Change.

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> [27.01.2014]

IPCC 2003: Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC GPG LULUCF).

<http://www.ipcc-nggip.iges.or.jp/public/gp/lulucf/gp/lulucf.htm> [27.01.2014]

IPCC 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change.

<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm> [27.01.2014]

Jandl, R., Lindner, M., Vesterdal, L., Bauwens, B., Baritz, R., Hagedorn, F., Johnson, D. W., Minkinen, K., Byrne, K. A. 2007: How strongly can forest management influence soil carbon sequestration? *Geoderma* 137: 253-268.

<http://dx.doi.org/10.1016/j.geoderma.2006.09.003>

Jarvis, P.G., Linder, S. 2000: Constraints to growth of boreal forests. *Nature* 405: 904-905.

<http://dx.doi.org/10.1038/35016154>

Karhu, K., Wall, A., Vanhala, P., Liski, J., Esala, M., Regina, K. 2011: Effects of afforestation and deforestation on boreal soil carbon stocks - Comparison of measured C stocks with Yass07 model results. *Geoderma* 164: 33-45.

<http://dx.doi.org/10.1016/j.geoderma.2011.05.008>

Karl, T., Guenther, A., Lindinger, C., Jordan, A., Fall, R., Lindinger, W. 2001: Eddy covariance measurements of oxygenated volatile organic compound fluxes from crop harvesting using a redesigned proton-transfer-reaction mass spectrometer. *Journal of Geophysical Research* 106 D20: 24157-24167.

<http://dx.doi.org/10.1029/2000JD000112>

Kaufmann, E. 2001: Estimation of standing timber, growth and cut. In: Brassel, P., Lischke, H. (eds.): *Swiss National Forest Inventory: Methods and Models of the Second Assessment*. Swiss Federal Research Institute WSL, Birmensdorf: 162-196.

<http://www.lfi.ch/publikationen/publ/methods/methods.pdf> [28.02.2013]

Kaufmann, E. 2005: Personal communication from Edgar Kaufmann (WSL, Zürich) to Esther Thürig (FOEN, Bern), 12.12.2005.

www.bafu.admin.ch/ghginv-ref

Kaufmann, U. 2013: Energieverbrauch stationäre Motoren und Gasturbinen ab 1990. Excel data file for internal use on behalf of the Federal Office for the Environment, Bern. Eicher + Pauli, Liestal, 17.12.2008.

www.bafu.admin.ch/ghginv-ref

de Keizer et al. 2007: See above under character D (de Keizer)

Keller, M. (Red.), 2005: Schweizerisches Landesforstinventar. Anleitungen für die Feldaufnahmen der Erhebungen 2004-2007. [Field manual for the third Swiss national forest inventory 2004-2007]. Swiss Federal Research Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland.

<http://www.lfi.ch/publikationen/publ/anleitung3.php> [28.02.2013]

Keller, J., Andreani-Aksoyoglu, S., Joss, U. 1995: Inventory of natural emissions in Switzerland. In: Power, H., Moussiopoulos, N., Brebbia C.A. (eds.): *Air Pollution III, Volume 2, Air Pollution Engineering and Management*: 339-346. Computational Mechanics Publications.

www.bafu.admin.ch/ghginv-ref

Keller, A., Rossier, N., Desaulles, A. 2005: Schwermetallbilanzen von Landwirtschaftspartellen der Nationalen Bodenbeobachtung. NABO – Nationales Bodenbeobachtungsnetz der Schweiz. [Heavy-metal balances of agricultural soil monitoring sites. NABO – Swiss Soil Monitoring Network]. Schriftenreihe der FAL 54, Zürich-Reckenholz.

www.bafu.admin.ch/ghginv-ref

Keller, A., Desaulles, A., Schwab, P., Weisskopf, P., Scheid, S., Oberholzer, H.-R. 2006: Monitoring soil quality in the long term: examples from the Swiss Soil Monitoring Network. *Mitteilungen der Österreichischen Bodenkundlichen Gesellschaft, Heft 73*: 5-12.

www.bafu.admin.ch/ghginv-ref

Keller, A. 2013: Details on NABO soil data uncertainties. Written communication from Armin Keller (Agroscope) to Daniel Bretscher (ART), 28.01.2013.

www.bafu.admin.ch/ghginv-ref

Keller/INFRAS 2013: Data were provided by M. Keller to FOEN. E-Mail 07.05.2013 from M.Keller (INFRAS) to B. Müller (FOEN).

Knighton, W.B., Herndon, S.C., Miake-Lye, R.C. 2009: Aircraft Engine Speciated Organic Gases: Speciation of Unburned Organic Gases in Aircraft. US EPA-420-R-09-902. May 2009
<http://www.epa.gov/nonroad/aviation/420r09902.pdf> [28.01.2014]

Koehlerei, 2014: Geschichte der Holzköhlerei in Romoos. <http://www.koehlerei.ch/> [02.04.2014]

Köck, K., Leifeld, J., Fuhrer, J. 2013: A model-based inventory of sinks and sources of CO₂ in agricultural soils in Switzerland: development of a concept. Agroscope, Zürich.

www.bafu.admin.ch/ghginv-ref

König, G., Brunda, M., Puxbaum, H., Hewitt, C.N., Duckham, S.C., Rudolph, J. 1995: Relative contribution of oxygenated hydrocarbons to the total biogenic VOC emissions of selected mid-European agricultural and natural plant species. Atmospheric Environment 29: 861–874.

[http://dx.doi.org/10.1016/1352-2310\(95\)00026-U](http://dx.doi.org/10.1016/1352-2310(95)00026-U)

Kreuzer 2012: Wissenschaftlicher Schlussbericht zuhanden des BAFU und des BLW für das Projekt: Technische Massnahmen und deren Potenzial zur Reduktion der THG CH₄ und N₂O aus der Schweizer Tierhaltung. ETH 10.2. 2012.

www.bafu.admin.ch/ghginv-ref

Kuhn, P. 2011: Supply of data on tree species and age in the city of Bern. Written communication from Peter Kuhn (Stadtgärtnerei Bern) to Beat Rihm (Meteotest, Bern), 08.03.2012.

www.bafu.admin.ch/ghginv-ref

Külling, D. R., Dohme, F., Menzi, H., Sutter, F., Lischer, P., Kreuzer, M. 2002: Methane emissions of differently fed dairy cows and corresponding methane and nitrogen emissions from their manure during storage. Environmental Monitoring and Assessment 79 (2): 129-150.

Külling, D. R., Menzi, H., Sutter, F., Lischer, P., Kreuzer, M. 2003: Ammonia, nitrous oxide and methane emissions from differently stored dairy manure derived from grass- and hay-based rations. Nutrient Cycling in Agroecosystems 65 (1): 13-22.

Kupper, T., Bonjour, C., Achermann, B., Zaucker, F., Rihm, B., Menzi, H. 2013: Ammoniakemissionen in der Schweiz 1990-2010 und Prognose bis 2020. Hochschule für Agrar-, Forst- und Lebensmittelwissenschaften, Zollikofen. URL:

<http://www.agrammon.ch/dokumente-zum-download/>

Lamlom, S.H., Savidge, R.A. 2003: A reassessment of carbon content in wood: variation within and between 41 North American species. Biomass and Bioenergy 25: 381-388.

[http://dx.doi.org/10.1016/S0961-9534\(03\)00033-3](http://dx.doi.org/10.1016/S0961-9534(03)00033-3)

Leifeld, J., Bassin, S., Fuhrer, J. 2003: Carbon stocks and carbon sequestration potentials in agricultural soils in Switzerland. Schriftenreihe der FAL 44. Zürich-Reckenholz.

www.bafu.admin.ch/ghginv-ref

Leifeld, J., Bassin, S., Fuhrer, J. 2005: Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. Agriculture, Ecosystems & Environment 105 (1/2): 255-266.

<http://dx.doi.org/10.1016/j.agee.2004.03.006>

Leifeld, J., Fuhrer, J. 2005: Greenhouse gas emissions from Swiss agriculture since 1990: Implications for environmental policies to mitigate global warming. *Environmental Science & Policy* 8: 410-417.

<http://dx.doi.org/10.1016/j.envsci.2005.04.001>

Leifeld, J., Zimmermann, M., Fuhrer, J. 2007: Characterization of soil carbon stocks and site-specific sequestration potentials of agricultural soils. Extended Summary. Final Report, Contract Number: 810.03.0716 / 2003.C.04. Agroscope Reckenholz-Tänikon Research Station ART.

www.bafu.admin.ch/ghginv-ref

Leifeld, J., Reiser, R., Oberholzer, H.R. 2009: Consequences of conventional versus organic farming on soil carbon: Results from a 27-year field experiment. *Agronomy Journal* 101: 1204-1218.

<http://dx.doi.org/10.2134/agronj2009.0002>

Leupro 2012: Entwicklung und Prognose der Emissionsfaktoren Feuerungen für den Zeitraum 1990 – 2035.

www.bafu.admin.ch/ghginv-ref

Mathys, L., Thürig E. 2010: Baumbiomasse in der Landschaft. [Living biomass of trees in Non-Forest Land]. Final report, Sigmaplan and WSL on behalf of the Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

Meining, S., V.Wilpert, K., Schäffer, J., Schröter, H. 2008: Waldzustandsbericht 2008 der Forstlichen Versuchs- und Forschungsanstalt Baden- Württemberg. Freiburg .

Menzi, H., Frick, R., Kaufmann, R. 1997: Ammoniak-Emissionen in der Schweiz: Ausmass und technische Beurteilung des Reduktionspotenzials. [Emissions d'ammoniac en Suisse: amplitude et évaluation technique du potential de réduction]. Schriftenreihe der FAL 26. Zürich-Reckenholz. [in German, with English and French summary]

Meteotest 2013a: LULUCF and KP-LULUCF. Comparison of Activity Data. Documentation by Rihm, B., Meteotest, Bern on behalf of the Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

Meteotest 2013b: Revision of the Spatial Strata in the Activity Data - Sector LULUCF of the Swiss Greenhouse Gas Inventory. Documentation by Rihm, B., Meteotest, Bern on behalf of the Federal Office for the Environment, Bern. www.bafu.admin.ch/ghginv-ref

Meteotest 2014: Treibhausgasinventar Sektor LULUCF, Tabellen zur Berechnung der Treibhausgasemissionen im Sektor LULUCF in der Schweiz inklusive Begleitdokumentation. Internes Dokument. [GHG inventory, sector LULUCF. Spreadsheets to calculate the GHG emissions of the LULUCF sector in Switzerland inclusive accompanying documents. Internal document]. Meteotest, Bern.

MISTA 2013: Milchstatistik der Schweiz 2012. Schweizerischer Bauernverband (SBV), TSM Treuhand GmbH, Schweizer Milchproduzenten (SMP).

Moeri, A.C. 2007: Kohlenstoffvorräte in Schweizer Waldböden mit besonderer Berücksichtigung der organischen Auflage. Diplomarbeit bei der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft (WSL). Geographisches Institut der Universität Zürich.

www.bafu.admin.ch/ghginv-ref

Mohn, J. 2011: Bestimmung des Anteils biogener und fossiler CO₂ Emissionen aus Schweizer KVAs. Eidg. Materialprüfungsanstalt und Forschungsanstalt. Schlussbericht. Im Auftrag des Bundesamts für Umwelt.

www.bafu.admin.ch/ghginv-ref

Mohn, J. 2013: Bestimmung der N₂O und CH₄ Emissionen aus Schweizer KVA's. Eidg. Materialprüfungs- und Forschungsanstalt. Schlussbericht. Im Auftrag des Bundesamts für Umwelt.

Moller, H.B., Sommer, S.G., Ahring, B.K. 2004: Biological degradation and greenhouse gas emissions during pre-storage of liquid animal manure. *Journal of Environmental Quality* 33 (1): 27-36.

Monni, S., Peltoniemi, M., Palosuo, T., Lehtonen, A., Mäkipää, R., Savolainen, I. 2007: Uncertainty of forest carbon stock changes – implications to the total uncertainty of GHG inventory of Finland. *Climatic Change* 81: 391-413.
<http://dx.doi.org/10.1007/s10584-006-9140-4>

Nave, L. E., Vance, E. D., Swanston, C. W., Curtis, P. S. 2010: Harvest impacts on soil carbon storage in temperate forests. *Forest Ecology and Management* 259: 857-866.
<http://dx.doi.org/10.1016/j.foreco.2009.12.009>

Nowak, D., Crane, D. 2002: Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution* 116: 381–389. [http://dx.doi.org/10.1016/S0269-7491\(01\)00214-7](http://dx.doi.org/10.1016/S0269-7491(01)00214-7)

Nussbaum, M., Papritz, A., Baltensweiler, A., Walthert, L. 2012: Organic Carbon Stocks of Swiss Forest Soils. Final Report. Institute of Terrestrial Ecosystems, ETH Zürich and Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.
<http://dx.doi.org/10.3929/ethz-a-007555133>

Nussbaumer, T., Boogen, N. 2010: Emissionsfaktoren von Holzfeuerungen – Aktualisierung des Arbeitsblatts Emissionsfaktoren Feuerungen und Vorabklärungen zur Bestimmung des Kondensatanteils, Verenum im Auftrag des Bundesamts für Umwelt, (BAFU), 2010.

OcCC 2008: Das Klima ändert – was nun? Der neue UN-Klimabericht (IPCC 2007) und die wichtigsten Ergebnisse aus Sicht der Schweiz. The Advisory Body on Climate Change, Bern.
<http://proclimweb.scnat.ch/Products/OcCC-IPCC/OcCC-IPCC-lowres.pdf> [27.01.2014]

OE 2014: Liechtenstein's Greenhouse Gas Inventory 1990-2012, National Inventory Report under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol, Submission of April 2013 to the United Nations Framework Convention on Climate Change. Office of Environmental Protection (OEP). Principality of Liechtenstein. Vaduz, 15 April 2014.

Park, K.H., Thompson, A.G., Marinier, M., Clark, K., Wagner-Riddle, C. 2006: Greenhouse gas emissions from stored liquid swine manure in a cold climate. *Atmospheric Environment* 40 (4): 618-627.

Perruchoud, D., Kienast, F., Kaufmann, E., Bräker, O.U 1999: 20th Century Carbon Budget of Forest Soils in the Alps. *Ecosystems* 2: 320-337.
<http://dx.doi.org/10.1007/s100219900083>

Peter, S., Hartmann, M., Hediger, W. 2006: Entwicklung der landwirtschaftlichen Emissionen umweltrelevanter Stickstoffverbindungen. Schlussbericht, Dezember 2006. Schriftenreihe Info Agrar Wirtschaft 2006/1. ETH Zürich; Institute for Environmental Decisions IED.
www.bafu.admin.ch/ghginv-ref

Prasuhn, V., Braun, M. 1994: Abschätzung der Phosphor- und Stickstoffverluste aus diffusen Quellen in die Gewässer des Kantons Bern. [Estimation des pertes en phosphore et en azote dans les eaux du canton de Berne à partir de sources diffuses]. Schriftenreihe der FAC Liebefeld 17. [in German, with English and French summary]

Prasuhn, V., Mohni, R. 2003: GIS-gestützte Abschätzung der Phosphor- und Stickstoffeinträge aus diffusen Quellen in die Gewässer des Kantons Bern. FAL, interner Bericht z.H. Amt für Gewässerschutz und Abfallwirtschaft, Kanton Bern. Zürich-Reckenholz.

Prognos 2012a: Die Energieperspektiven für die Schweiz bis 2050 – Energienachfrage und Elektrizitätsangebot in der Schweiz 2000-2050. Prognos AG im Auftrag des Bundesamtes für Energie, Basel. http://www.bfe.admin.ch/themen/00526/00527/index.html?lang=de&dossier_id=05024 [16.02.2014]

Prognos 2013: CO₂-Emissionen 1990-2012 von Industrie- und Dienstleistungen, Endbericht / Kurzdokumentation zuhänden Bundesamt für Umwelt, Bern. Prognos, Basel. www.bafu.admin.ch/ghginv-ref

Quantis 2014: Methanemissionen der Schweizer Gaswirtschaft. Zeitreihe 1990 bis 2012. Schlussbericht. Quantis im Auftrag des Schweizerischen Vereins des Gas- und Wasserfaches SVGW und des Bundesamts für Umwelt BAFU. www.bafu.admin.ch/ghginv-ref

RAP 1999: Fütterungsempfehlungen und Nährwerttabellen für Wiederkäuer [Apports alimentaires recommandés et tables de la valeur nutritive des aliments pour les ruminants]. Landwirtschaftliche Lehrmittelzentrale, Zollikofen. Vierte Auflage.

Richard, L. 1995: Ecologie des mégaphorbiaies subalpines à aune vert de la Vanoise et des régions environnantes (seconde partie) – Phytoécologie. Trav Sci Parc Nat Vanoise 19: 131-160.

Rühr, N., Eugster, W. 2009: Soil respiration fluxes and carbon sequestration of two mountain forests in Switzerland. Report on behalf of the Federal Office for the Environment, Bern. www.bafu.admin.ch/ghginv-ref

Ruffner, H.P. 2005: Written communication from Hans Peter Ruffner (Forschungsanstalt Wädenswil) to Jens Leifeld (ART, Reckenholz), 19.12.2005. www.bafu.admin.ch/ghginv-ref

Rusch, S., Hagedorn, F., Zimmermann, S., Lüscher, P. 2009: Bodenkohlenstoff nach Windwurf – eine CO₂-Quelle? [Soil carbon after wind throw – a source of CO₂?]. Report on behalf of the Federal Office for the Environment. [German] www.bafu.admin.ch/ghginv-ref

SAEFL 1992: Abfallkonzept für die Schweiz, Schriftenreihe Umwelt Nr. 173. Swiss Agency for the Environment, Forests and Landscape, Bern.

SAEFL 1993: NABO - Nationales Bodenbeobachtungsnetz. Messresultate 1985-1991. [NABO - Réseau d'observation des sols: période d'observation 1985-1991]. Schriftenreihe Umwelt Nr. 200. [Cahier de l'environnement No 200]. Swiss Agency for the Environment, Forests and Landscape, Bern. <http://www.bafu.admin.ch/publikationen/publikation/00321/index.html?lang=de> [German] <http://www.bafu.admin.ch/publikationen/publikation/00321/index.html?lang=fr> [French] [09.04.2014]

SAEFL 1995: Emissionen des Strassenverkehrs 1950-2010. [Emissions polluantes du trafic routier de 1950 à 2010]. Schriftenreihe Umwelt Nr. 255. [Cahier de l'environnement No 255]. Swiss Agency for the Environment, Forests and Landscape, Bern. [available in German and French].

SAEFL 1996: Schadstoffemissionen und Treibstoffverbrauch des Offroad Sektors, Umwelt-Materialien Nr. 49. Elektrowatt Ingenieurunternehmung AG, Zürich; Technik Thermische Maschinen, Niederrohrdorf; Swiss Agency for the Environment, Forests and Landscape, Bern.

SAEFL 1996a: Luftschadstoff-Emissionen aus natürlichen Quellen der Schweiz. [Emissions polluantes dues aux sources naturelles en Suisse]. Schriftenreihe Umwelt Nr. 257. [Cahier de l'environnement No 257]. Swiss Agency for the Environment, Forests and Landscape, Bern. www.bafu.admin.ch/ghginv-ref

SAEFL 2000: Handbuch Emissionsfaktoren für stationäre Quellen. Ausgabe 2000, Reihe Vollzug Umwelt. Swiss Agency for the Environment, Forests and Landscape, Bern.

SAEFL 2000a: Nationales Boden-Beobachtungsnetz - Veränderungen von Schadstoffgehalten nach 5 und 10 Jahren. [Réseau national d'observation des sols - Variations des teneurs en polluants après 5 et 10 ans de suivi]. Schriftenreihe Umwelt Nr. 320. [Cahier de l'environnement No 320]. Swiss Agency for the Environment, Forests and Landscape, Bern.

<http://www.bafu.admin.ch/publikationen/publikation/00470/index.html?lang=de> [German]

<http://www.bafu.admin.ch/publikationen/publikation/00470/index.html?lang=fr> [French]

[28.02.2014]

SAEFL 2001: Bauabfälle Schweiz - Mengen, Perspektiven und Entsorgungswege. Umwelt-Materialien Nr. 131. Band 1: Kennwerte. Swiss Agency for the Environment, Forests and Landscape, Bern.

SAEFL 2004a: Handbook Emission Factors for Road Transport. Swiss Agency for the Environment, Forests and Landscape / INFRAS, CD ROM, Bern.

<http://www.bafu.admin.ch/luft/00596/00597/00605/index.html?lang=en> [28.02.2014]

SAEFL 2004b: Swiss national forest programme (Swiss NFP), Environmental documentation No. 363, Swiss Agency for the Environment, Forests and Landscape, Bern. 117 pp.

<http://www.bafu.admin.ch/publikationen/publikation/00527/index.html?lang=en> [28.02.2014]

SAEFL 2005h: Personal communication from R. Quartier (SAEFL, Bern) to J. Füssler (Ernst Basler + Partner, Zollikon), 23 February 2005.

www.bafu.admin.ch/ghginv-ref

SBV 2004: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2003.

[Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung 2003]. Swiss Farmers Union, Brugg. [available in German and French]

SBV 2007: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2006.

[Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung, 2006]. Swiss Farmers Union, Brugg. [available in German and French]

SBV 2007a: Données mensuelles sur l'agriculture. 67.8. August 2007. [Landwirtschaftliche Monatszahlen 67.8. August 2007]. Swiss Farmers Union, Brugg. [available in German and French]

www.bafu.admin.ch/ghginv-ref

SBV 2012: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2011.

[Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung, 2011]. Swiss Farmers Union, Brugg. [available in German and French]

<http://www.sbv-usb.ch/de/statistik/> [28.02.2014]

SBV 2013: Statistiques et évaluations concernant l'agriculture et l'alimentation, 2012.

[Statistische Erhebungen und Schätzungen über Landwirtschaft und Ernährung, 2012]. Swiss Farmers Union, Brugg. [available in German and French]

<http://www.sbv-usb.ch/de/statistik/> [28.09.2014]

Schmid, M., Neftel, A., Fuhrer, J. 2000: Lachgasemissionen aus der Schweizer Landwirtschaft. [Emissions de protoxyde d'azote de l'agriculture Suisse]. Schriftenreihe der FAL 33. Zürich-Reckenholz.

www.bafu.admin.ch/ghginv-ref

Schneider, M. 2010: Personal communication from Manuel Schneider (ART, Reckenholz) to Daniel Bretscher (ART, Reckenholz), 21.09.2010.

www.bafu.admin.ch/ghginv-ref

Schwager, S. 2005: Personal communication from Stefan Schwager (FOEN, Bern) to Andreas Liechti (FOEN, Bern), 23.12.2005.

www.bafu.admin.ch/ghginv-ref

- Schwarz, W. 2001:** Emissionen des Kältemittels R-134a aus mobilen Klimaanlageen. Jährliche Emissionsraten von bis zu sieben Jahre alten Pkw-Klimaanlagen. [Emission of Refrigerant R-134a from Mobile Air-Conditioning Systems Annual Rate of Emission from Passenger-Car Air-Conditioning Systems up to Seven Years Old]. Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 360 09 006), Frankfurt.
<http://www.oekorecherche.de/deutsch/berichte/volltext/vollR134a.pdf> [German] [30.01.2013]
<http://www.oekorecherche.de/english/berichte/volltext/MAC-LOSS-2001.pdf> [English] [30.01.2013]
- Schwarz, W., Wartmann, S. 2005:** Emissionen und Emissionsprognosen von H-FKW, FKW und SF₆ in Deutschland – Aktueller Stand und Entwicklung eines Systems zur jährlichen Ermittlung. Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 202 41 356), Frankfurt.
<http://www.umweltdaten.de/publikationen/fpdf-l/3000.pdf> [German] [30.01.2013]
- Schweizerischer Bundesrat 2008:** Verordnung des EVD über Ethoprogramme (Ethoprogrammverordnung), 25. Juni 2008 (Stand am 1. Oktober 2008). Der Schweiz. Bundesrat, Bern, Schweiz.
- SECO 2014:** GDP and expenditure-side components (yearly and quarterly data). State Secretariat for Economic Affairs SECO.
http://www.seco.admin.ch/themen/00374/00456/00458/index.html?lang=en&download=NHZLpZeg7t,lnp6l0NTU042l2Z6ln1ad1lZn4Z2qZpnO2Yug2Z6gpJCGdn15qWym162epYbg2c_JiKbNoKSn6A-- [10.02.2014]
- Seringas 2010:** Upgrading Seringas to V5.3.0, release notes, CDC Climat, Arcueil, France.
www.bafu.admin.ch/ghginv-ref
- SFCA 2013:** Mineralölsteuer: Versteuerte Mengen 2012.
http://www.ezv.admin.ch/zollinfo_firmen/04020/04256/04263/04521/04523/index.html?lang=de [11.02.2014]
- SFOE 1991:** Schweizerische Gesamtenergiestatistik 1990. Statistique globale suisse de l'énergie 1990. Swiss Federal Office of Energy, Bern.
- SFOE 2001:** Schweizerische Gesamtenergiestatistik 2000. Statistique globale suisse de l'énergie 2000. Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=fr&name=fr_166628788.pdf [German and French] [31.01.2014]
- SFOE 2012:** Schweizerische Gesamtenergiestatistik 2011. Statistique globale suisse de l'énergie 2011. Swiss Federal Office of Energy, Bern. In German and French.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_872595201.pdf [28.01.2014]
- SFOE 2013:** Schweizerische Gesamtenergiestatistik 2012. Statistique globale suisse de l'énergie 2012. Swiss Federal Office of Energy, Bern. In German and French.
[http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_551182741.pdf&endung=Schweizerische Gesamtenergiestatistik 2012](http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_551182741.pdf&endung=Schweizerische%20Gesamtenergiestatistik%202012) [28.01.2014]
- SFOE 2013a:** Schweizerische Statistik der erneuerbaren Energien. Ausgabe 2012. Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_219304565.pdf [11.02.2014]
- SFOE 2013b:** Schweizerische Holzenergiestatistik. Erhebung für das Jahr 2012. Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/php/modules/publikationen/stream.php?extlang=de&name=de_610124540.pdf [German, with French summary] [23.02.2014]

SFOE 2013c: Thermische Stromproduktion inklusive Wärmekraftkopplung (WKK) in der Schweiz. Ausgabe 2012. Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/themen/00526/00541/00543/index.html?lang=de&dossier_id=00774
[German, with French summary] [23.02.2014]

SFOE 2013d: Energieverbrauch in der Industrie und im Dienstleistungssektor. Ausgabe 2012. Swiss Federal Office of Energy, Bern.
http://www.bfe.admin.ch/themen/00526/00541/00543/index.html?lang=de&dossier_id=00775
[German, with French summary] [02.04.2014]

SFSO 1997: Digital terrain model („Geländedaten“, 100m-Raster). Swiss Federal Statistical Office, GEOSTAT, Neuchâtel.
<http://www.bfs.admin.ch/bfs/portal/fr/index/dienstleistungen/geostat/datenbeschreibung/gelaendedaten.html> [13.03.2013]

SFSO 2000a: Digital soil map 1:200'000 („Bodeneignungskarte“, BEK). Swiss Federal Statistical Office, GEOSTAT, Neuchâtel.
http://www.bfs.admin.ch/bfs/portal/fr/index/dienstleistungen/geostat/datenbeschreibung/digitale_bodeneignungskarte.html [17.01.2012]

SFSO 2002: Einblicke in die schweizerische Landwirtschaft. [Insights into Swiss Agriculture]. Swiss Federal Statistical Office, Neuchâtel. [available in German and French]

SFSO 2005: Swiss Land Use Statistics (Arealstatistik Schweiz). Supply of hectare-based data of the first survey (Arealstatistik 1979/85, ASCH1) and second survey (Arealstatistik 1992/97, ASCH2). Swiss Federal Statistical Office, Neuchâtel.
http://www.bfs.admin.ch/bfs/portal/fr/index/infothek/erhebungen_quellen/blank/blank/arealstatistik/01.html [13.03.2013] and
<http://www.bfs.admin.ch/bfs/portal/fr/index/infothek/nomenklaturen/blank/blank/arealstatistik/01.html> [13.03.2013]

SFSO 2006a: Arealstatistik 2004/09 - Kategorienkatalog. Bodenbedeckung, Bodennutzung. Sektion Geoinformatik, Swiss Federal Statistical Office, Neuchâtel.
www.bafu.admin.ch/ghginv-ref
See also: <http://www.bfs.admin.ch/bfs/portal/fr/index/themen/02/03.html> and
http://www.bfs.admin.ch/bfs/portal/fr/index/infothek/erhebungen_quellen/blank/blank/arealstatistik/01.html [13.03.2014]

SFSO 2009a: Volkswirtschaftliche Gesamtrechnung der Schweiz im Jahr 2008 (National Accounting 2008) Bruttoinlandprodukt (BIP), Swiss Federal Statistical Office, Neuchâtel, November 2009.
<http://www.bfs.admin.ch/bfs/portal/de/index/themen/04/22/publ.html?publicationID=3777>
[10.02.2014]

SFSO 2009c: Actualités OFS: Prestations du transport privé de personnes par la route, Séries chronologiques actualisées jusqu'en 2008, Neuchâtel 2010.

SFSO 2010c: Fahrzeugbewegungen und Fahrleistungen im Güterverkehr, bis 2008, Neuchâtel 2010.

SFSO 2013: Supply of provisional data of the AREA Land Use Statistics. Written communication from Felix Weibel and Jürg Burkhalter (SFSO, Neuchâtel) to Lukas Mathys (Sigmaplan, Bern), 16.07.2013. www.bafu.admin.ch/ghginv-ref

SFSO 2013a: Die Bevölkerung der Schweiz 2012. Swiss Federal Office. Neuchâtel.
<http://www.bfs.admin.ch/bfs/portal/de/index/themen/01/02/blank/dos/mittlere/01.html>
05.02.2014]

SFSO 2013b: Road vehicles in Switzerland – stock 2012:
<http://www.bfs.admin.ch/bfs/portal/de/index/themen/11/03/blank/02/01/01.html> [11.02.2013]

SFSO 2013c: Wood production in Switzerland 1975-2012. Swiss Federal Statistical Office, Neuchâtel. <http://www.agr-bfs.ch> [official text in German, English, French and Italian] [14.10.2013]

SFSO 2013d: STAT-TAB: Die interaktive Statistikdatenbank; Datenwürfel für Thema 07.2 – Landwirtschaft. Swiss Federal Statistical Office, Neuchâtel. http://www.pxweb.bfs.admin.ch/Database/German_07%20-%20Land-%20und%20Forstwirtschaft/07.2%20-%20Landwirtschaft/07.2%20-%20Landwirtschaft.asp?lang=1&prod=07&secprod=2&openChild=true [25.9.2013]

SHL 2010: Bestimmung Tiere der Rindviehkategorien ab 2009 für die Berechnung des Ammoniakinventars. Schweizerische Hochschule für Landwirtschaft SHL, Zollikofen.

Sigmaplan 2010a: Deforestation under Kyoto Protocol. Documentation of implementation. Internal documentation by Mathys, L., Sigmaplan, Bern on behalf of the Federal Office for the Environment, Bern. www.bafu.admin.ch/ghginv-ref

Sigmaplan 2012a: LULUCF and KP-LULUCF. Comparison of deforestation data. Final Report. Internal document. Sigmaplan, Bern. www.bafu.admin.ch/ghginv-ref

Sigmaplan 2014: Treibhausgasinventar Sektor LULUCF, Tabellen zur Berechnung der Aktivitätsdaten im Sektor LULUCF in der Schweiz inklusive Begleitdokumentation. Internes Dokument. [GHG inventory, sector LULUCF. Spreadsheets to calculate the activity data of the LULUCF sector in Switzerland inclusive accompanying documents. Internal document]. Sigmaplan, Bern.

Soliva, C.R. 2006: Report to the attention of IPCC about the data set and calculation method used to estimate methane formation from enteric fermentation of agricultural livestock population and manure management in Swiss agriculture. On behalf of the Federal Office for the Environment, Bern. ETH Zurich, Institute of Animal Science. www.bafu.admin.ch/ghginv-ref

Soliva, C.R. 2006a: Dokumentation der Berechnungsgrundlage von Methan aus der Verdauung und dem Hofdünger landwirtschaftlicher Nutztiere. Im Auftrag des Bundesamtes für Umwelt, Bern. ETH Zürich, Institut für Nutztierwissenschaften. www.bafu.admin.ch/ghginv-ref

Sommer, S.G., Petersen, S.O., Sorensen, P., Poulsen, H.D., Moller, H.B. 2007: Methane and carbon dioxide emissions and nitrogen turnover during liquid manure storage. Nutrient Cycling in Agroecosystems 78 (1): 27-36.

Spicer C.W., Holdren M.W., Riggins R.M., Lyon T.F. 1994: Chemical composition and photochemical reactivity of exhaust from aircraft turbine engines. Annales Geophysicae 12 (10-11): 944-955. <http://dx.doi.org/10.1007/s00585-994-0944-0> [28.01.2014]

Spiecker, H., 1999: Growth trends in European forests, Do we have sufficient knowledge? In Karjalainen T, Spiecker H, Laroussinie, O. (Eds) 1999. Causes and consequences of accelerating tree growth in Europe. Proceedings of the international seminar held in Nancy, France 14–16. May 1998. European Forest Institute Proceedings No. 27: 157-169.

Spiess, E. 2005: Die Stickstoffbilanz der Schweiz. In: Herzog, F., Richner, W. (eds.): Evaluation der Ökomassnahmen Bereich Stickstoff und Phosphor. Schriftenreihe der FAL 57. Zürich Reckenholz: 26-31.

Spirig, C., Neftel, A. 2002: Biogene VOC und Aerosole. Bedeutung der biogenen flüchtigen organischen Verbindungen für die Aerosolbildung. Schriftenreihe der FAL 42. Zürich-Reckenholz.

SQS 2008: IQNet and SQS ISO 9001:2000 Certificate. Certified area: National Inventory System. Field of activity: Greenhouse Gas Inventory and CO₂ Statistics. Registration Number: CH-34433. Issued on 04.01.2008

www.bafu.admin.ch/ghginv-ref

SQS 2010: IQNet and SQS ISO 9001:2008 Certificate. Certified area: National Inventory System and National Registry. Field of activity: Greenhouse Gas Inventory, CO₂ Statistics and Accounting of Kyoto Protocol Units. Registration Number: CH-34433. Issued on 01.12.2010

www.bafu.admin.ch/ghginv-ref

Stangle, R. 2004: Ingenieurbiologische Hangsicherungsmassnahmen in Wildbacheinzugsgebieten – Bestandesentwicklung und langfristige Wirksamkeit. Klagenfurt: Internationale Forschungsgesellschaft Interpraevent, Tagungspublikation Internationales Symposium Interpraevent 2004, Band 1, pp. 295–305.

Statistics Finland 2012: Greenhouse gas emissions in Finland 1990-2010. National Inventory Report under the UNFCCC and the Kyoto Protocol.

Stieger, J. 2014: Methane emissions at farm and regional scales: origin, magnitude and spatio-temporal variability. PhD Thesis ETHZ; Zürich, Switzerland.

Stricker, B. 2012: Energiebedarfswerte der Pferde, Esel und Ponys. Written communication from Brigitte Stricker (ALP-Haras, Agroscope Liebefeld-Posieux research station) to Daniel Bretscher (ART, Reckenholz), August-October 2012. www.bafu.admin.ch/ghginv-ref

Swiss Confederation 1983: Loi fédérale du 7 octobre 1983 sur la protection de l'environnement (Loi sur la protection de l'environnement, LPE). As at 01 August 2008.

http://www.admin.ch/ch/f/rs/c814_01.html [official text in German, French and Italian] [27.01.2014]

Swiss Confederation 1985: Ordonnance du 16 décembre 1985 sur la protection de l'air (OPair). [Swiss Federal Ordinance on Air Pollution Control]. As of 01 January 2009: Annex 5

http://www.admin.ch/ch/f/rs/c814_318_142_1.html [official text in German, French and Italian] [10.02.2014]

www.bafu.admin.ch/ghginv-ref [English; as of 28 March 2000]

Swiss Confederation 1991: Loi fédérale du 4 octobre 1991 sur les forêts (Loi sur les forêts, LFo). As at 01 January 2008.

http://www.admin.ch/ch/f/rs/c921_0.html [official text in German, French and Italian] [28.02.2013]

Swiss Confederation 1991a: Ordonnance du 21 janvier 1991 sur la protection des hauts-marais et des marais de transition d'importance nationale (Ordonnance sur les hauts-marais). As at 01 January 2008.

http://www.admin.ch/ch/f/rs/c451_32.html [official text in German, French and Italian] [13.03.2013]

Swiss Confederation 1992: Ordonnance du 30 novembre 1992 sur les forêts (OFo). As at 01 March 2011.

http://www.admin.ch/ch/f/rs/c921_01.html [official text in German, French and Italian] [05.04.2011]

Swiss Confederation 1994: Ordonnance du 7 septembre 1994 sur la protection des bas-marais d'importance nationale (Ordonnance sur les bas-marais). As at 01 February 2010.

http://www.admin.ch/ch/f/rs/c451_33.html [official text in German, French and Italian] [18.01.2012]

Swiss Confederation 1997: Ordonnance du 12 novembre 1997 sur la taxe d'incitation sur les composés organiques volatils (OCOV). As at 01 January 2009.

http://www.admin.ch/ch/f/rs/c814_018.html [official text in German, French and Italian] [10.02.2014]

Swiss Confederation 1998: Ordonnance du 1er juillet 1998 sur les atteintes portées aux sols (OSol) (Ordinance against deterioration of soils). As at 01 June 2012.

<http://www.admin.ch/opc/fr/classified-compilation/19981783/index.html> [official text in German, French and Italian] [04.04.2014]

Swiss Confederation 2005: Ordinance of 18 May 2005 on Risk Reduction related to the Use of certain particularly dangerous Substances, Preparations and Articles (Ordinance on Chemical Risk Reduction, ORRChem). SR Number 814.81. As on 1 August 2011.

http://www.admin.ch/ch/d/sr/814_81/index.html [official text in German, French and Italian] [10.02.2014]

<http://www.admin.ch/ch/e/rs/8/814.81.en.pdf> [unofficial translation in English] [30.1.2013]

Swiss Confederation 2011: Bundesgesetz über die Reduktion der CO₂-Emissionen (CO₂-Gesetz). As of 1.1.2013. http://www.admin.ch/ch/d/sr/c641_71.html

Swiss Confederation 2012: Verordnung über die Reduktion der CO₂-Emissionen (CO₂-Verordnung). As of 12.2.2013. http://www.admin.ch/ch/d/sr/c641_711.html

Swiss-TS 2013: Quality management according to ISO 9001:2008. Certificate. Certified area: National Inventory System and Reporting under the UNFCCC and the Kyoto Protocol. Registration Number: 13-299-113. Issued on 25.11.2013. www.bafu.admin.ch/ghginv-ref

Thürig, E., Palosuo, T., Bucher, J., Kaufmann, E. 2005: The impact of windthrow on carbon sequestration in Switzerland: a model-based assessment. Forest Ecology and Management 210: 337-350.

<http://dx.doi.org/10.1016/j.foreco.2005.02.030>

Thürig, E., Kaufmann, E., Schmid, S., Bugmann, H. 2005a: Treibhausgas Inventar: Waldkennzahlen und jährlicher Klimaeinfluss. Internal report to FOEN. Inclusive: Excel-Datei Klimafaktoren.berechnen.xls zur Berechnung des jährlichen klimakorrigierten Zuwachses.

www.bafu.admin.ch/ghginv-ref

Thürig, E., Weber, P., Lischke, H., Schmatz, D., Kaufmann, E., Zandt, H., Dobbertin, M., Zingg, A., Cherubini, P., Waldner, P. 2009: Verbesserung der Klimaabhängigkeit der Wachstumsfunktion für Szenarienanalysen der Waldentwicklung (Holznutzungspotential, Nachhaltigkeit) KLIWAWA. [Improvement of climate dependency of growth curves for analyzing forest development scenarios (sustainable Potential Wood Supply)]. Forschungsanstalt für Wald, Schnee und Landschaft WSL. Research project on behalf of the Federal Office for the Environment, Bern.

www.bafu.admin.ch/ghginv-ref

Thürig, E., Herold, A. 2013: Recalculation of emission factors in Swiss forests for the Swiss GHGI. Internal documentation of technical adjustments of data delivery and more recent data. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.

www.bafu.admin.ch/ghginv-ref

Thürig, E. 2014: NFI 2009–2012: Calculation of emission factors in Swiss forests for the Swiss GHGI. Internal Report commissioned by the Swiss Federal Office for the Environment.

www.bafu.admin.ch/ghginv-ref

Thuille, A., Schulze, E.D. 2006: Carbon dynamics in successional and afforested spruce stands in Thuringia and the Alps. Global Change Biology 12: 325-342.

<http://dx.doi.org/10.1111/j.1365-2486.2005.01078.x>

TTM 2006a: Emissionsfaktoren CH₄, TTM, A. Mayer 16.1.2006

www.bafu.admin.ch/ghginv-ref

TTM 2006b: Emissionsfaktoren N₂O, TTM, A. Mayer 16.1.2006

www.bafu.admin.ch/ghginv-ref

Tuomi, M., Thum, T., Järvinen, H., Fronzek, S., Berg, B., Harmon, M., Trofymow, J.A., Sevanto, S., Liski, J. 2009: Leaf litter decomposition - Estimates of global variability based on Yasso07 model. *Ecological Modelling* 220: 3362-3371.
<http://dx.doi.org/10.1016/j.ecolmodel.2009.05.016>

Tuomi, M., Laiho, R., Repo, A., Liski, J. 2011: Wood decomposition model for boreal forests. *Ecological Modelling* 222: 709-718.
<http://dx.doi.org/10.1016/j.ecolmodel.2010.10.025>

UBA 2005: Emissionen, Aktivitätsraten und Emissionsfaktoren von fluorierten Treibhausgasen (F-Gasen) in Deutschland für die Jahre 1995 – 2002. Texte: 14 / 05. Umwelt Bundes Amt. Berlin.
<http://www.umweltdaten.de/publikationen/fpdf-l/2902.pdf> [German] [30.01.2013]

UBA 2006: Einsatz von Sekundärbrennstoffen, Forschungsbericht 204 42 203/02, UBA-FB 000893; available for download from www.umweltbundesamt.de

UBA 2007: Daten von H-FKW, FKW und SF₆ für die nationale Emissionsberichterstattung gemäß Klimarahmenkonvention für die Berichtsjahre 2004 und 2005“ "Fluorierte Treibhausgase". Gutachten durch die Oeko-Recherche GmbH für das Umweltbundesamt (FKZ 205 41 217/01), Frankfurt.
<http://www.umweltdaten.de/publikationen/fpdf-l/3439.pdf> [German] [30.01.2013]

UBA/Ökorecherche 2012: Entsorgung von Kältemitteln aus stationären Anlagen, Emissionsfaktor der Entsorgung, Emissionsberichterstattung von F-Gasen in der Kälte- und Klimatechnik. UBA Umweltbundesamt Berlin – 24.10.12. Öko-Recherche Büro für Umweltforschung und -beratung GmbH, W. Schwarz, Barbara Gschrey. Umweltbundesamt Berlin - 24. Oktober 2012, Ergebnisse des Forschungsvorhabens 1. Gewerbekälte und 2. Industriekälte

Umweltbundesamt 2012: Austria's National Inventory Report 2012. Submission under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Anderl, M et al. REP-0381.
<http://www.umweltbundesamt.at/fileadmin/site/publikationen/REP0381.pdf> [27.03.2013]

UNECE 2003: Guidelines for Estimating and Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution. Air Pollution Studies No. 15. New York and Geneva.
<http://www.unece.org/env/documents/2003/eb/air/ece.eb.air.80.E.pdf> [27.01.2014]

UNFCCC 2003: Review of the Implementation of Commitments and of other Provisions of the Convention. National Communications: Greenhouse Gas Inventories from Parties Included in Annex I to the Convention. UNFCCC guidelines on reporting and review. FCCC/CP/2002/8.
<http://unfccc.int/resource/docs/cop8/08.pdf> [28.02.2014]

UNFCCC 2004: Report of the individual review of the greenhouse gas inventory of Switzerland submitted in 2004. FCCC/WEB/IRI/2004/CHE, 15 December 2004.
http://unfccc.int/files/national_reports/annex_i_ghg_inventories/inventory_review_reports/application/pdf/2004_irr_in-country_review_switzerland.pdf [27.01.2014]

UNFCCC 2006: Report of the individual review of the greenhouse gas inventory of Switzerland submitted in 2005. FCCC/ARR/2005/CHE, 11 April 2006.
<http://unfccc.int/resource/docs/2006/arr/che.pdf> [27.01.2014]

UNFCCC 2006a: Guidelines for national systems under Article 5, paragraph 1, of the Kyoto Protocol. Decision 19/CMP.1. FCCC/KP/CMP/2005/8/Add.3,
<http://unfccc.int/resource/docs/2005/cmp1/eng/08a03.pdf#page=14> [28.02.2014]

UNFCCC 2006b: Updated UNFCCC reporting guidelines on annual inventories following incorporation of the provisions of decision 14/CP.11. FCCC/SBSTA/2006/9.

<http://unfccc.int/resource/docs/2006/sbsta/eng/09.pdf> [27.01.2014]

UNFCCC 2007: Report of the individual review of the greenhouse gas inventory of Switzerland submitted in 2006. FCCC/ARR/2006/CHE, 31 July 2007.

<http://unfccc.int/resource/docs/2007/arr/che.pdf> [27.01.2014]

UNFCCC 2007a: Report of the review of the initial report of Switzerland. FCCC/IRR/2007/CHE, 17 August 2007.

<http://unfccc.int/resource/docs/2007/irr/che.pdf> [10.02.2014]

UNFCCC 2008: Kyoto Protocol Reference Manual on Accounting of Emissions and Assigned Amount.

http://unfccc.int/resource/docs/publications/08_unfccc_kp_ref_manual.pdf [27.01.2014]

UNFCCC 2009: Report of the individual review of the greenhouse gas inventories of Switzerland submitted in 2007 and 2008. FCCC/ARR/2008/CHE, 29 April 2009.

<http://unfccc.int/resource/docs/2009/arr/che.pdf> [27.01.2014]

UNFCCC 2009a: Annotated outline of the National Inventory Report including reporting elements under the Kyoto Protocol

http://unfccc.int/files/national_reports/annex_i_ghg_inventories/reporting_requirements/application/pdf/annotated_nir_outline.pdf [27.01.2014]

UNFCCC 2010: Report of the individual review of the annual submission of Switzerland submitted in 2009. FCCC/ARR/2009/CHE, 26 January 2010.

<http://unfccc.int/resource/docs/2010/arr/che.pdf> [27.01.2014]

UNFCCC 2011: Report of the individual review of the annual submission of Switzerland submitted in 2010. FCCC/ARR/2010/CHE, 23 May 2011.

<http://unfccc.int/resource/docs/2011/arr/che.pdf> [27.01.2014]

UNFCCC 2012: Report of the individual review of the annual submission of Switzerland submitted in 2011. FCCC/ARR/2011/CHE, 16 May 2012.

<http://unfccc.int/resource/docs/2012/arr/che.pdf> [27.01.2014]

UNFCCC 2013: Report of the individual review of the annual submission of Switzerland submitted in 2012. FCCC/ARR/2012/CHE, 31 May 2013.

<http://unfccc.int/resource/docs/2013/arr/che.pdf> [27.01.2014]

UNFCCC 2013a: Potential Problems and Further Questions from the ERT formulated in the course of the 2013 review of the greenhouse gas inventories of Party submitted in 2013. Bonn, 07 September 2013.

UNFCCC 2014: Report of the individual review of the annual submission of Switzerland submitted in 2013. FCCC/ARR/2013/CHE, 28 February 2014.

<http://unfccc.int/resource/docs/2014/arr/che.pdf> [27.01.2014]

USEPA 1995: Compilation of air pollutant emission factors, Chapter 10.7 Charcoal.

<http://www.epa.gov/ttn/chief/ap42/index.html> [25.03.2014]

UVEK 2003: Düngen mit Klärschlamm wird verboten, UVEK Eidgenössisches Departement für Umwelt, Verkehr, Energie, Kommunikation.

http://www.admin.ch/cp/d/3e816ebe_1@presse1.admin.ch.html [28.02.2013]

Van der Weerden, T.J., Jarvis, S.C. 1997: Ammonia emission factors for N fertilizers applied to two contrasting grassland soils. Environmental Pollution 95(2): 205-211.

[http://dx.doi.org/10.1016/S0269-7491\(96\)00099-1](http://dx.doi.org/10.1016/S0269-7491(96)00099-1)

- Van Miegroet, H., Olsson, M. 2011:** Ecosystem Disturbance and Soil Organic Carbon – A Review. In: Soil Carbon in Sensitive European Ecosystems. John Wiley & Sons, Ltd, Chichester: 85-117.
<http://dx.doi.org/10.1002/9781119970255.ch5>
- Vanninen, P., Mäkelä, A. 1999:** Fine root biomass of Scots pine stands differing in age and soil fertility in southern Finland. *Tree Physiology* 19: 823–830.
<http://dx.doi.org/10.1093/treephys/19.12.823>
- Vesterdal, L., Ritter, E., Gundersen, P. 2002:** Change in soil organic carbon following afforestation of former arable land. *Forest Ecology and Management* 169: 137-147.
[http://dx.doi.org/10.1016/S0378-1127\(02\)00304-3](http://dx.doi.org/10.1016/S0378-1127(02)00304-3)
- Vorreiter, L. 1949:** Holztechnologisches Handbuch; Band I: Allgemeines, Holzkunde, Holzschutz und Holzvergütung. Verlag Georg Fromme & Co., Wien.
- VSTB 2003:** Erläuterungen zur Zielvereinbarung. Verband Schweizerischer Trocknungsbetriebe VSTB. August 2003
- VTG 2013:** Consumption of aviation fuel and jet kerosene of Swiss military aircraft 1990-2012. Amilcare Santino Foglia (VTG, Dübendorf) to Anouk-Aimée Bass (FOEN, Bern),
www.bafu.admin.ch/ghqinv-ref
- VTT 2004:** Transit Bus Emission Study: Comparison of Emissions from Diesel and Natural Gas Buses. Research Report PRO3/P5150/04. VTT Processes, Espoo.
- Walther U., Menzi, H., Ryser, J.-P., Flisch, R., Jeangros, B., Maillard, A., Siegenthaler, A., Vuilloud, P.A. 1994:** Grundlagen für die Düngung im Acker- und Futterbau. *Agrarforschung* 1(7): 1-40.
www.bafu.admin.ch/ghqinv-ref]
- Warneke C., Luxembourg, S.L., de Gouw, J.A., Rinne, H.J.I., Guenther, A.B., Fall, R. 2002:** Disjunct eddy covariance measurements of oxygenated VOC fluxes from an Alfalfa field before and after cutting. *Journal of Geophysical Research* 107 D8: 4067.
<http://dx.doi.org/10.1029/2001JD000594>
- Widmer, A. 2006:** Written communication from Albert Widmer (Forschungsanstalt Wädenswil) to Jens Leifeld (ART, Reckenholz), 20.06.2006.
www.bafu.admin.ch/ghqinv-ref
- Wiesen, P., Kleffmann, J., Kurtenbach, R., Becker, K.H. 1994:** Nitrous Oxide and Methane Emissions from Aero Engines. *Geophysical Research Letters* 21: 2027-2030.
<http://dx.doi.org/10.1029/94GL01709>
- Wirth, C., Schulze, E. D., Schwalbe, G., Tomczyk, I., Weber, G.-E., Weller, E. 2004:** Dynamik der Kohlenstoffvorräte in den Wäldern Thüringens: Abschlussbericht zur 1. Phase des BMBF-Projektes "Modelluntersuchung zur Umsetzung des Kyoto-Protokolls". *Mitteilungen der Thüringer Landesanstalt für Wald, Jagd und Fischerei* 23.
- WSL 2013:** Swissfire database. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.
<http://www.wsl.ch/swissfire> [16.09.2013]
- Wutzler, T., Wirth, C., Schumacher, J., 2008:** Generic biomass functions for common beech (*Fagus sylvatica*) in Central Europe: predictions and components of uncertainty. *Canadian Journal of Forest Research* 38: 1661-1675.
<http://dx.doi.org/10.1139/X07-194>
- Xinmin, J. 2004:** Die Methanemissionen der Schweizer Gasindustrie. Abschätzung der Gasemissionen [Methane emissions from Swiss gas industry. Estimation of methane emissions]. *Gas, Wasser, Abwasser* 5/2004: 337-345.
www.bafu.admin.ch/ghqinv-ref

Zeitz, J. O., Soliva, C. R., Kreuzer, M. 2012: Swiss diet types for cattle: how accurately are they reflected by the Intergovernmental Panel on Climate Change default values? *Journal of Integrative Environmental Sciences*, 9:sup1,199-216.

<http://dx.doi.org/10.1080/1943815X.2012.709253>

Zimmermann, M., Leifeld, J., Schmidt, M.W.I., Smith, P., Fuhrer, J. 2007: Measured soil organic matter fractions can be related to pools in the RothC model. *European Journal of Soil Science* 58: 658–667.

<http://dx.doi.org/10.1111/j.1365-2389.2006.00855.x>

Zimmermann, S., Hiltbrunner, D. 2012: Turnover and stabilization of soil organic matter: effect of land-use change in alpine regions. Schlussbericht [German]. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.

www.bafu.admin.ch/ghqinv-ref

ZPK 2013: Annual Report of Federation of the Swiss Pulp, paper and board industry.

<http://www.zpk.ch/> [02.04.2014]

References to EMIS database comments

Table A - 1 Assignments of NFR Codes to titles of EMIS database comments. These internal documents will be made available, on request, to reviewers by the NIC.

| NFR Code CRF [UNECE] | EMIS Title | NFR Code CRF [UNECE] | EMIS Title |
|----------------------|--|----------------------|---|
| 1 A 1 a | Kehrichtverbrennungsanlagen | 2 G | Sprengen und Schiessen |
| 1 A 1 a | Sondermüllverbrennungsanlagen | 3 A 1 | Farben-Anwendung Bau |
| 1 A 1 a & 6 A 1 | Kehrichtdeponien | 3 A 2 | Farben-Anwendung andere industrielle |
| 1 A 1 a & 6 D | Vergärung IG (industriell-gewerblich) | 3 A 1 | Farben-Anwendung Haushalte** |
| 1 A 1 a & 6 D | Vergärung LW (landwirtschaftlich) | 3 A 1 | Farben-Anwendung Holz |
| 1 A 2 a | Eisengiessereien Kupolöfen | 3 A 2 | Farben-Anwendung andere nicht industrielle |
| 1 A 2 a | Stahl-Produktion Wärmeöfen** | 3 A 2 | Farben-Anwendung Autoreparatur |
| 1 A 2 b | Buntmetallgiessereien übriger Betrieb** | 3 B 1 | Elektronik-Reinigung |
| 1 A 2 b & 2 C 3 | Aluminium Produktion | 3 B 1 | Metallreinigung |
| 1 A 2 d | Zellulose-Produktion Feuerung* | 3 B 1 | Reinigung Industrie übrige |
| 1 A 2 fi & 2 A 3 | Feinkeramik Produktion* | 3 B 2 | Chemische Reinigung |
| 1 A 2 fi & 2 A 7 | Glas übrige Produktion* | 3 C | Druckfarben Produktion |
| 1 A 2 fi & 2 A 7 | Glaswolle Produktion Rohprodukt* | 3 C | Farben-Produktion |
| 1 A 2 fi & 2 A 7 | Hohlglass Produktion* | 3 C | Feinchemikalien-Produktion** |
| 1 A 2 fi | Kalkproduktion, Feuerung* | 3 C | Gummi-Verarbeitung |
| 1 A 2 fi | Mischgut Produktion | 3 C | Klebband-Produktion |
| 1 A 2 fi & 2 A 3 | Steinwolle Produktion | 3 C | Klebstoff-Produktion |
| 1 A 2 fi & 2 A 3 | Ziegeleien** | 3 C | Lösungsmittel-Umschlag und -Lager |
| 1 A 2 fi | Zementwerke Feuerung | 3 C | Pharmazeutische Produktion** |
| 1 A 2 fi | Faserplatten Produktion** | 3 C | Polyester-Verarbeitung |
| 1 A 3 a & 1 A 5 | Flugverkehr | 3 C | Polystyrol-Verarbeitung |
| 1 A 3 b i-vii | Strassenverkehr | 3 C | Polyurethan-Verarbeitung |
| 1 A 3 c | Schienerverkehr | 3 C | PVC-Verarbeitung |
| 1 A 3 e | Gasverteilung Netzverluste | 3 C | Gerben von Ledermaterialien |
| 1 A 4 c i | Gastrocknung | 3 D [3 D 3] | Korrosionsschutz im Freien |
| 1 A 4 div. | Off-Road | 3 D 1 [3 D 3] | Lachgasanwendung Spitäler** |
| 1 Energy Model*** | Energie New | 3 D 5 [3 D 2] | Reinigungs- und Lösemittel; Haushalte |
| 1A solid fuels/wood | Holzfeuerungen | 3 D 5 [3 D 2] | Spraydosen Haushalte** |
| 1 B 2 a iv | Raffinerie, Leckverluste | 3 D 5 [3 D 3] | Betonzusatzmittel-Anwendung |
| 1 B 2 a v | Benzinumschlag Tanklager | 3 D 5 [3 D 3] | Coiffeursalons |
| 1 B 2 a v | Benzinumschlag Tankstellen | 3 D 5 [3 D 1] | Druckereien |
| 1 B 2 c | Raffinerie, Abfackelung | 3 D 5 [3 D 3] | Entfernung von Farben und Lacken |
| 2 A 1 | Zementwerke Rohmaterial | 3 D 5 [3 D 3] | Entwachsung von Fahrzeugen |
| 2 A 1 | Zementwerke übriger Betrieb | 3 D 5 [3 D 3] | Fahrzeug-Unterbodenschutz |
| 2 A 2 | Kalkproduktion, Rohmaterial* | 3 D 5 [3 D 3] | Feuerwerke |
| 2 A 2 | Kalkproduktion, übriger Betrieb* | 3 D 5 [3 D 3] | Flugzeug-Enteisung |
| 2 A 5 | Dachpappen Produktion Emissionen aus Bitumen | 3 D 5 [3 D 3] | Gas-Anwendung |
| 2 A 5 | Dachpappen Produktion Voranstrich | 3 D 5 [3 D 3] | Gesundheitswesen, übrige** |
| 2 A 5 | Dachpappen Verlegung Bitumen | 3 D 5 [3 D 3] | Glaswolle Imprägnierung* |
| 2 A 5 | Dachpappen Verlegung Voranstrich | 3 D 5 [3 D 3] | Holzschutzmittel-Anwendung |
| 2 A 6 | Strassenbelagsarbeiten** | 3 D 5 [3 D 3] | Klebstoff-Anwendung |
| 2 A 7 | Gips-Produktion übriger Betrieb** | 3 D 5 [3 D 3] | Kosmetika-Produktion** |
| 2 B 1 | Ammoniak-Produktion* | 3 D 5 [3 D 3] | Kosmetik-Institute |
| 2 B 2 | Salpetersäure Produktion* | 3 D 5 [3 D 3] | Kühlschmiermittel-Verwendung |
| 2 B 4 | Graphit und Siliziumkarbid Produktion* | 3 D 5 [3 D 3] | Lachgasanwendung Haushalt** |
| 2 B 5 | Ammoniumnitrat-Produktion* | 3 D 5 [3 D 3] | Lösungsmittel-Emissionen IG nicht zugeordnet |
| 2 B 5 | Chlorgas-Produktion | 3 D 5 [3 D 3] | Medizinische Praxen** |
| 2 B 5 | Essigsäure-Produktion** | 3 D 5 [3 D 3] | Öl- und Fettgewinnung |
| 2 B 5 | Ethen-Produktion* | 3 D 5 [3 D 3] | Papier- und Karton-Produktion** |
| 2 B 5 | Formaldehyd-Produktion | 3 D 5 [3 D 3] | Parfum- und Aromen-Produktion** |
| 2 B 5 | PVC-Produktion | 3 D 5 [3 D 3] | Pflanzenschutzmittel-Verwendung |
| 2 B 5 | Salzsäure-Produktion** | 3 D 5 [3 D 3] | Pharma-Produkte im Haushalt |
| 2 B 5 | Schwefelsäure-Produktion* | 3 D 5 [3 D 3] | Reinigung Gebäude IGD** |
| 2 C 1 | Eisengiessereien Elektroschmelzöfen | 3 D 5 [3 D 3] | Schmierstoff-Verwendung |
| 2 C 1 | Eisengiessereien übriger Betrieb | 3 D 5 [3 D 3] | Spraydosen IndustrieGewerbe |
| 2 C 1 | Stahl-Produktion Elektroschmelzöfen** | 3 D 5 [3 D 3] | Tabakwaren Konsum |
| 2 C 1 | Stahl-Produktion übriger Betrieb** | 3 D 5 [3 D 3] | Tabakwaren Produktion** |
| 2 C 1 | Stahl-Produktion Walzwerke** | 3 D 5 [3 D 3] | Textilien-Produktion |
| 2 C 5 d | Verzinkereien | 3 D 5 [3 D 3] | Wissenschaftliche Laboratorien |
| 2 C 5 e | Buntmetallgiessereien Elektroöfen** | 3 D 5 [3 D 3] | Steinwolle-Imprägnierung |
| 2 C 5 e | Batterie-Recycling** | 4 div.*** | Landwirtschaft |
| 2 D 1 | Zellulose Produktion übriger Betrieb* | 6 B 1 [6 B] | Kläranlagen Industriell |
| 2 D 1 | Faserplatten Produktion** | 6 B 2 [6 B] | Kläranlagen Kommunal |
| 2 D 1 | Spanplatten Produktion* | 6 C [6 C d] | Krematorien |
| 2 D 2 | Bierbrauereien | 6 C 1 [6 C e] | Abfallverbrennung Land- und Forstwirtschaft |
| 2 D 2 | Branntwein Produktion | 6 C 2 [6 C a] | Spitalabfallverbrennung |
| 2 D 2 | Brot Produktion | 6 C 2 [6 C b] | Kabelabbrand |
| 2 D 2 | Fleischräuchereien | 6 C 2 [6 C b] | Klärschlammverbrennung |
| 2 D 2 | Kaffeeröstereien | 6 C 2 [6 C c] | Abfallverbrennung illegal |
| 2 D 2 | Müllereien | 6 D | Kompostierung Industrie |
| 2 D 2 | Wein Produktion | 6 D | Shredder Anlagen |
| 2 D 2 | Zucker Produktion | 6 D | Biogasaufbereitung (Methanverlust) |
| 2 D 3 | Holzkohle Produktion | 7 C 1 | Kompostierung, Verbreitung als Dünger im Haushalt |
| 2 F div. | Synthetische Gase | 7 D | Brand- und Feuerschäden Immobilien |
| 2 D 3 | Holzbearbeitung | 7 D | Brand- und Feuerschäden Motorfahrzeuge |

* confidential process

** confidential EMIS comment

** work in progress

Annexes

Annex 1: Key Category Analysis (KCA)

A1.1 Methodology

The key category analysis is performed according to the IPCC Good Practice Guidance (IPCC 2000, chapter 7): A Tier 1 level and trend assessment is applied with the proposed threshold of 95%. A Tier 2 key category analysis has also been carried out for this submission with the proposed threshold of 90% of the sum of all level assessments weighted with their relative source uncertainty. All main source categories have been disaggregated into sub-sources (e.g. 2A, 2B, 2C etc.) and gases (CO₂, CH₄, N₂O, HFCs, PFCs, and SF₆).

For some important sources, an even more detailed level of disaggregation has been used in order to clearly identify and isolate the most important sources.

This is the case for the important source category 1A Energy Fuel Combustion sources, where the source categories have been disaggregated further to the level of sub-categories as for example 1A1 Fuel Combustion – Energy Industries, 1A2 Fuel Combustion – Manufacturing Industries, etc. Even further disaggregation has been realized for the source category Transport (1A3) and Other Sectors (1A4):

The source Transport (1A3) has been further split into Civil Aviation (1A3a), Road Transportation (1A3b), Railways (1A3c), and Navigation (1A3d).

A more detailed disaggregation has also been carried out for Other Sectors (1A4) which has been split into Commercial/Institutional (1A4a), Residential (1A4b) and Agriculture/Forestry (1A4c).

Also the categories Mineral Products (2A) and Metal Production (2C) have been disaggregated into source categories. Consumption of Halocarbons and SF₆ (2F) has been split into its source categories 2F1 to 2F9.

Agricultural Soils (4D) have been split into its source categories 4D1 to 4D4.

Uncertainty data have been taken from the uncertainty analysis, where the disaggregation of source and sink categories is in accordance with the key category analysis.

A1.2 KCA Tier 1 2012 without LULUCF categories.

Results of Key Category Analysis Tier 1 – Level and Trend

Table A - 2 Key category analysis Tier 1 2012 (without LULUCF) regarding level and trend.

| Tier 1 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | | | |
|---|--|----------------------------------|--|--|---------------|--------------------|---|--------------------------------------|---|-------------------------|-------------------------|----------------------------------|---------------------------------|---------------------------------|
| No. | A | | | | | B Direct GHG | C | | D Year 2012 Estimate [Gg CO2 eq] | E-L Level Assessm | E-T Trend Assessm | F-T % Contrib. in Trend | M Result level assessm | N Result trend assessm |
| | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | | | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | | | | | | |
| 1 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO2 | 11335.27 | 9016.58 | 17.53% | 0.04013 | 9.5% | KC level | KC trend | |
| 2 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 14.33% | 0.05183 | 12.3% | KC level | KC trend | |
| 3 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CO2 | 2587.68 | 6767.05 | 13.15% | 0.08494 | 20.1% | KC level | KC trend | |
| 4 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 5.91% | 0.02881 | 6.8% | KC level | KC trend | |
| 5 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 5.28% | 0.02471 | 5.8% | KC level | KC trend | |
| 6 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 5.15% | 0.02527 | 6.0% | KC level | KC trend | |
| 7 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 5.13% | 0.01900 | 4.5% | KC level | KC trend | |
| 8 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH4 | 2635.45 | 2496.98 | 4.85% | 0.00132 | 0.3% | KC level | - | |
| 9 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 4.07% | 0.02102 | 5.0% | KC level | KC trend | |
| 10 | 2A1 | 2. Industrial | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 1787.11 | 3.47% | 0.01336 | 3.2% | KC level | KC trend | |
| 11 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 2.88% | 0.01044 | 2.5% | KC level | KC trend | |
| 12 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 1143.10 | 2.22% | 0.00342 | 0.8% | KC level | KC trend | |
| 13 | 2F1 | 2. Industrial | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | | HFC | 0.02 | 1137.81 | 2.21% | 0.02274 | 5.4% | KC level | KC trend | |
| 14 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 1.56% | 0.00261 | 0.6% | KC level | KC trend | |
| 15 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 674.93 | 1.31% | 0.00250 | 0.6% | KC level | KC trend | |
| 16 | 4B | 4. Agriculture | B. Manure Management | | | CH4 | 671.61 | 646.11 | 1.26% | 0.00014 | 0.0% | KC level | - | |
| 17 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CO2 | 547.34 | 540.01 | 1.05% | 0.00015 | 0.0% | KC level | - | |
| 18 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CO2 | 289.73 | 498.74 | 0.97% | 0.00434 | 1.0% | KC level | KC trend | |
| 19 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.88% | 0.01432 | 3.4% | KC level | KC trend | |
| 20 | 4B | 4. Agriculture | B. Manure Management | | | N2O | 454.68 | 335.81 | 0.65% | 0.00213 | 0.5% | KC level | KC trend | |
| 21 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Other Fuels | CO2 | 134.15 | 288.60 | 0.56% | 0.00316 | 0.7% | KC level | KC trend | |
| 22 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 240.28 | 0.47% | 0.00121 | 0.3% | KC level | - | |
| 23 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 220.79 | 0.43% | 0.00192 | 0.5% | KC level | KC trend | |
| 24 | 1B2 | 1. Energy | B. Fugitive Emissions from 2. Oil and Natural Gas | | | CH4 | 263.72 | 169.45 | 0.33% | 0.00174 | 0.4% | - | KC trend | |
| 25 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 158.26 | 0.31% | 0.01021 | 2.4% | - | KC trend | |
| 26 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 155.28 | 0.30% | 0.00389 | 0.9% | - | KC trend | |
| 27 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CO2 | 252.55 | 136.65 | 0.27% | 0.00218 | 0.5% | - | KC trend | |
| 28 | 2F9 | 2. Industrial | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 135.91 | 0.26% | 0.00117 | 0.3% | - | - | |
| 29 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | | CO2 | 111.93 | 121.14 | 0.24% | 0.00025 | 0.1% | - | - | |
| 30 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO2 | 203.58 | 114.80 | 0.22% | 0.00166 | 0.4% | - | KC trend | |
| 31 | 6D | 6. Waste | D. Other | | | CH4 | 29.94 | 113.76 | 0.22% | 0.00169 | 0.4% | - | KC trend | |
| 32 | 2B | 2. Industrial | B. Chemical Industry | | | CO2 | 111.22 | 110.47 | 0.21% | 0.00005 | 0.0% | - | - | |
| 33 | 2A3 | 2. Industrial | A. Mineral Products; Limestone and Dolomite Use, Emissions | | | CO2 | 150.39 | 98.48 | 0.19% | 0.00096 | 0.2% | - | - | |
| 34 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CO2 | 0.00 | 83.59 | 0.16% | 0.00167 | 0.4% | - | KC trend | |
| 35 | 2F9 | 2. Industrial | F. Consumption of Halocarbons and SF6; Other | | | HFC | 0.00 | 76.14 | 0.15% | 0.00152 | 0.4% | - | - | |
| 36 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | N2O | 5.74 | 66.06 | 0.13% | 0.00121 | 0.3% | - | - | |
| 37 | 2A2 | 2. Industrial | A. Mineral Products; Lime Production-CO2 | | | CO2 | 53.35 | 54.26 | 0.11% | 0.00005 | 0.0% | - | - | |
| 38 | 2B | 2. Industrial | B. Chemical Industry | | | N2O | 68.13 | 53.57 | 0.10% | 0.00025 | 0.1% | - | - | |
| 39 | 1A3e | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CO2 | 31.42 | 45.44 | 0.09% | 0.00030 | 0.1% | - | - | |
| 40 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 44.62 | 0.09% | 0.00125 | 0.3% | - | - | |
| 41 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | | CO2 | 28.69 | 39.69 | 0.08% | 0.00024 | 0.1% | - | - | |
| 42 | 1B2 | 1. Energy | B. Fugitive Emissions from 2. Oil and Natural Gas | | | CO2 | 84.62 | 39.29 | 0.08% | 0.00086 | 0.2% | - | - | |
| 43 | 2F8 | 2. Industrial | F. Consumption of Halocarbons and SF6; Electrical Eq. | | | SF6 | 64.04 | 36.81 | 0.07% | 0.00051 | 0.1% | - | - | |
| 44 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 33.78 | 0.07% | 0.00123 | 0.3% | - | - | |
| 45 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CO2 | 54.59 | 33.60 | 0.07% | 0.00039 | 0.1% | - | - | |
| 46 | 2C | 2. Industrial | C. Metal Production; Magnesium Foundries | | | SF6 | 0.00 | 31.72 | 0.06% | 0.00063 | 0.2% | - | - | |
| 47 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | N2O | 142.38 | 27.52 | 0.05% | 0.00222 | 0.5% | - | KC trend | |
| 48 | 6C | 6. Waste | C. Waste Incineration | | | N2O | 19.06 | 25.86 | 0.05% | 0.00015 | 0.0% | - | - | |
| 49 | 6D | 6. Waste | D. Other | | | N2O | 5.82 | 25.55 | 0.05% | 0.00040 | 0.1% | - | - | |
| 50 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | N2O | 20.85 | 20.96 | 0.04% | 0.00001 | 0.0% | - | - | |
| 51 | 4D4 | 4. Agriculture | D. Agricultural Soils; Use of sewage sludge as fertilizers | | | N2O | 28.30 | 20.85 | 0.04% | 0.00013 | 0.0% | - | - | |
| 52 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | N2O | 27.72 | 19.75 | 0.04% | 0.00014 | 0.0% | - | - | |
| 53 | 2F7 | 2. Industrial | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | | SF6 | 0.00 | 19.55 | 0.04% | 0.00039 | 0.1% | - | - | |
| 54 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CH4 | 97.47 | 18.82 | 0.04% | 0.00152 | 0.4% | - | - | |
| 55 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | N2O | 25.94 | 18.69 | 0.04% | 0.00013 | 0.0% | - | - | |
| 56 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CO2 | 41.45 | 18.48 | 0.04% | 0.00044 | 0.1% | - | - | |
| 57 | 2F2 | 2. Industrial | F. Consumption of Halocarbons and SF6; Hard Foam | | | HFC | 0.00 | 13.27 | 0.03% | 0.00027 | 0.1% | - | - | |
| 58 | 2F4 | 2. Industrial | F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and Other | | | HFC | 0.00 | 13.22 | 0.03% | 0.00026 | 0.1% | - | - | |
| 59 | 2F9 | 2. Industrial | F. Consumption of Halocarbons and SF6; Other | | | PFC | 0.00 | 13.10 | 0.03% | 0.00026 | 0.1% | - | - | |
| 60 | 7 | 7. Other | | | | CO2 | 10.96 | 12.92 | 0.03% | 0.00005 | 0.0% | - | - | |
| 61 | 6C | 6. Waste | C. Waste Incineration | | | CO2 | 54.10 | 12.34 | 0.02% | 0.00080 | 0.2% | - | - | |
| 62 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | N2O | 13.92 | 11.76 | 0.02% | 0.00004 | 0.0% | - | - | |
| 63 | 2C1 | 2. Industrial | C. Metal Production; Steel Production | | | CO2 | 9.20 | 9.89 | 0.02% | 0.00002 | 0.0% | - | - | |
| 64 | 6B | 6. Waste | B. Wastewater Handling | | | CH4 | 4.65 | 9.28 | 0.02% | 0.00010 | 0.0% | - | - | |
| 65 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | N2O | 10.79 | 9.07 | 0.02% | 0.00003 | 0.0% | - | - | |
| 66 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | N2O | 11.70 | 7.80 | 0.02% | 0.00007 | 0.0% | - | - | |
| 67 | 2A7 | 2. Industrial | A. Mineral Products; Other non-specified-CO2 | | | CO2 | 15.30 | 7.68 | 0.01% | 0.00014 | 0.0% | - | - | |
| 68 | 2F5 | 2. Industrial | F. Consumption of Halocarbons and SF6; Solvents | | | PFC | 0.00 | 7.29 | 0.01% | 0.00015 | 0.0% | - | - | |
| 69 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Biomass | N2O | 1.68 | 6.85 | 0.01% | 0.00010 | 0.0% | - | - | |
| 70 | 2F7 | 2. Industrial | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | | PFC | 0.00 | 6.54 | 0.01% | 0.00013 | 0.0% | - | - | |
| 71 | 6C | 6. Waste | C. Waste Incineration | | | CH4 | 11.58 | 6.14 | 0.01% | 0.00010 | 0.0% | - | - | |
| 72 | 2F1 | 2. Industrial | F. Consumption of Halocarbons and SF6; Refrigeration | | | PFC | 0.04 | 6.14 | 0.01% | 0.00012 | 0.0% | - | - | |
| 73 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CH4 | 3.24 | 6.10 | 0.01% | 0.00006 | 0.0% | - | - | |
| 74 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | N2O | 4.96 | 5.54 | 0.01% | 0.00001 | 0.0% | - | - | |
| 75 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Other Fuels | N2O | 2.28 | 5.24 | 0.01% | 0.00006 | 0.0% | - | - | |
| 76 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Gaseous Fuels | CH4 | 2.66 | 4.74 | 0.01% | 0.00004 | 0.0% | - | - | |
| 77 | 2F5 | 2. Industrial | F. Consumption of Halocarbons and SF6; Solvents | | | HFC | 0.00 | 4.59 | 0.01% | 0.00009 | 0.0% | - | - | |
| 78 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Biomass | CH4 | 9.74 | 4.13 | 0.01% | 0.00011 | 0.0% | - | - | |
| 79 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Gaseous Fuels | CH4 | 2.41 | 3.83 | 0.01% | 0.00003 | 0.0% | - | - | |
| 80 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | N2O | 0.00 | 3.47 | 0.01% | 0.00007 | 0.0% | - | - | |

Table A - 2 continued. Key category analysis Tier 1 2012 (without LULUCF) regarding level and trend.

| Tier 1 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | |
|---|--|---------------|---|--|-----------------|---|--|------------------|------------------|---------------------|-------------------------|-------------------------|
| A | | | | | B | C | D | E-L | E-T | F-T | M | N |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO ₂ eq] | Year 2012 Estimate [Gg CO ₂ eq] | Level Assessment | Trend Assessment | % Contrib. in Trend | Result level assessment | Result trend assessment |
| 81 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Biomass | N ₂ O | 1.45 | 3.23 | 0.01% | 0.00004 | 0.0% | - |
| 82 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | N ₂ O | 2.15 | 2.86 | 0.01% | 0.00002 | 0.0% | - |
| 83 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Solid Fuels | N ₂ O | 6.44 | 2.41 | 0.00% | 0.00008 | 0.0% | - |
| 84 | 2B | 1. Industrial | B. Chemical Industry | | | CH ₄ | 1.54 | 2.39 | 0.00% | 0.00002 | 0.0% | - |
| 85 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CH ₄ | 3.71 | 2.28 | 0.00% | 0.00003 | 0.0% | - |
| 86 | 1A3b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CH ₄ | 6.00 | 2.26 | 0.00% | 0.00007 | 0.0% | - |
| 87 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Biomass | CH ₄ | 2.46 | 1.56 | 0.00% | 0.00002 | 0.0% | - |
| 88 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | N ₂ O | 0.79 | 1.46 | 0.00% | 0.00001 | 0.0% | - |
| 89 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CH ₄ | 1.62 | 1.43 | 0.00% | 0.00000 | 0.0% | - |
| 90 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | CH ₄ | 3.06 | 1.40 | 0.00% | 0.00003 | 0.0% | - |
| 91 | 2C5 | 2. Industrial | F. Metal Production; Non-ferrous metals-CO ₂ | | CO ₂ | | 1.65 | 1.36 | 0.00% | 0.00000 | 0.0% | - |
| 92 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | N ₂ O | 2.49 | 1.35 | 0.00% | 0.00002 | 0.0% | - |
| 93 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Gaseous Fuels | N ₂ O | 0.59 | 1.15 | 0.00% | 0.00001 | 0.0% | - |
| 94 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | N ₂ O | 2.01 | 1.13 | 0.00% | 0.00002 | 0.0% | - |
| 95 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CH ₄ | 0.65 | 1.12 | 0.00% | 0.00001 | 0.0% | - |
| 96 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | CH ₄ | 2.32 | 1.04 | 0.00% | 0.00002 | 0.0% | - |
| 97 | 2G | 2. Industrial | F. Other | | CO ₂ | | 1.04 | 0.91 | 0.00% | 0.00000 | 0.0% | - |
| 98 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | N ₂ O | 0.66 | 0.82 | 0.00% | 0.00000 | 0.0% | - |
| 99 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Gaseous Fuels | N ₂ O | 0.55 | 0.82 | 0.00% | 0.00001 | 0.0% | - |
| 100 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | N ₂ O | 0.00 | 0.70 | 0.00% | 0.00001 | 0.0% | - |
| 101 | 1B2 | 1. Energy | B. Fugitive Emissions from Fuels | 2. Oil and Natural Gas | | N ₂ O | 0.62 | 0.68 | 0.00% | 0.00000 | 0.0% | - |
| 102 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CH ₄ | 0.49 | 0.67 | 0.00% | 0.00000 | 0.0% | - |
| 103 | 7 | 7. Other | | | | N ₂ O | 0.62 | 0.62 | 0.00% | 0.00000 | 0.0% | - |
| 104 | 7 | 7. Other | | | | CH ₄ | 0.55 | 0.57 | 0.00% | 0.00000 | 0.0% | - |
| 105 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CH ₄ | 1.38 | 0.55 | 0.00% | 0.00002 | 0.0% | - |
| 106 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | N ₂ O | 0.60 | 0.55 | 0.00% | 0.00000 | 0.0% | - |
| 107 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | CH ₄ | 0.58 | 0.54 | 0.00% | 0.00000 | 0.0% | - |
| 108 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | N ₂ O | 0.38 | 0.52 | 0.00% | 0.00000 | 0.0% | - |
| 109 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | CH ₄ | 0.33 | 0.43 | 0.00% | 0.00000 | 0.0% | - |
| 110 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Other Fuels | CH ₄ | 0.54 | 0.37 | 0.00% | 0.00000 | 0.0% | - |
| 111 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | N ₂ O | 0.21 | 0.35 | 0.00% | 0.00000 | 0.0% | - |
| 112 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | N ₂ O | 0.16 | 0.28 | 0.00% | 0.00000 | 0.0% | - |
| 113 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CH ₄ | 0.24 | 0.25 | 0.00% | 0.00000 | 0.0% | - |
| 114 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | N ₂ O | 0.29 | 0.18 | 0.00% | 0.00000 | 0.0% | - |
| 115 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | CH ₄ | 0.80 | 0.15 | 0.00% | 0.00001 | 0.0% | - |
| 116 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Solid Fuels | CH ₄ | 0.40 | 0.14 | 0.00% | 0.00000 | 0.0% | - |
| 117 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CH ₄ | 0.16 | 0.12 | 0.00% | 0.00000 | 0.0% | - |
| 118 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CH ₄ | 0.00 | 0.10 | 0.00% | 0.00000 | 0.0% | - |
| 119 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CH ₄ | 0.09 | 0.04 | 0.00% | 0.00000 | 0.0% | - |
| 120 | 1A3el | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CH ₄ | 0.06 | 0.03 | 0.00% | 0.00000 | 0.0% | - |
| 121 | 1A3el | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | N ₂ O | 0.02 | 0.03 | 0.00% | 0.00000 | 0.0% | - |
| 122 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | CH ₄ | 0.00 | 0.02 | 0.00% | 0.00000 | 0.0% | - |
| 123 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | CH ₄ | 0.01 | 0.02 | 0.00% | 0.00000 | 0.0% | - |
| 124 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | N ₂ O | 0.02 | 0.01 | 0.00% | 0.00000 | 0.0% | - |
| 125 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | CH ₄ | 0.01 | 0.01 | 0.00% | 0.00000 | 0.0% | - |
| 126 | 6D | 6. Waste | D. Other | | | CO ₂ | 0.00 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 127 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CO ₂ | 9.24 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 128 | 2C3 | 2. Industrial | F. Metal Production; Aluminium Production-CO ₂ | | | CO ₂ | 139.26 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 129 | 2C3 | 2. Industrial | F. Metal Production; Aluminium Production-PFC | | | PFC | 100.17 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 130 | 2C | 2. Industrial | F. Metal Production; Aluminium Foundries | | | SF ₆ | 0.00 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 131 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CO ₂ | 44.84 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 132 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CH ₄ | 0.10 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 133 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | N ₂ O | 0.24 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 134 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CO ₂ | 9.24 | 0.00 | 0.00% | 0.00000 | 0.0% | - |
| 135 | 6D | 6. Waste | D. Other | | | CO ₂ | 0.00 | 0.00 | 0.00% | 0.00000 | 0.0% | - |

A1.3 KCA Tier 1 2012 including LULUCF categories

Results of Key Category Analysis Tier 1 – Level and Trend

Table A - 3 Key category analysis Tier 1 2012 (with LULUCF) regarding level and trend.

| Tier 1 Key category analysis 2012 with LULUCF categories | | | | | | | | | | | | | |
|--|--|----------------------------------|--|---|-----------------|---|--|----------------------|----------------------|----------------------------|---------------------------|---------------------------|----------|
| No. | A IPCC Source Categories and fuels if applicable (with LULUCF categories) | | | | B Direct GHG | C Base Year 1990 Estimate [Gg CO2 eq] | D Year 2012 Estimate [Gg CO2 eq] | E-L Level Assessm | E-T Trend Assessm | F-T % Contrib. in Trend | M Result level assessm | N Result trend assessm | |
| 1 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO2 | 11335.27 | 9016.58 | 16.21% | 0.03753 | 9.2% | KC level | KC trend |
| 2 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 13.26% | 0.04828 | 11.8% | KC level | KC trend |
| 3 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CO2 | 2587.68 | 6767.05 | 12.16% | 0.07824 | 19.1% | KC level | KC trend |
| 4 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 5.46% | 0.02679 | 6.5% | KC level | KC trend |
| 5 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 4.88% | 0.02273 | 5.5% | KC level | KC trend |
| 6 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 4.76% | 0.02324 | 5.7% | KC level | KC trend |
| 7 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 4.75% | 0.01770 | 4.3% | KC level | KC trend |
| 8 | 4A | 4. Agriculture | A. Enteric Fermentation | | | CH4 | 2635.45 | 2496.98 | 4.49% | 0.00134 | 0.3% | KC level | KC trend |
| 9 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CO2 | -2416.89 | -2134.56 | 3.84% | 0.00409 | 1.0% | KC level | KC trend |
| 10 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 3.77% | 0.01934 | 4.7% | KC level | KC trend |
| 11 | 2A1 | 2. Industry | A. Mineral Products; Cement Production-CO2 | | | CO2 | 2524.68 | 1787.11 | 3.21% | 0.01244 | 3.0% | KC level | KC trend |
| 12 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 2.67% | 0.00959 | 2.3% | KC level | KC trend |
| 13 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | | N2O | 1351.48 | 1143.10 | 2.05% | 0.00322 | 0.8% | KC level | KC trend |
| 14 | 2F1 | 2. Industry | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | | HFC | 0.02 | 1137.81 | 2.05% | 0.02098 | 5.1% | KC level | KC trend |
| 15 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 1.45% | 0.00237 | 0.6% | KC level | KC trend |
| 16 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | | CO2 | 345.17 | 707.27 | 1.27% | 0.00683 | 1.7% | KC level | KC trend |
| 17 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | | N2O | 822.48 | 674.93 | 1.21% | 0.00234 | 0.6% | KC level | KC trend |
| 18 | 4B | 4. Agriculture | B. Manure Management | | | CH4 | 671.61 | 646.11 | 1.16% | 0.00016 | 0.0% | KC level | - |
| 19 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CO2 | 547.34 | 540.01 | 0.97% | 0.00012 | 0.0% | KC level | - |
| 20 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | | CO2 | -621.57 | -518.61 | 0.93% | 0.00161 | 0.4% | KC level | KC trend |
| 21 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CO2 | 289.73 | 498.74 | 0.90% | 0.00399 | 1.0% | KC level | KC trend |
| 22 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.82% | 0.01327 | 3.2% | KC level | KC trend |
| 23 | 4B | 4. Agriculture | B. Manure Management | | | N2O | 454.68 | 335.81 | 0.60% | 0.00198 | 0.5% | KC level | KC trend |
| 24 | 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | | CO2 | 382.71 | 302.93 | 0.54% | 0.00129 | 0.3% | KC level | - |
| 25 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Other Fuels | CO2 | 134.15 | 288.60 | 0.52% | 0.00291 | 0.7% | KC level | KC trend |
| 26 | 6B | 6. Waste | B. Wastewater Handling | | | N2O | 184.72 | 240.28 | 0.43% | 0.00111 | 0.3% | KC level | - |
| 27 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | | N2O | 128.10 | 220.79 | 0.40% | 0.00177 | 0.4% | KC level | KC trend |
| 28 | 1B2 | 1. Energy | B. Fugitive Emissions from Oil and Natural Gas | | | CH4 | 263.72 | 169.45 | 0.30% | 0.00162 | 0.4% | KC level | KC trend |
| 29 | 5C2 | 5. LULUCF | C. Grassland | 2. Land converted to Grassland | | CO2 | 59.85 | 169.07 | 0.30% | 0.00204 | 0.5% | - | KC trend |
| 30 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | | CH4 | 688.16 | 158.26 | 0.28% | 0.00945 | 2.3% | - | KC trend |
| 31 | 3 | 3. Solvent and Other Product Use | | | | CO2 | 360.04 | 155.28 | 0.28% | 0.00361 | 0.9% | - | KC trend |
| 32 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CO2 | 252.55 | 136.65 | 0.25% | 0.00202 | 0.5% | - | KC trend |
| 33 | 2F9 | 2. Industry | F. Consumption of Halocarbons and SF6; Other | | | SF6 | 79.58 | 135.91 | 0.24% | 0.00108 | 0.3% | - | - |
| 34 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CO2 | 107.09 | 134.74 | 0.24% | 0.00056 | 0.1% | - | - |
| 35 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | | CO2 | 111.93 | 121.14 | 0.22% | 0.00022 | 0.1% | - | - |
| 36 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO2 | 203.58 | 114.80 | 0.21% | 0.00154 | 0.4% | - | KC trend |
| 37 | 6D | 6. Waste | D. Other | | | CH4 | 29.94 | 113.76 | 0.20% | 0.00156 | 0.4% | - | KC trend |
| 38 | 5F2 | 5. LULUCF | F. Other Land | 2. Land converted to Other Land | | CO2 | 91.98 | 112.28 | 0.20% | 0.00042 | 0.1% | - | - |
| 39 | 2B | 2. Industry | B. Chemical Industry | | | CO2 | 111.22 | 110.47 | 0.20% | 0.00004 | 0.0% | - | - |
| 40 | 2A3 | 2. Industry | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | | CO2 | 150.39 | 98.48 | 0.18% | 0.00089 | 0.2% | - | - |
| 41 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CO2 | 0.00 | 83.59 | 0.15% | 0.00154 | 0.4% | - | KC trend |
| 42 | 2F9 | 2. Industry | F. Consumption of Halocarbons and SF6; Other | | | HFC | 0.00 | 76.14 | 0.14% | 0.00140 | 0.3% | - | KC trend |
| 43 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | N2O | 5.74 | 66.06 | 0.12% | 0.00111 | 0.3% | - | - |
| 44 | 2A2 | 2. Industry | A. Mineral Products; Lime Production-CO2 | | | CO2 | 53.35 | 54.26 | 0.10% | 0.00004 | 0.0% | - | - |
| 45 | 2B | 2. Industry | B. Chemical Industry | | | N2O | 68.13 | 53.57 | 0.10% | 0.00024 | 0.1% | - | - |
| 46 | 1A3e | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CO2 | 31.42 | 45.44 | 0.08% | 0.00027 | 0.1% | - | - |
| 47 | 3 | 3. Solvent and Other Product Use | | | | N2O | 110.14 | 44.62 | 0.08% | 0.00116 | 0.3% | - | - |
| 48 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | | CO2 | 28.69 | 39.69 | 0.07% | 0.00022 | 0.1% | - | - |
| 49 | 1B2 | 1. Energy | B. Fugitive Emissions from Oil and Natural Gas | | | CO2 | 84.62 | 39.29 | 0.07% | 0.00080 | 0.2% | - | - |
| 50 | 2F8 | 2. Industry | F. Consumption of Halocarbons and SF6; Electrical Eq. | | | SF6 | 64.04 | 36.81 | 0.07% | 0.00047 | 0.1% | - | - |
| 51 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 33.78 | 0.06% | 0.00114 | 0.3% | - | - |
| 52 | 5E1 | 5. LULUCF | E. Settlements | 1. Settlements remaining Settlements | | CO2 | 3.60 | 33.69 | 0.06% | 0.00056 | 0.1% | - | - |
| 53 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CO2 | 54.59 | 33.60 | 0.06% | 0.00036 | 0.1% | - | - |
| 54 | 2C | 2. Industry | C. Metal Production; Magnesium Foundries | | | SF6 | 0.00 | 31.72 | 0.06% | 0.00058 | 0.1% | - | - |
| 55 | 5D2 | 5. LULUCF | D. Wetlands | 2. Land converted to Wetlands | | CO2 | 20.06 | 28.95 | 0.05% | 0.00017 | 0.0% | - | - |
| 56 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | N2O | 142.38 | 27.52 | 0.05% | 0.00205 | 0.5% | - | KC trend |
| 57 | 6C | 6. Waste | C. Waste Incineration | | | N2O | 19.06 | 25.86 | 0.05% | 0.00013 | 0.0% | - | - |
| 58 | 6D | 6. Waste | D. Other | | | N2O | 5.82 | 25.55 | 0.05% | 0.00037 | 0.1% | - | - |
| 59 | 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | | CO2 | 43.33 | 22.26 | 0.04% | 0.00037 | 0.1% | - | - |
| 60 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | N2O | 20.85 | 20.96 | 0.04% | 0.00001 | 0.0% | - | - |
| 61 | 4D4 | 4. Agriculture | D. Agricultural Soils; Use of sewage sludge as fertilizers | | | N2O | 28.30 | 20.85 | 0.04% | 0.00012 | 0.0% | - | - |
| 62 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | N2O | 27.72 | 19.75 | 0.04% | 0.00013 | 0.0% | - | - |
| 63 | 2F7 | 2. Industry | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | | SF6 | 0.00 | 19.55 | 0.04% | 0.00036 | 0.1% | - | - |
| 64 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CH4 | 97.47 | 18.82 | 0.03% | 0.00141 | 0.3% | - | KC trend |
| 65 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | N2O | 25.94 | 18.69 | 0.03% | 0.00012 | 0.0% | - | - |
| 66 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CO2 | 41.45 | 18.48 | 0.03% | 0.00040 | 0.1% | - | - |
| 67 | 2F2 | 2. Industry | F. Consumption of Halocarbons and SF6; Hard Foam | | | HFC | 0.00 | 13.27 | 0.02% | 0.00024 | 0.1% | - | - |
| 68 | 2F4 | 2. Industry | F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and Other | | | HFC | 0.00 | 13.22 | 0.02% | 0.00024 | 0.1% | - | - |
| 69 | 2F9 | 2. Industry | F. Consumption of Halocarbons and SF6; Other | | | PFC | 0.00 | 13.10 | 0.02% | 0.00024 | 0.1% | - | - |
| 70 | 7 | 7. Other | | | | CO2 | 10.96 | 12.92 | 0.02% | 0.00004 | 0.0% | - | - |
| 71 | 6C | 6. Waste | C. Waste Incineration | | | CO2 | 54.10 | 12.34 | 0.02% | 0.00074 | 0.2% | - | - |
| 72 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Liquid Fuels | N2O | 13.92 | 11.76 | 0.02% | 0.00003 | 0.0% | - | - |
| 73 | 2C1 | 2. Industry | C. Metal Production; Steel Production | | | CO2 | 9.20 | 9.89 | 0.02% | 0.00002 | 0.0% | - | - |
| 74 | 6B | 6. Waste | B. Wastewater Handling | | | CH4 | 4.65 | 9.28 | 0.02% | 0.00009 | 0.0% | - | - |
| 75 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | N2O | 10.79 | 9.07 | 0.02% | 0.00003 | 0.0% | - | - |
| 76 | 5D2 | 5. LULUCF | D. Land converted to Wetlands | 5. (II) Non-CO2 emissions from drainage of soils and wetlands | | CH4 | 9.03 | 9.03 | 0.02% | 0.00000 | 0.0% | - | - |
| 77 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutional | Liquid Fuels | N2O | 11.70 | 7.80 | 0.01% | 0.00007 | 0.0% | - | - |
| 78 | 2A7 | 2. Industry | A. Mineral Products; Other non-specified-CO2 | | | CO2 | 15.30 | 7.68 | 0.01% | 0.00013 | 0.0% | - | - |
| 79 | 2F5 | 2. Industry | F. Consumption of Halocarbons and SF6; Solvents | | | PFC | 0.00 | 7.29 | 0.01% | 0.00013 | 0.0% | - | - |
| 80 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construction | Biomass | N2O | 1.68 | 6.85 | 0.01% | 0.00010 | 0.0% | - | - |

Table A – 3 continued. Key category analysis Tier 1 2012 (with LULUCF) regarding level and trend.

| Tier 1 Key category analysis 2012 with LULUCF categories | | | | | | | | | | | | | |
|--|---|---------------|--|--|----------------|-------------------------------------|--------------------------------|---------------|---------------|---------------------|----------------------|----------------------|---|
| A | | | | | B | C | D | E-L | E-T | F-T | M | N | |
| No. | IPCC Source Categories and fuels if applicable (with LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | Level Assessm | Trend Assessm | % Contrib. in Trend | Result level assessm | Result trend assessm | |
| 81 | 2F7 | 2. Industrial | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | PFC | 0.00 | 6.54 | 0.01% | 0.00012 | 0.0% | - | - | |
| 82 | 6C | 6. Waste | C. Waste Incineration | | CH4 | 11.58 | 6.14 | 0.01% | 0.00009 | 0.0% | - | - | |
| 83 | 2F1 | 2. Industrial | F. Consumption of Halocarbons and SF6; Refrigeration | | PFC | 0.04 | 6.14 | 0.01% | 0.00011 | 0.0% | - | - | |
| 84 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CH4 | 3.24 | 6.10 | 0.01% | 0.00005 | 0.0% | - | - |
| 85 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | N2O | 4.96 | 5.54 | 0.01% | 0.00001 | 0.0% | - | - |
| 86 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Other Fuels | N2O | 2.28 | 5.24 | 0.01% | 0.00006 | 0.0% | - | - |
| 87 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Gaseous Fuels | CH4 | 2.66 | 4.74 | 0.01% | 0.00004 | 0.0% | - | - |
| 88 | 2F5 | 2. Industrial | F. Consumption of Halocarbons and SF6; Solvents | | HFC | 0.00 | 4.59 | 0.01% | 0.00008 | 0.0% | - | - | |
| 89 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Instituti | Biomass | CH4 | 9.74 | 4.13 | 0.01% | 0.00010 | 0.0% | - | - |
| 90 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Instituti | Gaseous Fuels | CH4 | 2.41 | 3.83 | 0.01% | 0.00003 | 0.0% | - | - |
| 91 | 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | N2O | 5.58 | 3.58 | 0.01% | 0.00003 | 0.0% | - | - | |
| 92 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | N2O | 0.00 | 3.47 | 0.01% | 0.00006 | 0.0% | - | - |
| 93 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Instituti | Biomass | N2O | 1.45 | 3.23 | 0.01% | 0.00003 | 0.0% | - | - |
| 94 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | N2O | 2.15 | 2.86 | 0.01% | 0.00001 | 0.0% | - | - |
| 95 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Solid Fuels | N2O | 6.44 | 2.41 | 0.00% | 0.00007 | 0.0% | - | - |
| 96 | 2B | 2. Industrial | B. Chemical Industry | | CH4 | 1.54 | 2.39 | 0.00% | 0.00002 | 0.0% | - | - | |
| 97 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CH4 | 3.71 | 2.28 | 0.00% | 0.00002 | 0.0% | - | - |
| 98 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CH4 | 6.00 | 2.26 | 0.00% | 0.00007 | 0.0% | - | - |
| 99 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Biomass | CH4 | 2.46 | 1.56 | 0.00% | 0.00002 | 0.0% | - | - |
| 100 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | N2O | 0.79 | 1.46 | 0.00% | 0.00001 | 0.0% | - | - |
| 101 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CH4 | 1.62 | 1.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 102 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Instituti | Liquid Fuels | CH4 | 3.06 | 1.40 | 0.00% | 0.00003 | 0.0% | - | - |
| 103 | 2C5 | 2. Industrial | C. Metal Production; Non-ferrous metals-CO2 | | CO2 | 1.65 | 1.36 | 0.00% | 0.00000 | 0.0% | - | - | |
| 104 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | N2O | 2.49 | 1.35 | 0.00% | 0.00002 | 0.0% | - | - |
| 105 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Gaseous Fuels | N2O | 0.59 | 1.15 | 0.00% | 0.00001 | 0.0% | - | - |
| 106 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | N2O | 2.01 | 1.13 | 0.00% | 0.00002 | 0.0% | - | - |
| 107 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CH4 | 0.65 | 1.12 | 0.00% | 0.00001 | 0.0% | - | - |
| 108 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Liquid Fuels | CH4 | 2.32 | 1.04 | 0.00% | 0.00002 | 0.0% | - | - |
| 109 | 2G | 2. Industrial | G. Other | | CO2 | 1.04 | 0.91 | 0.00% | 0.00000 | 0.0% | - | - | |
| 110 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | N2O | 0.66 | 0.82 | 0.00% | 0.00000 | 0.0% | - | - |
| 111 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Instituti | Gaseous Fuels | N2O | 0.55 | 0.82 | 0.00% | 0.00001 | 0.0% | - | - |
| 112 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | N2O | 0.00 | 0.70 | 0.00% | 0.00001 | 0.0% | - | - |
| 113 | 1B2 | 1. Energy | B. Fugitive Emissions from | 2. Oil and Natural Gas | | N2O | 0.62 | 0.68 | 0.00% | 0.00000 | 0.0% | - | - |
| 114 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CH4 | 0.49 | 0.67 | 0.00% | 0.00000 | 0.0% | - | - |
| 115 | 7 | 7. Other | | | | N2O | 0.62 | 0.62 | 0.00% | 0.00000 | 0.0% | - | - |
| 116 | 7 | 7. Other | | | | CH4 | 0.55 | 0.57 | 0.00% | 0.00000 | 0.0% | - | - |
| 117 | 5D1 | 5. LULUCF | D. Wetlands | 1. Wetlands remaining Wetlands | | CO2 | 2.87 | 0.56 | 0.00% | 0.00004 | 0.0% | - | - |
| 118 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CH4 | 1.38 | 0.55 | 0.00% | 0.00001 | 0.0% | - | - |
| 119 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | N2O | 0.60 | 0.55 | 0.00% | 0.00000 | 0.0% | - | - |
| 120 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | CH4 | 0.58 | 0.54 | 0.00% | 0.00000 | 0.0% | - | - |
| 121 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | N2O | 0.38 | 0.52 | 0.00% | 0.00000 | 0.0% | - | - |
| 122 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | Biomass Burnin | CO2 | 25.36 | 0.51 | 0.00% | 0.00045 | 0.1% | - | - |
| 123 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | CH4 | 0.33 | 0.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 124 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CH4 | 20.90 | 0.42 | 0.00% | 0.00037 | 0.1% | - | - |
| 125 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Other Fuels | CH4 | 0.54 | 0.37 | 0.00% | 0.00000 | 0.0% | - | - |
| 126 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | N2O | 0.21 | 0.35 | 0.00% | 0.00000 | 0.0% | - | - |
| 127 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | N2O | 0.16 | 0.28 | 0.00% | 0.00000 | 0.0% | - | - |
| 128 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CH4 | 0.24 | 0.25 | 0.00% | 0.00000 | 0.0% | - | - |
| 129 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | N2O | 0.29 | 0.18 | 0.00% | 0.00000 | 0.0% | - | - |
| 130 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | CH4 | 0.80 | 0.15 | 0.00% | 0.00001 | 0.0% | - | - |
| 131 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Solid Fuels | CH4 | 0.40 | 0.14 | 0.00% | 0.00000 | 0.0% | - | - |
| 132 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CH4 | 0.16 | 0.12 | 0.00% | 0.00000 | 0.0% | - | - |
| 133 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CH4 | 0.00 | 0.10 | 0.00% | 0.00000 | 0.0% | - | - |
| 134 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | N2O | 4.77 | 0.09 | 0.00% | 0.00008 | 0.0% | - | - |
| 135 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CH4 | 0.09 | 0.04 | 0.00% | 0.00000 | 0.0% | - | - |
| 136 | 1A3ei | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CH4 | 0.06 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - |
| 137 | 1A3ei | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | N2O | 0.02 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - |
| 138 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | CH4 | 0.00 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 139 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | CH4 | 0.01 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 140 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | N2O | 0.02 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 141 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | CH4 | 0.01 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 142 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CH4 | 0.41 | 0.00 | 0.00% | 0.00001 | 0.0% | - | - |
| 143 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | N2O | 0.19 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - |
| 144 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CO2 | 44.84 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - |
| 145 | 2C3 | 2. Industrial | C. Metal Production; Aluminium Production-CO2 | | CO2 | 139.26 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - | |
| 146 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CO2 | 9.24 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - | |
| 147 | 6D | 6. Waste | D. Other | | CO2 | 0.00 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - | |
| 148 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CH4 | 0.10 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - |
| 149 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | N2O | 0.24 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - |
| 150 | 2C3 | 2. Industrial | C. Metal Production; Aluminium Production-PFC | | PFC | 100.17 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - | |
| 151 | 2C | 2. Industrial | C. Metal Production; Aluminium Foundries | | SF6 | 0.00 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - | |

A1.4 KCA Tier 2 2012 without LULUCF categories.

Results of Key Category Analysis Tier 2 – Level and Trend

Table A - 4 Key category analysis Tier 2 2012 (without LULUCF) regarding level and trend.

| Tier 2 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | | |
|---|--|----------------------------------|--|-----------------------------------|----------------|-------------------------------------|--------------------------------|---------------------------------|---------------------------------|---------------------|----------------------|----------------------|----------|
| A | | | | | B | C | D | E-L | E-T | F-T | M | N | |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | Level Assessm. with Uncertainty | Trend Assessm. with Uncertainty | % Contrib. in Trend | Result level assessm | Result trend assessm | |
| 1 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | N2O | 822.48 | 674.93 | 2.18% | 0.00415 | 8.1% | KC level | KC trend | |
| 2 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | N2O | 1351.48 | 1143.10 | 1.85% | 0.00285 | 5.6% | KC level | KC trend | |
| 3 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 1.67% | 0.00781 | 15.3% | KC level | KC trend |
| 4 | 4A | 4. Agriculture | A. Enteric Fermentation | | CH4 | 2635.45 | 2496.98 | 0.89% | 0.00024 | 0.5% | KC level | - | |
| 5 | 4B | 4. Agriculture | B. Manure Management | | CH4 | 671.61 | 646.11 | 0.68% | 0.00008 | 0.2% | KC level | - | |
| 6 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Gasoline | CO2 | 11335.27 | 9016.58 | 0.45% | 0.00103 | 2.0% | KC level | KC trend |
| 7 | 4B | 4. Agriculture | B. Manure Management | | N2O | 454.68 | 335.81 | 0.41% | 0.00135 | 2.6% | KC level | KC trend | |
| 8 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | N2O | 128.10 | 220.79 | 0.36% | 0.00163 | 3.2% | KC level | KC trend | |
| 9 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Diesel | CO2 | 2587.68 | 6767.05 | 0.29% | 0.00190 | 3.7% | KC level | KC trend |
| 10 | 2F1 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | HFC | 0.02 | 1137.81 | 0.27% | 0.00273 | 5.3% | KC level | KC trend | |
| 11 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Gaseous Fuels | CO2 | 1424.38 | 2649.60 | 0.26% | 0.00127 | 2.5% | KC level | KC trend |
| 12 | 6B | 6. Waste | B. Wastewater Handling | | N2O | 184.72 | 240.28 | 0.23% | 0.00061 | 1.2% | KC level | KC trend | |
| 13 | 6D | 6. Waste | D. Other | | CH4 | 29.94 | 113.76 | 0.22% | 0.00170 | 3.3% | KC level | KC trend | |
| 14 | 2F9 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Other | | SF6 | 79.58 | 135.91 | 0.21% | 0.00094 | 1.8% | KC level | KC trend | |
| 15 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Gaseous Fuels | CO2 | 1074.09 | 2096.41 | 0.20% | 0.00105 | 2.1% | KC level | KC trend |
| 16 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 0.20% | 0.00072 | 1.4% | KC level | KC trend |
| 17 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CH4 | 688.16 | 158.26 | 0.18% | 0.00596 | 11.7% | KC level | KC trend | |
| 18 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Other Fuels | CO2 | 134.15 | 288.60 | 0.18% | 0.00100 | 2.0% | KC level | KC trend |
| 19 | 3 | 3. Solvent and Other Product Use | | | CO2 | 360.04 | 155.28 | 0.15% | 0.00195 | 3.8% | KC level | KC trend | |
| 20 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Gaseous Fuels | CO2 | 987.24 | 1482.76 | 0.14% | 0.00052 | 1.0% | KC level | KC trend |
| 21 | 2F9 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Other | | HFC | 0.00 | 76.14 | 0.12% | 0.00122 | 2.4% | KC level | KC trend | |
| 22 | 1B2 | 1. Energy | B. Fugitive Emissions from F2. Oil and Natural Gas | | CH4 | 263.72 | 169.45 | 0.10% | 0.00052 | 1.0% | KC level | KC trend | |
| 23 | 2A1 | 2. Industrial Pro | A. Mineral Products; Cement Production-CO2 | | CO2 | 2524.68 | 1787.11 | 0.10% | 0.00038 | 0.7% | KC level | - | |
| 24 | 2A3 | 2. Industrial Pro | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | CO2 | 150.39 | 98.48 | 0.10% | 0.00049 | 1.0% | - | KC trend | |
| 25 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 0.08% | 0.00040 | 0.8% | - | KC trend |
| 26 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 0.07% | 0.00026 | 0.5% | - | - |
| 27 | 3 | 3. Solvent and Other Product Use | | | N2O | 110.14 | 44.62 | 0.07% | 0.00100 | 2.0% | - | KC trend | |
| 28 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.07% | 0.00111 | 2.2% | - | KC trend |
| 29 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CO2 | 289.73 | 498.74 | 0.05% | 0.00022 | 0.4% | - | - |
| 30 | 2B | 2. Industrial Pro | B. Chemical Industry | | N2O | 68.13 | 53.57 | 0.04% | 0.00010 | 0.2% | - | - | |
| 31 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Biomass | CH4 | 97.87 | 33.78 | 0.04% | 0.00078 | 1.5% | - | KC trend |
| 32 | 6D | 6. Waste | D. Other | | N2O | 5.82 | 25.55 | 0.04% | 0.00032 | 0.6% | - | - | |
| 33 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | N2O | 20.85 | 20.96 | 0.03% | 0.00001 | 0.0% | - | - |
| 34 | 4D4 | 4. Agriculture | D. Agricultural Soils; Use of sewage sludge as fertilizers | | N2O | 28.30 | 20.85 | 0.03% | 0.00011 | 0.2% | - | - | |
| 35 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | N2O | 27.72 | 19.75 | 0.03% | 0.00012 | 0.2% | - | - |
| 36 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Liquid Fuels | N2O | 25.94 | 18.69 | 0.03% | 0.00010 | 0.2% | - | - |
| 37 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Diesel | N2O | 5.74 | 66.06 | 0.03% | 0.00027 | 0.5% | - | - |
| 38 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Gasoline | N2O | 142.38 | 27.52 | 0.03% | 0.00111 | 2.2% | - | KC trend |
| 39 | 1B2 | 1. Energy | B. Fugitive Emissions from F2. Oil and Natural Gas | | CO2 | 84.62 | 39.29 | 0.02% | 0.00026 | 0.5% | - | - | |
| 40 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 0.02% | 0.00004 | 0.1% | - | - |
| 41 | 2B | 2. Industrial Pro | B. Chemical Industry | | CO2 | 111.22 | 110.47 | 0.02% | 0.00000 | 0.0% | - | - | |
| 42 | 2F9 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Other | | PFC | 0.00 | 13.10 | 0.02% | 0.00021 | 0.4% | - | - | |
| 43 | 6C | 6. Waste | C. Waste Incineration | | N2O | 19.06 | 25.86 | 0.02% | 0.00006 | 0.1% | - | - | |
| 44 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Liquid Fuels | N2O | 13.92 | 11.76 | 0.02% | 0.00003 | 0.1% | - | - |
| 45 | 2F7 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Semiconductor Manufactur | | SF6 | 0.00 | 19.55 | 0.02% | 0.00016 | 0.3% | - | - | |
| 46 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agricul | Liquid Fuels | CO2 | 547.34 | 540.01 | 0.01% | 0.00000 | 0.0% | - | - |
| 47 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Biomass | N2O | 10.79 | 9.07 | 0.01% | 0.00002 | 0.0% | - | - |
| 48 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Gasoline | CH4 | 97.47 | 18.82 | 0.01% | 0.00056 | 1.1% | - | KC trend |
| 49 | 2F2 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Hard Foam | | HFC | 0.00 | 13.27 | 0.01% | 0.00013 | 0.3% | - | - | |
| 50 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Liquid Fuels | N2O | 11.70 | 7.80 | 0.01% | 0.00006 | 0.1% | - | - |
| 51 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Biomass | N2O | 1.68 | 6.85 | 0.01% | 0.00008 | 0.2% | - | - |
| 52 | 2F4 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and | | HFC | 0.00 | 13.22 | 0.01% | 0.00011 | 0.2% | - | - | |
| 53 | 7 | 7. Other | | | CO2 | 10.96 | 12.92 | 0.01% | 0.00002 | 0.0% | - | - | |
| 54 | 6C | 6. Waste | C. Waste Incineration | | CO2 | 54.10 | 12.34 | 0.01% | 0.00032 | 0.6% | - | - | |
| 55 | 2C | 2. Industrial Pro | C. Metal Production; Magnesium Foundries | | SF6 | 0.00 | 31.72 | 0.01% | 0.00010 | 0.2% | - | - | |
| 56 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agricul | Liquid Fuels | N2O | 4.96 | 5.54 | 0.01% | 0.00001 | 0.0% | - | - |
| 57 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Natural Gas | CO2 | 0.00 | 83.59 | 0.01% | 0.00008 | 0.2% | - | - |
| 58 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Other Fuels | N2O | 2.28 | 5.24 | 0.01% | 0.00005 | 0.1% | - | - |
| 59 | 2C1 | 2. Industrial Pro | C. Metal Production; Steel Production | | CO2 | 9.20 | 9.89 | 0.01% | 0.00001 | 0.0% | - | - | |
| 60 | 6C | 6. Waste | C. Waste Incineration | | CH4 | 11.58 | 6.14 | 0.01% | 0.00006 | 0.1% | - | - | |
| 61 | 2F8 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Electrical Eq. | | SF6 | 64.04 | 36.81 | 0.01% | 0.00005 | 0.1% | - | - | |
| 62 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CO2 | 252.55 | 136.65 | 0.01% | 0.00005 | 0.1% | - | - |
| 63 | 6B | 6. Waste | B. Wastewater Handling | | CH4 | 4.65 | 9.28 | 0.01% | 0.00003 | 0.1% | - | - | |
| 64 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Natural Gas | N2O | 0.00 | 3.47 | 0.01% | 0.00006 | 0.1% | - | - |
| 65 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | | CO2 | 111.93 | 121.14 | 0.01% | 0.00001 | 0.0% | - | - |
| 66 | 2F7 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Semiconductor Manufactur | | PFC | 0.00 | 6.54 | 0.01% | 0.00005 | 0.1% | - | - | |
| 67 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Solid Fuels | CO2 | 54.59 | 33.60 | 0.01% | 0.00003 | 0.1% | - | - |
| 68 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Biomass | N2O | 1.45 | 3.23 | 0.01% | 0.00003 | 0.1% | - | - |
| 69 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | N2O | 2.15 | 2.86 | 0.00% | 0.00001 | 0.0% | - | - |
| 70 | 1A3ei | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CO2 | 31.42 | 45.44 | 0.00% | 0.00001 | 0.0% | - | - |
| 71 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | N2O | 2.49 | 1.35 | 0.00% | 0.00003 | 0.1% | - | - |
| 72 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Solid Fuels | N2O | 6.44 | 2.41 | 0.00% | 0.00006 | 0.1% | - | - |
| 73 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Gaseous Fuels | CH4 | 3.24 | 6.10 | 0.00% | 0.00002 | 0.0% | - | - |
| 74 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | N2O | 2.01 | 1.13 | 0.00% | 0.00002 | 0.0% | - | - |
| 75 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO2 | 203.58 | 114.80 | 0.00% | 0.00002 | 0.0% | - | - |
| 76 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Gaseous Fuels | CH4 | 2.66 | 4.74 | 0.00% | 0.00001 | 0.0% | - | - |
| 77 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Biomass | CH4 | 9.74 | 4.13 | 0.00% | 0.00003 | 0.1% | - | - |
| 78 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | N2O | 0.66 | 0.82 | 0.00% | 0.00001 | 0.0% | - | - |
| 79 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Gaseous Fuels | N2O | 0.79 | 1.46 | 0.00% | 0.00001 | 0.0% | - | - |
| 80 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Gaseous Fuels | CH4 | 2.41 | 3.83 | 0.00% | 0.00001 | 0.0% | - | - |

Table A – 4 continued. Key category analysis Tier 2 2012 (without LULUCF) regarding level and trend.

| Tier 2 Key category analysis 2012 without LULUCF categories | | | | | | | | | | | | | |
|---|--|-------------------|--|-----------------------------------|----------------|-------------------------------------|--------------------------------|---------------------------------|---------------------------------|---------------------|----------------------|----------------------|---|
| | A | | | | B | C | D | E-L | E-T | F-T | M | N | |
| No. | IPCC Source Categories and fuels if applicable (without LULUCF categories) | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | Year 2012 Estimate [Gg CO2 eq] | Level Assessm. with Uncertainty | Trend Assessm. with Uncertainty | % Contrib. in Trend | Result level assessm | Result trend assessm | |
| 81 | 2A2 | 2. Industrial Pro | A. Mineral Products; Lime Production-CO2 | | CO2 | 53.35 | 54.26 | 0.00% | 0.00000 | 0.0% | - | - | |
| 82 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Biomass | N2O | 0.00 | 0.70 | 0.00% | 0.00002 | 0.0% | - | - |
| 83 | 7 | 7. Other | | | N2O | 0.62 | 0.62 | 0.00% | 0.00000 | 0.0% | - | - | |
| 84 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Gaseous Fuels | CO2 | 41.45 | 18.48 | 0.00% | 0.00002 | 0.0% | - | - |
| 85 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Gaseous Fuels | N2O | 0.59 | 1.15 | 0.00% | 0.00001 | 0.0% | - | - |
| 86 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | CO2 | 28.69 | 39.69 | 0.00% | 0.00001 | 0.0% | - | - | |
| 87 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | N2O | 0.60 | 0.55 | 0.00% | 0.00000 | 0.0% | - | - |
| 88 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | N2O | 0.38 | 0.52 | 0.00% | 0.00000 | 0.0% | - | - |
| 89 | 2F1 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Refrigeration | | PFC | 0.04 | 6.14 | 0.00% | 0.00001 | 0.0% | - | - | |
| 90 | 2B | 2. Industrial Pro | B. Chemical Industry | | CH4 | 1.54 | 2.39 | 0.00% | 0.00001 | 0.0% | - | - | |
| 91 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Solid Fuels | CH4 | 3.71 | 2.28 | 0.00% | 0.00001 | 0.0% | - | - |
| 92 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Liquid Fuels | CH4 | 6.00 | 2.26 | 0.00% | 0.00002 | 0.0% | - | - |
| 93 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Gaseous Fuels | N2O | 0.55 | 0.82 | 0.00% | 0.00000 | 0.0% | - | - |
| 94 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Biomass | CH4 | 2.46 | 1.56 | 0.00% | 0.00000 | 0.0% | - | - |
| 95 | 7 | 7. Other | | | CH4 | 0.55 | 0.57 | 0.00% | 0.00000 | 0.0% | - | - | |
| 96 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Liquid Fuels | CH4 | 1.62 | 1.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 97 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Comm | Liquid Fuels | CH4 | 3.06 | 1.40 | 0.00% | 0.00001 | 0.0% | - | - |
| 98 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CH4 | 0.65 | 1.12 | 0.00% | 0.00000 | 0.0% | - | - |
| 99 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Liquid Fuels | CH4 | 2.32 | 1.04 | 0.00% | 0.00001 | 0.0% | - | - |
| 100 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Biomass | N2O | 0.21 | 0.35 | 0.00% | 0.00000 | 0.0% | - | - |
| 101 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | N2O | 0.16 | 0.28 | 0.00% | 0.00000 | 0.0% | - | - |
| 102 | 1B2 | 1. Energy | B. Fugitive Emissions from F | 2. Oil and Natural Gas | N2O | 0.62 | 0.68 | 0.00% | 0.00000 | 0.0% | - | - | |
| 103 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CH4 | 0.49 | 0.67 | 0.00% | 0.00000 | 0.0% | - | - |
| 104 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | CH4 | 0.58 | 0.54 | 0.00% | 0.00000 | 0.0% | - | - |
| 105 | 2A7 | 2. Industrial Pro | A. Mineral Products; Other non-specified-CO2 | | CO2 | 15.30 | 7.68 | 0.00% | 0.00000 | 0.0% | - | - | |
| 106 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CH4 | 0.24 | 0.25 | 0.00% | 0.00000 | 0.0% | - | - |
| 107 | 2F5 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Solvents | | PFC | 0.00 | 7.29 | 0.00% | 0.00000 | 0.0% | - | - | |
| 108 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Reside | Solid Fuels | N2O | 0.29 | 0.18 | 0.00% | 0.00000 | 0.0% | - | - |
| 109 | 2C5 | 2. Industrial Pro | C. Metal Production; Non-ferrous metals-CO2 | | CO2 | 1.65 | 1.36 | 0.00% | 0.00000 | 0.0% | - | - | |
| 110 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | CH4 | 0.33 | 0.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 111 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Diesel | CH4 | 1.38 | 0.55 | 0.00% | 0.00000 | 0.0% | - | - |
| 112 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Other Fuels | CH4 | 0.54 | 0.37 | 0.00% | 0.00000 | 0.0% | - | - |
| 113 | 2F5 | 2. Industrial Pro | F. Consumption of Halocarbons and SF6; Solvents | | HFC | 0.00 | 4.59 | 0.00% | 0.00000 | 0.0% | - | - | |
| 114 | 2G | 2. Industrial Pro | G. Other | | CO2 | 1.04 | 0.91 | 0.00% | 0.00000 | 0.0% | - | - | |
| 115 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Biomass | CH4 | 0.80 | 0.15 | 0.00% | 0.00000 | 0.0% | - | - |
| 116 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industr | Solid Fuels | CH4 | 0.40 | 0.14 | 0.00% | 0.00000 | 0.0% | - | - |
| 117 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CH4 | 0.16 | 0.12 | 0.00% | 0.00000 | 0.0% | - | - |
| 118 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Natural Gas | CH4 | 0.00 | 0.10 | 0.00% | 0.00000 | 0.0% | - | - |
| 119 | 1A3ei | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | N2O | 0.02 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - | |
| 120 | 1A3ei | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | CH4 | 0.06 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - | |
| 121 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Gaseous Fuels | CH4 | 0.09 | 0.04 | 0.00% | 0.00000 | 0.0% | - | - |
| 122 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Trans | Biomass | CH4 | 0.00 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 123 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agric | Gaseous Fuels | N2O | 0.02 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 124 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | CH4 | 0.01 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 125 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | CH4 | 0.01 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 126 | 6D | 6. Waste | D. Other | | CO2 | 0.00 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 127 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CO2 | 9.24 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 128 | 2C3 | 2. Industrial Pro | C. Metal Production; Aluminium Production-CO2 | | CO2 | 139.26 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 129 | 2C3 | 2. Industrial Pro | C. Metal Production; Aluminium Production-PFC | | PFC | 100.17 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 130 | 2C | 2. Industrial Pro | C. Metal Production; Aluminium Foundries | | SF6 | 0.00 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 131 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CO2 | 44.84 | 0.00 | 0.00% | - | 0.0% | - | - |
| 132 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CH4 | 0.10 | 0.00 | 0.00% | - | 0.0% | - | - |
| 133 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | N2O | 0.24 | 0.00 | 0.00% | - | 0.0% | - | - |
| 134 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CO2 | 9.24 | 0.00 | 0.00% | - | 0.0% | - | - | |
| 135 | 6D | 6. Waste | D. Other | | CO2 | 0.00 | 0.00 | 0.00% | - | 0.0% | - | - | |

A1.5 KCA Tier 2 2012 including LULUCF categories

Results of Key Category Analysis Tier 2 – Level and Trend

Table A - 5 Key category analysis Tier 2 2012 (with LULUCF) regarding level and trend.

| Tier 2 Key category analysis 2012 with LULUCF categories | | | | | | | | | | | | | |
|--|--|----------------------------------|---|--|-----------------|---|--|--|--|----------------------------|---------------------------|---------------------------|----------|
| No. | A IPCC Source Categories and fuels if applicable (with LULUCF categories) | | | | B Direct GHG | C Base Year 1990 Estimate [Gg CO2 eq] | D Year 2012 Estimate [Gg CO2 eq] | E-L Level Assessm. with Uncertainty | E-T Trend Assessm. with Uncertainty | F-T % Contrib. in Trend | M Result level assessm | N Result trend assessm | |
| 1 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | CO2 | 107.09 | 134.74 | 5.05% | 0.01165 | 15.8% | KC level | KC trend | |
| 2 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | CO2 | -2416.89 | -2134.56 | 2.41% | 0.00257 | 3.5% | KC level | KC trend | |
| 3 | 4D3 | 4. Agriculture | D. Agricultural Soils; Indirect Emissions | | N2O | 822.48 | 674.93 | 2.01% | 0.00389 | 5.3% | KC level | KC trend | |
| 4 | 4D1 | 4. Agriculture | D. Agricultural Soils; Direct Soil Emissions | | N2O | 1351.48 | 1143.10 | 1.71% | 0.00268 | 3.6% | KC level | KC trend | |
| 5 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | CO2 | 1519.73 | 2714.50 | 1.54% | 0.00719 | 9.7% | KC level | KC trend |
| 6 | 5B1 | 5. LULUCF | B. Cropland | 1. Cropland remaining Cropland | CO2 | 345.17 | 707.27 | 1.40% | 0.00751 | 10.2% | KC level | KC trend | |
| 7 | 4A | 4. Agriculture | A. Enteric Fermentation | | CH4 | 2635.45 | 2496.98 | 0.82% | 0.00025 | 0.3% | KC level | - | |
| 8 | 4B | 4. Agriculture | B. Manure Management | | CH4 | 671.61 | 646.11 | 0.63% | 0.00009 | 0.1% | KC level | - | |
| 9 | 5A2 | 5. LULUCF | A. Forest Land | 2. Land converted to Forest Land | CO2 | -621.57 | -518.61 | 0.59% | 0.00101 | 1.4% | KC level | KC trend | |
| 10 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CO2 | 11335.27 | 9016.58 | 0.42% | 0.00097 | 1.3% | KC level | KC trend |
| 11 | 4B | 4. Agriculture | B. Manure Management | | N2O | 454.68 | 335.81 | 0.38% | 0.00126 | 1.7% | KC level | KC trend | |
| 12 | 4D2 | 4. Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | N2O | 128.10 | 220.79 | 0.34% | 0.00150 | 2.0% | KC level | KC trend | |
| 13 | 5E2 | 5. LULUCF | E. Settlements | 2. Land converted to Settlements | CO2 | 382.71 | 302.93 | 0.27% | 0.00065 | 0.9% | KC level | KC trend | |
| 14 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CO2 | 2587.68 | 6767.05 | 0.27% | 0.00175 | 2.4% | KC level | KC trend |
| 15 | 2F1 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Refrig. & AC Eq. | | HFC | 0.02 | 1137.81 | 0.25% | 0.00252 | 3.4% | KC level | KC trend | |
| 16 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuel | CO2 | 1424.38 | 2649.60 | 0.24% | 0.00117 | 1.6% | KC level | KC trend |
| 17 | 6B | 6. Waste | B. Wastewater Handling | | N2O | 184.72 | 240.28 | 0.22% | 0.00055 | 0.8% | KC level | KC trend | |
| 18 | 6D | 6. Waste | D. Other | | CH4 | 29.94 | 113.76 | 0.21% | 0.00157 | 2.1% | KC level | KC trend | |
| 19 | 5C2 | 5. LULUCF | C. Grassland | 2. Land converted to Grassland | CO2 | 59.85 | 169.07 | 0.21% | 0.00138 | 1.9% | KC level | KC trend | |
| 20 | 2F9 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Other | | SF6 | 79.58 | 135.91 | 0.20% | 0.00086 | 1.2% | KC level | KC trend | |
| 21 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Gaseous Fuel | CO2 | 1074.09 | 2096.41 | 0.19% | 0.00097 | 1.3% | KC level | KC trend |
| 22 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CO2 | 10248.79 | 7374.50 | 0.18% | 0.00067 | 0.9% | KC level | KC trend |
| 23 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CH4 | 688.16 | 158.26 | 0.17% | 0.00551 | 7.5% | KC level | KC trend | |
| 24 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Other Fuels | CO2 | 134.15 | 288.60 | 0.16% | 0.00092 | 1.2% | - | KC trend |
| 25 | 3 | 3. Solvent and Other Product Use | | | CO2 | 360.04 | 155.28 | 0.14% | 0.00180 | 2.4% | - | KC trend | |
| 26 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institution | Gaseous Fuel | CO2 | 987.24 | 1482.76 | 0.13% | 0.00048 | 0.7% | - | - |
| 27 | 2F9 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Other | | HFC | 0.00 | 76.14 | 0.11% | 0.00112 | 1.5% | - | KC trend | |
| 28 | 5F2 | 5. LULUCF | F. Other Land | 2. Land converted to Other Land | CO2 | 91.98 | 112.28 | 0.10% | 0.00021 | 0.3% | - | - | |
| 29 | 1B2 | 1. Energy | B. Fugitive Emissions fr | 2. Oil and Natural Gas | CH4 | 263.72 | 169.45 | 0.09% | 0.00049 | 0.7% | - | KC trend | |
| 30 | 2A1 | 2. Industrial Pr | A. Mineral Products; Cement Production-CO2 | | CO2 | 2524.68 | 1787.11 | 0.09% | 0.00035 | 0.5% | - | - | |
| 31 | 2A3 | 2. Industrial Pr | A. Mineral Products; Limestone and Dolomite Use, Emissions, CO2 | | CO2 | 150.39 | 98.48 | 0.09% | 0.00045 | 0.6% | - | - | |
| 32 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institution | Liquid Fuels | CO2 | 4606.43 | 3038.51 | 0.08% | 0.00037 | 0.5% | - | - |
| 33 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Liquid Fuels | CO2 | 3692.22 | 2640.18 | 0.07% | 0.00025 | 0.3% | - | - |
| 34 | 3 | 3. Solvent and Other Product Use | | | N2O | 110.14 | 44.62 | 0.06% | 0.00093 | 1.3% | - | KC trend | |
| 35 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Solid Fuels | CO2 | 1204.47 | 454.87 | 0.06% | 0.00103 | 1.4% | - | KC trend |
| 36 | 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | CO2 | 43.33 | 22.26 | 0.06% | 0.00053 | 0.7% | - | KC trend | |
| 37 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuel | CO2 | 289.73 | 498.74 | 0.04% | 0.00020 | 0.3% | - | - |
| 38 | 2B | 2. Industrial Pr | B. Chemical Industry | | N2O | 68.13 | 53.57 | 0.04% | 0.00010 | 0.1% | - | - | |
| 39 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | CH4 | 97.87 | 33.78 | 0.04% | 0.00072 | 1.0% | - | KC trend |
| 40 | 6D | 6. Waste | D. Other | | N2O | 5.82 | 25.55 | 0.04% | 0.00029 | 0.4% | - | - | |
| 41 | 5E1 | 5. LULUCF | E. Settlements | 1. Settlements remaining Settlements | CO2 | 3.60 | 33.69 | 0.03% | 0.00028 | 0.4% | - | - | |
| 42 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Other Fuels | N2O | 20.85 | 20.96 | 0.03% | 0.00001 | 0.0% | - | - |
| 43 | 4D4 | 4. Agriculture | D. Agricultural Soils; Use of sewage sludge as fertilizers | | N2O | 28.30 | 20.85 | 0.03% | 0.00010 | 0.1% | - | - | |
| 44 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | N2O | 27.72 | 19.75 | 0.03% | 0.00011 | 0.1% | - | - |
| 45 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | N2O | 25.94 | 18.69 | 0.03% | 0.00010 | 0.1% | - | - |
| 46 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | N2O | 5.74 | 66.06 | 0.03% | 0.00025 | 0.3% | - | - |
| 47 | 5D2 | 5. LULUCF | D. Wetlands | 2. Land converted to Wetlands | CO2 | 20.06 | 28.95 | 0.03% | 0.00009 | 0.1% | - | - | |
| 48 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | N2O | 142.38 | 27.52 | 0.02% | 0.00103 | 1.4% | - | KC trend |
| 49 | 1B2 | 1. Energy | B. Fugitive Emissions fr | 2. Oil and Natural Gas | CO2 | 84.62 | 39.29 | 0.02% | 0.00024 | 0.3% | - | - | |
| 50 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CO2 | 693.69 | 805.16 | 0.02% | 0.00003 | 0.0% | - | - |
| 51 | 2B | 2. Industrial Pr | B. Chemical Industry | | CO2 | 111.22 | 110.47 | 0.02% | 0.00000 | 0.0% | - | - | |
| 52 | 2F9 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Other | | PFC | 0.00 | 13.10 | 0.02% | 0.00019 | 0.3% | - | - | |
| 53 | 6C | 6. Waste | C. Waste Incineration | | N2O | 19.06 | 25.86 | 0.02% | 0.00005 | 0.1% | - | - | |
| 54 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Liquid Fuels | N2O | 13.92 | 11.76 | 0.02% | 0.00003 | 0.0% | - | - |
| 55 | 2F7 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | SF6 | 0.00 | 19.55 | 0.01% | 0.00014 | 0.2% | - | - | |
| 56 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CO2 | 547.34 | 540.01 | 0.01% | 0.00000 | 0.0% | - | - |
| 57 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Biomass | N2O | 10.79 | 9.07 | 0.01% | 0.00002 | 0.0% | - | - |
| 58 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Gasoline | CH4 | 97.47 | 18.82 | 0.01% | 0.00052 | 0.7% | - | KC trend |
| 59 | 2F2 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Hard Foam | | HFC | 0.00 | 13.27 | 0.01% | 0.00012 | 0.2% | - | - | |
| 60 | 5D2 | 5. LULUCF | D. Land converted to W(5)(I) Non-CO2 emissions from drainage of soils and wet | | CH4 | 9.03 | 9.03 | 0.01% | 0.00000 | 0.0% | - | - | |
| 61 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institution | Liquid Fuels | N2O | 11.70 | 7.80 | 0.01% | 0.00005 | 0.1% | - | - |
| 62 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Biomass | N2O | 1.68 | 6.85 | 0.01% | 0.00008 | 0.1% | - | - |
| 63 | 2F4 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Metered Dose Inhalers and Other | | HFC | 0.00 | 13.22 | 0.01% | 0.00010 | 0.1% | - | - | |
| 64 | 7 | 7. Other | | | CO2 | 10.96 | 12.92 | 0.01% | 0.00002 | 0.0% | - | - | |
| 65 | 6C | 6. Waste | C. Waste Incineration | | CO2 | 54.10 | 12.34 | 0.01% | 0.00030 | 0.4% | - | - | |
| 66 | 2C | 2. Industrial Pr | C. Metal Production; Magnesium Foundries | | SF6 | 0.00 | 31.72 | 0.01% | 0.00009 | 0.1% | - | - | |
| 67 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | N2O | 4.96 | 5.54 | 0.01% | 0.00001 | 0.0% | - | - |
| 68 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CO2 | 0.00 | 83.59 | 0.01% | 0.00008 | 0.1% | - | - |
| 69 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Construc | Other Fuels | N2O | 2.28 | 5.24 | 0.01% | 0.00004 | 0.1% | - | - |
| 70 | 2C1 | 2. Industrial Pr | C. Metal Production; Steel Production | | CO2 | 9.20 | 9.89 | 0.01% | 0.00001 | 0.0% | - | - | |
| 71 | 6C | 6. Waste | C. Waste Incineration | | CH4 | 11.58 | 6.14 | 0.01% | 0.00006 | 0.1% | - | - | |
| 72 | 2F8 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Electrical Eq. | | SF6 | 64.04 | 36.81 | 0.01% | 0.00005 | 0.1% | - | - | |
| 73 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | CO2 | 252.55 | 136.65 | 0.01% | 0.00005 | 0.1% | - | - | |
| 74 | 5B2 | 5. LULUCF | B. Cropland | 2. Land converted to Cropland | N2O | 5.58 | 3.58 | 0.01% | 0.00003 | 0.0% | - | - | |
| 75 | 6B | 6. Waste | B. Wastewater Handling | | CH4 | 4.65 | 9.28 | 0.01% | 0.00003 | 0.0% | - | - | |
| 76 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | N2O | 0.00 | 3.47 | 0.00% | 0.00005 | 0.1% | - | - |
| 77 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | CO2 | 111.93 | 121.14 | 0.00% | 0.00000 | 0.0% | - | - | |
| 78 | 2F7 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Semiconductor Manufacture | | PFC | 0.00 | 6.54 | 0.00% | 0.00005 | 0.1% | - | - | |
| 79 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CO2 | 54.59 | 33.60 | 0.00% | 0.00003 | 0.0% | - | - |
| 80 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institution | Biomass | N2O | 1.45 | 3.23 | 0.00% | 0.00003 | 0.0% | - | - |

Table A – 5 continued. Key category analysis Tier 2 2012 (with LULUCF) regarding level and trend.

| Tier 2 Key category analysis 2012 with LULUCF categories | | | | | | | | | | | | | | |
|--|---|------------------|--|---|----------------|-----|---------------|--|--------|--|--|-------------------------------|------------------------------|------------------------------|
| No. | A | | | | | B | C | | D | E-L Assessm. with Uncertainty | E-T Assessm. with Uncertainty | F-T % Contrib. in Trend | M Result level assessm | N Result trend assessm |
| | IPCC Source Categories and fuels if applicable (with LULUCF categories) | | | | | | Direct GHG | Base Year 1990 Estimate [Gg CO2 eq] | | | | | | |
| 81 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | N2O | | 2.15 | 2.86 | 0.00% | 0.00001 | 0.0% | - | - |
| 82 | 1A3el | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CO2 | | 31.42 | 45.44 | 0.00% | 0.00001 | 0.0% | - | - |
| 83 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | N2O | | 2.49 | 1.35 | 0.00% | 0.00003 | 0.0% | - | - |
| 84 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Solid Fuels | N2O | | 6.44 | 2.41 | 0.00% | 0.00006 | 0.1% | - | - |
| 85 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | CH4 | | 3.24 | 6.10 | 0.00% | 0.00002 | 0.0% | - | - |
| 86 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | N2O | | 2.01 | 1.13 | 0.00% | 0.00002 | 0.0% | - | - |
| 87 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CO2 | | 203.58 | 114.80 | 0.00% | 0.00002 | 0.0% | - | - |
| 88 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Gaseous Fuels | CH4 | | 2.66 | 4.74 | 0.00% | 0.00001 | 0.0% | - | - |
| 89 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutions | Biomass | CH4 | | 9.74 | 4.13 | 0.00% | 0.00003 | 0.0% | - | - |
| 90 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | N2O | | 0.66 | 0.82 | 0.00% | 0.00000 | 0.0% | - | - |
| 91 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Gaseous Fuels | N2O | | 0.79 | 1.46 | 0.00% | 0.00001 | 0.0% | - | - |
| 92 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutions | Gaseous Fuels | CH4 | | 2.41 | 3.83 | 0.00% | 0.00001 | 0.0% | - | - |
| 93 | 2A2 | 2. Industrial Pr | A. Mineral Products; Lime Production-CO2 | | CO2 | | | 53.35 | 54.26 | 0.00% | 0.00000 | 0.0% | - | - |
| 94 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | N2O | | 0.00 | 0.70 | 0.00% | 0.00002 | 0.0% | - | - |
| 95 | 7 | 7. Other | | | | N2O | | 0.62 | 0.62 | 0.00% | 0.00000 | 0.0% | - | - |
| 96 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CO2 | | 41.45 | 18.48 | 0.00% | 0.00002 | 0.0% | - | - |
| 97 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Gaseous Fuels | N2O | | 0.59 | 1.15 | 0.00% | 0.00001 | 0.0% | - | - |
| 98 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | | CO2 | | 28.69 | 39.69 | 0.00% | 0.00000 | 0.0% | - | - |
| 99 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | N2O | | 0.60 | 0.55 | 0.00% | 0.00000 | 0.0% | - | - |
| 100 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | N2O | | 0.38 | 0.52 | 0.00% | 0.00000 | 0.0% | - | - |
| 101 | 2F1 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Refrigeration | | PFC | | | 0.04 | 6.14 | 0.00% | 0.00001 | 0.0% | - | - |
| 102 | 2B | 1. Industrial Pr | B. Chemical Industry | | CH4 | | | 1.54 | 2.39 | 0.00% | 0.00000 | 0.0% | - | - |
| 103 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | CH4 | | 3.71 | 2.28 | 0.00% | 0.00001 | 0.0% | - | - |
| 104 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Liquid Fuels | CH4 | | 6.00 | 2.26 | 0.00% | 0.00002 | 0.0% | - | - |
| 105 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutions | Gaseous Fuels | N2O | | 0.55 | 0.82 | 0.00% | 0.00000 | 0.0% | - | - |
| 106 | 5D1 | 5. LULUCF | D. Wetlands | 1. Wetlands remaining Wetlands | | CO2 | | 2.87 | 0.56 | 0.00% | 0.00004 | 0.1% | - | - |
| 107 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Biomass | CH4 | | 2.46 | 1.56 | 0.00% | 0.00000 | 0.0% | - | - |
| 108 | 7 | 7. Other | | | | CH4 | | 0.55 | 0.57 | 0.00% | 0.00000 | 0.0% | - | - |
| 109 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Liquid Fuels | CH4 | | 1.62 | 1.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 110 | 1A4a | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Commercial/Institutions | Liquid Fuels | CH4 | | 3.06 | 1.40 | 0.00% | 0.00001 | 0.0% | - | - |
| 111 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | Biomass Burn | CO2 | | 25.36 | 0.51 | 0.00% | 0.00032 | 0.4% | - | - |
| 112 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | CH4 | | 0.65 | 1.12 | 0.00% | 0.00000 | 0.0% | - | - |
| 113 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Liquid Fuels | CH4 | | 2.32 | 1.04 | 0.00% | 0.00001 | 0.0% | - | - |
| 114 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | CH4 | | 20.90 | 0.42 | 0.00% | 0.00026 | 0.4% | - | - |
| 115 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | N2O | | 0.21 | 0.35 | 0.00% | 0.00000 | 0.0% | - | - |
| 116 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Gaseous Fuels | N2O | | 0.16 | 0.28 | 0.00% | 0.00000 | 0.0% | - | - |
| 117 | 1B2 | 1. Energy | B. Fugitive Emissions from | 2. Oil and Natural Gas | | N2O | | 0.62 | 0.68 | 0.00% | 0.00000 | 0.0% | - | - |
| 118 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Liquid Fuels | CH4 | | 0.49 | 0.67 | 0.00% | 0.00000 | 0.0% | - | - |
| 119 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gasoline | CH4 | | 0.58 | 0.54 | 0.00% | 0.00000 | 0.0% | - | - |
| 120 | 2A7 | 2. Industrial Pr | A. Mineral Products; Other non-specified-CO2 | | CO2 | | | 15.30 | 7.68 | 0.00% | 0.00000 | 0.0% | - | - |
| 121 | 1A3a | 1. Energy | A. Fuel Combustion | 3. Transport; Civil Aviation | | CH4 | | 0.24 | 0.25 | 0.00% | 0.00000 | 0.0% | - | - |
| 122 | 2F5 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Solvents | | PFC | | | 0.00 | 7.29 | 0.00% | 0.00000 | 0.0% | - | - |
| 123 | 1A4b | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Residential | Solid Fuels | N2O | | 0.29 | 0.18 | 0.00% | 0.00000 | 0.0% | - | - |
| 124 | 2C5 | 2. Industrial Pr | C. Metal Production; Non-ferrous metals-CO2 | | CO2 | | | 1.65 | 1.36 | 0.00% | 0.00000 | 0.0% | - | - |
| 125 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Biomass | CH4 | | 0.33 | 0.43 | 0.00% | 0.00000 | 0.0% | - | - |
| 126 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Diesel | CH4 | | 1.38 | 0.55 | 0.00% | 0.00000 | 0.0% | - | - |
| 127 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Other Fuels | CH4 | | 0.54 | 0.37 | 0.00% | 0.00000 | 0.0% | - | - |
| 128 | 2F5 | 2. Industrial Pr | F. Consumption of Halocarbons and SF6; Solvents | | HFC | | | 0.00 | 4.59 | 0.00% | 0.00000 | 0.0% | - | - |
| 129 | 2G | 2. Industrial Pr | G. Other | | CO2 | | | 1.04 | 0.91 | 0.00% | 0.00000 | 0.0% | - | - |
| 130 | 5A1 | 5. LULUCF | A. Forest Land | 1. Forest Land remaining Forest Land | | N2O | | 4.77 | 0.09 | 0.00% | 0.00006 | 0.1% | - | - |
| 131 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Biomass | CH4 | | 0.80 | 0.15 | 0.00% | 0.00000 | 0.0% | - | - |
| 132 | 1A2 | 1. Energy | A. Fuel Combustion | 2. Manufacturing Industries and Constr | Solid Fuels | CH4 | | 0.40 | 0.14 | 0.00% | 0.00000 | 0.0% | - | - |
| 133 | 1A5 | 1. Energy | A. Fuel Combustion | 5. Other | Liquid Fuels | CH4 | | 0.16 | 0.12 | 0.00% | 0.00000 | 0.0% | - | - |
| 134 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Natural Gas | CH4 | | 0.00 | 0.10 | 0.00% | 0.00000 | 0.0% | - | - |
| 135 | 1A3el | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | N2O | | 0.02 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - |
| 136 | 1A3el | 1. Energy | A. Fuel Combustion | 3. Transport; Other non-specified | | CH4 | | 0.06 | 0.03 | 0.00% | 0.00000 | 0.0% | - | - |
| 137 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | CH4 | | 0.09 | 0.04 | 0.00% | 0.00000 | 0.0% | - | - |
| 138 | 1A3b | 1. Energy | A. Fuel Combustion | 3. Transport; Road Transportation | Biomass | CH4 | | 0.00 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 139 | 1A4c | 1. Energy | A. Fuel Combustion | 4. Other Sectors; Agriculture/Forestry | Gaseous Fuels | N2O | | 0.02 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 140 | 1A3d | 1. Energy | A. Fuel Combustion | 3. Transport; Navigation | Gas/Diesel Oil | CH4 | | 0.01 | 0.02 | 0.00% | 0.00000 | 0.0% | - | - |
| 141 | 1A3c | 1. Energy | A. Fuel Combustion | 3. Transport; Railways | Liquid Fuels | CH4 | | 0.01 | 0.01 | 0.00% | 0.00000 | 0.0% | - | - |
| 142 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | CH4 | | 0.41 | 0.00 | 0.00% | 0.00001 | 0.0% | - | - |
| 143 | 5C1 | 5. LULUCF | C. Grassland | 1. Grassland remaining Grassland | | N2O | | 0.19 | 0.00 | 0.00% | 0.00000 | 0.0% | - | - |
| 144 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CO2 | | 44.84 | 0.00 | 0.00% | - | 0.0% | - | - |
| 145 | 2C3 | 2. Industrial Pr | C. Metal Production; Aluminium Production-CO2 | | CO2 | | | 139.26 | 0.00 | 0.00% | - | 0.0% | - | - |
| 146 | 6A | 6. Waste | A. Solid Waste Disposal on Land | | CO2 | | | 9.24 | 0.00 | 0.00% | - | 0.0% | - | - |
| 147 | 6D | 6. Waste | D. Other | | CO2 | | | 0.00 | 0.00 | 0.00% | - | 0.0% | - | - |
| 148 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | CH4 | | 0.10 | 0.00 | 0.00% | - | 0.0% | - | - |
| 149 | 1A1 | 1. Energy | A. Fuel Combustion | 1. Energy Industries | Solid Fuels | N2O | | 0.24 | 0.00 | 0.00% | - | 0.0% | - | - |
| 150 | 2C3 | 2. Industrial Pr | C. Metal Production; Aluminium Production-PFC | | PFC | | | 100.17 | 0.00 | 0.00% | - | 0.0% | - | - |
| 151 | 2C | 2. Industrial Pr | C. Metal Production; Aluminium Foundries | | SF6 | | | 0.00 | 0.00 | 0.00% | - | 0.0% | - | - |

Annex 2: Sulphur Dioxide (SO₂)

Table A - 6 Sulphur content and SO₂ emission factors. For explanations see next page.

| year | maximum legal limit of sulphur content | | | | | |
|------|--|-----------------|----------------|--------------------|--------------------|-----------|
| | Diesel oil ppm | Gasoline ppm | Gas oil ppm | Natural gas ppm | Res. fuel oil % | Coal % |
| 1990 | 1400 | 200 | 2000 | 190 | 1.0 | 1.0 |
| 1991 | 1300 | 200 | 2000 | 190 | 1.0 | 1.0 |
| 1992 | 1200 | 200 | 2000 | 190 | 1.0 | 1.0 |
| 1993 | 1000 | 200 | 2000 | 190 | 1.0 | 1.0 |
| 1994 | 500 | 200 | 2000 | 190 | 1.0 | 1.0 |
| 2000 | 350 | 150 | 2000 | 190 | 1.0 | 1.0 |
| 2005 | 50 | 50 | 2000 | 190 | 1.0 | 1.0 |
| 2008 | 50 | 50 | 1000 | 190 | 1.0 | 1.0 |
| 2009 | 10 | 50 | 1000 | 190 | 1.0 | 1.0 |
| 2010 | 10 | 10 | 1000 | 190 | 1.0 | 1.0 |
| 2011 | 10 | 10 | 1000 | 190 | 1.0 | 1.0 |
| 2012 | 10 | 10 | 1000 | 190 | 1.0 | 1.0 |

| year | Effective sulphur content | | | | | |
|------|---------------------------|-----------------|----------------|--------------------|--------------------|-----------|
| | Diesel oil ppm | Gasoline ppm | Gas oil ppm | Natural gas ppm | Res. fuel oil % | Coal % |
| 1990 | 1400 | 200 | 1600 | 11.6 | 0.97 | 0.9 |
| 1991 | 1300 | 200 | 1300 | 11.6 | 0.89 | 0.9 |
| 1992 | 1200 | 200 | 1200 | 11.6 | 0.86 | 0.9 |
| 1993 | 1000 | 200 | 1000 | 11.6 | 0.87 | 0.9 |
| 1994 | 434 | 200 | 1350 | 11.6 | 0.77 | 0.9 |
| 1995 | 341 | 200 | 1170 | 11.6 | 0.78 | 0.9 |
| 1996 | 372 | 200 | 1160 | 11.6 | 0.78 | 0.9 |
| 1997 | 353 | 200 | 1250 | 11.6 | 0.70 | 0.9 |
| 1998 | 402 | 200 | 926 | 11.6 | 0.83 | 0.9 |
| 1999 | 443 | 200 | 650 | 11.6 | 0.62 | 0.9 |
| 2000 | 272 | 142 | 680 | 11.6 | 0.66 | 0.9 |
| 2001 | 250 | 121 | 830 | 11.6 | 0.82 | 0.9 |
| 2002 | 235 | 101 | 798 | 11.6 | 0.78 | 0.9 |
| 2003 | 200 | 81 | 700 | 11.6 | 0.79 | 0.9 |
| 2004 | 10 | 8.0 | 700 | 11.6 | 0.76 | 0.9 |
| 2005 | 10 | 8.0 | 799 | 11.6 | 0.78 | 0.9 |
| 2006 | 10 | 8.0 | 699 | 11.6 | 0.74 | 0.9 |
| 2007 | 10 | 8.0 | 630 | 11.6 | 0.71 | 0.9 |
| 2008 | 10 | 8.0 | 641 | 11.6 | 0.67 | 0.9 |
| 2009 | 10 | 8.0 | 539 | 11.6 | 0.64 | 0.9 |
| 2010 | 10 | 8.0 | 509 | 11.6 | 0.60 | 0.9 |
| 2011 | 10 | 8.0 | 477 | 11.6 | 0.60 | 0.9 |
| 2012 | 10 | 8.0 | 445 | 11.6 | 0.60 | 0.9 |

| year | Effective SO ₂ emission factor | | | | | |
|------|---|----------|---------|-------------|---------------|------|
| | Diesel oil | Gasoline | Gas oil | Natural gas | Res. fuel oil | Coal |
| | kg/TJ | | | | | |
| 1990 | 65.4 | 9.4 | 75.1 | 0.50 | 473 | 350 |
| 1991 | 60.7 | 9.4 | 61.0 | 0.50 | 432 | 350 |
| 1992 | 56.1 | 9.4 | 56.3 | 0.50 | 417 | 350 |
| 1993 | 46.7 | 9.4 | 46.9 | 0.50 | 422 | 350 |
| 1994 | 20.3 | 9.4 | 63.4 | 0.50 | 374 | 350 |
| 1995 | 15.9 | 9.4 | 54.9 | 0.50 | 377 | 350 |
| 1996 | 17.4 | 9.4 | 54.5 | 0.50 | 379 | 350 |
| 1997 | 16.5 | 9.4 | 58.7 | 0.50 | 340 | 350 |
| 1998 | 18.8 | 9.4 | 43.5 | 0.50 | 403 | 350 |
| 1999 | 20.7 | 9.4 | 30.5 | 0.50 | 301 | 350 |
| 2000 | 12.7 | 6.7 | 31.9 | 0.50 | 320 | 350 |
| 2001 | 11.7 | 5.7 | 39.0 | 0.50 | 398 | 350 |
| 2002 | 11.0 | 4.8 | 37.5 | 0.50 | 398 | 350 |
| 2003 | 9.3 | 3.8 | 32.9 | 0.50 | 383 | 350 |
| 2004 | 0.5 | 0.4 | 32.9 | 0.50 | 369 | 350 |
| 2005 | 0.5 | 0.4 | 37.5 | 0.50 | 379 | 350 |
| 2006 | 0.5 | 0.4 | 32.8 | 0.50 | 361 | 350 |
| 2007 | 0.5 | 0.4 | 29.6 | 0.50 | 344 | 350 |
| 2008 | 0.5 | 0.4 | 30.1 | 0.50 | 326 | 350 |
| 2009 | 0.5 | 0.4 | 25.3 | 0.50 | 309 | 350 |
| 2010 | 0.5 | 0.4 | 23.9 | 0.50 | 291 | 350 |
| 2011 | 0.5 | 0.4 | 22.4 | 0.50 | 291 | 350 |
| 2012 | 0.5 | 0.4 | 20.9 | 0.50 | 291 | 350 |

Explanation to Table A - 6

- For liquid and solid fuels the SO₂ emission factors are determined by the sulphur content. The upmost lines in Table A - 6 “maximum legal limit on sulphur content” show the maximum values as defined in the Federal Ordinance on Air Pollution Control OAPC (Swiss Confederation 1985).
- The lines in the middle part of Table A - 6 contain the effective sulphur contents. They are based on measurements: Summary and annual reports of the Swiss Petroleum Association (EV), reports by the Federal Administration of Customs (OZD) since 2000.
- The lines at the bottom part of Table A - 6 give the emission factors in kg/TJ. They are calculated from the sulphur content S, the net calorific value NCV and the quotient of the molar masses of S and SO₂

$$EF_{SO_2} = \frac{M_{SO_2}}{M_S} \cdot \frac{S}{NCV} = 2 \cdot \frac{S}{NCV}$$

- Coal: Note that the legal limit of sulphur content depends on the size of the heat capacity of the combustion system. The value shown in the table above (1%, 350 kg/TJ SO₂) holds for heat capacity below 1 MW; see OAPC Annex 4, §513 (Swiss Confederation 1985). For larger capacities the value is 3% (OAPC Annex 5, §2, Swiss Confederation 1985). For industrial combustion plants, the limit for the exhaust emissions actually sets the corresponding maximum sulphur content to 1.4% (500 kg/TJ).
- Residual fuel oil: OAPC Annex 5, §11, lit.2 sets 2.8% for the legal limit. Simultaneously, OAPC dispenses from emission control measurements if residual fuel oil is used with sulphur content of maximum 1% (see OAPC Annex 3, §421, lit. 2, Swiss Confederation 1985), which holds for most combustion plants.

Annex 3: Other detailed methodological descriptions for individual source or sink categories

A3.1 Sector Energy

A3.1.1 Swiss Energy Flow

The diagrams show a summary of the Swiss energy flow 2012 and 1990 as published by the Swiss Federal Office of Energy (SFOE 2013). Diagram languages are German and French.




Fig. 5 Detailliertes Energieflussdiagramm der Schweiz 2012 (in TJ)
Flux énergétique détaillé de la Suisse en 2012 (en TJ)

Figure A - 1 Energy flow in Switzerland 2012 in TJ (SFOE 2013)




Figure A - 2 Energy flow in Switzerland 1990 in TJ (SFOE 1991)

A3.1.2 Emissions from Fuel Consumption: Disaggregation of Fuel Consumption

See Chapter 3.2.5.

A3.1.3 Emission from Manufacturing Industries and Construction

The emission factors of precursors in the manufacturing industries and construction sector are given below. Emission factors for greenhouse gases are given in 3.2.7.

Table A - 7 Emission factors of precursors from Manufacturing Industries and Construction

| 1A2 Emission factors (mix of bottom-up and top-down approach (modelling)) | NO _x | CO | NMVOC | SO ₂ |
|--|-----------------|-------|-------|-----------------|
| | kg/TJ | kg/TJ | kg/TJ | kg/TJ |
| 1A2a Iron and Steel | | | | |
| Gas oil | 33 | 8 | 2 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | 125 | 13 | 4 | 291 |
| Petroleum coke | NO | NO | NO | NO |
| Bituminous coal | 14 | 2'282 | 8 | 311 |
| Lignite | NO | NO | NO | NO |
| Natural gas | 36 | 5 | 2 | 0.5 |
| 1A2b Non-Ferrous Metals | | | | |
| Gas oil | 33 | 182 | 37 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | NO | NO | NO | NO |
| Petroleum coke | NO | NO | NO | NO |
| Bituminous coal | NO | NO | NO | NO |
| Lignite | NO | NO | NO | NO |
| Natural gas | 19 | 9 | 2 | 0.5 |
| 1A2c Chemicals | | | | |
| Gas oil | 33 | 8 | 2 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | 125 | 13 | 4 | 291 |
| Petroleum coke | NO | NO | NO | NO |
| Bituminous coal | NO | NO | NO | NO |
| Lignite | NO | NO | NO | NO |
| Natural gas | 19 | 9 | 2 | 0.5 |
| 1A2d Pulp, Paper and Print | | | | |
| Gas oil | 33 | 8 | 2 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | 125 | 13 | 4 | 291 |
| Petroleum coke | NO | NO | NO | NO |
| Bituminous coal | NO | NO | NO | NO |
| Lignite | NO | NO | NO | NO |
| Natural gas | 19 | 9 | 2 | 0.5 |
| 1A2e Food Processing, Beverages and Tobacco | | | | |
| Gas oil | 33 | 8 | 2 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | NO | NO | NO | NO |
| Petroleum coke | NO | NO | NO | NO |
| Bituminous coal | NO | NO | NO | NO |
| Lignite | NO | NO | NO | NO |
| Natural gas | 19 | 9 | 2 | 0.5 |
| 1A2fi Other | | | | |
| Gas oil | 33 | 8 | 7 | 22 |
| Liquefied petroleum gas | 33 | 8 | 2 | 22 |
| Residual fuel oil | 125 | 13 | 4 | 291 |
| Petroleum coke | 200 | 100 | 0 | 500 |
| Bituminous coal | 200 | 100 | 10 | 500 |
| Lignite | 213 | 100 | 10 | 500 |
| Natural gas | 19 | 9 | 3 | 0.5 |
| Biomass | 137 | 282 | 7 | 15 |
| Other fuels (fossil waste incineration in cement industry) | IE | IE | IE | IE |
| Fuels, not itemized (fiber construction board, fine ceramics, glass, glass wool, bottle glas, lime, asphalt, rock wool, brick, cement) | 253 | 455 | 20 | 127 |
| 1A2fii Construction and industrial machinery | NO _x | CO | NMVOC | SO ₂ |
| | kg/h | kg/h | kg/h | kg/h |
| Diesel and gasoline | 440 | 807 | 75 | 0.5 |

A3.1.4 Civil Aviation

This paragraph contains further information to the emission modelling. More complete information is provided in FOCA (2006-2012) and on request for reviewers by FOCA.

Emission factors

Table A - 8 Aircraft cruise factors, used for cruise emission calculation (extract of list of 881 aircraft) GKL_ICAO = ICAO seat categories. Mass emissions are given in kilograms or grams per nautical mile (NM).

| Aircraft Cruise _Factors | | | | | | |
|--------------------------|----------|-------------------|------------|-----------|----------|---------|
| Aircraft_ICAO | GKL_ICAO | Cruise_D_Source | kg_fuel_NM | kg_NOx_NM | g_VOC_NM | g_CO_NM |
| AA1 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AA5 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AC11 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AC14 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AC50 | 0 | P001FOCA | 0.77 | 0.021 | 4.14 | 364.17 |
| AC68 | 0 | P001FOCA | 0.77 | 0.0075 | 4.14 | 364.17 |
| AC6T | 1 | FOCAINV95-03.2T | 1.58 | 0.021 | 0.87 | 2.9 |
| AC90 | 1 | FOCAINV95-03.2T | 1.58 | 0.021 | 0.87 | 2.9 |
| AC95 | 1 | FOCAINV95-03.2T | 1.58 | 0.021 | 0.87 | 2.9 |
| AEST | 0 | P001FOCA | 0.77 | 0.021 | 4.14 | 364.17 |
| AJET | 0 | FOCAEDBJ014 | 2.92 | 0.0146 | 8.53 | 63 |
| ALO2 | 0 | FOCAHeli | 1.91 | 0.024 | 0.42 | 2.1 |
| ALO3 | 0 | FOCAHeli | 1.91 | 0.024 | 0.42 | 2.1 |
| AN12 | 0 | AN26*2 | 5.36 | 0.0062 | 143 | 348 |
| AN2 | 0 | FOCA/91/DC3 | 0.82 | 0.0002 | 13.7 | 1000 |
| AN22 | 6 | FOCAINV95-03.2T*2 | 3.16 | 0.042 | 1.74 | 5.8 |
| AN24 | 2 | AN26 | 2.68 | 0.0031 | 71.7 | 174 |
| AN26 | 1 | 500 | 2.68 | 0.0031 | 71.7 | 174 |
| AN72 | 2 | FOCAINV95-03.2J | 6.4 | 0.1 | 0.83 | 10 |
| AR7 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AR7A | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS02 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS16 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS20 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS24 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS25 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS26 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| AS2T | 0 | FOCAEDBT758 | 0.95 | 0.005 | 1.8 | 12 |
| AS30 | 0 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| AS32 | 1 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| AS33 | 0 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| AS35 | 0 | FOCAHeli | 1.91 | 0.024 | 0.42 | 2.1 |
| AS50 | 0 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| AS55 | 0 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| AS65 | 0 | FOCAHeli*2 | 3.82 | 0.048 | 0.82 | 4.2 |
| ASK1 | 0 | P002FOCA | 0.21 | 0.0098 | 1.79 | 61.7 |
| ASTA | 0 | FOCAINV95-03.B | 3.016 | 0.046 | 0.3 | 2.8 |
| ASTR | 0 | FOCAINV95-03.B | 3.016 | 0.046 | 0.3 | 2.8 |
| ASTRA | 0 | FOCAINV95-03.B | 3.016 | 0.046 | 0.3 | 2.8 |
| AT42 | 1 | FOCAINV95-03.2T | 1.58 | 0.021 | 0.87 | 2.9 |
| AT43 | 1 | 500 | 1.6 | 0.013 | 0 | 15 |

Activity data

LTO-cycle times (minutes). ICAO standard cycle times were originally designed for emissions certification, not for emissions modelling. Today, they do generally not match real world aircraft LTO operations. Swiss FOCA has therefore adjusted some of the ICAO standard cycle times for different aircraft categories. For jets, the mean time for taxi-in and taxi-out at Swiss airports has been determined 20 minutes instead of the standard 26 minutes.

Table A - 9 For jets, business jets, turboprops, piston engines and helicopters, the times in mode are shown and are based on ICAO, US EPA and Swiss FOCA data. "Type" is a classification variable. J = Jet, T = Turboprop, P = Piston, H = Helicopter, B = Business jet, SJ = Supersonic Jet. The number in "Type" stands for the number of engines. For Jet Aircraft, the cycle times and associated thrust settings still lead to an overestimation of LTO emissions (FOCA 2007b).

| LTO Cycle | | | | |
|-----------|---------------|---------------|---------------|-----------|
| Type | Time_Take_Off | Time_Climbout | Time_Approach | Zeit_Taxi |
| 1J | 0.7 | 2.2 | 4 | 20 |
| 1T | 0.5 | 2.5 | 4.5 | 13 |
| 1P | 0.3 | 2.5 | 3 | 12 |
| 1H | 0 | 6.5 | 6.5 | 7 |
| 2B | 0.4 | 0.5 | 1.6 | 13 |
| 3B | 0.4 | 0.5 | 1.6 | 13 |
| 2T | 0.5 | 2.5 | 4.5 | 13 |
| 4T | 0.5 | 2.5 | 4.5 | 13 |
| 2J | 0.7 | 2.2 | 4 | 20 |
| 3J | 0.7 | 2.2 | 4 | 20 |
| 4J | 0.7 | 2.2 | 4 | 20 |
| 2P | 0.3 | 2.5 | 3 | 12 |
| 3P | 0.3 | 2.5 | 3 | 12 |
| 4P | 0.3 | 2.5 | 3 | 12 |
| 2H | 0 | 6.5 | 6.5 | 7 |
| 4SJ | 1.2 | 2 | 2.3 | 20 |
| 3H | 0 | 6.5 | 6.5 | 7 |
| 4H | 0 | 6.5 | 6.5 | 7 |
| 4B | 0.4 | 0.5 | 1.6 | 13 |

Table A - 10 Aircraft-Engine Combinations and associated codes for SWISS FOCA emissions database. (Extract from list of more than 26'000 individual aircraft)

| Aircraft Engine Combinations | | | | | | | |
|------------------------------|--------------------------------------|-------------------|----------|------|------|-------------|---------------|
| Engine Name | Aircraft Name | Aircraft Registr. | No. Eng. | Code | Type | Aircr. ICAO | Source |
| V2527-A5 | AIRBUS A320-232 | ECHXA | 2 | J220 | 2J | A320 | 1IA003 |
| CF34-3B1 | BOMBARDIER CRJ200ER (CL-600-2B19) | ECHXM | 2 | J090 | 2J | CRJ2 | 1GE034 |
| CFM56-3C1 | BOEING 737-4K5 | ECHXT | 2 | J022 | 2J | B734 | 1CM007 |
| TPE331-11U-611G | FAIRCHILD (SWEARIN-GEN) SA227AC METR | ECHXY | 2 | T310 | 2T | SW4 | FOI |
| CFM56-5B4/P | AIRBUS A320-214 | ECHYC | 2 | J067 | 2J | A320 | 3CM026 |
| CFM56-5B4/P | AIRBUS A320-214 | ECHYD | 2 | J067 | 2J | A320 | 3CM026 |
| CF34-3B1 | BOMBARDIER CRJ200ER (CL-600-2B19) | ECHYG | 2 | J090 | 2J | CRJ2 | 1GE034 |
| CFEC-FE738-1-1B | DASSAULT FALCON 2000 | ECHYI | 2 | B130 | 2B | F2TH | FOI-Honeywell |
| GA TPE331-11U-612G | | ECHZH | 2 | T310 | 2T | FA3 | FOI |
| CF34-3B1 | BOMBARDIER CRJ200ER (CL-600-2B19) | ECHZR | 2 | J090 | 2J | CRJ2 | 1GE034 |
| CFM56-7B27B1 | BOEING 737-86Q (WINGLETS) | ECHZS | 2 | J075 | 2J | B738 | 3CM034 |
| CFM56-5B4/P | AIRBUS A320-214 | ECHZU | 2 | J067 | 2J | A320 | 3CM026 |
| CF34-3B1 | BOMBARDIER CRJ200ER (CL-600-2B19) | ECIAA | 2 | J090 | 2J | CRJ2 | 1GE034 |
| FJ44-1A | CESSNA 525 CITATIONJET | ECIAB | 2 | B001 | 2B | C525 | FOCA |
| CFM56-5B4/P | AIRBUS A320-214 | ECIAG | 2 | J067 | 2J | A320 | 3CM026 |
| V2527-A5 | AIRBUS A320-232 | ECIAZ | 2 | J220 | 2J | A320 | 1IA003 |
| BRBR700-710A2-20 | BOMBARDIER BD-700-1A10 GLOBAL EX-PRE | ECIBD | 2 | J854 | 2J | GLEX | 4BR009 |
| PT6A-60A | BEECH-CRAFT KING AIR 350 (RAYTHEON B | ECIBK | 2 | T738 | 2T | B350 | FOI |
| CF34-3B1 | BOMBARDIER CRJ200ER (CL-600-2B19) | ECIBM | 2 | J090 | 2J | CRJ2 | 1GE034 |
| CFM56-7B27B1 | BOEING 737-81Q (WINGLETS) | ECICD | 2 | J075 | 2J | B738 | 3CM034 |
| CFM56-5B4/P | AIRBUS A320-214 | ECICK | 2 | J067 | 2J | A320 | 3CM026 |

Emissions

The output of the FOCA emission modelling consists of tables with the following structure:

Table A - 11 Extract of the output file of FOCA emission and fuel consumption modelling. Upper part: LTO, lower part: cruise (example for 2004). Emissions and fuel consumption in tons.

| Airport | Distance | Type Traffic | Movements | Type | Aircraft ICAO | Engine Name | Fuel (LTO) tons | Emissions (LTO) in tons | | | | | |
|---------|------------|--------------|-----------|------|---------------|--------------|-----------------|-------------------------|------------------|-----------------|-----------------|-------|--------|
| | Km | | No. | | | | | CO ₂ | H ₂ O | SO ₂ | NO _x | VOC | CO |
| LSGG | 181501.69 | Taxi | 165 | 2B | C550 | JT15D-4 | 5673.492 | 17871.5 | 6978.395 | 5.673 | 26.04 | 139 | 359.2 |
| LSGG | 164165.197 | Taxi | 77 | 2J | B752 | RB211-535E4 | 47470.5 | 149532.1 | 58388.72 | 47.47 | 554.91 | 0 | 361.47 |
| LSGG | 133166.837 | Taxi | 118 | 2B | F2TH | CFE738-1-1B | 6164.2728 | 19417.46 | 7582.056 | 6.164 | 87.539 | 40.59 | 185.53 |
| LSGG | 117228.943 | Taxi | 99 | 3B | F900 | TFE731-60-1C | 5668.542 | 17855.91 | 6972.307 | 5.669 | 46.937 | 28.13 | 163.44 |
| LSGG | 114258.902 | Taxi | 134 | 2B | LJ45 | TFE731-20R | 4725.108 | 14884.09 | 5811.883 | 4.725 | 31.31 | 53.62 | 169.01 |
| LSGG | 112510.267 | Taxi | 100 | 2B | F2TH | CFE738-1-1B | 5223.96 | 16455.47 | 6425.471 | 5.224 | 74.186 | 34.4 | 157.23 |
| LSGG | 107945.477 | Taxi | 96 | 2B | C560 | JT15D-5D | 3795.3216 | 11955.26 | 4668.246 | 3.795 | 16.959 | 271.6 | 287.98 |
| LSGG | 181501.69 | Taxi | 165 | 2B | C550 | JT15D-4 | 307732.68 | 969357.9 | 378511.2 | 307.7 | 4513 | 29.43 | 274.71 |
| LSGG | 164165.197 | Taxi | 77 | 2J | B752 | RB211-535E4 | 673698.47 | 2122150 | 828649.1 | 673.7 | 7986.4 | 647.8 | 1038.2 |
| LSGG | 133166.837 | Taxi | 118 | 2B | F2TH | CFE738-1-1B | 225781.85 | 711212.8 | 277711.7 | 225.8 | 3311.2 | 21.59 | 201.55 |
| LSGG | 117228.943 | Taxi | 99 | 3B | F900 | TFE731-60-1C | 298139.18 | 939138.4 | 366711.2 | 298.1 | 4372.3 | 28.52 | 266.14 |
| LSGG | 114258.902 | Taxi | 134 | 2B | LJ45 | TFE731-20R | 193723.81 | 610230 | 238280.3 | 193.7 | 2841 | 18.53 | 172.93 |
| LSGG | 106761.289 | Taxi | 100 | 2B | F2TH | CFE738-1-1B | 181011.75 | 570187 | 222644.4 | 181 | 2654.6 | 17.31 | 161.58 |
| LSGG | 103217.159 | Taxi | 96 | 2B | C560 | JT15D-5D | 175002.74 | 551258.6 | 215253.4 | 175 | 2566.5 | 16.74 | 156.22 |

A3.1.5 Road Transportation

Emission factors

The derivation of the emission factors for road vehicles is described in detail in INFRAS 2010 (this report is available in English). Some important features of the emission factor methodologies are summarised in this paragraph.

The emission factors have to be differentiated according to the vehicle categories. Each category contains a number of vehicle classes, which differ by emission concepts. The next table illustrates the classes of the passenger cars. Similar “segmentations” hold for the other vehicle categories too. Emission factors for vehicle classes are combined to average emission factors for vehicles categories weighted according to the fleet composition, which varies from year to year (see below).

Table A - 12 Vehicle segmentation of the passenger cars. Each segment is subdivided into three cubic capacities: <1.4 litre, 1.4-2.0 litres, > 2.0 litres (INFRAS 2010).

| Fuel type | Vehicle segment |
|-----------|-------------------------------|
| Gasoline | <ECE |
| | AGV82 (CH) |
| | PreEuro 3WayCat <1987 |
| | PreEuro 3WayCat 1987-90 |
| | ECE-15'00 |
| | ECE-15'01/02 |
| | ECE-15'03 |
| | Euro-1 |
| | Euro-2 |
| | Euro-3 |
| | Euro-4 |
| | Euro-5 |
| | Euro-6 |
| Diesel | <1986 |
| | 1986-1988 |
| | Euro-1 |
| | Euro-2 |
| | Euro-3 |
| | Euro-4 |
| | Euro-5 Diesel Particle Filter |
| | Euro-6 Diesel Particle Filter |

The emission factors published in the handbook (CD ROM, INFRAS 2010) are classified by “traffic situations”. The scheme (see Table below) distinguishes the traffic situations along 4 dimensions: urban/rural areas, 5 functional road types, speed limit and 4 levels of service. This leads to the definition of 276 different traffic situations in total. A traffic situation is primarily characterised by the type of road which induces a typical driving behaviour. (Because driving behaviour is not independent of the amount of traffic on that particular road, on the same segment different driving patterns may exist.) For the handbook several typical traffic situations have been defined, based on driving behaviour studies in Germany and in Switzerland (see e.g. SAEFL 1995, Chp. 4).

Table A - 13 Traffic situation-scheme in HBEFA 3.1 (INFRAS 2010). Every traffic situation is characterised by a typical driving pattern (i.e. a speed-time curve)

| | | | Speed Limit [km/h] | | | | | | | | | | | | |
|-------|--------------------------------|---------------------|--------------------|----|----|----|----|----|----|-----|-----|-----|-----|------|--|
| Area | Road type | Levels of service | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 | 110 | 120 | 130 | >130 | |
| Rural | Motorway-Nat. | 4 levels of service | | | | | | | | | | | | | |
| | Semi-Motorway | 4 levels of service | | | | | | | | | | | | | |
| | TrunkRoad/Primary-Nat. | 4 levels of service | | | | | | | | | | | | | |
| | Distributor/Secondary | 4 levels of service | | | | | | | | | | | | | |
| | Distributor/Secondary(sinuous) | 4 levels of service | | | | | | | | | | | | | |
| | Local/Collector | 4 levels of service | | | | | | | | | | | | | |
| | Local/Collector(sinuous) | 4 levels of service | | | | | | | | | | | | | |
| | Access-residential | 4 levels of service | | | | | | | | | | | | | |
| Urban | Motorway-Nat. | 4 levels of service | | | | | | | | | | | | | |
| | Motorway-City | 4 levels of service | | | | | | | | | | | | | |
| | TrunkRoad/Primary-Nat. | 4 levels of service | | | | | | | | | | | | | |
| | TrunkRoad/Primary-City | 4 levels of service | | | | | | | | | | | | | |
| | Distributor/Secondary | 4 levels of service | | | | | | | | | | | | | |
| | Local/Collector | 4 levels of service | | | | | | | | | | | | | |
| | Access-residential | 4 levels of service | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | |

Traffic situations are defined independently of vehicle categories (LDV, HDV, 2-wheelers). But behind the same traffic situation each vehicle category may know its own “driving pattern” which may be expressed as a speed curve (i.e. speed time series). Emission factors originally are derived for these underlying driving patterns based on measurements performed on laboratory test benches. Emission factors per traffic situation are then calculated by attributing the driving patterns to different traffic situations (based on statistical analysis).

Emission factors for Switzerland are shown in the next table (FOEN 2010i, updated based on Prognos 2012a). They represent weighted averages over all traffic situations. The year indicates the date when the corresponding vehicle class appears in the market. E.g. “Euro-3” standard came into force on 1 Jan, 2001, but the first vehicles with Euro-3 standard already appeared in 1999.

Table A - 14 Mean emission factors of passenger cars (PC), light duty vehicles (LDV), heavy duty vehicles (HDV), coaches, urban buses (Bus) and Motorcycles (MC) in grams per kilometre, incl. cold starts and evaporation. (FOEN 2010i, updated based on Prognos 2012a). CO₂ (rep.) refers to the fossil part, CO₂ (total) includes fossil and biomass.

| Pollutant | Year | PC | LDV | HDV | Coach | Bus | MC |
|--|------|-------|-------|--------|--------|--------|-------|
| grams per vehicle kilometre, incl. cold starts and evaporation | | | | | | | |
| CH ₄ | 1990 | 0.080 | 0.090 | 0.020 | 0.017 | 0.053 | 0.236 |
| CH ₄ | 1995 | 0.050 | 0.065 | 0.018 | 0.016 | 0.046 | 0.159 |
| CH ₄ | 2000 | 0.033 | 0.039 | 0.013 | 0.014 | 0.034 | 0.120 |
| CH ₄ | 2005 | 0.021 | 0.020 | 0.009 | 0.011 | 0.018 | 0.103 |
| CH ₄ | 2010 | 0.013 | 0.012 | 0.004 | 0.007 | 0.005 | 0.092 |
| CH ₄ | 2015 | 0.009 | 0.006 | 0.002 | 0.004 | 0.003 | 0.082 |
| CO | 1990 | 9.58 | 20.16 | 2.39 | 2.09 | 5.99 | 14.70 |
| CO | 1995 | 5.46 | 14.60 | 2.18 | 2.01 | 5.68 | 14.14 |
| CO | 2000 | 3.59 | 8.86 | 1.77 | 1.84 | 4.64 | 13.62 |
| CO | 2005 | 2.53 | 4.39 | 1.61 | 1.73 | 2.92 | 11.68 |
| CO | 2010 | 1.64 | 2.09 | 1.44 | 1.70 | 1.26 | 8.09 |
| CO | 2015 | 1.19 | 1.28 | 1.27 | 1.63 | 1.03 | 5.63 |
| CO ₂ (rep.) | 1990 | 234 | 249 | 803 | 871 | 1'194 | 82 |
| CO ₂ (rep.) | 1995 | 237 | 252 | 799 | 860 | 1'199 | 90 |
| CO ₂ (rep.) | 2000 | 230 | 254 | 759 | 833 | 1'162 | 92 |
| CO ₂ (rep.) | 2005 | 217 | 246 | 790 | 823 | 1'127 | 94 |
| CO ₂ (rep.) | 2010 | 195 | 237 | 768 | 812 | 1'078 | 96 |
| CO ₂ (rep.) | 2015 | 169 | 224 | 739 | 794 | 1'045 | 91 |
| CO ₂ (total) | 1990 | 234 | 249 | 803 | 871 | 1'194 | 82 |
| CO ₂ (total) | 1995 | 237 | 252 | 799 | 860 | 1'199 | 90 |
| CO ₂ (total) | 2000 | 230 | 255 | 760 | 834 | 1'163 | 92 |
| CO ₂ (total) | 2005 | 217 | 246 | 793 | 826 | 1'131 | 94 |
| CO ₂ (total) | 2010 | 199 | 240 | 777 | 821 | 1'094 | 99 |
| CO ₂ (total) | 2015 | 180 | 232 | 760 | 817 | 1'079 | 98 |
| HC | 1990 | 1.58 | 2.02 | 0.85 | 0.70 | 2.20 | 3.69 |
| HC | 1995 | 0.92 | 1.38 | 0.74 | 0.66 | 1.93 | 2.65 |
| HC | 2000 | 0.57 | 0.77 | 0.56 | 0.60 | 1.42 | 2.08 |
| HC | 2005 | 0.36 | 0.38 | 0.38 | 0.47 | 0.73 | 1.64 |
| HC | 2010 | 0.23 | 0.19 | 0.16 | 0.27 | 0.22 | 1.19 |
| HC | 2015 | 0.17 | 0.12 | 0.07 | 0.16 | 0.12 | 0.85 |
| N ₂ O | 1990 | 0.009 | 0.005 | 0.008 | 0.008 | 0.003 | 0.002 |
| N ₂ O | 1995 | 0.012 | 0.007 | 0.009 | 0.008 | 0.003 | 0.002 |
| N ₂ O | 2000 | 0.011 | 0.009 | 0.009 | 0.008 | 0.003 | 0.002 |
| N ₂ O | 2005 | 0.005 | 0.007 | 0.008 | 0.007 | 0.002 | 0.002 |
| N ₂ O | 2010 | 0.003 | 0.006 | 0.030 | 0.014 | 0.001 | 0.002 |
| N ₂ O | 2015 | 0.002 | 0.005 | 0.041 | 0.023 | 0.002 | 0.002 |
| NMHC | 1990 | 1.504 | 1.930 | 0.827 | 0.681 | 2.151 | 3.451 |
| NMHC | 1995 | 0.871 | 1.320 | 0.721 | 0.640 | 1.880 | 2.489 |
| NMHC | 2000 | 0.538 | 0.735 | 0.543 | 0.582 | 1.383 | 1.964 |
| NMHC | 2005 | 0.343 | 0.362 | 0.373 | 0.459 | 0.714 | 1.538 |
| NMHC | 2010 | 0.217 | 0.180 | 0.155 | 0.265 | 0.209 | 1.096 |
| NMHC | 2015 | 0.163 | 0.116 | 0.069 | 0.156 | 0.110 | 0.773 |
| NO _x | 1990 | 2.147 | 4.167 | 22.457 | 22.929 | 33.896 | 0.294 |
| NO _x | 1995 | 1.608 | 3.485 | 20.751 | 21.648 | 32.841 | 0.392 |
| NO _x | 2000 | 1.301 | 3.067 | 18.148 | 19.938 | 29.997 | 0.423 |
| NO _x | 2005 | 0.968 | 2.595 | 15.051 | 17.360 | 24.703 | 0.444 |
| NO _x | 2010 | 0.679 | 2.117 | 9.512 | 13.381 | 18.000 | 0.400 |
| NO _x | 2015 | 0.535 | 1.857 | 6.213 | 10.142 | 13.576 | 0.352 |
| SO ₂ | 1990 | 0.040 | 0.093 | 0.714 | 0.774 | 1.061 | 0.010 |
| SO ₂ | 1995 | 0.031 | 0.041 | 0.173 | 0.186 | 0.260 | 0.011 |
| SO ₂ | 2000 | 0.022 | 0.034 | 0.131 | 0.144 | 0.201 | 0.008 |
| SO ₂ | 2005 | 0.001 | 0.001 | 0.005 | 0.005 | 0.007 | 0.000 |
| SO ₂ | 2010 | 0.001 | 0.001 | 0.005 | 0.005 | 0.007 | 0.001 |
| SO ₂ | 2015 | 0.001 | 0.001 | 0.005 | 0.005 | 0.006 | 0.001 |

Activity Data

Activity data for the emission model are the mileages of the vehicle categories per traffic situation. To that aim, three steps must be carried out.

1. Vehicle turnover: The vehicle fleet is built up for each year accounting for the stock changes. This vehicle turnover is modelled on the basis of new registrations and by applying survival probabilities. Trends in traffic volume per vehicle category, including structural changes (size distributions, shares of diesel vehicles) are then combined to draw the continual substitution of older technologies by new ones constantly altering the fleet composition or mileage by emission concepts in all vehicle categories (see following Figure).
2. The total mileage is calculated by vehicle stock multiplied with the specific mileage per vehicle and annum. The latter data are derived from household surveys and from specific odometer readings during vehicle inspections (ARE 2002).
3. Assignment of the mileage to the traffic situations for all vehicle categories. This step requires the adoption of the traffic model: Each road segment carries its mileage and its traffic, which allows the assignment sought.




Figure A - 3 Mileage composition by emission concept (in million vehicle kilometres per year), FOEN 2010i.

Modelling hot exhaust emissions

As a next step in the modelling process, the mileage classified by vehicle segments and traffic situations is multiplied with the emission factors resulting in hot exhaust emissions.

The results do not yet contain the emissions from tank tourism. For this purpose a special procedure is carried out (described in section 3.2.8.2 b), providing the fuel consumption of tank tourism. From that, the emissions are calculated by multiplication with mean emission factors.

Cold start and evaporative emissions

The handbook also contains emission factors for modelling cold start excess emissions and evaporative emissions (diurnal and hot/warm soak and running losses). For a technical description the reader may be referred to INFRAS (2010).

Results show that for CO₂ the hot exhaust emissions contribute to 97% of the total. Only 3% stem from cold start excess emissions. For CH₄ however, the picture is much different. Only about 40% of the emission total is hot exhaust. More than 59% are cold start excess emissions, the rest results evaporative emissions. For N₂O, no cold start emissions or evaporative emissions are taken into account due to lack of data.

A3.1.6 Off-road Vehicles

Methodology

The emissions of the whole off-road sector underwent a complete revision for Submission 2010. The emissions are calculated with a Tier 2 method. Activity data and emission factors were updated for Submission 2010; for the present Submission, activity data have again been recalculated following the update of the latest figures on population and economy (Prognos 2012a). The modelling is carried out in a database (INFRAS 2008) that is structured in analogy to the on-road database (INFRAS 2010). The off-road sector has been allocated to 1A2, 1A3 and 1A4, with only military offroad remaining in 1A5.

The modelling of the emission and of the fuel consumption are carried out by using the formula

$$E_{i,j,t,\tau}^g = N_{i,j,t} \cdot T_{i,j,t} \cdot \omega_{t-\tau} \cdot P_{i,j} \cdot L_{i,j} \cdot v_{t-\tau} \cdot \varepsilon_{i,j,\tau}^g$$

E: Emission and fuel consumption

N: number of vehicles

T: average operating hours per year

ω : age dependency

P: motor power in kW

L: load factor

v: degradation factor (due to aging)

ε : emission factor in g/kWh

indices: g: gas (CH₄, N₂O, CO, NO_x, SO₂) and fuel consumption,

i off-road family (railway, navigation etc.),

j size class,

t: year (1980, 1985, 1990, 1995, 2000, ... , 2020)

τ : year of construction (note: $t - \tau$ = age of vehicle)

Note that the emissions are only calculated in steps of 5 years. Emissions for years in-between like 1991, 1992 etc. are interpolated linearly.

Emission and fuel consumption factors for off-road vehicles

The CO₂ emission factors are derived from fuel type and fuel consumption (see tables below). The emission factors for CH₄ and N₂O are only specified by the fuel type.

Table A - 15 CH₄ (TTM 2006a) and N₂O (TTM 2006b) emission factors used in the off-road model (INFRAS 2008).

| Gas | Diesel | Gasoline | |
|------------------|--------|----------|----------|
| | | 4-stroke | 2-stroke |
| | | mg/kWh | |
| CH ₄ | 6 | 500 | 4000 |
| N ₂ O | 30 | 50 | -- |

The values differ from default values (IPCC 1996, vol III, tbl 1-7, 1-8, conversion factor used: 1 g/kWh = 278 kg/TJ): For CH₄ IPCC recommends 18 mg/kWh for diesel oil, 72 mg/kWh for gasoline four-stroke, 210 mg/kWh gasoline two-stroke. For N₂O IPCC gives 2 mg/kWh (diesel oil and gasoline four-stroke) and 6 mg/kWh (gasoline two-stroke).

Table A - 16 Emission and consumption factors for diesel engines (without ships and rail vehicles). PreEU-A etc. indicate emission standards.

| Basic emission factors of diesel engines (g/kWh) | | | | | | |
|--|------------------|-----------------|-------------------|--------------------|---------------------|----------------------|
| power class | PreEU-A <1996 | PreEU-B 1996 | EU-I 2002/2003 | EU-II 2003/2004 | EU-III 2007/2008 | EU-IIIB 2011/2012 |
| Carbon monoxide (CO) | | | | | | |
| <18 kW | 6.71 | 6.71 | 2.90 | 2.90 | 2.90 | 2.90 |
| 18-37 kW | 6.71 | 6.71 | 2.76 | 2.42 | 2.06 | 1.76 |
| 37-75 kW | 4.68 | 4.68 | 1.87 | 1.63 | 1.39 | 1.19 |
| 75-130 kW | 3.62 | 3.62 | 1.28 | 1.01 | 0.86 | 0.73 |
| >130 kW | 3.62 | 3.62 | 1.04 | 0.91 | 0.77 | 0.66 |
| VOC | | | | | | |
| <18 kW | 2.28 | 2.28 | 1.60 | 1.00 | 0.59 | 0.59 |
| 18-37 kW | 2.41 | 2.41 | 0.92 | 0.56 | 0.37 | 0.37 |
| 37-75 kW | 1.33 | 1.33 | 0.65 | 0.46 | 0.33 | 0.24 |
| 75-130 kW | 0.91 | 0.91 | 0.45 | 0.35 | 0.28 | 0.17 |
| >130 kW | 0.91 | 0.91 | 0.43 | 0.3 | 0.22 | 0.17 |
| Nitrogen oxides (NO_x) | | | | | | |
| <18 kW | 10.31 | 8.2 | 5.95 | 5.95 | 5.95 | 5.95 |
| 18-37 kW | 10.31 | 8.2 | 6.34 | 6.34 | 6.34 | 6.34 |
| 37-75 kW | 12.4 | 9.87 | 8.95 | 6.56 | 3.90 | 3.39 |
| 75-130 kW | 12.52 | 9.96 | 8.44 | 5.67 | 3.32 | 2.97 |
| >130 kW | 12.52 | 9.96 | 8.19 | 5.66 | 3.38 | 1.80 |
| Fuel consumption (FC) | | | | | | |
| <18 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 18-37 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 37-75 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 75-130 kW | 223 | 223 | 223 | 223 | 223 | 223 |
| >130 kW | 223 | 223 | 223 | 223 | 223 | 223 |

Table A - 17 Emission and consumption factors for gasoline four-stroke engines. PreEU-A etc. indicate emission standards.

| Basic emission factors of equipment with 4-stroke gasoline engines (g/kWh). | | | | | |
|---|------------------|-----------------|-----------------|--------------|--------------------|
| power class | PreEU-A <1995 | PreEU-B 1995 | PreEU-C 2000 | EU-I 2004 | EU-II 2005-2009 |
| Carbon monoxide (CO) | | | | | |
| <66 ccm | 470 | 470 | 470 | 467 | 467 |
| 66-100 ccm | 470 | 470 | 470 | 467 | 467 |
| 100-225 ccm | 470 | 470 | 470 | 467 | 467 |
| >225 ccm | 470 | 470 | 470 | 467 | 467 |
| VOC | | | | | |
| <66 ccm | 60 | 60 | 60 | 41 | 41 |
| 66-100 ccm | 40 | 40 | 40 | 32 | 32 |
| 100-225 ccm | 20 | 20 | 20 | 12 | 12 |
| >225 ccm | 20 | 20 | 20 | 10 | 9 |
| Nitrogen oxides (NO_x) | | | | | |
| <66 ccm | 1.5 | 2 | 3 | 4.5 | 4.5 |
| 66-100 ccm | 1.5 | 2 | 3 | 3.6 | 3.6 |
| 100-225 ccm | 3.5 | 3.5 | 3.5 | 2.8 | 2.8 |
| >225 ccm | 3.5 | 3.5 | 3.5 | 2.2 | 1.9 |
| Fuel consumption (FC) | | | | | |
| <66 ccm | 500 | 500 | 500 | 480 | 480 |
| 66-100 ccm | 480 | 480 | 480 | 470 | 470 |
| 100-225 ccm | 460 | 460 | 460 | 450 | 450 |
| >225 ccm | 460 | 460 | 460 | 450 | 450 |

Table A - 18 Emission and consumption factors for gasoline two-stroke engines. PreEU-A etc. indicate emission standards.

| Basic emission factors of equipment with 2-stroke gasoline engines (g/kWh) | | | | | |
|--|------------------|-----------------|-----------------|--------------|--------------------|
| gas/fuel consumption | PreEU-A <1995 | PreEU-B 1995 | PreEU-C 2000 | EU-I 2004 | EU-II 2009/2011 |
| Carbon monoxide (CO) | 650 | 640 | 620 | 600 | 600 |
| VOC | 260 | 250 | 150 | 100 | 41 |
| Nitrogen oxides (NO _x) | 1.5 | 2.0 | 3.0 | 4.8 | 4.5 |
| Fuel consumption (FC) | 660 | 650 | 550 | 500 | 440 |

Table A - 19 Emission and consumption factors for rail vehicles with diesel engines. PreEU etc. indicate emission standards.

| Basic emission factors of rail vehicles (g/kWh) | | | | | |
|---|----------------|---------------|----------------|----------------------|-----------------|
| power class | PreEU <2000 | UIC I 2000 | UIC II 2003 | EU IIIa 2006/2009 | EU IIIb 2012 |
| Carbon monoxide (CO) | | | | | |
| <560 kW | 4 | 3 | 2.5 | 2.5 | 2.5 |
| >560 kW | 4 | 3 | 3 | 3 | 3 |
| VOC | | | | | |
| <560 kW | 1.6 | 0.8 | 0.6 | 0.4 | 0.17 |
| >560 kW | 1.6 | 0.8 | 0.8 | 0.5 | 0.4 |
| Nitrogen oxides (NO_x) | | | | | |
| <560 kW | 13 | 12 | 6 | 3.2 | 1.8 |
| >560 kW | 16 | 12 | 9.5 | 5.4 | 3.2 |
| Fuel consumption (FC) | | | | | |
| <560 kW | 223 | 223 | 223 | 223 | 223 |
| >560 kW | 223 | 223 | 223 | 223 | 223 |

Table A - 20 Emission and consumption factors for ships with diesel engines. PreSAV etc. indicate emission standards.

| Basic emission factors of diesel engine ships (g/kWh) | | | | | | |
|---|-----------------|-------------|--------------|---------------|------------------|------------------|
| power class | PreSAV <1995 | SAV 1995 | EU I 2003 | EU II 2008 | EU III-a 2009 | EU III-b 2012 |
| Carbon monoxide (CO) | | | | | | |
| <18 kW | 6.7 | 6.7 | 6.7 | 6.7 | 6.7 | 5.5 |
| 18-37 kW | 6.7 | 6.7 | 6.7 | 6.7 | 6.7 | 5 |
| 37-75 kW | 5.9 | 5.9 | 5.9 | 4.5 | 4.5 | 4.5 |
| 75-130 kW | 5 | 5 | 4.5 | 4.5 | 4.5 | 4.5 |
| >130 kW | 5 | 5 | 4.5 | 4.5 | 4.5 | 3.5 |
| VOC | | | | | | |
| <18 kW | 10 | 7.2 | 5 | 3 | 2 | 1 |
| 18-37 kW | 10 | 7.2 | 5 | 3 | 2 | 1 |
| 37-75 kW | 10 | 5.4 | 1.2 | 1.2 | 1.1 | 0.6 |
| 75-130 kW | 10 | 4.1 | 1.2 | 0.9 | 0.8 | 0.4 |
| 130-399 kW | 5 | 3.6 | 1.2 | 0.9 | 0.8 | 0.4 |
| 300-560 kW | 5 | 3.2 | 1.2 | 0.9 | 0.8 | 0.4 |
| >560 kW | 5 | 2.8 | 1.2 | 0.9 | 0.8 | 0.4 |
| Nitrogen oxides (NO_x) | | | | | | |
| <18 kW | 10.3 | 10.3 | 10.3 | 10.3 | 10.3 | 8 |
| 18-37 kW | 10.3 | 10.3 | 10.3 | 10.3 | 10.3 | 8 |
| 37-75 kW | 12.4 | 12.4 | 8.3 | 6.3 | 5.7 | 3.3 |
| 75-130 kW | 12.5 | 12.5 | 8.3 | 6.3 | 5.7 | 3.3 |
| >130 kW | 12.5 | 12.5 | 8.3 | 6.3 | 5.7 | 2 |
| Fuel consumption (FC) | | | | | | |
| <18 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 18-37 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 37-75 kW | 248 | 248 | 248 | 248 | 248 | 248 |
| 75-130 kW | 223 | 223 | 223 | 223 | 223 | 223 |
| >130 kW | 223 | 223 | 223 | 223 | 223 | 223 |

Table A - 21 Emission and consumption factors for boats with diesel engines. PreSAV etc. indicate emission standards.

| Basic emission factors of diesel engine boats (g/kWh) | | | | |
|--|----------------------------|---------------------|----------------------|-----------------------|
| power class | PreSAV <1995 | SAV 1995 | EU-I 2007 | EU II 2012 |
| Carbon monoxide (CO) | | | | |
| <4.4 kW | 6.7 | 6.7 | 4.5 | 4 |
| 4.4-7.4 kW | 6.7 | 6.7 | 4.5 | 4 |
| 7.4-37 kW | 6.7 | 6.7 | 4.5 | 4 |
| 37-74 kW | 5.9 | 5.9 | 4.5 | 4 |
| 74-100 kW | 5 | 5 | 4.5 | 4 |
| >100 kW | 5 | 3.6 | 3.6 | 3.5 |
| VOC | | | | |
| <4.4 kW | 10 | 10 | 2.4 | 1.5 |
| 4.4-7.4 kW | 10 | 10 | 2.1 | 1.5 |
| 7.4-37 kW | 10 | 2 | 1.7 | 1.5 |
| 37-74 kW | 10 | 1.4 | 1.4 | 1.3 |
| 74-100 kW | 10 | 1.2 | 1.2 | 1 |
| >100 kW | 5 | 1.2 | 1.2 | 1 |
| Nitrogen oxides (NO_x) | | | | |
| <4.4 kW | 13 | 11 | 8.8 | 6 |
| 4.4-7.4 kW | 13 | 11 | 8.8 | 6 |
| 7.4-37 kW | 13 | 11 | 8.8 | 6 |
| 37-74 kW | 13 | 11 | 8.8 | 6 |
| 74-100 kW | 13 | 11 | 8.8 | 6 |
| >100 kW | 13 | 11 | 8.8 | 6 |
| Fuel consumption (FC) | | | | |
| <4.4 kW | 400 | 400 | 400 | 400 |
| 4.4-7.4 kW | 400 | 400 | 400 | 400 |
| 7.4-37 kW | 400 | 380 | 380 | 380 |
| 37-74 kW | 380 | 350 | 350 | 350 |
| 74-100 kW | 400 | 330 | 330 | 330 |
| >100 kW | 300 | 300 | 300 | 300 |

Table A - 22 Emission and consumption factors for boats with gasoline engines. PreSAV etc. indicate emission standards.

| Basic emission factors of gasoline engine boats (g/kWh) | | | | | | |
|---|--------------------------|-------------|----------------|--------------------------|-------------|------------|
| power class | 2-stroke gasoline engine | | | 4-stroke gasoline engine | | |
| | PreSAV <1995 | SAV 1995 | SAV/EU 2007 | PreSAV <1995 | SAV 2007 | EU 2007 |
| Carbon monoxide (CO) | | | | | | |
| <4.4 kW | 645 | 315 | 315 | 350 | 315 | 315 |
| 4.4-7.4 kW | 645 | 200 | 225 | 350 | 200 | 225 |
| 7.4-37 kW | 645 | 100 | 162 | 350 | 100 | 162 |
| 37-74 kW | 645 | 65 | 144 | 350 | 65 | 144 |
| 74-100 kW | 645 | 55 | 141 | 350 | 55 | 141 |
| >100 kW | 645 | 45 | 139 | 350 | 45 | 139 |
| VOC | | | | | | |
| <4.4 kW | 260 | 22 | 25 | 25 | 22 | 25 |
| 4.4-7.4 kW | 260 | 12 | 13 | 20 | 12 | 13 |
| 7.4-37 kW | 260 | 6 | 8 | 20 | 6 | 8 |
| 37-74 kW | 260 | 4 | 6 | 20 | 4 | 6 |
| 74-100 kW | 260 | 3.3 | 5 | 20 | 3.3 | 5 |
| >100 kW | 260 | 2.1 | 5 | 20 | 2.1 | 5 |
| Nitrogen oxides (NO_x) | | | | | | |
| <4.4 kW | 15 | 13 | 13 | 3.5 | 13 | 13 |
| 4.4-7.4 kW | 15 | 9.3 | 9.3 | 3.5 | 9.3 | 9.3 |
| 7.4-37 kW | 15 | 9.3 | 9.3 | 3.5 | 9.3 | 9.3 |
| 37-74 kW | 15 | 9.3 | 9.3 | 3.5 | 9.3 | 9.3 |
| 74-100 kW | 15 | 9.3 | 9.3 | 3.5 | 9.3 | 9.3 |
| >100 kW | 15 | 9.6 | 9.6 | 3.5 | 9.6 | 9.6 |
| Fuel consumption (FC) | | | | | | |
| <4.4 kW | 700 | 400 | 400 | 400 | 400 | 400 |
| 4.4-7.4 kW | 700 | 400 | 400 | 400 | 400 | 400 |
| 7.4-37 kW | 650 | 380 | 380 | 380 | 380 | 380 |
| 37-74 kW | 650 | 380 | 380 | 380 | 380 | 380 |
| 74-100 kW | 650 | 380 | 380 | 380 | 380 | 380 |
| >100 kW | 650 | 380 | 380 | 380 | 380 | 380 |

Table A - 23 Emission and consumption factors (FC) for ships with steam engines (gas oil). steam 1 etc. indicate emission standards.

| Basic emission factors of steam (gas oil) engine ships (g/kWh) | | | | | | | |
|--|------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| pollutant | steam 1 <1950 | steam 2 1950 | steam 3 1980 | steam 4 1990 | steam 5 1995 | steam 6 2005 | steam 7 2005 |
| CO | 0.3 | 0.3 | 0.3 | 0.09 | 0.09 | 0.09 | 0.09 |
| HC | 0.45 | 0.45 | 0.45 | 0.33 | 0.33 | 0.33 | 0.33 |
| NO _x | 2.34 | 2.34 | 2.34 | 1.77 | 1.56 | 1.26 | 1.03 |
| PM | 0.033 | 0.024 | 0.015 | 0.009 | 0.006 | 0.006 | 0.006 |
| FC | 1406 | 1012 | 787 | 703 | 703 | 703 | 703 |

Activity data off-road vehicles

The activity data are described in detail in INFRAS (2008). Aggregated numbers are shown in the following tables.

Table A - 24 Number of vehicles per off-road family (INFRAS 2008)

| Family | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 |
|--------------|------------------|------------------|------------------|------------------|------------------|------------------|
| | no. of vehicles | | | | | |
| Construction | 56'070 | 52'443 | 47'995 | 47'354 | 48'162 | 48'770 |
| Industry | 13'947 | 18'372 | 22'748 | 22'748 | 23'739 | 24'499 |
| Agriculture | 324'567 | 324'047 | 337'869 | 339'948 | 359'496 | 372'900 |
| Forestry | 13'844 | 13'357 | 13'055 | 12'749 | 12'548 | 12'144 |
| Garden/Hobby | 659'828 | 719'118 | 779'052 | 763'881 | 786'481 | 800'506 |
| Navigation | 93'395 | 89'042 | 82'674 | 82'647 | 86'790 | 90'134 |
| Railway | 1'300 | 1'305 | 1'255 | 1'255 | 1'318 | 1'370 |
| Military | 1'340 | 1'340 | 1'340 | 1'340 | 1'408 | 1'462 |
| Sum | 1'164'291 | 1'219'024 | 1'285'988 | 1'271'922 | 1'319'942 | 1'351'784 |

Table A - 25 Operating hours per vehicle per year and (Table A- 1million) operating hours per off-road family (INFRAS 2008).

| Family | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 |
|--------------|-----------------------------------|------|------|------|------|------|
| | operating hours per veh. per year | | | | | |
| Construction | 299 | 353 | 383 | 386 | 387 | 388 |
| Industry | 628 | 648 | 660 | 660 | 660 | 659 |
| Agriculture | 119 | 118 | 112 | 108 | 104 | 100 |
| Forestry | 199 | 201 | 203 | 202 | 202 | 201 |
| Garden/Hobby | 22 | 25 | 27 | 27 | 27 | 28 |
| Navigation | 40 | 39 | 40 | 40 | 40 | 40 |
| Railway | 612 | 627 | 616 | 616 | 616 | 616 |
| Military | 51 | 53 | 54 | 52 | 49 | 47 |

| Family | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 |
|--------------|-------------------------|--------------|--------------|--------------|---------------|---------------|
| | mio. of operating hours | | | | | |
| Construction | 16.75 | 18.52 | 18.38 | 18.26 | 18.65 | 18.95 |
| Industry | 8.76 | 11.90 | 15.01 | 15.01 | 15.66 | 16.16 |
| Agriculture | 38.77 | 38.21 | 37.68 | 36.57 | 37.26 | 37.45 |
| Forestry | 2.76 | 2.68 | 2.64 | 2.57 | 2.53 | 2.44 |
| Garden/Hobby | 14.42 | 17.71 | 21.09 | 20.78 | 21.52 | 22.04 |
| Navigation | 3.72 | 3.46 | 3.34 | 3.33 | 3.49 | 3.61 |
| Railway | 0.80 | 0.82 | 0.77 | 0.77 | 0.81 | 0.84 |
| Military | 0.07 | 0.07 | 0.07 | 0.07 | 0.07 | 0.07 |
| Sum | 86.04 | 93.37 | 98.99 | 97.36 | 100.00 | 101.56 |

A3.2 Industrial Processes

Illustrative Example of modelling Mobile Air-Conditioning / Cars

Table A - 26 Model structure and assumptions for calculating emissions from mobile air conditioning in cars. The example represents data for the year 2012. The basic parameters have not changed for the present inventory.

| Parameters for Car Air-Conditioning | | |
|--|---------------------------------|--------|
| Initial Charge in kg | 1994 | 0.81 |
| | 2002 | 0.7 |
| | Other years inter-/extrapolated | |
| All units are imported with refrigerants charged | | |
| Emission factor 1995 | Annual loss | 8.5% |
| | Share recharged regularly | 6.0% |
| | Share not recharged | 2.5% |
| Charge at end of life | | 58.0% |
| Disposal emissions | Up to 1999 | 100.0% |
| | From 2000 onwards | 50.0% |
| Export of second hand cars | | 50.0% |
| Reuse of recovered refrigerant (estimate value) | | 80.0% |
| Servicing emission factor | | 10.0% |
| Product lifetime | | 15 |
| Market growth rate | | 0.5% |

| Year | New registered vehicles | Vehicles in use | Disposed vehicles | AC units in new registered cars | | | Stock - AC units in use | | Disposal | Initial charge |
|------|-------------------------|-------------------------|-------------------|---------------------------------|--------------------------|--------------------|------------------------------------|-----------------|--------------------|----------------|
| | (VSAI, EFKO) | (B. f. Statistik/Astra) | | Portion of vehicles with AC [%] | R134a as refrigerant [%] | AC units with R134 | Portion of vehicles with R134a [%] | Units with R134 | Units AC with R134 | kg /vehicle |
| 1989 | 335'094 | 2'895'842 | | 5 | 0 | 0 | 0 | 0 | 0 | 0.85 |
| 1990 | 327'456 | 2'985'399 | 237'899 | 6 | 0 | 0 | 0 | 0 | 0 | 0.84 |
| 1991 | 314'824 | 3'057'800 | 242'423 | 7 | 10 | 2'204 | 0 | 2'204 | 0 | 0.83 |
| 1992 | 296'009 | 3'091'230 | 262'579 | 9 | 30 | 7'992 | 0 | 10'196 | 0 | 0.83 |
| 1993 | 262'814 | 3'109'524 | 244'520 | 14 | 66 | 24'284 | 1 | 34'480 | 0 | 0.82 |
| 1994 | 270'009 | 3'165'043 | 214'490 | 19 | 90 | 46'172 | 3 | 80'652 | 0 | 0.81 |
| 1995 | 272'897 | 3'229'169 | 208'771 | 24 | 100 | 65'495 | 5 | 146'147 | 0 | 0.78 |
| 1996 | 269'529 | 3'268'073 | 230'625 | 38 | 100 | 102'421 | 8 | 248'568 | 0 | 0.77 |
| 1997 | 272'441 | 3'323'421 | 217'093 | 52 | 100 | 141'669 | 12 | 390'237 | 0 | 0.76 |
| 1998 | 297'336 | 3'383'275 | 237'482 | 68 | 100 | 202'188 | 18 | 592'426 | 0 | 0.75 |
| 1999 | 317'985 | 3'467'275 | 233'985 | 75 | 100 | 238'489 | 24 | 830'914 | 0 | 0.73 |
| 2000 | 315'398 | 3'545'247 | 237'426 | 77 | 100 | 242'856 | 30 | 1'073'771 | 0 | 0.72 |
| 2001 | 317'126 | 3'629'713 | 232'660 | 85 | 100 | 269'557 | 37 | 1'343'328 | 0 | 0.71 |
| 2002 | 295'109 | 3'704'822 | 220'000 | 87 | 100 | 256'745 | 43 | 1'600'073 | 0 | 0.70 |
| 2003 | 271'541 | 3'754'000 | 222'363 | 89 | 100 | 241'671 | 49 | 1'841'744 | 0 | 0.70 |
| 2004 | 269'211 | 3'811'351 | 211'860 | 91 | 100 | 244'982 | 55 | 2'086'726 | 0 | 0.70 |
| 2005 | 259'426 | 3'863'807 | 206'970 | 92 | 100 | 238'672 | 60 | 2'325'398 | 0 | 0.70 |
| 2006 | 269'421 | 3'899'917 | 233'311 | 96 | 100 | 258'644 | 66 | 2'582'409 | 1'633 | 0.70 |
| 2007 | 284'674 | 3'955'787 | 228'804 | 96 | 100 | 273'287 | 72 | 2'849'519 | 6'178 | 0.70 |
| 2008 | 288'525 | 4'030'965 | 213'347 | 96 | 100 | 276'984 | 77 | 3'106'789 | 19'713 | 0.70 |
| 2009 | 266'018 | 4'051'569 | 245'414 | 96 | 100 | 255'377 | 82 | 3'320'201 | 41'966 | 0.70 |
| 2010 | 294'239 | 4'119'370 | 226'438 | 96 | 100 | 282'469 | 86 | 3'548'325 | 54'345 | 0.70 |
| 2011 | 327'896 | 4'209'300 | 237'966 | 96 | 100 | 314'780 | 90 | 3'772'678 | 90'427 | 0.70 |
| 2012 | 328'139 | 4'254'725 | 282'714 | 96 | 100 | 315'013 | 93 | 3'940'680 | 147'011 | 0.70 |

| R 134a | Activity | | | Emissions | | | | Recharge | |
|--------|---------------------|-------|----------|----------------------|----------|----------|-------|----------------|------------------|
| | Input with vehicles | Stock | Disposed | Stock incl. Recharge | Disposal | Recharge | Total | import in bulk | recovered/reused |
| | [t] | [t] | [t] | [t] | [t] | [t] | [t] | [t] | [t] |
| 1990 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0.1 | 0.0 |
| 1992 | 7 | 8 | 0 | 0 | 0 | 0 | 0 | 0.3 | 0.0 |
| 1993 | 20 | 28 | 0 | 2 | 0 | 0 | 2 | 1.1 | 0.0 |
| 1994 | 37 | 65 | 0 | 4 | 0 | 0 | 4 | 2.8 | 0.0 |
| 1995 | 51 | 115 | 0 | 8 | 0 | 0 | 8 | 5.4 | 0.0 |
| 1996 | 79 | 191 | 0 | 14 | 0 | 1 | 14 | 9.2 | 0.0 |
| 1997 | 107 | 294 | 0 | 23 | 0 | 2 | 23 | 14.6 | 0.0 |
| 1998 | 151 | 437 | 0 | 35 | 0 | 4 | 35 | 21.9 | 0.0 |
| 1999 | 175 | 599 | 0 | 49 | 0 | 5 | 49 | 31.1 | 0.0 |
| 2000 | 176 | 757 | 0 | 65 | 0 | 8 | 65 | 40.7 | 0.0 |
| 2001 | 192 | 924 | 0 | 82 | 0 | 11 | 82 | 50.4 | 0.0 |
| 2002 | 180 | 1'072 | 0 | 100 | 0 | 15 | 100 | 59.9 | 0.0 |
| 2003 | 169 | 1'201 | 0 | 114 | 0 | 18 | 114 | 68.2 | 0.0 |
| 2004 | 171 | 1'326 | 0 | 125 | 0 | 18 | 125 | 75.8 | 0.0 |
| 2005 | 167 | 1'444 | 0 | 137 | 0 | 19 | 137 | 83.1 | 0.0 |
| 2006 | 181 | 1'571 | 1 | 146 | 0 | 18 | 146 | 90.5 | 0.2 |
| 2007 | 191 | 1'706 | 3 | 156 | 1 | 17 | 157 | 98.3 | 0.6 |
| 2008 | 194 | 1'839 | 9 | 168 | 2 | 17 | 170 | 106.4 | 1.9 |
| 2009 | 179 | 1'947 | 20 | 178 | 5 | 17 | 183 | 113.6 | 4.0 |
| 2010 | 198 | 2'061 | 25 | 188 | 6 | 18 | 195 | 120.2 | 5.0 |
| 2011 | 220 | 2'188 | 41 | 200 | 10 | 19 | 210 | 127.5 | 8.1 |
| 2012 | 221 | 2'296 | 65 | 210 | 16 | 19 | 226 | 134.5 | 13.0 |

A3.3 Agriculture

Additional data for estimating enteric fermentation emission factors for cattle

Table A - 27 Data for estimating enteric fermentation emission factors for cattle. Reference: IPCC 1997c, p 4.31 – 4.33

| Data for estimating enteric fermentation emission factors for cattle in Switzerland | | | | | | | | | | |
|---|------------------|---------------------------|---------------------------------------|--|-----------------------------|-----------------|----------------------------|---|---|---|
| Type | Age ^a | Weight ^a kg | Weight Gain ^a kg/day | Feeding Situation / Further Specification ^a | Milk ^b kg/day | Work hrs/day | Pregnant ^a % | Digesti- bility of Feed ^d % | CH ₄ Conver- sion ^d % | Em. Factor kg/head/ year ^e |
| Mature Dairy Cattle | NA | 650 | 0 | | 16.1 - 22.6 ^c | 0 | 305 days of lactation | 60 | 6.00 | 101.52 - 122.83 |
| Mature Non-Dairy Cattle | NA | 550 | 0 | | 8.2 | 0 | | 60 | 6.00 | 80.71 |
| Fattening Calves | 0-98 days | 60-200 | 1.43 | Rations of unskimmed milk and supplement feed when life weight exceeds 100 kg. Rations are apportioned on two servings per day. | 0 | 0 | 0 | 65 | 0.00 | 0.00 |
| Pre-Weaned Calves | 0-10 month | 60-325 | 1 | "Natura beef" production, milk from mother cow and additional feed. | 0 | 0 | 0 | 65 | 6.00 | 18.03 |
| Breeding Calves | 0-4 month | 50-120 | 0.8 | Feeding plan for a dismission with 14 to 15 weeks. Milk, feed concentrate (100kg in total), hay (80 kg in total). | 0 | 0 | 0 | 65 | 6.00 | 26.58 |
| Breeding Cattle (4-12 months) | 4-12 | 120-300 | 0.8 | Premature race (Milk-race) | 0 | 0 | 0 | 60 | 6.00 | |
| Breeding Cattle (> 1 year) | 12-28/30 | 300-600 | 0.8 | Premature race (Milk-race) | 0 | 0 | 0 | 60 | 6.00 | 50.79 |
| Fattening Calves (0-4 months) | 0-4 month | 70-175 | 0.86 | Diet based on milk or milk-powder and feed concentrate, hay and/or silage | 0 | 0 | 0 | 65 | 6.00 | 40.78 |
| Fattening Cattle (4-12 months) | 4-12 month | 175-550 | 1.3 | Feeding recommendations for fattening steers, concentrate based | 0 | 0 | 0 | 60 | 6.00 | |

^a data source: RAP 1999 and calculations according to Soliva 2006

^b Milk production in kg/day is calculated by dividing the average annual milk production per head by 305 days (lactation period).

^c data source: Swiss farmers union (SBV 2011).

^d data source: IPCC 1997c and IPCC 2000

^e For better comparability emission factors of young cattle have been converted to kg/head/year although the time span of most of the individual categories is less than 365 days.

Additional data for estimating manure management CH₄ emission factors

Table A - 28 Data for estimating manure management CH₄ emission factors. Reference: IPCC 1997c, Tables B-1 to B-7.

| Data for estimating Manure Management CH ₄ emission factors in Switzerland | | | | | | | |
|---|---------------------------|---|----------------------------|------------------------|---------------------------------|-----------------------|---|
| Type | Weight kg ^a | Digestibilit y of Feed % ^b | Energy Intake MJ/day | Feed Intake kg/day | % Ash Dry Basis ^b | VS kg/head/ day | B ₀ m ³ CH ₄ /kg VS ^b |
| Mature Dairy Cattle | 650 | 60 | 258 - 312 | 15.89 ^c | 8 | 5.15-6.24 | 0.24 |
| Mature Non-Dairy Cattle | 550 | 60 | 205.1 | 10.96 ^c | 8 | 4.09 | 0.24 |
| Fattening Calves | 60 – 200 | 65 | 47.6 | 2.02 ^a | 8 | 0.83 | 0.17 |
| Pre-Weaned Calves | 60 – 325 | 65 | 55.7 | 2.98 ^a | 8 | 0.97 | 0.17 |
| Breeding Calves | 50 – 120 | 65 | 26.9 | 1.5 ^a | 8 | 0.47 | 0.17 |
| Breeding Cattle (4-12 months) | 120 – 300 | 60 | 89.2 | 4.88 ^a | 8 | 1.78 | 0.17 |
| Breeding Cattle (> 1 year) | 300 – 600 | 60 | 129.1 | 7.78 ^a | 8 | 2.57 | 0.17 |
| Fattening Calves (0-4 months) | 70 – 175 | 65 | 55.6 | 3.27 ^a | 8 | 0.97 | 0.17 |
| Fattening Cattle (4-12 months) | 175 – 550 | 60 | 124.6 | 6.82 ^a | 8 | 2.48 | 0.17 |
| Sheep | Not determined | 60 | 21 - 24 | 1.09-1.24 ^c | 8 | 0.40 ^b | 0.19 |
| Goats | Not determined | 60 | 25 - 28 | 1.21-1.26 ^c | 8 | 0.28 ^b | 0.17 |
| Horses | Not determined | 70 | 107 - 108 | 7.73-7.83 ^c | 4 | 1.72 ^b | 0.33 |
| Mules and Asses | Not determined | 70 | 39 - 40 | Not estimated | 4 | 0.94 ^b | 0.33 |
| Swine | Not determined | 75 | 26 - 32 | Not estimated | 2 | 0.50 ^b | 0.45 |
| Poultry | Not determined | Not estimated | 1.2 - 1.6 ^d | Not estimated | Not estimated | 0.10 ^b | 0.32 |

^a RAP 1999

^b IPCC 1997c and IPCC 2000

^c Flisch et al. 2009

^d based on metabolizable energy (ME)

Additional data for N₂O emission calculation of agricultural soils

Table A - 29 Additional data for N₂O emission calculation of agricultural soils.

| 2012 | Total crop production Crop(O) and Crop(BF) (kg DM) | Nitrogen incorporated with crop residues F(CR) (t N) | N ₂ O emissions from crop residues (t N ₂ O) | N fixed per kg crop DM (kg N/kg crop) | N fixed (kg N) | N ₂ O emissions from N fixation (t N ₂ O) |
|------------------------------|--|--|--|---------------------------------------|----------------|---|
| 1. Cereals | | | | | | |
| Wheat | 426'771'400 | 1'788 | 35.12 | | | |
| Barley | 156'827'550 | 800 | 15.70 | | | |
| Maize | 124'897'300 | 1'191 | 23.39 | | | |
| Oats | 7'450'250 | 46 | 0.91 | | | |
| Rye | 8'896'100 | 38 | 0.74 | | | |
| Other: | | | | | | |
| Triticale | 47'007'550 | 230 | 4.53 | | | |
| Spelt | 12'203'450 | 112 | 2.19 | | | |
| Mix of Fodder Cereals | 844'050 | 4 | 0.08 | | | |
| Mix of Bread Cereals | 209'100 | 1 | 0.02 | | | |
| 2. Pulse | | | | | | |
| Dry Beans | 804'100 | 32 | 0.63 | 0.0521 | 41'861 | 0.82 |
| Peas (Eiweisserbsen) | 11'139'250 | 328 | 6.44 | 0.0424 | 471'780 | 9.27 |
| Soybeans | 2'635'000 | 109 | 2.13 | 0.0671 | 176'700 | 3.47 |
| Other: | | | | | | |
| Leguminous Vegetables | 2'771'489 | 402 | 7.90 | 0.1170 | 324'149 | 6.37 |
| 3. Tuber and Root | | | | | | |
| Potatoes | 96'694'400 | 353 | 6.94 | | | |
| Other: | | | | | | |
| Fodder Beet | 10'864'000 | 87 | 1.71 | | | |
| Sugar Beet | 368'007'420 | 3'312 | 65.06 | | | |
| 5. Other | | | | | | |
| Fruit | 51'304'640 | 205 | 4.03 | | | |
| Grass | 6'261'698'258 | 22'299 | 438.02 | 0.0051 | 32'117'271 | 630.87 |
| Green Corn | 109'609'171 | 50 | 0.98 | | | |
| Non-Leguminous Vegetables | 55'165'829 | 514 | 10.09 | | | |
| Rape | 59'760'000 | 873 | 17.14 | | | |
| Renewable Energy Crops | 4'176'000 | 61 | 1.20 | | | |
| Silage Corn | 644'759'829 | 370 | 7.27 | | | |
| Sunflowers | 7'820'000 | 166 | 3.25 | | | |
| Tobacco | 1'071'000 | 28 | 0.55 | | | |
| Vine | 25'430'600 | 153 | 3.00 | | | |
| Total Non-leguminous | 2'219'769'639 | 10'381 | 203.91 | | | |
| Total Leguminous | 17'349'839 | 870 | 17.10 | | 1'014'490 | 19.9 |
| Total excluding grass | 2'237'119'478 | 11'251 | 221.01 | | 1'014'490 | 19.9 |
| Total including grass | 8'498'817'736 | 33'551 | 659.03 | | 33'131'761 | 650.8 |

Table A - 30 Additional data for N₂O emission calculation of agricultural soils.

| 2012 | Residue/ Crop ratio (kg/kg) | Dry matter (dm) fraction of residue (kg/kg) | Nitrogen content of residues (kg/kg) |
|---------------------------|-----------------------------|---|--------------------------------------|
| 1. Cereals | | | |
| Wheat | 1.15 | 0.85 | 0.0037 |
| Barley | 1.00 | 0.85 | 0.0051 |
| Maize | 1.11 | 0.85 | 0.0086 |
| Oats | 1.27 | 0.85 | 0.0049 |
| Rye | 1.17 | 0.85 | 0.0036 |
| Other : | | | |
| Triticale | 1.25 | 0.85 | 0.0039 |
| Spelt | 1.56 | 0.85 | 0.0059 |
| Mix of Fodder Cereals | 1.00 | 0.85 | 0.0051 |
| Mix of Bread Cereals | 1.17 | 0.85 | 0.0037 |
| 2. Pulse | | | |
| Dry Beans | 1.13 | 0.85 | 0.0353 |
| Peas (Eiweisserbsen) | 1.25 | 0.85 | 0.0235 |
| Soybeans | 1.00 | 0.85 | 0.0412 |
| Other: | | | |
| Leguminous Vegetables | 3.87 | 0.16 | 0.0328 |
| 3. Tuber and Root | | | |
| Potatoes | 0.47 | 0.13 | 0.0127 |
| Other : | | | |
| Fodder Beet | 0.41 | 0.15 | 0.0233 |
| Sugar Beet | 0.67 | 0.15 | 0.0220 |
| 5. Other | | | |
| Fruit | NA | 0.17 | 0.0040 |
| Grass | 0.26 | NA | 0.0215 |
| Green Corn | 0.05 | 0.32 | 0.0091 |
| Non-Leguminous Vegetables | 0.46 | 0.13 | 0.0230 |
| Rape | 1.86 | 0.85 | 0.0083 |
| Renewable Energy Crops | 1.86 | 0.85 | 0.0083 |
| Silage Corn | 0.05 | 0.32 | 0.0115 |
| Sunflowers | 2.00 | 0.60 | 0.0150 |
| Tobacco | 1.18 | NA | 0.0221 |
| Vine | NA | 0.20 | 0.0060 |

Annex 4: CO₂ Reference Approach and comparison with Sectoral Approach, and relevant information on the national energy balance

Reviewers have repeatedly asked for explanations of the apparent differences between the energy data held by the International Energy Agency (IEA) and the data reported in the reference approach. In order to clarify the pertaining issues, the reasons for the major differences are given below. Data for the year 2010 are used to illustrate the description.

General remarks

The **net calorific values** used by IEA differ from those used in the GHG inventory. In order to avoid differences caused by the conversion with different NCV, the comparison between IEA and the reference approach is made in Gg.

Stock changes as reported by IEA are only including primary stocks (IEA 2005), while the reporting in the reference approach includes secondary and tertiary stocks. This results in a particularly large difference for gas oil, as retailers and end-consumers hold considerable amounts of heating fuel on stock. The IEA subsumes secondary and tertiary stock changes under statistical difference.

All data regarding liquid fuel consumption reported by the IEA includes fuel consumption in **Liechtenstein** (Geographical coverage in IEA 2012). For reporting purposes under the UNFCCC, consumption of Liechtenstein is subtracted.

Data sources used for the comparison shown in table A-33 below are:

- Switzerland's greenhouse gas inventory 1990-2011, submission of 15. April 2013, CRF-table 1.A(b), (FOEN 2013).
- Energy statistics of OECD countries (2012 Edition), (IEA 2012).

Liquid fuels

The total amount of liquid fuel consumption as reported in the greenhouse gas inventory is 11'052 Gg. There is a difference of 13 Gg (0.1%) between CRF and IEA. This difference is primarily caused by the different methodology used for aviation bunkers (see below).

Crude oil

Crude oil in the reference approach contains additives, while IEA lists them separately (data in italics in table A-33). The difference between CRF and IEA is smaller than 0.1% if the sum of additives, refinery feedstocks and crude oil is considered.

Gasoline

The comparison is made for motor gasoline only. Aviation gasoline is included under aviation fuels. Gasoline reported by IEA includes gasoline used in Liechtenstein (LIE), which is subtracted for reporting under the UNFCCC. The difference between CRF and IEA is approximately 0.1%, if the consumption of LIE is taken into account.

Aviation fuels

The different aviation fuels are aggregated in the greenhouse gas inventory. For comparison of IEA and reference approach, all aviation fuels are summed up. The difference between IEA and reference approach if considering the apparent final consumption is 12 Gg (approximately 1% of imports). This difference is largely due to a different methodology used to estimate international bunker. Aviation bunkers have to be reported monthly to the IEA. As

the tier 3 approach used for the greenhouse gas inventory is not available on a monthly basis, the international bunker fuel estimate of IEA consists of the total consumption at the two international airports in Zurich and Geneva, while all remaining fuel use is considered domestic. The reporting in the national greenhouse gas inventory is based on a much more detailed approach, where information on single flights is taken into account. Due to the different approach, the numbers are somewhat different. However, the order of magnitude is the same, and the information in the inventory is based on a higher-tier method and presumably more accurate.

Diesel and gas oil

The IEA numbers include diesel and gas oil used in Liechtenstein. Furthermore, stock changes are reported differently in the CRF and by the IEA. Secondary and tertiary stock changes are subsumed under statistical difference by the IEA, while they are included in the stock change reported in the reference approach. If the statistical difference is taken into account, the difference in the apparent consumption is less than 0.1%.

Residual fuel oil

Data agree between IEA and UNFCCC. It seems as if there is a rounding error in the imported amounts, leading to an apparent difference of 1 Gg. According to the foreign trade statistics, 33'693 t of residual fuel oil had been imported in 2010.

Bitumen

Bitumen is a main feedstock in the greenhouse gas inventory. Data between IEA and the reference approach compare well. Again, small differences are likely due to the use of rounded values, leading to apparent differences of the order of 1-2 Gg.

Petroleum coke

There are considerable differences (26 Gg) in the reported numbers for petroleum coke import. The reason for this apparent difference is that for IEA, all petroleum coke is reported together. In the greenhouse gas inventory, however, only the petroleum coke used as a fuel is reported under petroleum coke, while calcined petroleum coke is reported together with "other oil" as feedstocks. This is largely a consequence of the treatment of fuels and feedstocks in the Swiss energy statistic (SFOE 2012).

Lubricants

There are small differences between IEA and the reference approach, as the data reported to the IEA comprises a slightly different set of customs tariff headings for lubricants to the one used for the Swiss energy statistic. The substances not reported under lubricants in the reference approach are reported under other oil.

Liquefied petroleum gas (LPG)

The reporting of liquefied petroleum gas in the greenhouse gas inventory includes white spirit and lamp oil. As for petroleum coke, IEA numbers include fuels that are used as feedstocks, while in the reference approach, only liquefied petroleum gas, white spirit and lamp oil used as fuels are reported under liquefied petroleum gas. The difference in apparent consumption between IEA and the reference approach is 3 Gg (0.03% of total liquid fuel consumption).

Other oil products

In the greenhouse gas inventory, all other oil products are reported together, while IEA has a finer degree of disaggregation. As already mentioned above, the share of petroleum coke that is used as a feedstock is reported under other oil in the greenhouse gas inventory. Therefore, the difference between IEA and the reference approach corresponds largely to the difference in apparent consumption of petroleum coke.

Solid fuels

Solid fuels play only a minor role in Switzerland (246 Gg) and are reported in good agreement.

Gaseous fuels

In the greenhouse gas inventory, the amount of gas reported under 1B2b Fugitive emissions is subtracted from the total gas import as reported by IEA, as this gas is not used for energy purposes. Taking this into account the difference is of the order of 2 TJ.

Table A - 31 Comparison of the IEA energy statistic with the Reference Approach for the year 2010. Numbers in italics are fuels that are reported in a finer disaggregation in the IEA energy statistic than in the Reference Approach. Numbers in bold aggregate the data to the level of disaggregation used in the Reference Approach.

| CRF vs. IEA (2010) | Import | | Export | | Bunker | | Stock change | | Stat.diff. | LIE | Consumption | |
|--|---------|---------|--------|------|--------|--------|--------------|-------|------------|-----|----------------|----------------|
| Gg | IEA | CRF | IEA | CRF | IEA | CRF | IEA | CRF | IEA | CRF | IEA | CRF |
| Crude oil | 4'488 | 4'546 | | | | | 0 | 1 | 0 | | 4'488 | 4'547 |
| Refinery feedstocks | 3 | | | | | | 1 | | 2 | | 6 | |
| Additives/blending components | 51 | | | | | | -1 | | 2 | | 52 | |
| | | | | | | | | | | | 4'546 | 4'547 |
| Motor gasoline | 1'850 | 1'838 | | | | | -9 | -6 | 4 | 15 | 1'830 | 1'832 |
| Aviation gasoline | 7 | | | | | | -2 | | -1 | | 4 | |
| Kerosene type jet fuel | 1'354 | 1'362 | | | -1'367 | -1'352 | | 2 | 6 | | -7 | 12 |
| Other Kerosene | 3 | | | | | | | | | | 3 | |
| | | | | | | | | | | | 0 | 12 |
| Gas/diesel oil | 3'510 | 3'485 | -21 | -39 | -10 | -11 | 38 | 1'072 | 1'020 | 27 | 4'510 | 4'507 |
| Fuel oil | 33 | 34 | -323 | -316 | | | -17 | -17 | 7 | | -300 | -299 |
| Liquefied petroleum gases (LPG) | 50 | 54 | -24 | -25 | | | | | | 0.1 | 26 | 29 |
| White spirit & SBP | 7 | | | | | | | | -1 | | 6 | |
| | | | | | | | | | | | 32 | 29 |
| Bitumen | 317 | 318 | -2 | -2 | | | | | | | 315 | 317 |
| Lubricants | 86 | 72 | -38 | -16 | | | | | 7 | | 55 | 56 |
| Petroleum coke | 73 | 47 | | | | | | | | | 73 | 47 |
| Naphtha | 1 | | | | | | 5 | | -1 | | 5 | |
| Paraffin waxes | 1 | | | | | | | | | | 1 | |
| Non-specified oil products / other oil | 4 | 63 | - | -23 | | | - | -6 | | | 4 | 33 |
| | | | | | | | | | | | 10 | 33 |
| Liquid fuels | | | | | | | | | | | 11'039 | 11'052 |
| Anthracite | 7 | | | | | | | | | | 7 | |
| Other bituminous coal | 123 | 152 | | | | | 36 | 32 | | | 159 | 184 |
| Lignite | 66 | 62 | | | | | -4 | | | | 62 | 62 |
| Coke oven coke | 18 | | | | | | | | | | 18 | |
| Solid fuels | | | | | | | | | | | 246 | 246 |
| Natural gas (TJ, NCV) | 126'014 | 125'627 | | | | | | | | | 126'014 | 125'627 |
| Fugitive emissions (TJ, NCV) | | 389 | | | | | | | | | | 389 |
| Gaseous fuels | | | | | | | | | | | 126'014 | 126'016 |

Annex 5: Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded

No supplementary information to the statements given in Chapter 1.8 Completeness Assessment.

Annex 6: Additional information to be considered as part of the NIR submission (where relevant) or other useful reference information

A6.1 Independent verification of the National Swiss Inventory for F-gases

Introduction:

Since 2000 the Swiss Federal Laboratories for Materials Science and Technology (Empa) performs continuous measurements of halogenated greenhouse gases at the high-Alpine site of Jungfraujoch (3580 m asl). These measurements are used for the independent estimation of hydrofluorocarbon (HFC) emissions from Switzerland and neighbouring countries and can serve as a verification tool for their Swiss emissions. For this verification the so-called tracer-ratio method is applied, where Swiss HFC pollution events are scaled to concurrent pollution events of carbon monoxide (CO) and then multiplied by the Swiss CO emission inventory. Other methods that rely on atmospheric observations are also being developed at Empa for future usage. Similar approaches are also used for independent verification of greenhouse gas emissions in the United Kingdom (UK MetOffice – using measurements from Mace Head, Ireland) and in Australia (CSIRO – using measurements from Cape Grim, Tasmania).

Method description:

For yearly estimates of Swiss emissions of HFCs based on data from Jungfraujoch, only periods are used when the air masses at the high-Alpine station of Jungfraujoch are predominantly influenced by emissions from Switzerland. The number of events which can be used each year depends on the meteorological conditions and is between 7-15 days per year. The process to select these periods is shown in Figure A-4 and is shortly described here. First, the trajectories from the COSMO-model from MeteoSwiss are screened for periods when the Jungfraujoch site has been under the influence of air masses which were within the Swiss boundary layer for the last 48 hours. Second, for these periods mixing ratios of HFCs are compared with those of CO. Periods which show a concurrent increase for both groups of compounds are selected for the independent verification of Swiss emissions, as this is taken as an indication of thorough mixing of Swiss emissions during the transport to the measurement site. Third, the emissions are calculated for each case/day using the formula in Figure A-4. The resulting emissions are only used for the annual emission estimate if they are within three standard deviations of the average (Grubbs test). This criterion is met by approximately 90% of the selected data. Finally, annual emissions are estimated as the median from these individual cases. These annual estimates are merged to a 3-year annual average centred over a 3-year period (e.g. the estimate for 2009 emissions is calculated by using data from 2008–2010). The error of the estimates based on data from Jungfraujoch has been assessed by using the range of 25%-75% percentiles of the estimates from single pollution events, since 2009. For estimates between 2001-2008 the average of the 2009-2011 errors has been taken. An additional absolute error could occur if the Swiss emissions of CO are over/underestimated by the inventory. This would linearly be transmitted to the emissions of the fluorinated greenhouse gases.




Figure A - 4 Description of the procedure to estimate annual emissions of HFCs from Switzerland by using continuous measurements of HFCs at Jungfraujoch (Switzerland).

Results and Discussion:

In the following, Swiss emissions of five HFCs (HFC-134a, HFC-125, HFC-152a, HFC-143a, HFC-32) are estimated based on data from Jungfraujoch and are compared to the emission estimate of the Swiss greenhouse gas inventory. Further emission estimates of other HFC's and other fluorinated greenhouse gases (e.g. SF_6) will be added in future National Inventory Reports (NIR) upon availability.

HFC-134a

HFC-134a is the most important anthropogenic HFC. Its main source is the diffuse emission from its usage as cooling agent in mobile air conditioners (MACs). Further relevant applications are the usage as propellant and the usage as cooling agent in the industrial and commercial refrigeration. The stock of HFC-134a in MACs and the related emissions have been steadily increasing over the past years. The stabilization of the total emissions between 2005 and 2010 is related to the decreasing HFC use as propellant and optimizations in the industrial and commercial refrigeration. Increasing tendencies are found again in the inventory 2012 due to the still growing stock of HFC-134a in refrigeration and air-conditioning equipment and due to new applications using HFC-134a for research (i.e. as tracer gas). Estimated emissions based on measurements at Jungfraujoch agree fairly well with the emission estimates of the Swiss greenhouse gas inventory. The emissions according to the inventory are slightly higher than the ones based on measurements. But in recent years, the data agree within the estimated uncertainty of 25%, except for the last 3 years. Differences in the last 3 years might occur from the conservative model assumption regarding research use (100% loss assumed, increasing consumption in research of HFC-134a 2010-2012).




Figure A - 5 Comparison of HFC-134a emissions from Switzerland: Inventory and estimates from measurements at Jungfraujoch.

HFC-125

HFC-125 is mainly used as cooling agent in air conditioners and commercial refrigeration equipment. Estimated emissions from Jungfraujoch measurement data are in fairly good agreement with emissions provided by the inventory. Although, in recent years, the inventory emissions seem to slightly exceed the estimates based on data from Jungfraujoch.




Figure A - 6 Comparison of HFC-125 emissions from Switzerland: Inventory and estimates from measurements at Jungfraujoch.

HFC-152a

HFC-152a is mainly used as a blowing agent. It has been used in open-cell polyurethane (PU) foams, in closed cell PU-Sprays and closed-cell extruded polystyrene (XPS) foams. In open cell foams, 100% of emissions are related to the blowing process. In closed cell foams a portion of the blowing agent remains in the product, emissions occur continuously over the lifetime, depending on the cell- and molecular-structure of the blowing agent. Unlike for other blowing agents, experts assume that within the first year of the foam lifetime 95-100% of HFC-152a is emitted. The emissions of the first year are commonly allocated to the country of production (according to UNFCCC good practice guidance). These assumptions and allocation are also applied for the model used in the Swiss inventory for estimating HFC-152a emissions under source category 2F2 Foam Blowing.

HFC-152a emissions from foams in the inventory are mainly related to the production and consumption of PU-Spray (Swiss production HFC free since 2008). Most of other foam products are imported and consequently these emissions are allocated to the country of origin. The reported decrease in the inventory since 2003 reflects the replacement of HFC-152a in PU-Spray.

Up to the year 2002 estimated emissions from Jungfrauoch measurement data are lower than reported in the inventory and from then onwards they are higher. This can be explained by the UNFCCC practice to allocate HFC-152a emissions of the first year to the country of production of foams (which is except for PU-Spray mainly outside Switzerland). However, in reality a fraction of these first year emissions actually occur during usage of the products (e.g. for insulation) in Switzerland and therefore are reflected in the measurements but are not reflected by definition in the inventory¹⁹. Emissions estimated from Jungfrauoch show a consistent negative trend related to the partial phase-out of HFC-152a from the foam-blowing applications.




Figure A - 7 Comparison of HFC-152a emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

HFC-143a and HFC-32

HFC-143a and HFC-32 are mainly used as cooling agent mixtures in commercial refrigeration and stationary air conditioners (together with HFC-134a and/or HFC-125). Estimated emissions from Jungfrauoch measurement data are consistently slightly lower than emissions provided by the inventory. However, they normally agree within the uncertainty of 40% reached for these two compounds by the Jungfrauoch-based estimates.

¹⁹ Nonetheless it is important to apply the UNFCCC approach in the inventory as otherwise double counting may occur when allocating the total emissions to the country of origin and the country of product use.




Figure A - 8 Comparison of HFC-143a emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.




Figure A - 9 Comparison of HFC-32 emissions from Switzerland: Inventory and estimates from measurements at Jungfrauoch.

Annex 7: Supplementary Information on the Uncertainty Analysis

A7.1 Approach for the Monte Carlo Simulations

As a first step, the shape and extent of the probability distributions were derived for the activity data and emission factors, based on measured data, literature or expert judgement. The mean value of the probability distributions was set equal to the value of the GHG inventory. In most cases, normal distributions were assumed, for three agricultural categories, triangular distributions are applied.

As a second step, emissions were calculated as emission factor multiplied by the corresponding activity data. For those cases where the activity data or emission factor for a specific source category were not available as well as for all non key categories, emissions were modelled directly, with the mean value set equal to the value of the GHG inventory and an adequate probability distribution of the emissions.

In a third step, the correlations were chosen. Correlations may have a significant effect on the overall inventory uncertainty. The more the source categories are differentiated the more correlations can be considered. The choice was restricted to categories with relevant amounts of uncertainty. For consistency reasons, Crystal Ball software adjusted a few of the correlation coefficients by an average of 0.10.

The Monte Carlo simulation then provided information on the simulated distribution, on the 2.5 and 97.5 percentiles of emissions, on the uncertainty of the national total emission in 2012 and in the base year 1990 as well as on the trend uncertainty 1990–2012.

A7.2 Assumptions for the Monte Carlo Simulations

A7.2.1 Assumptions for the Probability Distributions

For almost all source and sink categories, normal distributions have been chosen. The important exceptions are the agricultural source categories 4B and 4D shown in the table below.

Table A - 32 Probability distribution assigned to activity data and emission factors (1990 and 2012) of categories that are not normally distributed. For the remaining categories, normal probability distributions have been assigned to the emission uncertainties.

| IPCC Source Category | | | | Gas | Probability distribution | |
|----------------------|-------------|--|---------------------|-----|--------------------------|------------|
| | | | | | AD | EF |
| 4B | Agriculture | B. Manure Management | liquid | N2O | normal | triangular |
| 4B | Agriculture | B. Manure Management | solid | N2O | normal | triangular |
| 4D2 | Agriculture | D. Agricultural Soils; Pasture, Range and Paddock Manure | | N2O | normal | triangular |
| 4D3 | Agriculture | D. Agricultural Soils; Indirect Emissions | deposition | N2O | normal | triangular |
| 4D3 | Agriculture | D. Agricultural Soils; Indirect Emissions | leaching and runoff | N2O | normal | triangular |

A7.2.2 Assumptions for the Correlation Coefficients

Since there are no quantitative correlations available, only three values 1 and ± 0.5 have been used if any are assumed:

- “strong” positive correlations are set to $r = 1.0$ (like perfect correlations),
- “weak” correlations are set to $r = \pm 0.5$.

For modelling of the level uncertainty, the following assumption has been made:

- CO₂ emission factors of liquid fuels are strongly and positively correlated ($r = 1.0$) for large sources (more than 1000 Gg CO₂ eq. The restriction is not relevant since correlations have hardly any influence for small sources). The same holds for gaseous fuels.
- Activity data of liquid and gaseous fuels from the categories 1A2, 1A4a and 1A4b are negatively correlated ($r = -0.5$), since the total amount is well known but the partitioning into the different categories is not. Therefore, if the amount is overestimated in one category it is underestimated in one of the other categories.
- Activity data of 4A (Enteric Fermentation) and 4B (Manure Management) are positively correlated ($r=0.5$) since they are both based on livestock numbers.

For modelling of the trend uncertainty, the following assumptions have been made:

- CO₂ emission factors of each source of gasoline and gas oil are strongly and positively correlated ($r = 1.0$) between 1990 and 2012.
- Activity data/emissions of the major sources (1A2: CO₂, 1A3: CO₂, 1A4: CO₂, 4A: CH₄, 4B: CH₄, 2F: HFC) are medium and positively correlated between 1990 and 2012 ($r = 0.5$).

Table A - 33 Correlation coefficients of activity data . "b_": base year 1990." t_": 2012.

| | b_AD_1A2Gaseous FuelsCO2 (MC) | t_AD_1A2Gaseous FuelsCO2 (MC) | b_AD_1A2Liquid FuelsCO2 (MC) | t_AD_1A2Liquid FuelsCO2 (MC) | b_AD_1A3bDieselCO2 (MC) | t_AD_1A3bDieselCO2 (MC) | b_AD_1A3bGasolineCO2 (MC) | t_AD_1A3bGasolineCO2 (MC) | b_AD_1A4aGaseous FuelsCO2 (MC) | t_AD_1A4aGaseous FuelsCO2 (MC) | b_AD_1A4aLiquid FuelsCO2 (MC) | t_AD_1A4aLiquid FuelsCO2 (MC) | b_AD_1A4bGaseous FuelsCO2 (MC) | t_AD_1A4bGaseous FuelsCO2 (MC) | b_AD_1A4bLiquid FuelsCO2 (MC) | t_AD_1A4bLiquid FuelsCO2 (MC) | b_EM_2F1HFC (MC) | t_EM_2F1HFC (MC) | b_AD_4ACH4 (MC) | t_AD_4ACH4 (MC) | b_AD_4BCH4 (MC) | t_AD_4BCH4 (MC) |
|--------------------------------|-------------------------------|-------------------------------|------------------------------|------------------------------|-------------------------|-------------------------|---------------------------|---------------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|------------------|------------------|-----------------|-----------------|-----------------|-----------------|
| b_AD_1A2Gaseous FuelsCO2 (MC) | 1.0 | 0.5 | | | | | | | -0.5 | | | | -0.5 | | | | | | | | | |
| t_AD_1A2Gaseous FuelsCO2 (MC) | | 1.0 | | | | | | | -0.5 | | | | -0.5 | | | | | | | | | |
| b_AD_1A2Liquid FuelsCO2 (MC) | | | 1.0 | 0.5 | | | | | | | -0.5 | | | | -0.5 | | | | | | | |
| t_AD_1A2Liquid FuelsCO2 (MC) | | | | 1.0 | | | | | | | -0.5 | | | | -0.5 | | | | | | | |
| b_AD_1A3bDieselCO2 (MC) | | | | | 1.0 | 0.5 | | | | | | | | | | | | | | | | |
| t_AD_1A3bDieselCO2 (MC) | | | | | | 1.0 | | | | | | | | | | | | | | | | |
| b_AD_1A3bGasolineCO2 (MC) | | | | | | | 1.0 | 0.5 | | | | | | | | | | | | | | |
| t_AD_1A3bGasolineCO2 (MC) | | | | | | | | 1.0 | | | | | | | | | | | | | | |
| b_AD_1A4aGaseous FuelsCO2 (MC) | | | | | | | | | 1.0 | 0.5 | | | -0.5 | | | | | | | | | |
| t_AD_1A4aGaseous FuelsCO2 (MC) | | | | | | | | | | 1.0 | | | -0.5 | | | | | | | | | |
| b_AD_1A4aLiquid FuelsCO2 (MC) | | | | | | | | | | | 1.0 | 0.5 | | | -0.5 | | | | | | | |
| t_AD_1A4aLiquid FuelsCO2 (MC) | | | | | | | | | | | | 1.0 | | | -0.5 | | | | | | | |
| b_AD_1A4bGaseous FuelsCO2 (MC) | | | | | | | | | | | | | 1.0 | 0.5 | | | | | | | | |
| t_AD_1A4bGaseous FuelsCO2 (MC) | | | | | | | | | | | | | | 1.0 | | | | | | | | |
| b_AD_1A4bLiquid FuelsCO2 (MC) | | | | | | | | | | | | | | | 1.0 | 0.5 | | | | | | |
| t_AD_1A4bLiquid FuelsCO2 (MC) | | | | | | | | | | | | | | | | 1.0 | | | | | | |
| b_EM_2F1HFC (MC) | | | | | | | | | | | | | | | | | 1.0 | 0.5 | | | | |
| t_EM_2F1HFC (MC) | | | | | | | | | | | | | | | | | | 1.0 | | | | |
| b_AD_4ACH4 (MC) | | | | | | | | | | | | | | | | | | | 1.0 | 0.5 | 0.5 | |
| t_AD_4ACH4 (MC) | | | | | | | | | | | | | | | | | | | | 1.0 | 0.5 | |
| b_AD_4BCH4 (MC) | | | | | | | | | | | | | | | | | | | | | 1.0 | 0.5 |
| t_AD_4BCH4 (MC) | | | | | | | | | | | | | | | | | | | | | | 1.0 |

Table A - 34 Correlation coefficients for CO₂ emission factors of category 1A1.

| | b_EF_1A1Other FuelsCO ₂ (MC) | t_EF_1A1Other FuelsCO ₂ (MC) |
|---|---|---|
| b_EF_1A1Other FuelsCO ₂ (MC) | 1.0 | 0.5 |
| t_EF_1A1Other FuelsCO ₂ (MC) | | 1.0 |

Table A - 35 Correlation coefficients for emission factors of agricultural categories 4A, 4B and 4D.

| | t_EF_4D40N2O (MC) | b_EF_4D40N2O (MC) | t_EF_4D3leaching and runoffN2O (MC) | b_EF_4D3leaching and runoffN2O (MC) | t_EF_4D3depositionN2O (MC) | b_EF_4D3depositionN2O (MC) | t_EF_4D20N2O (MC) | b_EF_4D20N2O (MC) | t_EF_4D1organic soilsN2O (MC) | b_EF_4D1organic soilsN2O (MC) | t_EF_4D1fertilizerN2O (MC) | b_EF_4D1fertilizerN2O (MC) | t_EF_4BsolidN2O (MC) | b_EF_4BsolidN2O (MC) | t_EF_4BliquidN2O (MC) | b_EF_4BliquidN2O (MC) | t_EF_4BCH4 (MC) | b_EF_4BCH4 (MC) | t_EF_4ACH4 (MC) | b_EF_4ACH4 (MC) | |
|--|-------------------|-------------------|-------------------------------------|-------------------------------------|----------------------------|----------------------------|-------------------|-------------------|-------------------------------|-------------------------------|----------------------------|----------------------------|----------------------|----------------------|-----------------------|-----------------------|-----------------|-----------------|-----------------|-----------------|--|
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |

Table A - 36 Correlation coefficients for activity data of the agricultural categories 4A and 4B.

| | b_AD_4ACH4 (MC) | t_AD_4ACH4 (MC) | b_AD_4BCH4 (MC) | t_AD_4BCH4 (MC) | b_AD_4BsolidN2O (MC) | t_AD_4BsolidN2O (MC) |
|----------------------|-----------------|-----------------|-----------------|-----------------|----------------------|----------------------|
| b_AD_4ACH4 (MC) | 1.0 | 0.5 | 0.5 | | | |
| t_AD_4ACH4 (MC) | | 1.0 | | 0.5 | | |
| b_AD_4BCH4 (MC) | | | 1.0 | 0.5 | 0.5 | |
| t_AD_4BCH4 (MC) | | | | 1.0 | | 0.5 |
| b_AD_4BsolidN2O (MC) | | | | | 1.0 | 0.5 |
| t_AD_4BsolidN2O (MC) | | | | | | 1.0 |

A7.3 Comments to the Results of Monte Carlo Simulations

A7.3.1 Relation between simulated and inventory values

The Monte Carlo method simulates a probability distribution of the Swiss greenhouse gas emissions from which all relevant statistical parameters can be derived (mean, standard deviation and percentiles). The simulated mean value may slightly differ from the reported CRF value.

The discrepancy between simulated and reported values becomes apparent when mean numbers in Figure 1-4 are compared to reported numbers in the CRF-tables. Note that it is not a relevant issue for the uncertainty analysis but is rather confusing for readers and reviewers who carefully study the numbers. For transparency reasons, the numbers are explained in Table A - 37.

The absolute percentiles generated by the simulation are firstly expressed as relative numbers (the simulated mean is set to 100%). Then the relative numbers are transferred to the numbers reported in the CRF-tables, and they are applied to derive the absolute uncertainties.

Table A - 37 Mean values, 2.5 and 97.5 percentiles of the Monte Carlo simulation and corresponding values of the reported emissions (as of CRF-tables).

| Year | Parameters | Unit | Emission (without LULUCF) | Lower bound 2.5 percentile | Upper bound 97.5 percentile | Lower uncertainty | Upper uncertainty |
|------|-------------------------|-----------------------|------------------------------|-------------------------------|--------------------------------|----------------------|----------------------|
| 1990 | simulated values | | | | | | |
| | absolute | Gg CO ₂ eq | 53'357 | 51'314 | 55'748 | -2'043 | 2'392 |
| | relative | % | 100.0% | 96.2% | 104.5% | -3.8% | 4.5% |
| | values of CRF | | | | | | |
| | absolute | Gg CO ₂ eq | 52'890 | 50'865 | 55'261 | -2'025 | 2'371 |
| | relative | % | 100.0% | 96.2% | 104.5% | -3.8% | 4.5% |
| 2012 | simulated values | | | | | | |
| | absolute | Gg CO ₂ eq | 51'825 | 49'945 | 53'944 | -1'880 | 2'119 |
| | relative | % | 100.0% | 96.4% | 104.1% | -3.6% | 4.1% |
| | values of CRF | | | | | | |
| | absolute | Gg CO ₂ eq | 51'449 | 49'582 | 53'552 | -1'867 | 2'103 |
| | relative | % | 100.0% | 96.4% | 104.1% | -3.6% | 4.1% |

A7.3.2 Tornado Diagram

The following chart shows the results of a sensitivity analysis, depicting the most important sources of uncertainty including LULUCF. These can either be emission factors, activity data or emissions. The bars depict the amount of uncertainty introduced compared to total emissions (on x-axis), and the the numbers attached to the bars indicate the value of the corresponding emission factor or activity data.

The figure shows that the very high uncertainty of the emissions of category 5C1 Grasland remaining Grasland (CO₂) cause the most important contribution to the total uncertainty (see bar on top of the Tornado plot). Further important contributions stem from 5A1 Forestland remaining Forestland (CO₂), 4D1 Direct Soil Emission, Synthetic Fertilizers (N₂O), 4D3 Indirect Emissions, Leaching and Runoff (N₂O).




Figure A - 10 Most important sources of uncertainty in 2012 (incl. LULUCF). Explanations see text above figure. Abbrev.: The letter "t" refers to 2012, "EF" emission factor, "AD" activity data, "EM" emission. x-axis: National total of CO₂ eq emissions in 2012. Numbers attached to the bars indicate the values of emissions factor, activity or emission data in original units (Gg CO₂eq, kg/TJ, ha etc.).

Annex 8: Supplementary Information under Article 7, paragraph 1 of the Kyoto Protocol

No supplementary information under this item.